

Third Edition

Organic Chemistry

Janice Gorzynski Smith

COMMON FUNCTIONAL GROUPS

Type of Compound	General Structure	Example	Functional Group	Type of Compound	General Structure	Example	Functional Group
Acid chloride			-COCl	Aromatic compound			phenyl group
Alcohol			-OH hydroxy group	Carboxylic acid			-COOH carboxy group
Aldehyde			C=O carbonyl group	Ester			-COOR
Alkane	R-H	CH ₃ CH ₃	--	Ether			-OR alkoxy group
Alkene			double bond	Ketone			C=O carbonyl group
Alkyl halide			-X halo group	Nitrile	R-C≡N:	CH ₃ -C≡N:	-C≡N cyano group
Alkyne	-C≡C-	H-C≡C-H	triple bond	Sulfide			-SR alkylthio group
Amide			-CONH ₂ , -CONHR, -CONR ₂	Thiol			-SH mercapto group
Amine	R-NH ₂ or R ₂ NH or R ₃ N	CH ₃ -NH ₂	-NH ₂ amino group	Thioester			-COSR
Anhydride							

Organic Chemistry

Third Edition

Janice Gorzynski Smith

University of Hawai'i at Mānoa





*F*or Megan Sarah

About the Author

Janice Gorzynski Smith was born in Schenectady, New York, and grew up following the Yankees, listening to the Beatles, and water skiing on Sacandaga Reservoir. She became interested in chemistry in high school, and went on to major in chemistry at Cornell University where she received an A.B. degree *summa cum laude*. Jan earned a Ph.D. in Organic Chemistry from Harvard University under the direction of Nobel Laureate E. J. Corey, and she also spent a year as a National Science Foundation National Needs Postdoctoral Fellow at Harvard. During her tenure with the Corey group she completed the total synthesis of the plant growth hormone gibberellic acid.

Following her postdoctoral work, Jan joined the faculty of Mount Holyoke College where she was employed for 21 years. During this time she was active in teaching organic chemistry lecture and lab courses, conducting a research program in organic synthesis, and serving as department chair. Her organic chemistry class was named one of Mount Holyoke's "Don't-miss courses" in a survey by *Boston* magazine. After spending two sabbaticals amidst the natural beauty and diversity in Hawai'i in the 1990s, Jan and her family moved there permanently in 2000. She is currently a faculty member at the University of Hawai'i at Mānoa, where she teaches the two-semester organic chemistry lecture and lab courses. In 2003, she received the Chancellor's Citation for Meritorious Teaching.

Jan resides in Hawai'i with her husband Dan, an emergency medicine physician. She has four children: Matthew and Zachary, age 14 (margin photo on page 163); Jenna, a student at Temple University's Beasley School of Law; and Erin, an emergency medicine physician and co-author of the *Student Study Guide/Solutions Manual* for this text. When not teaching, writing, or enjoying her family, Jan bikes, hikes, snorkels, and scuba dives in sunny Hawai'i, and time permitting, enjoys travel and Hawaiian quilting.



The author (far right) and her family from the left: husband Dan, and children Zach, Erin, Jenna, and Matt.

Contents in Brief

	Prologue	1
1	Structure and Bonding	6
2	Acids and Bases	54
3	Introduction to Organic Molecules and Functional Groups	81
4	Alkanes	113
5	Stereochemistry	159
6	Understanding Organic Reactions	196
7	Alkyl Halides and Nucleophilic Substitution	228
8	Alkyl Halides and Elimination Reactions	278
9	Alcohols, Ethers, and Epoxides	312
10	Alkenes	358
11	Alkynes	399
12	Oxidation and Reduction	426
13	Mass Spectrometry and Infrared Spectroscopy	463
14	Nuclear Magnetic Resonance Spectroscopy	494
15	Radical Reactions	538
16	Conjugation, Resonance, and Dienes	571
17	Benzene and Aromatic Compounds	607
18	Electrophilic Aromatic Substitution	641
19	Carboxylic Acids and the Acidity of the O–H Bond	688
20	Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction	721
21	Aldehydes and Ketones—Nucleophilic Addition	774
22	Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution	825
23	Substitution Reactions of Carbonyl Compounds at the α Carbon	880
24	Carbonyl Condensation Reactions	916
25	Amines	949
26	Carbon–Carbon Bond-Forming Reactions in Organic Synthesis	1002
27	Carbohydrates	1027
28	Amino Acids and Proteins	1074
29	Lipids	1119
30	Synthetic Polymers	1148
	Appendices	A-1
	Glossary	G-1
	Credits	C-1
	Index	I-1

Contents

Preface xviii
Acknowledgments xxiii
List of *How To's* xxv
List of Mechanisms xxvii
List of Selected Applications xxx

Prologue 1

What Is Organic Chemistry? 1
Some Representative Organic Molecules 2
Ginkgolide B—A Complex Organic Compound from the Ginkgo Tree 4



1 Structure and Bonding 6

1.1 The Periodic Table 7
1.2 Bonding 10
1.3 Lewis Structures 12
1.4 Lewis Structures Continued 17
1.5 Resonance 18
1.6 Determining Molecular Shape 23
1.7 Drawing Organic Structures 27
1.8 Hybridization 32
1.9 Ethane, Ethylene, and Acetylene 36
1.10 Bond Length and Bond Strength 40
1.11 Electronegativity and Bond Polarity 42
1.12 Polarity of Molecules 44
1.13 L-Dopa—A Representative Organic Molecule 45
 Key Concepts 46
 Problems 47




2 Acids and Bases 54

2.1 Brønsted–Lowry Acids and Bases 55
2.2 Reactions of Brønsted–Lowry Acids and Bases 56
2.3 Acid Strength and pK_a 58
2.4 Predicting the Outcome of Acid–Base Reactions 61
2.5 Factors That Determine Acid Strength 62
2.6 Common Acids and Bases 70
2.7 Aspirin 71
2.8 Lewis Acids and Bases 72
 Key Concepts 74
 Problems 75



3 Introduction to Organic Molecules and Functional Groups 81

- 3.1 Functional Groups 82
- 3.2 An Overview of Functional Groups 83
- 3.3 Intermolecular Forces 87
- 3.4 Physical Properties 90
- 3.5 Application: Vitamins 97
- 3.6 Application of Solubility: Soap 98
- 3.7 Application: The Cell Membrane 100
- 3.8 Functional Groups and Reactivity 102
- 3.9 Biomolecules 104
 - Key Concepts* 105
 - Problems* 106



4 Alkanes 113

- 4.1 Alkanes—An Introduction 114
- 4.2 Cycloalkanes 118
- 4.3 An Introduction to Nomenclature 119
- 4.4 Naming Alkanes 120
- 4.5 Naming Cycloalkanes 125
- 4.6 Common Names 127
- 4.7 Fossil Fuels 128
- 4.8 Physical Properties of Alkanes 129
- 4.9 Conformations of Acyclic Alkanes—Ethane 129
- 4.10 Conformations of Butane 134
- 4.11 An Introduction to Cycloalkanes 137
- 4.12 Cyclohexane 138
- 4.13 Substituted Cycloalkanes 141
- 4.14 Oxidation of Alkanes 147
- 4.15 Lipids—Part 1 149
 - Key Concepts* 151
 - Problems* 153



5 Stereochemistry 159

- 5.1 Starch and Cellulose 160
- 5.2 The Two Major Classes of Isomers 162
- 5.3 Looking Glass Chemistry—Chiral and Achiral Molecules 163
- 5.4 Stereogenic Centers 166
- 5.5 Stereogenic Centers in Cyclic Compounds 168
- 5.6 Labeling Stereogenic Centers with *R* or *S* 170
- 5.7 Diastereomers 175
- 5.8 Meso Compounds 177
- 5.9 *R* and *S* Assignments in Compounds with Two or More Stereogenic Centers 179
- 5.10 Disubstituted Cycloalkanes 180

- 5.11 Isomers—A Summary 181
- 5.12 Physical Properties of Stereoisomers 182
- 5.13 Chemical Properties of Enantiomers 186
 - Key Concepts 188
 - Problems 190




6 Understanding Organic Reactions 196

- 6.1 Writing Equations for Organic Reactions 197
- 6.2 Kinds of Organic Reactions 198
- 6.3 Bond Breaking and Bond Making 200
- 6.4 Bond Dissociation Energy 203
- 6.5 Thermodynamics 206
- 6.6 Enthalpy and Entropy 209
- 6.7 Energy Diagrams 210
- 6.8 Energy Diagram for a Two-Step Reaction Mechanism 213
- 6.9 Kinetics 215
- 6.10 Catalysts 218
- 6.11 Enzymes 219
 - Key Concepts 220
 - Problems 222



7 Alkyl Halides and Nucleophilic Substitution 228

- 7.1 Introduction to Alkyl Halides 229
- 7.2 Nomenclature 230
- 7.3 Physical Properties 231
- 7.4 Interesting Alkyl Halides 232
- 7.5 The Polar Carbon–Halogen Bond 234
- 7.6 General Features of Nucleophilic Substitution 235
- 7.7 The Leaving Group 236
- 7.8 The Nucleophile 238
- 7.9 Possible Mechanisms for Nucleophilic Substitution 242
- 7.10 Two Mechanisms for Nucleophilic Substitution 243
- 7.11 The S_N2 Mechanism 244
- 7.12 Application: Useful S_N2 Reactions 250
- 7.13 The S_N1 Mechanism 252
- 7.14 Carbocation Stability 256
- 7.15 The Hammond Postulate 258
- 7.16 Application: S_N1 Reactions, Nitrosamines, and Cancer 261
- 7.17 When Is the Mechanism S_N1 or S_N2 ? 262
- 7.18 Vinyl Halides and Aryl Halides 267
- 7.19 Organic Synthesis 267
 - Key Concepts 270
 - Problems 271



8 Alkyl Halides and Elimination Reactions 278

- 8.1 General Features of Elimination 279
 - 8.2 Alkenes—The Products of Elimination Reactions 281
 - 8.3 The Mechanisms of Elimination 285
 - 8.4 The E2 Mechanism 285
 - 8.5 The Zaitsev Rule 288
 - 8.6 The E1 Mechanism 291
 - 8.7 S_N1 and E1 Reactions 294
 - 8.8 Stereochemistry of the E2 Reaction 295
 - 8.9 When Is the Mechanism E1 or E2? 298
 - 8.10 E2 Reactions and Alkyne Synthesis 299
 - 8.11 When Is the Reaction S_N1, S_N2, E1, or E2? 300
- Key Concepts* 304
Problems 305



9 Alcohols, Ethers, and Epoxides 312

- 9.1 Introduction 313
 - 9.2 Structure and Bonding 314
 - 9.3 Nomenclature 314
 - 9.4 Physical Properties 318
 - 9.5 Interesting Alcohols, Ethers, and Epoxides 319
 - 9.6 Preparation of Alcohols, Ethers, and Epoxides 321
 - 9.7 General Features—Reactions of Alcohols, Ethers, and Epoxides 323
 - 9.8 Dehydration of Alcohols to Alkenes 324
 - 9.9 Carbocation Rearrangements 328
 - 9.10 Dehydration Using POCl₃ and Pyridine 330
 - 9.11 Conversion of Alcohols to Alkyl Halides with HX 331
 - 9.12 Conversion of Alcohols to Alkyl Halides with SOCl₂ and PBr₃ 335
 - 9.13 Tosylate—Another Good Leaving Group 338
 - 9.14 Reaction of Ethers with Strong Acid 341
 - 9.15 Reactions of Epoxides 343
 - 9.16 Application: Epoxides, Leukotrienes, and Asthma 347
 - 9.17 Application: Benzo[a]pyrene, Epoxides, and Cancer 349
- Key Concepts* 349
Problems 351



10 Alkenes 358

- 10.1 Introduction 359
- 10.2 Calculating Degrees of Unsaturation 360
- 10.3 Nomenclature 362
- 10.4 Physical Properties 365
- 10.5 Interesting Alkenes 366
- 10.6 Lipids—Part 2 366
- 10.7 Preparation of Alkenes 369
- 10.8 Introduction to Addition Reactions 370

- 10.9 Hydrohalogenation—Electrophilic Addition of HX 371
- 10.10 Markovnikov's Rule 374
- 10.11 Stereochemistry of Electrophilic Addition of HX 376
- 10.12 Hydration—Electrophilic Addition of Water 378
- 10.13 Halogenation—Addition of Halogen 379
- 10.14 Stereochemistry of Halogenation 381
- 10.15 Halohydrin Formation 383
- 10.16 Hydroboration–Oxidation 385
- 10.17 Keeping Track of Reactions 390
- 10.18 Alkenes in Organic Synthesis 391
 - Key Concepts 393
 - Problems 394



11 Alkynes 399

- 11.1 Introduction 400
- 11.2 Nomenclature 401
- 11.3 Physical Properties 402
- 11.4 Interesting Alkynes 402
- 11.5 Preparation of Alkynes 404
- 11.6 Introduction to Alkyne Reactions 405
- 11.7 Addition of Hydrogen Halides 406
- 11.8 Addition of Halogen 409
- 11.9 Addition of Water 409
- 11.10 Hydroboration–Oxidation 412
- 11.11 Reaction of Acetylide Anions 414
- 11.12 Synthesis 417
 - Key Concepts 419
 - Problems 421



12 Oxidation and Reduction 426

- 12.1 Introduction 427
- 12.2 Reducing Agents 428
- 12.3 Reduction of Alkenes 428
- 12.4 Application: Hydrogenation of Oils 432
- 12.5 Reduction of Alkynes 434
- 12.6 The Reduction of Polar C–X σ Bonds 437
- 12.7 Oxidizing Agents 438
- 12.8 Epoxidation 439
- 12.9 Dihydroxylation 442
- 12.10 Oxidative Cleavage of Alkenes 444
- 12.11 Oxidative Cleavage of Alkynes 446
- 12.12 Oxidation of Alcohols 447
- 12.13 Green Chemistry 450
- 12.14 Application: The Oxidation of Ethanol 451
- 12.15 Sharpless Epoxidation 451
 - Key Concepts 454
 - Problems 457



13 Mass Spectrometry and Infrared Spectroscopy 463

- 13.1 Mass Spectrometry 464
- 13.2 Alkyl Halides and the M + 2 Peak 468
- 13.3 Fragmentation 469
- 13.4 Other Types of Mass Spectrometry 472
- 13.5 Electromagnetic Radiation 474
- 13.6 Infrared Spectroscopy 476
- 13.7 IR Absorptions 478
- 13.8 IR and Structure Determination 485
 - Key Concepts 487
 - Problems 488



14 Nuclear Magnetic Resonance Spectroscopy 494

- 14.1 An Introduction to NMR Spectroscopy 495
- 14.2 ^1H NMR: Number of Signals 498
- 14.3 ^1H NMR: Position of Signals 502
- 14.4 The Chemical Shift of Protons on sp^2 and sp Hybridized Carbons 505
- 14.5 ^1H NMR: Intensity of Signals 507
- 14.6 ^1H NMR: Spin-Spin Splitting 508
- 14.7 More Complex Examples of Splitting 513
- 14.8 Spin-Spin Splitting in Alkenes 516
- 14.9 Other Facts About ^1H NMR Spectroscopy 517
- 14.10 Using ^1H NMR to Identify an Unknown 519
- 14.11 ^{13}C NMR Spectroscopy 522
- 14.12 Magnetic Resonance Imaging (MRI) 527
 - Key Concepts 527
 - Problems 528



15 Radical Reactions 538

- 15.1 Introduction 539
- 15.2 General Features of Radical Reactions 540
- 15.3 Halogenation of Alkanes 541
- 15.4 The Mechanism of Halogenation 542
- 15.5 Chlorination of Other Alkanes 545
- 15.6 Chlorination versus Bromination 546
- 15.7 Halogenation as a Tool in Organic Synthesis 548
- 15.8 The Stereochemistry of Halogenation Reactions 549
- 15.9 Application: The Ozone Layer and CFCs 551
- 15.10 Radical Halogenation at an Allylic Carbon 552
- 15.11 Application: Oxidation of Unsaturated Lipids 556
- 15.12 Application: Antioxidants 557
- 15.13 Radical Addition Reactions to Double Bonds 558
- 15.14 Polymers and Polymerization 560
 - Key Concepts 563
 - Problems 564




16 Conjugation, Resonance, and Dienes 571

- 16.1 Conjugation 572
- 16.2 Resonance and Allylic Carbocations 574
- 16.3 Common Examples of Resonance 575
- 16.4 The Resonance Hybrid 577
- 16.5 Electron Delocalization, Hybridization, and Geometry 578
- 16.6 Conjugated Dienes 580
- 16.7 Interesting Dienes and Polyenes 581
- 16.8 The Carbon–Carbon σ Bond Length in 1,3-Butadiene 581
- 16.9 Stability of Conjugated Dienes 583
- 16.10 Electrophilic Addition: 1,2- Versus 1,4-Addition 584
- 16.11 Kinetic Versus Thermodynamic Products 586
- 16.12 The Diels–Alder Reaction 588
- 16.13 Specific Rules Governing the Diels–Alder Reaction 590
- 16.14 Other Facts About the Diels–Alder Reaction 595
- 16.15 Conjugated Dienes and Ultraviolet Light 597
 - Key Concepts 599
 - Problems 601



17 Benzene and Aromatic Compounds 607

- 17.1 Background 608
- 17.2 The Structure of Benzene 609
- 17.3 Nomenclature of Benzene Derivatives 610
- 17.4 Spectroscopic Properties 613
- 17.5 Interesting Aromatic Compounds 614
- 17.6 Benzene's Unusual Stability 615
- 17.7 The Criteria for Aromaticity—Hückel's Rule 617
- 17.8 Examples of Aromatic Compounds 620
- 17.9 What Is the Basis of Hückel's Rule? 626
- 17.10 The Inscribed Polygon Method for Predicting Aromaticity 629
- 17.11 Buckminsterfullerene—Is It Aromatic? 632
 - Key Concepts 633
 - Problems 633



18 Electrophilic Aromatic Substitution 641


- 18.1 Electrophilic Aromatic Substitution 642
- 18.2 The General Mechanism 642
- 18.3 Halogenation 644
- 18.4 Nitration and Sulfonation 646
- 18.5 Friedel–Crafts Alkylation and Friedel–Crafts Acylation 647
- 18.6 Substituted Benzenes 654
- 18.7 Electrophilic Aromatic Substitution of Substituted Benzenes 657
- 18.8 Why Substituents Activate or Deactivate a Benzene Ring 659
- 18.9 Orientation Effects in Substituted Benzenes 661

- 18.10 Limitations on Electrophilic Substitution Reactions with Substituted Benzenes 665
- 18.11 Disubstituted Benzenes 666
- 18.12 Synthesis of Benzene Derivatives 668
- 18.13 Halogenation of Alkyl Benzenes 669
- 18.14 Oxidation and Reduction of Substituted Benzenes 671
- 18.15 Multistep Synthesis 675
 - Key Concepts 678
 - Problems 680



19 Carboxylic Acids and the Acidity of the O–H Bond 688

- 19.1 Structure and Bonding 689
- 19.2 Nomenclature 690
- 19.3 Physical Properties 692
- 19.4 Spectroscopic Properties 693
- 19.5 Interesting Carboxylic Acids 694
- 19.6 Aspirin, Arachidonic Acid, and Prostaglandins 696
- 19.7 Preparation of Carboxylic Acids 697
- 19.8 Reactions of Carboxylic Acids—General Features 699
- 19.9 Carboxylic Acids—Strong Organic Brønsted–Lowry Acids 700
- 19.10 Inductive Effects in Aliphatic Carboxylic Acids 703
- 19.11 Substituted Benzoic Acids 705
- 19.12 Extraction 707
- 19.13 Sulfonic Acids 709
- 19.14 Amino Acids 710
 - Key Concepts 713
 - Problems 714



20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction 721

- 20.1 Introduction 722
- 20.2 General Reactions of Carbonyl Compounds 723
- 20.3 A Preview of Oxidation and Reduction 726
- 20.4 Reduction of Aldehydes and Ketones 727
- 20.5 The Stereochemistry of Carbonyl Reduction 729
- 20.6 Enantioselective Carbonyl Reductions 731
- 20.7 Reduction of Carboxylic Acids and Their Derivatives 733
- 20.8 Oxidation of Aldehydes 738
- 20.9 Organometallic Reagents 739
- 20.10 Reaction of Organometallic Reagents with Aldehydes and Ketones 742
- 20.11 Retrosynthetic Analysis of Grignard Products 746
- 20.12 Protecting Groups 748
- 20.13 Reaction of Organometallic Reagents with Carboxylic Acid Derivatives 750
- 20.14 Reaction of Organometallic Reagents with Other Compounds 753
- 20.15 α,β -Unsaturated Carbonyl Compounds 755
- 20.16 Summary—The Reactions of Organometallic Reagents 758

- 20.17 Synthesis 759
 Key Concepts 762
 Problems 765



21 Aldehydes and Ketones—Nucleophilic Addition 774

- 21.1 Introduction 775
21.2 Nomenclature 776
21.3 Physical Properties 779
21.4 Spectroscopic Properties 780
21.5 Interesting Aldehydes and Ketones 783
21.6 Preparation of Aldehydes and Ketones 784
21.7 Reactions of Aldehydes and Ketones—General Considerations 785
21.8 Nucleophilic Addition of H^- and R^- —A Review 789
21.9 Nucleophilic Addition of CN^- 790
21.10 The Wittig Reaction 792
21.11 Addition of 1° Amines 797
21.12 Addition of 2° Amines 800
21.13 Addition of H_2O —Hydration 802
21.14 Addition of Alcohols—Acetal Formation 804
21.15 Acetals as Protecting Groups 808
21.16 Cyclic Hemiacetals 809
21.17 An Introduction to Carbohydrates 812
 Key Concepts 813
 Problems 815



22 Carboxylic Acids and Their Derivatives— Nucleophilic Acyl Substitution 825

- 22.1 Introduction 826
22.2 Structure and Bonding 828
22.3 Nomenclature 830
22.4 Physical Properties 834
22.5 Spectroscopic Properties 835
22.6 Interesting Esters and Amides 836
22.7 Introduction to Nucleophilic Acyl Substitution 838
22.8 Reactions of Acid Chlorides 842
22.9 Reactions of Anhydrides 844
22.10 Reactions of Carboxylic Acids 845
22.11 Reactions of Esters 850
22.12 Application: Lipid Hydrolysis 853
22.13 Reactions of Amides 855
22.14 Application: The Mechanism of Action of β -Lactam Antibiotics 856
22.15 Summary of Nucleophilic Acyl Substitution Reactions 857
22.16 Natural and Synthetic Fibers 858
22.17 Biological Acylation Reactions 860
22.18 Nitriles 862
 Key Concepts 867
 Problems 870



23 Substitution Reactions of Carbonyl Compounds at the α Carbon 880

- 23.1 Introduction 881
- 23.2 Enols 881
- 23.3 Enolates 884
- 23.4 Enolates of Unsymmetrical Carbonyl Compounds 889
- 23.5 Racemization at the α Carbon 891
- 23.6 A Preview of Reactions at the α Carbon 892
- 23.7 Halogenation at the α Carbon 892
- 23.8 Direct Enolate Alkylation 897
- 23.9 Malonic Ester Synthesis 900
- 23.10 Acetoacetic Ester Synthesis 903
 - Key Concepts* 906
 - Problems* 908



24 Carbonyl Condensation Reactions 916

- 24.1 The Aldol Reaction 917
- 24.2 Crossed Aldol Reactions 921
- 24.3 Directed Aldol Reactions 925
- 24.4 Intramolecular Aldol Reactions 926
- 24.5 The Claisen Reaction 928
- 24.6 The Crossed Claisen and Related Reactions 930
- 24.7 The Dieckmann Reaction 932
- 24.8 The Michael Reaction 934
- 24.9 The Robinson Annulation 936
 - Key Concepts* 940
 - Problems* 941



25 Amines 949

- 25.1 Introduction 950
- 25.2 Structure and Bonding 950
- 25.3 Nomenclature 952
- 25.4 Physical Properties 954
- 25.5 Spectroscopic Properties 955
- 25.6 Interesting and Useful Amines 956
- 25.7 Preparation of Amines 960
- 25.8 Reactions of Amines—General Features 966
- 25.9 Amines as Bases 966
- 25.10 Relative Basicity of Amines and Other Compounds 968
- 25.11 Amines as Nucleophiles 975
- 25.12 Hofmann Elimination 977
- 25.13 Reaction of Amines with Nitrous Acid 980
- 25.14 Substitution Reactions of Aryl Diazonium Salts 982
- 25.15 Coupling Reactions of Aryl Diazonium Salts 986
- 25.16 Application: Synthetic Dyes 988

25.17 Application: Sulfa Drugs 990

Key Concepts 991

Problems 994



26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis 1002

26.1 Coupling Reactions of Organocuprate Reagents 1003

26.2 Suzuki Reaction 1005

26.3 Heck Reaction 1009

26.4 Carbenes and Cyclopropane Synthesis 1012

26.5 Simmons–Smith Reaction 1014

26.6 Metathesis 1015

Key Concepts 1020

Problems 1021



27 Carbohydrates 1027

27.1 Introduction 1028

27.2 Monosaccharides 1028

27.3 The Family of D-Aldoses 1034

27.4 The Family of D-Ketoses 1035

27.5 Physical Properties of Monosaccharides 1036

27.6 The Cyclic Forms of Monosaccharides 1036

27.7 Glycosides 1042

27.8 Reactions of Monosaccharides at the OH Groups 1046

27.9 Reactions at the Carbonyl Group—Oxidation and Reduction 1047

27.10 Reactions at the Carbonyl Group—Adding or Removing One Carbon Atom 1049

27.11 The Fischer Proof of the Structure of Glucose 1053

27.12 Disaccharides 1056

27.13 Polysaccharides 1059

27.14 Other Important Sugars and Their Derivatives 1061

Key Concepts 1066

Problems 1068



28 Amino Acids and Proteins 1074

28.1 Amino Acids 1075

28.2 Synthesis of Amino Acids 1078

28.3 Separation of Amino Acids 1081

28.4 Enantioselective Synthesis of Amino Acids 1085

28.5 Peptides 1086

28.6 Peptide Sequencing 1090

28.7 Peptide Synthesis 1094

28.8 Automated Peptide Synthesis 1099

28.9 Protein Structure 1101

28.10 Important Proteins 1106*Key Concepts 1111**Problems 1113***29 Lipids 1119****29.1 Introduction 1120****29.2 Waxes 1121****29.3 Triacylglycerols 1122****29.4 Phospholipids 1126****29.5 Fat-Soluble Vitamins 1128****29.6 Eicosanoids 1129****29.7 Terpenes 1132****29.8 Steroids 1138***Key Concepts 1143**Problems 1144***30 Synthetic Polymers 1148****30.1 Introduction 1149****30.2 Chain-Growth Polymers—Addition Polymers 1150****30.3 Anionic Polymerization of Epoxides 1156****30.4 Ziegler–Natta Catalysts and Polymer Stereochemistry 1157****30.5 Natural and Synthetic Rubbers 1159****30.6 Step-Growth Polymers—Condensation Polymers 1160****30.7 Polymer Structure and Properties 1164****30.8 Green Polymer Synthesis 1166****30.9 Polymer Recycling and Disposal 1169***Key Concepts 1172**Problems 1173***Appendix A** pK_a Values for Selected Compounds A-1**Appendix B** Nomenclature A-3**Appendix C** Bond Dissociation Energies for Some Common Bonds A-7**Appendix D** Reactions that Form Carbon–Carbon Bonds A-9**Appendix E** Characteristic IR Absorption Frequencies A-10**Appendix F** Characteristic NMR Absorptions A-11**Appendix G** General Types of Organic Reactions A-13**Appendix H** How to Synthesize Particular Functional Groups A-15

Glossary G-1

Credits C-1

Index I-1

Preface

My goal in writing *Organic Chemistry* was to create a text that showed students the beauty and logic of organic chemistry by giving them a book that they would *use*. This text is based on lecture notes and handouts that were developed in my own organic chemistry courses over my 30-year teaching career. I have followed two guiding principles: use relevant and interesting applications to illustrate chemical phenomena, and present the material in a student-friendly fashion using bulleted lists, solved problems, and extensive illustrations and summaries. *Organic Chemistry* is my attempt to simplify and clarify a course that intimidates many students—to make organic chemistry interesting, relevant, and accessible to *all* students, both chemistry majors and those interested in pursuing careers in biology, medicine, and other disciplines, without sacrificing the rigor they need to be successful in the future.

The Basic Features

- **Style** This text is different—by design. Today’s students rely more heavily on visual imagery to learn than ever before. The text uses less prose and more diagrams, equations, tables, and bulleted summaries to introduce and reinforce the major concepts and themes of organic chemistry.
- **Content** *Organic Chemistry* accents basic themes in an effort to keep memorization at a minimum. Relevant examples from everyday life are used to illustrate concepts, and this material is integrated throughout the chapter rather than confined to a boxed reading. Each topic is broken down into small chunks of information that are more manageable and easily learned. Sample problems are used as a tool to illustrate stepwise problem solving. Exceptions to the rule and older, less useful reactions are omitted to focus attention on the basic themes.
- **Organization** *Organic Chemistry* uses functional groups as the framework within which chemical reactions are discussed. Thus, the emphasis is placed on the reactions that different functional groups undergo, not on the reactions that prepare them. Moreover, similar reactions are grouped together so that parallels can be emphasized. These include acid–base reactions (Chapter 2), oxidation and reduction (Chapters 12 and 20), radical reactions (Chapter 15), and reactions of organometallic reagents (Chapter 20).

By introducing one new concept at a time, keeping the basic themes in focus, and breaking complex problems down into small pieces, I have found that many students find organic chemistry an intense but learnable subject. Many, in fact, end the year-long course surprised that they have actually *enjoyed* their organic chemistry experience.

Organization and Presentation

For the most part, the overall order of topics in the text is consistent with the way most instructors currently teach organic chemistry. There are, however, some important differences in the way topics are presented to make the material logical and more accessible. This can especially be seen in the following areas.

- **Review material** Chapter 1 presents a healthy dose of review material covering Lewis structures, molecular geometry and hybridization, bond polarity, and types of bonding. While many of these topics are covered in general chemistry courses, they are presented here from an organic chemist’s perspective. I have found that giving students a firm grasp of these fundamental concepts helps tremendously in their understanding of later material.
- **Acids and bases** Chapter 2 on acids and bases serves two purposes. It gives students experience with curved arrow notation using some familiar proton transfer reactions. It also illustrates how some fundamental concepts in organic structure affect a reaction, in this case an acid–base reaction. Since many mechanisms involve one or more acid–base reactions, I emphasize proton transfer reactions early and come back to this topic often throughout the text.

- **Functional groups** Chapter 3 uses the functional groups to introduce important properties of organic chemistry. Relevant examples—PCBs, vitamins, soap, and the cell membrane—illustrate basic solubility concepts. In this way, practical topics that are sometimes found in the last few chapters of an organic chemistry text (and thus often omitted because instructors run out of time) are introduced early so that students can better grasp why they are studying the discipline.
- **Stereochemistry** Stereochemistry (the three-dimensional structure of molecules) is introduced early (Chapter 5) and reinforced often, so students have every opportunity to learn and understand a crucial concept in modern chemical research, drug design, and synthesis.
- **Modern reactions** While there is no shortage of new chemical reactions to present in an organic chemistry text, I have chosen to concentrate on new methods that introduce a particular three-dimensional arrangement in a molecule, so-called asymmetric or enantioselective reactions. Examples include Sharpless epoxidation (Chapter 12), CBS reduction (Chapter 20), and enantioselective synthesis of amino acids (Chapter 28).
- **Grouping reactions** Since certain types of reactions have their own unique characteristics and terminology that make them different from the basic organic reactions, I have grouped these reactions together in individual chapters. These include acid–base reactions (Chapter 2), oxidation and reduction (Chapters 12 and 20), radical reactions (Chapter 15), and reactions of organometallic reagents (Chapter 20). I have found that focusing on a group of reactions that share a common theme helps students to better see their similarities.
- **Synthesis** Synthesis, one of the most difficult topics for a beginning organic student to master, is introduced in small doses, beginning in Chapter 7 and augmented with a detailed discussion of retrosynthetic analysis in Chapter 11. In later chapters, special attention is given to the retrosynthetic analysis of compounds prepared by carbon–carbon bond-forming reactions (for example, Sections 20.11 and 21.10C).
- **Spectroscopy** Since spectroscopy is such a powerful tool for structure determination, four methods are discussed over two chapters (Chapters 13 and 14).
- **Key Concepts** End-of-chapter summaries succinctly summarize the main concepts and themes of the chapter, making them ideal for review prior to working the end-of-chapter problems or taking an exam.


New to the Third Edition

- In response to reviewer feedback, **new sections** have been added on fragmentation patterns in mass spectrometry (Section 13.3) and peptide sequencing (Section 28.6). In addition, sections on splitting in NMR spectroscopy (Section 14.7) and substituent effects in substituted benzenes (Section 18.6) **have been rewritten** to clarify and focus the material. Some **mechanisms have been modified** by adding electron pairs to nucleophiles and leaving groups to more clearly indicate the course of the chemical reaction.
- **Twenty new NMR spectra** have been added in Chapters 14–25 to give students additional practice in this important type of analysis.
- Over **350 new problems** are included in the third edition. The majority of these problems are written at the intermediate level—more advanced than the easier drill problems, but not as complex as the challenge problems. Beginning with Chapter 11, there are additional multi-step synthesis problems that rely on reactions learned in earlier chapters.
- The **interior design has been modified** to tidy margins, and **art labeling has been simplified**, so students can focus more clearly on the important concepts in a section.
- **New micro-to-macro illustrations** are included on hydrogen bonding in DNA (Chapter 3), the production of ethanol from corn (Chapter 9), partial hydrogenation of vegetable oils (Chapter 12), artificial sweeteners (Chapter 27), and insulin (Chapter 28). Several 3-D illustrations of proteins have been added to Chapter 28 as well. The depiction of enzymes as biological catalysts in Chapter 6 has been redone to use an actual reaction—the conversion of the lactose in milk to glucose and galactose.
- **New health-related and environmental applications** are included in margin notes and problems. Topics include the health benefits of omega-3 fatty acids, α -hydroxy acids in skin care products, drugs such as Benadryl that contain ammonium salts, chloroethane as a local anesthetic, rebaudioside A (trade name Truvia), a sweetening agent isolated from a plant source, and many others.

Tools to Make Learning Organic Chemistry Easier


Illustrations

Organic Chemistry is supported by a well-developed illustration program. Besides traditional skeletal (line) structures and condensed formulas, there are numerous ball-and-stick molecular models and electrostatic potential maps to help students grasp the three-dimensional structure of molecules (including stereochemistry) and to better understand the distribution of electronic charge.




Micro-to-Macro Illustrations

Unique to *Organic Chemistry* are micro-to-macro illustrations, where line art and photos combine with chemical structures to reveal the underlying molecular structures giving rise to macroscopic properties of common phenomena. Examples include starch and cellulose (Chapter 5), adrenaline (Chapter 7), partial hydrogenation of vegetable oil (Chapter 12), and dopamine (Chapter 25).




Spectra

Over 100 spectra created specifically for *Organic Chemistry* are presented throughout the text. The spectra are color-coded by type and generously labeled. Mass spectra are green; infrared spectra are red; and proton and carbon nuclear magnetic resonance spectra are blue.



Mechanisms

Curved arrow notation is used extensively to help students follow the movement of electrons in reactions. Where appropriate, mechanisms are presented in parts to promote a better conceptual understanding.



Problem Solving

Sample Problems

Sample Problems show students how to solve organic chemistry problems in a logical, stepwise manner. More than 800 follow-up problems are located throughout the chapters to test whether students understand concepts covered in the Sample Problems.

How To's

How To's provide students with detailed instructions on how to work through key processes.

Applications and Summaries

Key Concept Summaries

Succinct summary tables reinforcing important principles and concepts are provided at the end of each chapter.


Margin Notes

Margin notes are placed carefully throughout the chapters, providing interesting information relating to topics covered in the text. Some margin notes are illustrated with photos to make the chemistry more relevant.




Canola, soybeans, and flaxseed are excellent dietary sources of linolenic acid, an essential fatty acid. Oils derived from omega-3 fatty acids (Problem 10.12) are currently thought to be especially beneficial for individuals at risk of developing coronary artery disease.

Sample Problem 15.4 Draw the products formed when **A** is treated with NBS + $h\nu$.



Solution

Hydrogen abstraction at the allylic C forms a resonance-stabilized radical (with two different resonance structures) that reacts with Br_2 to form two constitutional isomers as products.




Problem 15.20 Draw all constitutional isomers formed when each alkene is treated with NBS + $h\nu$.




HOW TO Name an Ester ($\text{RCO}_2\text{R}'$) Using the IUPAC System

Example Give a systematic name for each ester:




Step [1] Name the R' group bonded to the oxygen atom as an alkyl group.

- The name of the alkyl group, ending in the suffix *-yl*, becomes the *first* part of the ester name.



Step [2] Name the acyl group ($\text{RCO}-$) by changing the *-ic acid* ending of the parent carboxylic acid to the suffix *-ate*.

- The name of the acyl group becomes the *second* part of the name.



Answer: ethyl acetate

Answer: *tert*-butyl cyclohexanecarboxylate

KEY CONCEPTS

Alkenes


General Facts About Alkenes

- Alkenes contain a carbon-carbon double bond consisting of a stronger σ bond and a weaker π bond. Each carbon is sp^2 hybridized and trigonal planar (10.1).
- Alkenes are named using the suffix *-ene* (10.3).
- Alkenes with different groups on each end of the double bond exist as a pair of diastereomers, identified by the prefixes *E* and *Z* (10.3B).
- Alkenes have weak intermolecular forces, giving them low mp 's and bp 's, and making them water insoluble. A *cis* alkene is more polar than a *trans* alkene, giving it a slightly higher boiling point (10.4).
- Because a π bond is electron rich and much weaker than a σ bond, alkenes undergo addition reactions with electrophiles (10.8).

Stereochemistry of Alkene Addition Reactions (10.8)


A reagent XY adds to a double bond in one of three different ways:

- Syn addition**— X and Y add from the same side.




- Syn addition occurs in **hydroboration**.

- Anti addition**— X and Y add from opposite sides.



- Anti addition occurs in **halogenation** and **halohydrin formation**.


- Both syn and anti addition** occur when carbocations are intermediates.



- Syn and anti addition occur in **hydrohalogenation** and **hydration**.


Addition Reactions of Alkenes

[1] Hydrohalogenation—Addition of HX ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) (10.9–10.11)




- The mechanism has two steps.
- Carbocations are formed as intermediates.
- Carbocation rearrangements are possible.
- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- Syn and anti addition occur.

[2] Hydration and related reactions (Addition of H_2O or ROH) (10.12)



- For both reactions:
- The mechanism has three steps.
 - Carbocations are formed as intermediates.
 - Carbocation rearrangements are possible.
 - Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
 - Syn and anti addition occur.



Supplements for the Instructor and Student

The following items may accompany this text. Please consult your McGraw-Hill representative for policies, prices, and availability as some restrictions may apply.

McGraw-Hill Connect™ Chemistry is a web-based assignment and assessment platform that gives students the means to better connect with their course work, their instructors, and the important concepts that they will need to know for success now and in the future.



With *Connect Chemistry*, instructors can deliver assignments, quizzes, and tests online.

A majority of questions from the text are presented in an auto-gradable format and tied to the text's learning objectives. Instructors can edit existing questions and author entirely new problems. Track individual student performance—by question, assignment, or in relation to the class overall—with detailed grade reports. Integrate grade reports easily with Learning Management Systems (LMS) such as WebCT and Blackboard.

By choosing *Connect Chemistry*, instructors are providing their students with a powerful tool for improving academic performance and truly mastering course material. *Connect Chemistry* allows students to practice important skills at their own pace and on their own schedule. Importantly, students' assessment results and instructors' feedback are all saved online—so students can continually review their progress and plot their course to success.

Like *Connect Chemistry*, *Connect Chemistry Plus* provides students with online assignments and assessments, plus 24/7 online access to an eBook—an online edition of the text—to aid them in successfully completing their work, wherever and whenever they choose.

McGraw-Hill Presentation Center allows instructors to build instructional materials wherever, whenever, and however you want! Presentation Center is an online digital library containing assets such as photos, artwork, PowerPoints, and other media types that can be used to create customized lectures, visually enhanced tests and quizzes, compelling course websites, or attractive printed support materials. The McGraw-Hill Presentation Center library includes thousands of assets from many McGraw-Hill titles. This ever-growing resource gives instructors the power to utilize assets specific to an adopted textbook as well as content from all other books in the library. The Presentation Center can be accessed from the instructor side of your textbook's ARIS website, and the Presentation Center's dynamic search engine allows you to explore by discipline, course, textbook chapter, asset type, or keyword. Simply browse, select, and download the files you need to build engaging course materials. All assets are copyright McGraw-Hill Higher Education, but can be used by instructors for classroom purposes.

Brownstone's Diploma testing software serves up over 1,200 test questions to accompany *Organic Chemistry*. Diploma's software allows you to quickly create a customized test using McGraw-Hill's supplied questions, or by authoring your own questions. Diploma is a downloadable application that allows you to create your tests without an Internet connection—just download the software and question files directly to your computer.

Student Study Guide/Solutions Manual Written by Janice Gorzynski Smith and Erin Smith Berk, the Student Study Guide/Solutions Manual provides step-by-step solutions to all in-chapter and end-of-chapter problems. Each chapter begins with an overview of key concepts and includes key rules and summary tables.

Acknowledgments

When I started working on the first edition of *Organic Chemistry* in the fall of 1999, I had no sense of the magnitude of the task, or any idea of just how many people I would rely upon to complete it. Fortunately, I have had the steadfast support of a dedicated team of publishing professionals at McGraw-Hill.

I am especially thankful for the opportunity to work with three terrific women who have transformed the ideas and manuscript pages of the last two editions of *Organic Chemistry* into stunning texts—Tami Hodge (Senior Sponsoring Editor), Donna Nemmers (Senior Developmental Editor), and Jayne Klein (Senior Project Manager). All aspects of this project—from devising the overall plan for the third edition to obtaining valuable reviews to setting a workable production schedule—have been carried out with skill and efficiency. I couldn't ask for a better team of individuals with which to work.

Thanks also go out to Ryan Blankenship, who has recently assumed the role of Publisher for my project. Senior Marketing Manager Todd Turner has provided me with many valuable insights that result from his many contacts with current and potential users. I also appreciate the work of Laurie Jansen (Designer) and Carrie Burger (Photo Researcher) who are responsible for the visually pleasing appearance of this edition. Thanks are again due to Professor Spencer Knapp and his crew at Rutgers University, who prepared the many new spectra that appear in the third edition, and to freelance Developmental Editor John Murdzek for his meticulous editing and humorous insights on my project.

Organic Chemistry is complemented with useful supplements prepared by qualified and dedicated individuals. Special thanks go to Kathleen Halligan of York College of Pennsylvania who authored the instructor's test bank, and Layne Morsch of The University of Illinois, Springfield who prepared the PowerPoint lecture outlines. I am also grateful for the keen eyes of Matthew Dintzner of DePaul University, Michael Kurz of the University of Texas–San Antonio, and Margaret Ruth Leslie of Kent State University for their careful accuracy checking of the Test Bank and PowerPoint Lecture Outlines to accompany this text.

My immediate family has experienced the day-to-day demands of living with a busy author. Thanks go to my husband Dan and my children Erin, Jenna, Matthew, and Zachary, all of whom keep me grounded during the time-consuming process of writing and publishing a textbook. Erin, co-author of the Student Study Guide/Solutions Manual, continued this important task this year in the midst of planning a wedding, completing a residency in emergency medicine, and settling into a new home and profession.

Among the many others that go unnamed but who have profoundly affected this work are the thousands of students I

have been lucky to teach over the last 30 years. I have learned so much from my daily interactions with them, and I hope that the wider chemistry community can benefit from this experience by the way I have presented the material in this text.

This third edition has evolved based on the helpful feedback of many people who reviewed the second edition, class-tested the book, and attended focus groups or symposiums. These many individuals have collectively provided constructive improvements to the project.

Heba Abourahma, *Indiana University of Pennsylvania*
Madeline Adamczeski, *San José City College*
Sheikh Ahmed, *West Virginia University*
Jung-Mo Ahn, *University of Texas, Dallas*
Thomas Albright, *University of Houston*
Scott E. Allen, *University of Tampa*
Steven W. Anderson, *University of Wisconsin, Whitewater*
Mark E. Arant, *University of Louisiana, Monroe*
Thurston E. Banks, *Tennessee Technological University*
Debra L. Bautista, *Eastern Kentucky University*
David Bergbreiter, *Texas A&M University*
John M. Berger, *Montclair State University*
David Berkowitz, *University of Nebraska, Lincoln*
Steve Bertman, *Western Michigan University*
Silas C. Blackstock, *The University of Alabama*
James R. Blanton, *The Citadel*
David L. Boatright, *University of West Georgia*
Chad Booth, *Texas State University, San Marcos*
Ned Bowden, *University of Iowa*
Kathleen Brunke, *Christopher Newport University*
Christopher S. Callam, *The Ohio State University*
Suzanne Carpenter, *Armstrong Atlantic State University*
Steven Castle, *Brigham Young University*
Hamish S. Christie, *The University of Arizona*
Allen Clauss, *University of Wisconsin, Madison*
Barry A. Coddens, *Northwestern University*
Sergio Cortes, *University of Texas, Dallas*
James Ricky Cox, *Murray State University*
Jason P. Cross, *Temple University*
Peter de Lijser, *California State University, Fullerton*
Amy M. Deveau, *University of New England*
Brahmadeo Dewprashad, *Borough of Manhattan Community College*
Matthew Dintzner, *DePaul University*
Pamela S. Doyle, *Essex County College*
Nicholas Drapela, *Oregon State University*
Norma Kay Dunlap, *Middle Tennessee State University*
Ihasn Erden, *San Francisco State University*

John Michael Ferguson, *University of Central Oklahoma*
 David Flanigan, *Hillsborough Community College*
 David C. Forbes, *University of South Alabama*
 John W. Francis, *Columbus State Community College*
 Lee Friedman, *University of Maryland, College Park*
 Anne Gaquere, *University of West Georgia*
 Bob Gawley, *University of Arkansas*
 Jose L. Gonzalez–Roman, *Georgia Perimeter College, Decatur*
 Anne Elizabeth V. Gorden, *Auburn University*
 Steven M. Graham, *Saint John’s University*
 Dennis J. Gravert, *Saint Mary’s University of Minnesota*
 Ray A. Gross, Jr., *Prince George’s Community College*
 Stephen M. Gross, *Creighton University*
 Greg Hale, *University of Texas, Arlington*
 Kathleen M. Halligan, *York College of Pennsylvania*
 Scott Handy, *Middle Tennessee State University*
 Kenn Harding, *Texas A&M University*
 Jill Harp, *Winston Salem State University*
 Paul Higgs, *Barry University*
 Ed Hilinski, *Florida State University*
 Nadene Houser–Archfield, *Prince George’s Community College*
 Michael T. Huggins, *University of West Florida*
 Thomas G. Jackson, *University of South Alabama*
 Peter A. Jacobi, *Dartmouth College*
 Tamera S. Jahnke, *Missouri State University*
 David Andrew Jeffrey, *Georgia State University*
 Hima S. Joshi, *California Polytechnic State University,
 San Luis Obispo*
 Eric Kantorowski, *California Polytechnic State University,
 San Luis Obispo*
 Steven Kass, *University of Minnesota*
 Mushtaq Khan, *Union County College*
 Rebecca Kissling, *SUNY, Binghamton*
 Vera Kolb, *University of Wisconsin, Parkside*
 Grant Krow, *Temple University*
 Michael Kurz, *University of Texas, San Antonio*
 Michael Langohr, *Tarrant County College District*
 Michael S. Leonard, *Washington & Jefferson College*
 Chunmei Li, *Stephen F. Austin State University*
 Harriet A. Lindsay, *Eastern Michigan University*
 Robert D. Long, *Eastern New Mexico University*
 Douglas A. Loy, *The University of Arizona*
 Mitch Malachowski, *University of San Diego*
 Ned H. Martin, *University of North Carolina, Wilmington*
 Michael M. McCormick, *Boise State University*
 Owen McDougal, *Boise State University*
 Matt McIntosh, *University of Arkansas*
 Keith T. Mead, *Mississippi State University*
 Thomas Minehan, *California State University, Northridge*
 James A. Miranda, *California State University, Sacramento*
 Miguel O. Mitchell, *Salisbury University*
 Thomas W. Nalli, *Winona State University*

Donna J. Nelson, *University of Oklahoma*
 Dallas New, *University of Central Oklahoma*
 Jacqueline A. Nikles, *University of Alabama, Birmingham*
 William J. Nixon, Jr., *St. Petersburg College*
 David Allan Owen, *Murray State University*
 Anne B. Padias, *The University of Arizona*
 Daniel Palleros, *University of California, Santa Cruz*
 James W. Pavlik, *Worcester Polytechnic Institute*
 Otto Phanstiel, *University of Central Florida*
 Charles U. Pittman, Jr., *Mississippi State University*
 John R. Pollard, *The University of Arizona*
 Daniel P. Predecki, *Shippensburg University*
 Michael B. Ramey, *Appalachian State University*
 Michael Rathke, *Michigan State University*
 Partha S. Ray, *University of West Georgia*
 J. Ty Redd, *Southern Utah University*
 J. Michael Robinson, *The University of Texas, Permian Basin*
 Tomislav Rovis, *Colorado State University*
 Lev Ryzhkov, *Towson University*
 Raymond Sadeghi, *University of Texas, San Antonio*
 Robert Sammelson, *Ball State University*
 Jason M. Serin, *Glendale Community College*
 Heather Sklenicka, *Rochester Community and Technical
 College*
 Irina P. Smoliakova, *University of North Dakota*
 David Spurgeon, *The University of Arizona*
 Laurie S. Starkey, *California State Polytechnic University,
 Pomona*
 Chad Stearman, *Missouri State University*
 Jonathan M. Stoddard, *California State University, Fullerton*
 Robert Stolow, *Tufts University*
 Todd Swanson, *Gustavus Adolphus College*
 Richard Tarkka, *University of Central Arkansas*
 Eric S. Tillman, *Bucknell University*
 Eric L. Trump, *Emporia State University*
 Ken Walsh, *University of Southern Indiana*
 Don Warner, *Boise State University*
 Arlon A. Widder, *Georgia Perimeter College*
 Milton J. Wieder, *Metropolitan State College, Denver*
 Viktor Zhdankin, *University of Minnesota, Duluth*

Although every effort has been made to make this text and its accompanying Student Study Guide/Solutions Manual as error-free as possible, some errors undoubtedly remain and for them, I am solely responsible. Please feel free to email me about any inaccuracies, so that subsequent editions may be further improved.

With much aloha,

Janice Gorzynski Smith
jgsmith@hawaii.edu

List of *How To's*

How To boxes provide detailed instructions for key procedures that students need to master. Below is a list of each *How To* and where it is presented in the text.

- Chapter 1 Structure and Bonding**
How To Draw a Lewis Structure 13
- Chapter 2 Acids and Bases**
How To Determine the Relative Acidity of Protons 69
- Chapter 4 Alkanes**
How To Name an Alkane Using the IUPAC System 121
How To Name a Cycloalkane Using the IUPAC System 125
How To Draw a Newman Projection 132
How To Draw the Chair Form of Cyclohexane 139
How To Draw the Two Conformations for a Substituted Cyclohexane 142
How To Draw Two Conformations for a Disubstituted Cyclohexane 145
- Chapter 5 Stereochemistry**
How To Assign *R* or *S* to a Stereogenic Center 172
How To Find and Draw All Possible Stereoisomers for a Compound with Two Stereogenic Centers 176
- Chapter 7 Alkyl Halides and Nucleophilic Substitution**
How To Name an Alkyl Halide Using the IUPAC System 230
- Chapter 9 Alcohols, Ethers, and Epoxides**
How To Name an Alcohol Using the IUPAC System 315
- Chapter 10 Alkenes**
How To Name an Alkene 362
How To Assign the Prefixes *E* and *Z* to an Alkene 364
- Chapter 11 Alkynes**
How To Develop a Retrosynthetic Analysis 418
- Chapter 13 Mass Spectrometry and Infrared Spectroscopy**
How To Use MS and IR for Structure Determination 486
- Chapter 14 Nuclear Magnetic Resonance Spectroscopy**
How To Determine the Number of Protons Giving Rise to an NMR Signal 508
How To Use ¹H NMR Data to Determine a Structure 520
- Chapter 16 Conjugation, Resonance, and Dienes**
How To Draw the Product of a Diels–Alder Reaction 590
- Chapter 17 Benzene and Aromatic Compounds**
How To Use the Inscribed Polygon Method to Determine the Relative Energies of MOs for Cyclic, Completely Conjugated Compounds 629
- Chapter 18 Electrophilic Aromatic Substitution**
How To Determine the Directing Effects of a Particular Substituent 661
- Chapter 21 Aldehydes and Ketones—Nucleophilic Addition**
How To Determine the Starting Materials for a Wittig Reaction Using Retrosynthetic Analysis 795
- Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution**
How To Name an Ester (RCO₂R') Using the IUPAC System 831
How To Name a 2° or 3° Amide 831
- Chapter 24 Carbonyl Condensation Reactions**
How To Synthesize a Compound Using the Aldol Reaction 921
How To Synthesize a Compound Using the Robinson Annulation 939

Chapter 25 Amines

How To Name 2° and 3° Amines with Different Alkyl Groups 952

Chapter 27 Carbohydrates

How To Draw a Haworth Projection from an Acyclic Aldohexose 1039

Chapter 28 Amino Acids and Proteins

How To Use (*R*)- α -Methylbenzylamine to Resolve a Racemic Mixture of Amino Acids 1083

How To Synthesize a Dipeptide from Two Amino Acids 1095

How To Synthesize a Peptide Using the Merrifield Solid Phase Technique 1099

List of Mechanisms

Mechanisms are the key to understanding the reactions of organic chemistry. For this reason, great care has been given to present mechanisms in a detailed, step-by-step fashion. The list below indicates when each mechanism in the text is presented for the first time.

- Chapter 7 Alkyl Halides and Nucleophilic Substitution**
 - 7.1 The S_N2 Mechanism 245
 - 7.2 The S_N1 Mechanism 252
- Chapter 8 Alkyl Halides and Elimination Reactions**
 - 8.1 The E2 Mechanism 285
 - 8.2 The E1 Mechanism 291
- Chapter 9 Alcohols, Ethers, and Epoxides**
 - 9.1 Dehydration of 2° and 3° ROH—An E1 Mechanism 326
 - 9.2 Dehydration of a 1° ROH—An E2 Mechanism 327
 - 9.3 A 1,2-Methyl Shift—Carbocation Rearrangement During Dehydration 329
 - 9.4 Dehydration Using POCl₃ + Pyridine—An E2 Mechanism 331
 - 9.5 Reaction of a 1° ROH with HX—An S_N2 Mechanism 333
 - 9.6 Reaction of 2° and 3° ROH with HX—An S_N1 Mechanism 333
 - 9.7 Reaction of ROH with SOCl₂ + Pyridine—An S_N2 Mechanism 336
 - 9.8 Reaction of ROH with PBr₃—An S_N2 Mechanism 336
 - 9.9 Mechanism of Ether Cleavage in Strong Acid—
(CH₃)₃COCH₃ + HI → (CH₃)₃CI + CH₃I + H₂O 342
- Chapter 10 Alkenes**
 - 10.1 Electrophilic Addition of HX to an Alkene 373
 - 10.2 Electrophilic Addition of H₂O to an Alkene—Hydration 379
 - 10.3 Addition of X₂ to an Alkene—Halogenation 380
 - 10.4 Addition of X and OH—Halohydrin Formation 383
 - 10.5 Addition of H and BH₂—Hydroboration 386
- Chapter 11 Alkynes**
 - 11.1 Electrophilic Addition of HX to an Alkyne 407
 - 11.2 Addition of X₂ to an Alkyne—Halogenation 409
 - 11.3 Tautomerization in Acid 410
 - 11.4 Hydration of an Alkyne 411
- Chapter 12 Oxidation and Reduction**
 - 12.1 Addition of H₂ to an Alkene—Hydrogenation 430
 - 12.2 Dissolving Metal Reduction of an Alkyne to a Trans Alkene 436
 - 12.3 Reduction of RX with LiAlH₄ 437
 - 12.4 Epoxidation of an Alkene with a Peroxyacid 440
 - 12.5 Oxidation of an Alcohol with CrO₃ 448
 - 12.6 Oxidation of a 1° Alcohol to a Carboxylic Acid 449
- Chapter 15 Radical Reactions**
 - 15.1 Radical Halogenation of Alkanes 543
 - 15.2 Allylic Bromination with NBS 554
 - 15.3 Radical Addition of HBr to an Alkene 559
 - 15.4 Radical Polymerization of CH₂=CHZ 563
- Chapter 16 Conjugation, Resonance, and Dienes**
 - 16.1 Electrophilic Addition of HBr to a 1,3-Diene—1,2- and 1,4-Addition 585
- Chapter 18 Electrophilic Aromatic Substitution**
 - 18.1 General Mechanism—Electrophilic Aromatic Substitution 643
 - 18.2 Bromination of Benzene 645
 - 18.3 Formation of the Nitronium Ion (⁺NO₂) for Nitration 646

- 18.4 Formation of the Electrophile $^+\text{SO}_3\text{H}$ for Sulfonation 647
- 18.5 Formation of the Electrophile in Friedel–Crafts Alkylation—Two Possibilities 649
- 18.6 Friedel–Crafts Alkylation Using a 3° Carbocation 649
- 18.7 Formation of the Electrophile in Friedel–Crafts Acylation 649
- 18.8 Friedel–Crafts Alkylation Involving Carbocation Rearrangement 651
- 18.9 A Rearrangement Reaction Beginning with a 1° Alkyl Chloride 651
- 18.10 Benzylic Bromination 670
- Chapter 20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction**
- 20.1 Nucleophilic Addition—A Two-Step Process 724
- 20.2 Nucleophilic Substitution—A Two-Step Process 725
- 20.3 LiAlH_4 Reduction of RCHO and $\text{R}_2\text{C}=\text{O}$ 728
- 20.4 Reduction of RCOCl and RCOOR' with a Metal Hydride Reagent 735
- 20.5 Reduction of an Amide to an Amine with LiAlH_4 737
- 20.6 Nucleophilic Addition of $\text{R}''\text{MgX}$ to RCHO and $\text{R}_2\text{C}=\text{O}$ 743
- 20.7 Reaction of $\text{R}''\text{MgX}$ or $\text{R}''\text{Li}$ with RCOCl and RCOOR' 751
- 20.8 Carboxylation—Reaction of RMgX with CO_2 754
- 20.9 1,2-Addition to an α,β -Unsaturated Carbonyl Compound 756
- 20.10 1,4-Addition to an α,β -Unsaturated Carbonyl Compound 756
- Chapter 21 Aldehydes and Ketones—Nucleophilic Addition**
- 21.1 General Mechanism—Nucleophilic Addition 786
- 21.2 General Mechanism—Acid-Catalyzed Nucleophilic Addition 787
- 21.3 Nucleophilic Addition of ^-CN —Cyanohydrin Formation 791
- 21.4 The Wittig Reaction 794
- 21.5 Imine Formation from an Aldehyde or Ketone 798
- 21.6 Enamine Formation from an Aldehyde or Ketone 800
- 21.7 Base-Catalyzed Addition of H_2O to a Carbonyl Group 803
- 21.8 Acid-Catalyzed Addition of H_2O to a Carbonyl Group 803
- 21.9 Acetal Formation—Part [1] Formation of a Hemiacetal 806
- 21.10 Acetal Formation—Part [2] Formation of the Acetal 806
- 21.11 Acid-Catalyzed Cyclic Hemiacetal Formation 810
- 21.12 A Cyclic Acetal from a Cyclic Hemiacetal 811
- Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution**
- 22.1 General Mechanism—Nucleophilic Acyl Substitution 839
- 22.2 Conversion of Acid Chlorides to Anhydrides 843
- 22.3 Conversion of Acid Chlorides to Carboxylic Acids 843
- 22.4 Conversion of an Anhydride to an Amide 844
- 22.5 Conversion of Carboxylic Acids to Acid Chlorides 846
- 22.6 Fischer Esterification—Acid-Catalyzed Conversion of Carboxylic Acids to Esters 848
- 22.7 Conversion of Carboxylic Acids to Amides with DCC 850
- 22.8 Acid-Catalyzed Hydrolysis of an Ester to a Carboxylic Acid 851
- 22.9 Base-Promoted Hydrolysis of an Ester to a Carboxylic Acid 852
- 22.10 Amide Hydrolysis in Base 856
- 22.11 Hydrolysis of a Nitrile in Base 864
- 22.12 Reduction of a Nitrile with LiAlH_4 865
- 22.13 Reduction of a Nitrile with DIBAL-H 866
- 22.14 Addition of Grignard and Organolithium Reagents (R-M) to Nitriles 866
- Chapter 23 Substitution Reactions of Carbonyl Compounds at the α Carbon**
- 23.1 Tautomerization in Acid 883
- 23.2 Tautomerization in Base 884
- 23.3 Acid-Catalyzed Halogenation at the α Carbon 893
- 23.4 Halogenation at the α Carbon in Base 894
- 23.5 The Haloform Reaction 895
- Chapter 24 Carbonyl Condensation Reactions**
- 24.1 The Aldol Reaction 918
- 24.2 Dehydration of β -Hydroxy Carbonyl Compounds with Base 920
- 24.3 The Intramolecular Aldol Reaction 927
- 24.4 The Claisen Reaction 929
- 24.5 The Dieckmann Reaction 933
- 24.6 The Michael Reaction 935

- 24.7 The Robinson Annulation—Part [A] Michael Addition to Form a 1,5-Dicarbonyl Compound 937
- 24.8 The Robinson Annulation—Part [B] Intramolecular Aldol Reaction to Form a 2-Cyclohexenone 937
- Chapter 25 Amines**
- 25.1 The E2 Mechanism for the Hofmann Elimination 978
- 25.2 Formation of a Diazonium Salt from a 1° Amine 981
- 25.3 Formation of an *N*-Nitrosamine from a 2° Amine 982
- 25.4 Azo Coupling 987
- Chapter 26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis**
- 26.1 Suzuki Reaction 1008
- 26.2 Heck Reaction 1011
- 26.3 Formation of Dichlorocarbene 1013
- 26.4 Addition of Dichlorocarbene to an Alkene 1013
- 26.5 Simmons–Smith Reaction 1015
- 26.6 Olefin Metathesis: $2 \text{RCH}=\text{CH}_2 \rightarrow \text{RCH}=\text{CHR} + \text{CH}_2=\text{CH}_2$ 1017
- Chapter 27 Carbohydrates**
- 27.1 Glycoside Formation 1043
- 27.2 Glycoside Hydrolysis 1044
- Chapter 28 Amino Acids and Proteins**
- 28.1 Formation of an α -Amino Nitrile 1080
- 28.2 Edman Degradation 1092
- Chapter 29 Lipids**
- 29.1 Biological Formation of Geranyl Diphosphate 1135
- 29.2 Biological Formation of Farnesyl Diphosphate 1136
- 29.3 Isomerization of Geranyl Diphosphate to Neryl Diphosphate 1137
- Chapter 30 Synthetic Polymers**
- 30.1 Radical Polymerization of $\text{CH}_2=\text{CHPh}$ 1151
- 30.2 Forming Branched Polyethylene During Radical Polymerization 1153
- 30.3 Cationic Polymerization of $\text{CH}_2=\text{CHZ}$ 1154
- 30.4 Anionic Polymerization of $\text{CH}_2=\text{CHZ}$ 1154
- 30.5 Ziegler–Natta Polymerization of $\text{CH}_2=\text{CH}_2$ 1158

List of Selected Applications

Applications make any subject seem more relevant and interesting—for nonmajors and majors alike. The following is a list of the most important biological, medicinal, and environmental applications that have been integrated throughout *Organic Chemistry*. Each chapter opener showcases a current application relating to the chapter's topic. (Code: G = general; M = medicinal; B = biological; E = environmental)

Prologue

- G, E Examples of simple organic compounds—methane, a component in natural gas; ethanol, the alcohol in beer and wine; and trichlorofluoromethane, a refrigerant and aerosol propellant implicated in ozone destruction
- M Some complex organic compounds that are useful drugs—the antibiotic amoxicillin, the antidepressant fluoxetine (Prozac), and AZT, a drug used to treat HIV
- B Ginkgolide B, principal component of extracts from the ginkgo tree, *Ginkgo biloba*

Chapter 1 Structure and Bonding

- M L-Dopa, the drug of choice for the treatment of Parkinson's disease (Opener, Section 1.13)
- M Fosamax, a drug used to prevent bone loss in women (Section 1.4B)

Chapter 2 Acids and Bases

- M The acid–base chemistry of aspirin, the most widely used over-the-counter drug (Opener, Section 2.7)
- M Pseudoephedrine, the nasal decongestant in Sudafed (Section 2.5, Problem 2.18)

Chapter 3 Introduction to Organic Molecules and Functional Groups

- B Vitamin C, a water-soluble vitamin needed in the formation of the protein collagen (Opener)
- B How geckos stick to walls and ceilings (Section 3.3B)
- E Solubility principles and the pollutants MTBE and PCBs in the environment (Section 3.4C)
- B How structure explains the fat solubility of vitamin A and the water solubility of vitamin C (Section 3.5)
- G How soap cleans away dirt (Section 3.6)
- B The structure of the cell membrane (Section 3.7A)
- M How ionophores like the antibiotic valinomycin transport ions across a cell membrane (Section 3.7B)
- B Hydrogen bonding in DNA, deoxyribonucleic acid, the high molecular weight compound that stores the genetic information of an organism (Section 3.9)

Chapter 4 Alkanes

- G Alkanes, the major constituents of petroleum, which is refined to produce gasoline, diesel fuel, and home heating oil (Opener, Section 4.7)
- E The combustion of alkanes, the concentration of atmospheric carbon dioxide, and global warming (Section 4.14B)
- B An introduction to lipids, biomolecules whose properties can be explained by understanding alkane chemistry; cholesterol in the cell membrane (Section 4.15)

Chapter 5 Stereochemistry

- M The importance of the three-dimensional structure in the pain reliever (*S*)-naproxen (Opener)
- B How differences in the three-dimensional structure of starch and cellulose affect their shape and function (Section 5.1)
- M The three-dimensional structure of thalidomide, the anti-nausea drug that caused catastrophic birth defects (Section 5.5)
- M How mirror image isomers can have drastically different properties—the analgesic ibuprofen, the antidepressant fluoxetine, and the anti-inflammatory agent naproxen (Section 5.13)
- B The sense of smell—How mirror image isomers can smell differently (Section 5.13)

Chapter 6 Understanding Organic Reactions

- B Energy changes in the metabolism of glucose and the combustion of isooctane, a high-octane component of gasoline (Opener, Section 6.4)
- B Enzymes, biological catalysts (Section 6.11)

Chapter 7 Alkyl Halides and Nucleophilic Substitution

- B The biological synthesis of adrenaline, the hormone secreted in response to a strenuous or challenging activity (Opener, Section 7.12)
- E CFCs and DDT, two polyhalogenated compounds once widely used, now discontinued because of adverse environmental effects (Section 7.4)

- B S-Adenosylmethionine (SAM), a nutritional supplement used by the cell in key nucleophilic substitutions that synthesize amino acids, hormones, and neurotransmitters (Section 7.12)
- B How nitrosamines, compounds formed in cured meats preserved with sodium nitrite, are thought to be cancer-causing (Section 7.16)
- M The importance of organic synthesis in preparing useful drugs such as aspirin and taxol, an anticancer drug used to treat breast cancer (Section 7.19)
- Chapter 8 Alkyl Halides and Elimination Reactions**
- E DDE, a degradation product of the pesticide DDT (Opener, Section 8.1)
- B, M Elimination reactions in the synthesis of a prostaglandin, an antimalarial drug, and a female sex hormone (Section 8.4)
- Chapter 9 Alcohols, Ethers, and Epoxides**
- B Palytoxin, a toxic component isolated from marine soft corals of the genus *Palythoa* (Opener, Problem 9.80)
- G, E Ethanol, a gasoline additive and renewable fuel source that can be produced from the fermentation of carbohydrates in grains (Section 9.5)
- M The design of asthma drugs that block the synthesis of leukotrienes, highly potent molecules that contribute to the asthmatic response (Section 9.16)
- B The metabolism of polycyclic aromatic hydrocarbons (PAHs) to carcinogens that disrupt normal cell function resulting in cancer or cell death (Section 9.17)
- Chapter 10 Alkenes**
- B Fats and oils—the properties of saturated and unsaturated fatty acids (Opener, Section 10.6)
- G Ethylene, the starting material for preparing the polymer polyethylene and many other simple compounds used to make a variety of other polymers (Section 10.5)
- B Omega-3 fatty acids, highly unsaturated fatty acids thought to be beneficial for individuals at risk of developing coronary artery disease (Section 10.6, Problem 10.12)
- B The synthesis of the female sex hormone estrone (Section 10.15B)
- M The synthesis of artemisinin, an antimalarial drug isolated from qinghao, a Chinese herbal remedy (Section 10.16)
- Chapter 11 Alkynes**
- M Oral contraceptives (Opener, Section 11.4)
- M Synthetic hormones mifepristone and Plan B, drugs that prevent pregnancy (Section 11.4)
- Chapter 12 Oxidation and Reduction**
- B The metabolism of ethanol, the alcohol in alcoholic beverages (Opener, Section 12.14)
- B The partial hydrogenation of vegetable oils and the formation of “trans fats” (Section 12.4)
- B The use of disparlure, a sex pheromone, in controlling the spread of gypsy moths (Section 12.8)
- G Blood alcohol screening (Section 12.12)
- E Green chemistry—environmentally benign oxidation reactions (Section 12.13)
- B The synthesis of insect pheromones using asymmetric epoxidation (Section 12.15)
- Chapter 13 Mass Spectrometry and Infrared Spectroscopy**
- M Infrared spectroscopy and the structure determination of penicillin (Opener, Section 13.8)
- M Using instrumental analysis to detect THC, the active component in marijuana, and other drugs (Section 13.4B)
- B Mass spectrometry and high molecular weight biomolecules (Section 13.4C)
- Chapter 14 Nuclear Magnetic Resonance Spectroscopy**
- M Modern spectroscopic methods and the structure of the hormone melatonin (Opener, Problem 14.26)
- M Magnetic resonance imaging (MRI) and medicine (Section 14.12)
- Chapter 15 Radical Reactions**
- G Polystyrene, a common synthetic polymer used in packaging materials and beverage cups (Opener)
- E Ozone destruction and CFCs (Section 15.9)
- B The oxidation of unsaturated lipids by radical reactions (Section 15.11)
- M, B Two antioxidants—naturally occurring vitamin E and synthetic BHT (Section 15.12)
- G The formation of useful polymers from monomers by radical reactions (Section 15.14)
- Chapter 16 Conjugation, Resonance, and Dienes**
- M Lycopene, a highly unsaturated red pigment found in tomatoes, watermelon, and other fruits (Opener, Section 16.7)
- B The Diels–Alder reaction and the synthesis of tetrodotoxin, a toxin isolated from the puffer fish (Section 16.12)
- M The synthesis of steroids by Diels–Alder reactions (Section 16.14C)
- G Why lycopene and other highly conjugated compounds are colored (Section 16.15A)
- G How sunscreens work (Section 16.15B)

Chapter 17 Benzene and Aromatic Compounds

- B, M Capsaicin, the spicy component of hot peppers and active ingredient in topical creams for the treatment of chronic pain (Opener)
- G Polycyclic aromatic hydrocarbons (PAHs), constituents of cigarette smoke and diesel exhaust (Section 17.5)
- M Examples of common drugs that contain an aromatic ring—Zoloft, Valium, Novocain, Viracept, Viagra, and Claritin (Section 17.5)
- B Histamine and scombroid fish poisoning (Section 17.8)
- G Diamond, graphite, and buckminsterfullerene (Section 17.11)

Chapter 18 Electrophilic Aromatic Substitution

- M The synthesis of the hallucinogen LSD (Opener, Section 18.5D)
- M, E Examples of biologically active aryl chlorides—the drugs bupropion and chlorpheniramine, and 2,4-D and 2,4,5-T, herbicide components of the defoliant Agent Orange (Section 18.3)
- M Benzocaine, the active ingredient in the over-the-counter topical anesthetic Orajel (Section 18.14C)

Chapter 19 Carboxylic Acids and the Acidity of the O–H Bond

- G Hexanoic acid, the foul-smelling carboxylic acid in ginkgo seeds (Opener, Problem 19.51)
- B GHB (4-hydroxybutanoic acid), an illegal recreational intoxicant used as a “date rape” drug (Section 19.5)
- M, B How NSAIDs block the synthesis of prostaglandins to prevent inflammation (Section 19.6)
- B An introduction to amino acids, the building blocks of proteins; why vegetarians must have a balanced diet (Section 19.14)

Chapter 20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

- B The use of juvenile hormone mimics to control certain insect populations; the use of organometallic reagents to synthesize the C₁₈ juvenile hormone (Opener, Section 20.10C)
- B, M Reduction reactions in the synthesis of the analgesic ibuprofen and the perfume component muscone (Section 20.4)
- M The synthesis of the long-acting bronchodilator salmeterol (Section 20.6A)
- B Biological oxidation–reduction reactions with the coenzymes NADH and NAD⁺ (Section 20.6B)
- B The synthesis of the marine neurotoxin ciguatoxin CTX3C (Section 20.7)
- M The use of organometallic reagents to synthesize the oral contraceptive ethinylestradiol (Section 20.10C)

Chapter 21 Aldehydes and Ketones—Nucleophilic Addition

- M Digoxin, a naturally occurring drug isolated from the woolly foxglove plant and used to treat congestive heart failure (Opener, Problem 21.40)
- B Naturally occurring cyanohydrin derivatives—linamarin, from cassava root; and amygdalin, often called laetrile, from apricot, peach, and wild cherry pits (Section 21.9B)
- B The use of the Wittig reaction in the synthesis of β -carotene, the orange pigment in carrots (Section 21.10B)
- B The role of rhodopsin in the chemistry of vision (Section 21.11B)

Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

- G Nylon, the first synthetic fiber (Opener)
- M, B Compounds that contain an ester—vitamin C; cocaine, addictive stimulant from the leaves of the coca plant; and FK506, an immunosuppressant (Section 22.6)
- M, B Useful amides—proteins, the polyamide met-enkephalin, the anticancer drug Gleevec, the penicillin antibiotics, and the cephalosporin antibiotics (Section 22.6)
- G The synthesis of the insect repellent DEET (Section 22.8)
- M The use of acylation in the synthesis of aspirin, acetaminophen (the active ingredient in Tylenol), and heroin (Section 22.9)
- B The hydrolysis of triacylglycerols in the metabolism of lipids (Section 22.12A)
- G Olestra, a fake fat (Section 22.12A)
- G The synthesis of soap (Section 22.12B)
- M The mechanism of action of β -lactam antibiotics like penicillin (Section 22.14)
- G Natural and synthetic fibers—nylon and polyesters (Section 22.16)
- B Biological acylation reactions (Section 22.17)
- M Cholesteryl esters in plaque, the deposits that form on the walls of arteries (Section 22.17)

Chapter 23 Substitution Reactions of Carbonyl Compounds at the α Carbon

- M The synthesis of tamoxifen, an anticancer drug used in the treatment of breast cancer (Opener, Section 23.8C)
- M The synthesis of the antimalarial drug quinine by an intramolecular substitution reaction (Section 23.7C)

Chapter 24 Carbonyl Condensation Reactions

- M The synthesis of the anti-inflammatory agent ibuprofen (Opener, Problem 24.19)
- B The synthesis of periplanone B, sex pheromone of the female American cockroach (Section 24.3)
- B Synthesis of *ar*-turmerone, a component of turmeric, a principal ingredient in curry powder (Section 24.3)

- B The synthesis of the steroid progesterone by an intramolecular aldol reaction (Section 24.4)
- B The synthesis of the female sex hormone estrone by a Michael reaction (Section 24.8)

Chapter 25 Amines

- B Caffeine, an alkaloid found in coffee, tea, and cola beverages (Opener)
- M Histamine, antihistamines, and antiulcer drugs like Tagamet (cimetidine) (Section 25.6B)
- B Naturally occurring alkaloids—atropine from the poisonous nightshade plant, nicotine from tobacco, and coniine from hemlock (Section 25.6B)
- B, M Biologically active derivatives of 2-phenylethylamine—adrenaline, noradrenaline, methamphetamine, mescaline, and dopamine (Section 25.6C)
- B, M The neurotransmitter serotonin and widely used antidepressants called SSRIs (selective serotonin reuptake inhibitors) (Section 25.6C)
- M The synthesis of methamphetamine (Section 25.7C)
- M Drugs such as the antihistamine diphenhydramine, sold as water-soluble ammonium salts (Section 25.9)
- G Azo dyes (Section 25.15)
- G Perkin's mauveine and synthetic dyes (Section 25.16A)
- G How dyes bind to fabric (Section 25.16B)
- M Sulfa drugs (Section 25.17)

Chapter 26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis

- B, E Bombykol, the sex pheromone of the female silkworm moth (Opener, Section 26.2B)
- E Pyrethrin I, a biodegradable insecticide isolated from chrysanthemums (Section 26.4, Problem 26.33)
- M Ring-closing metathesis and the synthesis of epothilone A, an anticancer drug, and Sch38516, an antiviral agent (Section 26.6)

Chapter 27 Carbohydrates

- B Lactose, the carbohydrate in milk (Opener)
- B Glucose, the most common simple sugar (Section 27.6)
- B, M Naturally occurring glycosides—salicin from willow bark and solanine, isolated from the deadly nightshade plant (Section 27.7C)
- G Rebaudioside A (trade name Truvia), a sweet glycoside from the stevia plant (Section 27.7C)
- B Common disaccharides—maltose from malt, lactose from milk, and sucrose, common table sugar (Section 27.12)
- G Artificial sweeteners (Section 27.12C)
- B Common polysaccharides—cellulose, starch, and glycogen (Section 27.13)
- B, M Glucosamine, an over-the-counter remedy used for osteoarthritis, and chitin, the carbohydrate that gives rigidity to crab shells (Section 27.14A)
- B *N*-Glycosides and the structure of DNA (Section 27.14B)

Chapter 28 Amino Acids and Proteins

- B Myoglobin, the protein that stores oxygen in tissues (Opener, Section 28.10C)
- B The naturally occurring amino acids (Section 28.1)
- B The preparation of polypeptides and proteins using automated peptide synthesis—the Merrifield method (Section 28.8)
- B The structure of spider silk (Section 28.9B)
- M The structure of insulin (Section 28.9C)
- B β -Keratin, the protein in hair (Section 28.10A)
- B Collagen, the protein in connective tissue (Section 28.10B)
- B The globular protein hemoglobin; the structure of sickle cell hemoglobin (Section 28.10C)

Chapter 29 Lipids

- B Cholesterol, the most prominent steroid (Opener, Section 29.8B)
- B Triacylglycerols, the components of fats and oils (Section 29.3)
- B Energy storage and the metabolism of fats (Section 29.3)
- B The phospholipids in cell membranes (Section 29.4)
- B Fat-soluble vitamins—A, D, E, and K (Section 29.5)
- B The eicosanoids, a group of biologically active lipids that includes the prostaglandins and leukotrienes (Section 29.6)
- M Vioxx, Bextra, and Celebrex—anti-inflammatory drugs (Section 29.6)
- B, M The structure of steroids—cholesterol, female sex hormones, male sex hormones, adrenal cortical steroids, anabolic steroids, and oral contraceptives (Section 29.8)
- M Cholesterol and cholesterol-lowering drugs atorvastatin (Lipitor) and simvastatin (Zocor) (Section 29.8B)

Chapter 30 Synthetic Polymers

- G Polyethylene terephthalate, an easily recycled synthetic polymer used in transparent soft drink containers (Opener, Sections 30.6B and 30.9A)
- G Polyethylene, the plastic in milk jugs and plastic bags, and other chain-growth polymers (Section 30.2)
- G Using Ziegler–Natta catalysts to make high-density polyethylene (Section 30.4)
- B Natural and synthetic rubber (Section 30.5)
- G The synthesis of step-growth polymers—polyamides such as nylon and Kevlar, polyesters such as Dacron, polyurethanes such as spandex, and polycarbonates such as Lexan (Section 30.6)
- M Dissolving sutures (Section 30.6B)
- G Epoxy resins (Section 30.6E)
- E Green polymer synthesis—environmentally benign methods for preparing polymers (Section 30.8)
- E Polymer recycling (Section 30.9A)
- E Biodegradable polymers (Section 30.9B)

Prologue



What is organic chemistry?
Some representative organic molecules
Ginkgolide B—A complex organic compound from the ginkgo tree

Organic chemistry. You might wonder how a discipline that conjures up images of eccentric old scientists working in basement laboratories is relevant to you, a student in the twenty-first century.

Consider for a moment the activities that occupied your past 24 hours. You likely showered with soap, drank a caffeinated beverage, ate at least one form of starch, took some medication, read a newspaper, listened to a CD, and traveled in a vehicle that had rubber tires and was powered by fossil fuels. If you did any *one* of these, your life was touched by organic chemistry.

What Is Organic Chemistry?

- Organic chemistry is the chemistry of compounds that contain the element carbon.

It is one branch in the entire field of chemistry, which encompasses many classical subdisciplines including inorganic, physical, and analytical chemistry, and newer fields such as bioinorganic chemistry, physical biochemistry, polymer chemistry, and materials science.

Organic chemistry was singled out as a separate discipline for historical reasons. Originally, it was thought that compounds in living things, termed *organic compounds*, were fundamentally different from those in nonliving things, called *inorganic compounds*. Although we have known for more than 150 years that this distinction is artificial, the name *organic* persists. Today the term refers to the study of the compounds that contain carbon, many of which, incidentally, are found in living organisms.

It may seem odd that a whole discipline is devoted to the study of a single element in the periodic table, when more than 100 elements exist. It turns out, though, that there are far more organic compounds than any other type. **Organic chemicals affect virtually every facet of our lives, and for this reason, it is important and useful to know something about them.**

Clothes, foods, medicines, gasoline, refrigerants, and soaps are composed almost solely of organic molecules. Some, like cotton, wool, or silk are naturally occurring; that is, they can be isolated directly from natural sources. Others, such as nylon and polyester, are synthetic, meaning they are produced by chemists in the laboratory. By studying the principles and concepts of organic chemistry, you can learn more about compounds such as these and how they affect the world around you.

Realize, too, what organic chemistry has done for us. Organic chemistry has made available both comforts and necessities that were previously nonexistent, or reserved for only the wealthy. We have seen an enormous increase in life span, from 47 years in 1900 to over 70 years currently. To a large extent this is due to the isolation and synthesis of new drugs to fight infections and the availability of vaccines for childhood diseases. Chemistry has also given us the tools to control insect populations that spread disease, and there is more food for all because of fertilizers, pesticides, and herbicides. Our lives would be vastly different today without the many products that result from organic chemistry (Figure 1).

Figure 1

Products of organic chemistry
used in medicine

a. Oral contraceptives



b. Plastic syringes



c. Antibiotics





d. Synthetic heart valves




- Organic chemistry has given us contraceptives, plastics, antibiotics, and the knitted material used in synthetic heart valves.

Some Representative Organic Molecules

Perhaps the best way to appreciate the variety of organic molecules is to look at a few. Three simple organic compounds are **methane**, **ethanol**, and **trichlorofluoromethane**.



- **Methane**, the simplest of all organic compounds, contains one carbon atom. Methane—the main component of natural gas—occurs widely in nature. Like other hydrocarbons—organic compounds that contain only carbon and hydrogen—**methane is combustible**; that is, it burns in the presence of oxygen. Methane is the product of the anaerobic (without air) decomposition of organic matter by bacteria. The natural gas we use today was formed by the decomposition of organic material millions of years ago. Hydrocarbons such as methane are discussed in Chapter 4.
- **Ethanol**, the alcohol present in beer, wine, and other alcoholic beverages, is formed by the fermentation of sugar, quite possibly the oldest example of organic synthesis. Ethanol can also be made in the lab by a totally different process, but **the ethanol produced in the lab**




is identical to the ethanol produced by fermentation. Alcohols including ethanol are discussed in Chapter 9.

- **Trichlorofluoromethane** is a member of a class of molecules called chlorofluorocarbons or CFCs, which contain one or two carbon atoms and several halogens. Trichlorofluoromethane is an unusual organic molecule in that **it contains no hydrogen atoms**. Because it has a low molecular weight and is easily vaporized, trichlorofluoromethane has been used as an aerosol propellant and refrigerant. It and other CFCs have been implicated in the destruction of the stratospheric ozone layer, as is discussed in Chapter 15.

Because more complicated organic compounds contain many carbon atoms, organic chemists have devised a shorthand to draw them. Keep in mind the following when examining these structures:


- **Each solid line represents a two-electron covalent bond.**
- **When no atom is drawn at the corner of a ring, an organic chemist assumes it to be carbon.**

For example, in the six-membered ring drawn, there is one carbon atom at each corner of the hexagon.




Three complex organic molecules that are important medications are **amoxicillin**, **fluoxetine**, and **AZT**.


- **Amoxicillin** is one of the most widely used antibiotics in the penicillin family. The discovery and synthesis of such antibiotics in the twentieth century have made routine the treatment of infections that were formerly fatal. You were likely given some amoxicillin to treat an ear infection when you were a child. The penicillin antibiotics are discussed in Chapter 22.



- **Fluoxetine** is the generic name for the antidepressant **Prozac**. Prozac was designed and synthesized by chemists in the laboratory, and is now produced on a large scale in chemical factories. Because it is safe and highly effective in treating depression, Prozac is widely prescribed. Over 40 million individuals worldwide have used Prozac since 1986.




- **AZT**, the abbreviation for **azidodeoxythymidine**, is a drug that treats human immunodeficiency virus (HIV), the virus that causes acquired immune deficiency syndrome (AIDS). Also known by its generic name **zidovudine**, AZT represents a chemical success to a different challenge: synthesizing agents that combat viral infections.




Other complex organic compounds having interesting properties are **capsaicin** and **DDT**.

- **Capsaicin**, one member of a group of compounds called *vanilloids*, is responsible for the characteristic spiciness of hot peppers. It is the active ingredient in pepper sprays used for personal defense and topical creams used for pain relief.



- **DDT**, the abbreviation for **dichlorodiphenyltrichloroethane**, is a pesticide once called “miraculous” by Winston Churchill because of the many lives it saved by killing disease-carrying mosquitoes. DDT use is now banned in the United States and many developed countries because it is a nonspecific insecticide that persists in the environment.



What are the common features of these organic compounds?

- **All organic compounds contain carbon atoms and most contain hydrogen atoms.**
- **All the carbon atoms have four bonds.** A stable carbon atom is said to be *tetravalent*.
- Other elements may also be present. **Any atom that is not carbon or hydrogen is called a heteroatom.** Common heteroatoms include N, O, S, P, and the halogens.
- Some compounds have **chains** of atoms and some compounds have **rings**.

These features explain why there are so many organic compounds: **Carbon forms four strong bonds with itself and other elements. Carbon atoms combine together to form rings and chains.**

Ginkgolide B—A Complex Organic Compound from the Ginkgo Tree

Let's complete this discussion with **ginkgolide B** ($C_{20}H_{24}O_{10}$), a complex organic compound isolated from the ginkgo tree *Ginkgo biloba*, the oldest seed-producing plant that currently lives on earth (Figure 2). Also called the maidenhair tree, *Ginkgo biloba* has existed for over 280 million years, and fossil records indicate that it has undergone little significant evolutionary change for eons. Extracts from the roots, bark, leaves, and seeds have been used in traditional Chinese



- Hydrogen atoms bonded to ring carbons are omitted in the structure of ginkgolide B, a convention described in Section 1.7.

medicine to treat asthma and improve blood circulation. Today, ginkgo extracts comprise the most widely taken herbal supplements, used by some individuals to enhance memory and treat dementia. Recent findings of the National Institutes of Health, however, have cast doubt on its efficacy in providing any long-term improvement in cognitive function.

In 1932 ginkgolide B was one of four components isolated from ginkgo extracts, and its structure was determined in 1967. Although its rigid ring system of 20 carbons contained in a compact three-dimensional shape made it a challenging molecule to prepare in the laboratory, Professor E. J. Corey and co-workers at Harvard University reported the synthesis of ginkgolide B in the laboratory in 1988.

In this introduction, we have seen a variety of molecules that have diverse structures. They represent a minuscule fraction of the organic compounds currently known and the many thousands that are newly discovered or synthesized each year. The principles you learn in organic chemistry will apply to all of these molecules, from simple ones like methane and ethanol, to complex ones like capsaicin and ginkgolide B. It is these beautiful molecules, their properties, and their reactions that we will study in organic chemistry.

WELCOME TO THE WORLD OF ORGANIC CHEMISTRY!

1

Structure and Bonding

- 1.1 The periodic table
- 1.2 Bonding
- 1.3 Lewis structures
- 1.4 Lewis structures continued
- 1.5 Resonance
- 1.6 Determining molecular shape
- 1.7 Drawing organic structures
- 1.8 Hybridization
- 1.9 Ethane, ethylene, and acetylene
- 1.10 Bond length and bond strength
- 1.11 Electronegativity and bond polarity
- 1.12 Polarity of molecules
- 1.13 L-Dopa—A representative organic molecule



L-Dopa, also called levodopa, was first isolated from seeds of the broad bean plant *Vicia faba* in 1913. Since 1967 it has been the drug of choice for the treatment of Parkinson's disease, a debilitating illness that results from the degeneration of neurons that produce the neurotransmitter dopamine in the brain. L-Dopa is an oral medication that is transported to the brain by the bloodstream, where it is converted to dopamine. Since L-dopa must be taken in large doses with some serious side effects, today it is often given with other drugs that lessen its negative impact on an individual. In Chapter 1, we learn about the structure, bonding, and properties of organic molecules like L-dopa.

Before examining organic molecules in detail, we must review some important features about structure and bonding learned in previous chemistry courses. We will discuss these concepts primarily from an organic chemist's perspective, and spend time on only the particulars needed to understand organic compounds.


Important topics in Chapter 1 include drawing Lewis structures, predicting the shape of molecules, determining what orbitals are used to form bonds, and how electronegativity affects bond polarity. Equally important is Section 1.7 on drawing organic molecules, both shorthand methods routinely used for simple and complex compounds, as well as three-dimensional representations that allow us to more clearly visualize them.

1.1 The Periodic Table

All matter is composed of the same building blocks called **atoms**. There are two main components of an atom.

- The **nucleus** contains positively charged **protons** and uncharged **neutrons**. Most of the mass of the atom is contained in the nucleus.
- The **electron cloud** is composed of negatively charged **electrons**. The electron cloud comprises most of the volume of the atom.

Schematic of an atom



The charge on a proton is equal in magnitude but opposite in sign to the charge on an electron. In a neutral atom, the **number of protons in the nucleus equals the number of electrons**. This quantity, called the **atomic number**, is unique to a particular element. For example, every neutral carbon atom has an atomic number of six, meaning it has six protons in its nucleus and six electrons surrounding the nucleus.

In addition to neutral atoms, we will also encounter **charged ions**.

- A **cation** is positively charged and has fewer electrons than its neutral form.
- An **anion** is negatively charged and has more electrons than its neutral form.

The number of neutrons in the nucleus of a particular element can vary. **Isotopes** are two atoms of the same element having a different number of neutrons. The **mass number** of an atom is the total number of protons and neutrons in the nucleus. Isotopes have different mass numbers.


Isotopes of carbon and hydrogen are sometimes used in organic chemistry, as we will see in Chapter 14.

- The most common isotope of hydrogen has one proton and no neutrons in the nucleus, but 0.02% of hydrogen atoms have one proton and one neutron. This isotope of hydrogen is called **deuterium**, and is sometimes symbolized by the letter **D**.
- Most carbon atoms have six protons and six neutrons in the nucleus, but 1.1% have six protons and seven neutrons.

The **atomic weight** is the weighted average of the mass of all isotopes of a particular element, reported in atomic mass units (amu).

Each atom is identified by a one- or two-letter abbreviation that is the characteristic symbol for that element. Carbon is identified by the single letter **C**. Sometimes the atomic number is indicated as a subscript to the left of the element symbol, and the mass number is indicated as a superscript, as shown in Figure 1.1.

an s orbital (in the same shell) because its electron density is farther away from the nucleus. A p orbital is filled with electrons only after an s orbital of the same shell is full.



Let's now look at the elements in the first and second rows of the periodic table.

The First Row

The first row of the periodic table is formed by adding electrons to the first shell of orbitals around the nucleus. There is only one orbital in the first shell, called the **1s orbital**.

- **Remember:** Each orbital can have a maximum of two electrons.

As a result, there are **two elements in the first row**, one having one electron added to the $1s$ orbital, and one having two. The element hydrogen (H) has what is called a $1s^1$ configuration with one electron in the $1s$ orbital, and helium (He) has a $1s^2$ configuration with two electrons in the $1s$ orbital.




The Second Row

Every element in the second row has a filled first shell of electrons. Thus, all second-row elements have a $1s^2$ configuration. These electrons in the inner shell of orbitals are called **core electrons**, and are not usually involved in the chemistry of a particular element.

Each element in the second row of the periodic table has four orbitals available to accept additional electrons:

- **one 2s orbital**, the s orbital in the second shell
- **three 2p orbitals**, all dumbbell-shaped and perpendicular to each other along the x , y , and z axes



Because each of the four orbitals in the second shell can hold two electrons, there is a **maximum capacity of eight electrons** for elements in the second row. The second row of the periodic table consists of eight elements, obtained by adding electrons to the $2s$ and three $2p$ orbitals.

group number	→ 1A	2A	3A	4A	5A	6A	7A	8A
second row	→ Li	Be	B	C	N	O	F	Ne
number of valence electrons	→ 1	2	3	4	5	6	7	8

The outermost electrons are called **valence electrons**. The valence electrons are more loosely held than the electrons closer to the nucleus, and as such, they participate in chemical reactions. **The group number of a second-row element reveals its number of valence electrons.** For example, carbon in group 4A has four valence electrons, and oxygen in group 6A has six.

Problem 1.1 While the most common isotope of nitrogen has a mass number of 14 (nitrogen-14), a radioactive isotope of nitrogen has a mass number of 13 (nitrogen-13). Nitrogen-13 is used in PET (positron emission tomography) scans by physicians to monitor brain activity and diagnose dementia. For each isotope, give the following information: (a) the number of protons; (b) the number of neutrons; (c) the number of electrons in the neutral atom; and (d) the group number.

Problem 1.2 Consider the three atoms: [1] $^{31}_{15}\text{P}$; [2] $^{19}_{9}\text{F}$; and [3] ^2_1H . For each atom give the following information: (a) the atomic number; (b) the total number of electrons in the neutral atom; (c) the number of valence electrons; and (d) the group number.

1.2 Bonding

Until now our discussion has centered on individual atoms, but it is more common in nature to find two or more atoms joined together.

- **Bonding is the joining of two atoms in a stable arrangement.**

Bonding may occur between atoms of the same or different elements. Bonding is a favorable process because it always leads to **lowered energy and increased stability**. Joining two or more elements forms **compounds**. Although only about 100 elements exist, more than 30 million compounds are known. Examples of compounds include hydrogen gas (H_2), formed by joining two hydrogen atoms, and methane (CH_4), the simplest organic compound, formed by joining a carbon atom with four hydrogen atoms.

One general rule governs the bonding process.

- **Through bonding, atoms attain a complete outer shell of valence electrons.**

Alternatively, because the noble gases in column 8A of the periodic table are especially stable as atoms having a filled shell of valence electrons, the general rule can be restated.

- **Through bonding, atoms attain a stable noble gas configuration of electrons.**

What does this mean for first- and second-row elements? **A first-row element like hydrogen can accommodate two electrons around it.** This would make it like the noble gas helium at the end of the same row. **A second-row element is most stable with eight valence electrons around it** like neon. Elements that behave in this manner are said to follow the **octet rule**.

There are two different kinds of bonding: **ionic bonding** and **covalent bonding**.

- **Ionic bonds result from the transfer of electrons from one element to another.**
- **Covalent bonds result from the sharing of electrons between two nuclei.**


The type of bonding is determined by the location of an element in the periodic table. An ionic bond generally occurs when elements on the **far left** side of the periodic table combine with elements on the **far right** side, ignoring the noble gases, which form bonds only rarely. **The resulting ions are held together by extremely strong electrostatic interactions.** A positively charged cation formed from the element on the left side attracts a negatively charged anion formed from the element on the right side. The resulting **salts** are seen in many of the inorganic compounds with which you are familiar. Sodium chloride (NaCl) is common table salt, and potassium iodide (KI) is an essential nutrient added to make iodized salt.



Atoms readily form ionic bonds when they can attain a noble gas configuration by gaining or losing just one or two electrons.


Ionic compounds form extended crystal lattices that maximize the positive and negative electrostatic interactions. In NaCl, each positively charged Na^+ ion is surrounded by six negatively charged Cl^- ions, and each Cl^- ion is surrounded by six Na^+ ions.

NaCl—An ionic crystalline lattice



Lithium fluoride, LiF, is an example of an ionic compound.

- The element **lithium**, located in group 1A of the periodic table, has just one valence electron in its second shell. If this electron is lost, lithium forms the cation Li^+ having no electrons in the second shell. However, it will have a stable electronic arrangement with two electrons in the first shell like helium.
- The element **fluorine**, located in group 7A of the periodic table, has seven valence electrons. By gaining one it forms the anion F^- , which has a filled valence shell (an octet of electrons), like neon.
- Thus, lithium fluoride is a stable ionic compound.



- The transfer of electrons forms stable salts composed of cations and anions.

A **compound** may have either ionic or covalent bonds. A **molecule** has only covalent bonds.

The second type of bonding, **covalent bonding**, occurs with elements like carbon in the middle of the periodic table, which would otherwise have to gain or lose several electrons to form an ion with a complete valence shell. A **covalent bond is a two-electron bond**, and a compound with covalent bonds is called a **molecule**. Covalent bonds also form between two elements from the same side of the table, such as two hydrogen atoms or two chlorine atoms. H_2 , Cl_2 , and CH_4 are all examples of covalent molecules.

Problem 1.3


Label each bond in the following compounds as ionic or covalent.

- a. F_2 b. LiBr c. CH_3CH_3 d. NaNH_2

Problem 1.4

An element like fluorine forms either ionic or covalent bonds, depending on the identity of the element to which it bonds. What type of bonding is observed in each compound: (a) NaF , a toothpaste ingredient added to strengthen tooth enamel; (b) CFC_l_3 , a chlorofluorocarbon once widely used as an aerosol propellant? Explain why a difference is observed.

How many covalent bonds will a particular atom typically form? As you might expect, it depends on the location of the atom in the periodic table. In the first row, **hydrogen forms one covalent bond** using its one valence electron. When two hydrogen atoms are joined in a bond, each has a filled valence shell of two electrons.



Second-row elements can have no more than eight valence electrons around them. For neutral molecules, two consequences result.

- **Atoms with one, two, three, or four valence electrons form one, two, three, or four bonds**, respectively, in neutral molecules.
- **Atoms with five or more valence electrons** form enough bonds to give an octet. This results in the following simple equation:

$$\boxed{\text{predicted number of bonds}} = 8 - \text{number of valence electrons}$$

For example, B has three valence electrons, so it forms three bonds, as in BF_3 . N has five valence electrons, so it also forms three bonds ($8 - 5 = 3$ bonds), as in NH_3 .

These guidelines are used in Figure 1.3 to summarize the usual number of bonds formed by the common atoms in organic compounds. Notice that when second-row elements form fewer than four bonds their octets consist of both **bonding (shared) electrons** and **nonbonding (unshared) electrons**. Unshared electrons are also called **lone pairs**.

Nonbonded pair of electrons = unshared pair of electrons = lone pair

Problem 1.5 How many covalent bonds are predicted for each atom?


- a. O b. Al c. Br d. Si

1.3 Lewis Structures

Lewis structures are electron dot representations for molecules. There are three general rules for drawing Lewis structures.

1. Draw only the valence electrons.
2. Give every second-row element no more than eight electrons.
3. Give each hydrogen two electrons.

To draw a Lewis structure for a diatomic molecule like **HF**, recall that hydrogen has one valence electron and fluorine has seven. H and F each donate one electron to form a two-electron bond. The resulting molecule gives both H and F a filled valence shell.



The letter **X** is often used to represent one of the halogens in group 7A: F, Cl, Br, or I.

In a Lewis structure, a **solid line** indicates a two-electron covalent bond.

Figure 1.3

Summary: The usual number of bonds of common neutral atoms

			\downarrow nonbonded electron pair			
	—H	$\begin{array}{c} \\ \text{—C—} \\ \end{array}$	$\begin{array}{c} \\ \text{—}\ddot{\text{N}}\text{—} \\ \end{array}$	$\begin{array}{c} \\ \text{—}\ddot{\text{O}}\text{—} \\ \end{array}$	$\begin{array}{c} \\ \text{—}\ddot{\text{X}}\text{—} \\ \end{array}$	X = F, Cl, Br, I
number of bonds →	1	4	3	2	1	
number of nonbonded electron pairs →	0	0	1	2	3	


1.3A A Procedure for Drawing Lewis Structures

Drawing a Lewis structure for larger molecules is easier if you follow a stepwise procedure.


HOW TO Draw a Lewis Structure

Step [1] Arrange atoms next to each other that you think are bonded together.

- Always place hydrogen atoms and halogen atoms on the periphery because H and X (X = F, Cl, Br, and I) form only one bond each.



- As a first approximation, place no more atoms around an atom than the number of bonds it usually forms.



- In truth, the proper arrangement of atoms may not be obvious, or more than one arrangement may be possible (Section 1.4A). Even in many simple molecules, the connectivity between atoms must be determined experimentally.

Step [2] Count the electrons.

- Count the number of valence electrons from all atoms.
- Add one electron for each negative charge.
- Subtract one electron for each positive charge.
- This sum gives the total number of electrons that must be used in drawing the Lewis structure.

Step [3] Arrange the electrons around the atoms.

- Place a bond between every two atoms, giving two electrons to each H and no more than eight to any second-row atom.
- Use all remaining electrons to fill octets with lone pairs.
- If all valence electrons are used and an atom does not have an octet, form multiple bonds, as shown in Sample Problem 1.3.

Step [4] Assign formal charges to all atoms.


- Formal charges are discussed in Section 1.3C.

Sample Problems 1.1 and 1.2 illustrate how to draw Lewis structures in some simple organic molecules.

Sample Problem 1.1 Draw a Lewis structure for methane, CH_4 .

Solution


Step [1] Arrange the atoms.



Step [2] Count the electrons.

$$\begin{array}{r}
 1 \text{ C} \times 4 \text{ e}^- = 4 \text{ e}^- \\
 4 \text{ H} \times 1 \text{ e}^- = 4 \text{ e}^- \\
 \hline
 \mathbf{8 \text{ e}^- \text{ total}}
 \end{array}$$

Step [3] Add the bonds and lone pairs.



Adding four two-electron bonds around carbon uses all eight valence electrons, and so there are no lone pairs. To check whether a Lewis structure is valid, we must answer YES to three questions:

- Have all the electrons been used?
- Is each H surrounded by two electrons?
- Is each second-row element surrounded by no more than eight electrons?

The answer to all three questions is YES, so the Lewis structure drawn for CH_4 is valid.


Sample Problem 1.2

Draw a Lewis structure for methanol, a compound with molecular formula CH_4O .

Solution


Step [1]

Arrange the atoms.



Step [2]

Count the electrons.




Step [3]

Add the bonds and lone pairs.

Add bonds first...

...then lone pairs.



only 10 electrons used

In Step [3], placing bonds between all atoms uses only 10 electrons, and the O atom does not yet have a complete octet. To complete the structure, give the O atom two nonbonded electron pairs. This uses all 14 electrons, giving every H two electrons and every second-row element eight. We have now drawn a valid Lewis structure.

Problem 1.6

Draw a valid Lewis structure for each species:

- a. CH_3CH_3 b. CH_5N c. CH_3^- d. CH_3Cl

1.3B Multiple Bonds

Sample Problem 1.3 illustrates two examples when multiple bonds are needed in Lewis structures.


Sample Problem 1.3

Draw a Lewis structure for each compound. Assume the atoms are arranged as follows:

- a. ethylene, C_2H_4 b. acetylene, C_2H_2
- $$\begin{array}{c} \text{H C C H} \\ \text{H H} \end{array}$$

Solution


- a. **Ethylene, C_2H_4 :** Follow Steps [1] to [3] to draw a Lewis structure. After placing five bonds between the atoms and adding the two remaining electrons as a lone pair, one C still has no octet.



Step [3] Add the bonds and lone pairs.

Add bonds first...

...then lone pairs.



The number of electrons “owned” by an atom is determined by its number of bonds and lone pairs.


- An atom “owns” *all* of its unshared electrons and *half* of its shared electrons.

$$\boxed{\text{number of electrons owned}} = \boxed{\text{number of unshared electrons}} + \frac{1}{2} \boxed{\text{number of shared electrons}}$$

The number of electrons “owned” by different carbon atoms is indicated in the following examples:



- C shares eight electrons.
- C “owns” **four** electrons.



- Each C shares eight electrons.
- Each C “owns” **four** electrons.




- C shares six electrons.
- C has two unshared electrons.
- C “owns” **five** electrons.

Sample Problem 1.4 illustrates how formal charge is calculated on the atoms of a polyatomic ion. **The sum of the formal charges on the individual atoms equals the net charge on the molecule or ion.**

Sample Problem 1.4

Determine the formal charge on each atom in the ion H_3O^+ .



Solution

For each atom, two steps are needed:

Step [1] Determine the number of electrons an atom “owns.”




Step [2] Subtract this sum from its number of valence electrons.

O atom	H atoms
[1] number of electrons “owned” by O $2 + \frac{1}{2}(6) = 5$	[1] number of electrons “owned” by each H $0 + \frac{1}{2}(2) = 1$
[2] formal charge on O $6 - 5 = \boxed{+1}$	[2] formal charge on each H $1 - 1 = \boxed{0}$

The formal charge on each H is 0. The formal charge on oxygen is +1. The overall charge on the ion is the sum of all of the formal charges; $0 + 0 + 0 + 1 = +1$.


Problem 1.8

Calculate the formal charge on each second-row atom:



Problem 1.9

Draw a Lewis structure for each ion:



Sometimes it is easier to count bonds, rather than shared electrons when determining formal charge.

$$\frac{1}{2}[\text{number of shared electrons}] = \text{number of bonds}$$

When you first add formal charges to Lewis structures, use the procedure in Sample Problem 1.4. With practice, you will notice that certain bonding patterns always result in the same formal charge. For example, any N atom with four bonds (and, thus no lone pairs) has a +1 formal charge. Table 1.1 lists the bonding patterns and resulting formal charges for carbon, nitrogen, and oxygen.

A shortcut method to determine the number of bonds in a Lewis structure is given in the *Student Study Guide/Solutions Manual* (page 1–4).

Table 1.1 Formal Charge Observed with Common Bonding Patterns for C, N, and O


Atom	Number of valence electrons	Formal charge		
		+1	0	-1
C	4	$\begin{array}{c} + \\ \\ \text{---C---} \\ \end{array}$	$\begin{array}{c} \\ \text{---C---} \\ \end{array}$	$\begin{array}{c} \text{---}\ddot{\text{C}}\text{---} \\ \\ - \end{array}$
N	5	$\begin{array}{c} \\ \text{---N}^+ \\ \end{array}$	$\begin{array}{c} \cdot \\ \\ \text{---}\ddot{\text{N}}\text{---} \\ \end{array}$	$\begin{array}{c} \text{---}\ddot{\text{N}}\text{---} \\ \\ - \end{array}$
O	6	$\begin{array}{c} \cdot \\ \text{---}\ddot{\text{O}}^+ \\ \end{array}$	$\begin{array}{c} \cdot \\ \text{---}\ddot{\text{O}}\text{---} \\ \end{array}$	$\begin{array}{c} \text{---}\ddot{\text{O}}\text{---} \\ \\ - \end{array}$

1.4 Lewis Structures Continued

The discussion of Lewis structures concludes with the introduction of isomers and exceptions to the octet rule.

1.4A Isomers

In drawing a Lewis structure for a molecule with several atoms, sometimes more than one arrangement of atoms is possible for a given molecular formula. For example, there are two acceptable arrangements of atoms for the molecular formula $\text{C}_2\text{H}_6\text{O}$.



Both are valid Lewis structures, and both molecules exist. One is called ethanol, and the other, dimethyl ether. These two compounds are called **isomers**.

- **Isomers** are different molecules having the same molecular formula.

Ethanol and dimethyl ether are **constitutional isomers** because they have the same molecular formula, but the **connectivity of their atoms is different**. For example, ethanol has one C–C bond and one O–H bond, whereas dimethyl ether has two C–O bonds. A second class of isomers, called **stereoisomers**, is introduced in Section 4.13B.

Problem 1.10

Draw Lewis structures for each molecular formula.


- a. $\text{C}_2\text{H}_4\text{Cl}_2$ (two isomers) b. $\text{C}_3\text{H}_8\text{O}$ (three isomers) c. C_3H_6 (two isomers)

1.4B Exceptions to the Octet Rule

Most of the common elements in organic compounds—**C, N, O, and the halogens**—follow the octet rule. Hydrogen is a notable exception, because it accommodates only two electrons in bonding. Additional exceptions include boron and beryllium (second-row elements in groups 3A and 2A, respectively), and elements in the third row (particularly phosphorus and sulfur).

Elements in Groups 2A and 3A


Elements in groups 2A and 3A of the periodic table, such as beryllium and boron, do not have enough valence electrons to form an octet in a neutral molecule. Lewis structures for BeH_2 and BF_3 show that these atoms have only four and six electrons, respectively, around the central atom. There is nothing we can do about this! There simply aren't enough electrons to form an octet.



Alendronic acid, sold as a sodium salt under the trade name of **Fosamax**, is used to prevent osteoporosis in women. Osteoporosis decreases bone density, as shown by comparing normal bone (top) with brittle bone (bottom).



four electrons around Be




six electrons around B

Because the Be and B atoms each have less than an octet of electrons, these molecules are highly reactive.

Elements in the Third Row


A second exception to the octet rule occurs with some elements located in the third row and later in the periodic table. These elements have empty *d* orbitals available to accept electrons, and thus they may have **more than eight electrons** around them. For organic chemists, the two most common elements in this category are phosphorus and sulfur, which can have 10 or even 12 electrons around them. Examples of these phosphorus and sulfur compounds include the following:

10 electrons around S




dimethyl sulfoxide
(abbreviated as DMSO)

12 electrons around S



sulfuric acid


10 electrons around each P



alendronic acid

1.5 Resonance

Some molecules can't be adequately represented by a single Lewis structure. For example, two valid Lewis structures can be drawn for the anion $(\text{HCONH})^-$. One structure has a negatively charged N atom and a C–O double bond; the other has a negatively charged O atom and a C–N double bond. These structures are called **resonance structures** or **resonance forms**. A **double-headed arrow** is used to separate two resonance structures.



- **Resonance structures** are two Lewis structures having the *same* placement of atoms but a *different* arrangement of electrons.

Which resonance structure is an accurate representation for $(\text{HCONH})^-$? **The answer is neither of them.** The true structure is a composite of both resonance forms, and is called a **resonance hybrid**. The hybrid shows characteristics of *both* resonance structures.

Each resonance structure implies that electron pairs are localized in bonds or on atoms. In actuality, resonance allows certain electron pairs to be **delocalized** over two or more atoms, and this delocalization of electron density adds stability. **A molecule with two or more resonance structures is said to be resonance stabilized.** We will return to the resonance hybrid in Section 1.5C. First, however, we examine the general principles of resonance theory and learn how to interconvert two or more resonance structures.


1.5A An Introduction to Resonance Theory

Keep in mind the following basic principles of resonance theory.

- Resonance structures are not real. An individual resonance structure does not accurately represent the structure of a molecule or ion. Only the hybrid does.
- Resonance structures are *not* in equilibrium with each other. There is no movement of electrons from one form to another.
- Resonance structures are *not* isomers. Two isomers differ in the arrangement of *both* atoms and electrons, whereas resonance structures differ *only* in the *arrangement of electrons*.

Resonance structures are different ways of drawing the same compound. Two resonance structures are *not* different compounds.


For example, ions **A** and **B** are resonance structures because the atom position is the same in both compounds, but the location of an electron pair is different. In contrast, compounds **C** and **D** are isomers since the atom placement is different; **C** has an O–H bond, and **D** has an additional C–H bond.



Problem 1.11 Classify each pair of compounds as isomers or resonance structures.




Problem 1.12 Considering structures **A–D**, classify each pair of compounds as isomers, resonance structures, or neither: (a) **A** and **B**; (b) **A** and **C**; (c) **A** and **D**; (d) **B** and **D**.



1.5B Drawing Resonance Structures

To draw resonance structures, use the three rules that follow:

Rule [1] Two resonance structures differ in the position of multiple bonds and nonbonded electrons. The placement of atoms and single bonds always stays the same.



- b. **Two** curved arrows tell us to move **two** electron pairs. The second resonance structure has a formal charge of (-1) on O.


Move two electron pairs...




...then calculate formal charges.

This type of resonance-stabilized anion is called an **enolate anion**. Enolates are important intermediates in many organic reactions, and all of Chapters 23 and 24 is devoted to their preparation and reactions.

Problem 1.13 Follow the curved arrows to draw a second resonance structure for each species.



Problem 1.14 Use curved arrow notation to show how the first resonance structure can be converted to the second.



Two resonance structures can have exactly the same kinds of bonds, as they do in the carbocation in Sample Problem 1.5a, or they may have different types of bonds, as they do in the enolate in Sample Problem 1.5b. Either possibility is fine as long as the individual resonance structures are valid Lewis structures.

The ability to draw and manipulate resonance structures is an important skill that will be needed throughout your study of organic chemistry. With practice, you will begin to recognize certain common bonding patterns for which more than one Lewis structure can be drawn. For now, notice that two different resonance structures can be drawn in the following situations:

- When a lone pair is located on an atom directly bonded to a multiple bond.

lone pair adjacent to C=C



lone pair adjacent to C=O

- When an atom bearing a (+) charge is bonded to either a multiple bond or an atom with a lone pair.


(+) charge adjacent to a double bond



(+) charge adjacent to an atom with a lone pair



Problem 1.15 Draw a second resonance structure for each species.



1.5C The Resonance Hybrid


The **resonance hybrid** is the composite of all possible resonance structures. In the resonance hybrid, the electron pairs drawn in different locations in individual resonance structures are *delocalized*.

- The resonance hybrid is more stable than any resonance structure because it delocalizes electron density over a larger volume.

What does the hybrid look like? When all resonance forms are identical, as they were in the carbocation in Sample Problem 1.5a, each resonance form contributes **equally** to the hybrid.


When two resonance structures are different, the hybrid looks more like the “better” resonance structure. The “better” resonance structure is called the **major contributor** to the hybrid, and all others are **minor contributors**. The hybrid is the weighted average of the contributing resonance structures. What makes one resonance structure “better” than another? There are many factors, but for now, we will learn just two.

- A “better” resonance structure is one that has *more bonds* and *fewer charges*.



Comparing resonance structures **X** and **Y**, **X** is the major contributor because it has more bonds and fewer charges. Thus, the hybrid looks more like **X** than **Y**.

How can we draw a hybrid, which has delocalized electron density? First, we must determine what is different in the resonance structures. Two differences commonly seen are the **position of a multiple bond** and the **site of a charge**. The anion $(\text{HCONH})^-$ illustrates two conventions for drawing resonance hybrids.




Common symbols and conventions used in organic chemistry are listed on the inside back cover.


- **Double bond position.** Structure **A** has a C–O double bond, whereas structure **B** has a C–N double bond. A dashed line in the hybrid indicates partial double bond character between these atoms.
- **Location of the charge.** A negative charge resides on different atoms in **A** and **B**. The symbol δ^- (for a partial negative charge) indicates that the charge is delocalized on the N and O atoms in the hybrid.

This discussion of resonance is meant to serve as an introduction only. You will learn many more facets of resonance theory in later chapters. In Chapter 2, for example, the enormous effect of resonance on acidity is discussed.

Problem 1.16 Label the resonance structures in each pair as major, minor, or equal contributors to the hybrid. Then draw the hybrid.



Problem 1.17 Draw a second resonance structure for nitrous acid. Label each resonance structure as a major, minor, or equal contributor to the hybrid. Then draw the resonance hybrid.



1.6 Determining Molecular Shape


We can now use Lewis structures to determine the shape around a particular atom in a molecule. Consider the H_2O molecule. The Lewis structure tells us only which atoms are connected to each other, but it implies nothing about the geometry. What does the overall molecule look like? Is H_2O a bent or linear molecule? Two variables define a molecule's structure: **bond length** and **bond angle**.

1.6A Bond Length


Although the SI unit for bond length is the picometer (pm), the angstrom (\AA) is still widely used in the chemical literature; $1 \text{ \AA} = 10^{-10} \text{ m}$. As a result, $1 \text{ pm} = 10^{-2} \text{ \AA}$, and $95.8 \text{ pm} = 0.958 \text{ \AA}$.

Bond length is the average distance between the centers of two bonded nuclei. Bond lengths are typically reported in picometers (pm), where $1 \text{ pm} = 10^{-12} \text{ m}$. For example, the O–H bond length in H_2O is 95.8 pm. Average bond lengths for common bonds are listed in Table 1.2.

- Bond length *decreases* across a row of the periodic table as the size of the atom *decreases*.



- Bond length *increases* down a column of the periodic table as the size of an atom *increases*.



Increasing bond length

Learn these general trends. Often knowing such trends is more useful than learning a set of exact numbers, because we are usually interested in comparisons rather than absolute values.

Table 1.2 Average Bond Lengths

Bond	Length (pm)	Bond	Length (pm)	Bond	Length (pm)
H–H	74	H–F	92	C–F	133
C–H	109	H–Cl	127	C–Cl	177
N–H	101	H–Br	141	C–Br	194
O–H	96	H–I	161	C–I	213

1.6B Bond Angle

Bond angle determines the shape around any atom bonded to two other atoms. To determine the bond angle and shape around a given atom, we must first determine how many groups surround the atom. **A group is either an atom or a lone pair of electrons.** Then we use the **valence shell electron pair repulsion (VSEPR) theory** to determine the shape. VSEPR is based on the fact that electron pairs repel each other; thus:

- The most stable arrangement keeps these groups as far away from each other as possible.

A second-row element has only three possible arrangements, defined by the number of groups surrounding it.


To determine geometry:
[1] Draw a valid Lewis structure; [2] count groups around a given atom.

Number of groups	Geometry	Bond angle
• two groups	linear	180°
• three groups	trigonal planar	120°
• four groups	tetrahedral	109.5°

Let's examine several molecules to illustrate this phenomenon. In each example, we first need a valid Lewis structure, and then we merely count groups around a given atom to predict its geometry.

Two Groups Around an Atom

Any atom surrounded by only two groups is linear and has a bond angle of 180°. Two examples illustrating this geometry are **BeH₂** (beryllium hydride) and **HC≡CH** (acetylene). We consider each carbon atom in acetylene *separately*. Because each C is surrounded by two atoms and no lone pairs, each H–C–C bond angle in acetylene is 180°, and therefore all four atoms are linear.



Acetylene illustrates another important feature: **ignore multiple bonds in predicting geometry.** **Count only atoms and lone pairs.**

We will begin to represent molecules with models having balls for atoms and sticks for bonds, as in the ball-and-stick model of acetylene just shown. These representations are analogous to a set of molecular models. Balls are color-coded using accepted conventions: carbon (black), hydrogen (white or gray), oxygen (red), and so forth, as shown.





Three Groups Around an Atom

Any atom surrounded by three groups is trigonal planar and has bond angles of 120°. Two examples illustrating this geometry are **BF₃** (boron trifluoride) and **CH₂=CH₂** (ethylene). *Each* carbon atom of ethylene is surrounded by three atoms and no lone pairs, making *each* H–C–C bond angle 120°.

Most students in organic chemistry find that building models helps them visualize the shape of molecules. Invest in a set of models *now*.


Two trigonal planar molecules




Four Groups Around an Atom

Any atom surrounded by four groups is tetrahedral and has bond angles of approximately 109.5° . For example, the simple organic compound methane, CH_4 , has a central carbon atom with bonds to four hydrogen atoms, each pointing to a corner of a tetrahedron. This arrangement keeps four groups farther apart than a square planar arrangement in which all bond angles would be only 90° .

Tetrahedral arrangement




Square planar arrangement



How can we represent the three-dimensional geometry of a tetrahedron on a two-dimensional piece of paper? Place two of the bonds in the plane of the paper, one bond in front and one bond behind, using the following conventions:

- A *solid line* is used for a bond *in* the plane.
- A *wedge* is used for a bond in *front* of the plane.
- A *dashed line* is used for a bond *behind* the plane.




This is just one way to draw a tetrahedron for CH_4 . We can turn the molecule in many different ways, generating many equivalent representations. For example, all of the following are acceptable drawings for CH_4 .



Each drawing has two solid lines, one wedge, and one dashed line.


Finally, wedges and dashes are used for groups that are really aligned one behind another. It does not matter in the following two drawings whether the wedge or dash is skewed to the left or right, because the two H atoms are really aligned as shown in the three-dimensional model.




All carbons in stable molecules are **tetravalent**, but the geometry varies with the number of groups around the particular carbon.

Problem 1.18 Draw two different three-dimensional representations for CH_2Cl_2 (dichloromethane) using wedges and dashes.

Ammonia (NH_3) and water (H_2O) both have atoms surrounded by four groups, some of which are lone pairs. In NH_3 , the three H atoms and one lone pair around N point to the corners of a tetrahedron. The H–N–H bond angle of 107° is close to the theoretical tetrahedral bond angle of 109.5° . This molecular shape is referred to as a **trigonal pyramid**, because one of the groups around the N is a nonbonded electron pair, not another atom.




In H_2O , the two H atoms and two lone pairs around O point to the corners of a tetrahedron. The H–O–H bond angle of 105° is close to the theoretical tetrahedral bond angle of 109.5° . Water has a **bent** molecular shape, because two of the groups around oxygen are lone pairs of electrons.



In both NH_3 and H_2O the bond angle is somewhat smaller than the theoretical tetrahedral bond angle because of repulsion of the lone pairs of electrons. The bonded atoms are compressed into a smaller space with a smaller bond angle.

Predicting geometry based on counting groups is summarized in Figure 1.4.

Sample Problem 1.6 Determine the geometry around the indicated atom in each species.



Solution






Figure 1.4
Summary: Determining geometry based on the number of groups

Number of groups around an atom	Geometry	Bond angle	Examples
2	linear	180°	BeH_2 , $\text{HC}\equiv\text{CH}$
3	trigonal planar	120°	BF_3 , $\text{CH}_2=\text{CH}_2$
4	tetrahedral	109.5°	CH_4 , NH_3 , H_2O

Problem 1.19 Determine the geometry around all second-row elements in each compound.




Problem 1.20 Predict the indicated bond angles in each compound.



Problem 1.21

Using the principles of VSEPR theory, you can predict the geometry around any atom in any molecule, no matter how complex. Enanthotoxin is a poisonous compound isolated from a common variety of hemlock grown in England. Predict the geometry around the indicated atoms in enanthotoxin.



1.7 Drawing Organic Structures


Drawing organic molecules presents a special challenge. Because they often contain many atoms, we need shorthand methods to simplify their structures. The two main types of shorthand representations used for organic compounds are **condensed structures** and **skeletal structures**.

1.7A Condensed Structures

Condensed structures are most often used for compounds having a chain of atoms bonded together, rather than a ring. The following conventions are used:

- All of the atoms are drawn in, but the two-electron bond lines are generally omitted.
- Atoms are usually drawn next to the atoms to which they are bonded.
- Parentheses are used around similar groups bonded to the same atom.
- Lone pairs are omitted.


To interpret a condensed formula, it is usually best to start at the **left side** of the molecule and remember that the **carbon atoms must be tetravalent**. A carbon bonded to three H atoms becomes CH_3 ; a carbon bonded to two H atoms becomes CH_2 , and so forth. Two examples of condensed formulas for compounds having only carbon and hydrogen are given below.



Recall from the prologue:
A **heteroatom** is any atom that is not C or H.

Other examples of condensed structures with heteroatoms and carbon–carbon multiple bonds are given in Figure 1.5. You must learn how to convert a Lewis structure to a condensed structure, and vice versa.

Figure 1.5
Examples of condensed structures



Translating some condensed formulas is not obvious, and it will come only with practice. This is especially true for compounds containing a carbon–oxygen double bond. Some noteworthy examples in this category are given in Figure 1.6. While carbon–carbon double bonds are generally drawn in condensed structures, carbon–oxygen double bonds are usually omitted.

Sample Problem 1.7


Convert each condensed formula to a Lewis structure.

- a. $(\text{CH}_3)_2\text{CHOCH}_2\text{CH}_2\text{CH}_2\text{OH}$ b. $\text{CH}_3(\text{CH}_2)_2\text{CO}_2\text{C}(\text{CH}_3)_3$

Solution

Start at the left and proceed to the right, making sure that each carbon has four bonds. Give any heteroatom enough lone pairs to have an octet.

- a. $(\text{CH}_3)_2\text{CHOCH}_2\text{CH}_2\text{CH}_2\text{OH}$



- b. $\text{CH}_3(\text{CH}_2)_2\text{CO}_2\text{C}(\text{CH}_3)_3$




Figure 1.6
Condensed structures containing a C–O double bond



- Take special note of condensed structures with oxygen atoms. In these examples, the only way for all atoms to have an octet is by having a carbon–oxygen double bond.

Problem 1.22 Convert each condensed formula to a Lewis structure.

- a. $\text{CH}_3(\text{CH}_2)_4\text{CH}(\text{CH}_3)_2$ c. $(\text{CH}_3)_3\text{CCH}(\text{OH})\text{CH}_2\text{CH}_3$
 b. $(\text{CH}_3)_2\text{CHCH}(\text{CH}_2\text{CH}_3)_2$ d. $(\text{CH}_3)_2\text{CHCHO}$

Problem 1.23 Simplify each condensed structure by using parentheses around similar groups.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$ c. $\text{HOCH}_2-\overset{\text{CH}_2\text{OH}}{\underset{\text{H}}{\text{C}}}-\text{CH}_2\text{CH}_2\text{CH}_2-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_2-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_3$
 b. $\text{CH}_3\text{CH}_2\text{CH}_2-\overset{\text{CH}_2\text{CH}_3}{\underset{\text{H}}{\text{C}}}-\text{CH}_2\text{CH}_3$

Problem 1.24 During periods of strenuous exercise, the buildup of lactic acid $[\text{CH}_3\text{CH}(\text{OH})\text{CO}_2\text{H}]$ causes the aching feeling in sore muscles. Convert this condensed structure to a Lewis structure of lactic acid.

1.7B Skeletal Structures

Skeletal structures are used for organic compounds containing both rings and chains of atoms. Three important rules are used to draw them:

- Assume there is a carbon atom at the junction of any two lines or at the end of any line.
- Assume there are enough hydrogens around each carbon to make it tetravalent.
- Draw in all heteroatoms and the hydrogens directly bonded to them.

Carbon chains are drawn in a zigzag fashion, and rings are drawn as polygons, as shown for hexane and cyclohexane.






Figure 1.7 shows other examples of skeletal structures for a variety of Lewis structures, and Sample Problem 1.8 illustrates how to interpret the skeletal structure for a cyclic compound.

Figure 1.7
Examples of skeletal structures




Sample Problem 1.8



Draw a complete structure for vanillin showing all H atoms and lone pairs. Vanillin is the principal component of the extract of the vanilla bean.

**Solution**


Skeletal structures have a C atom at the junction of any two lines and at the end of any line. Each C must have enough H's to make it tetravalent. In structures that contain a -CHO group, the C atom is doubly bonded to the O atom and singly bonded to H. Each O atom needs two lone pairs to have a complete octet.

**Problem 1.25**




How many hydrogen atoms are present around each indicated carbon atom in the following molecules? Both compounds are active ingredients in some common sunscreens.

**Problem 1.26**

Convert each skeletal structure to a Lewis structure.




When heteroatoms are bonded to a carbon skeleton, the heteroatom is joined *directly* to the carbon to which it is bonded, with no H atoms in between. Thus, an OH group is drawn as OH or HO depending on where the OH is located. In contrast, when carbon appendages are bonded to a carbon skeleton, the H atoms will be drawn to the *right* of the carbon to which they are bonded regardless of the location.



1.7C Skeletal Structures with Charged Carbon Atoms


Take care in interpreting skeletal structures for positively and negatively charged carbon atoms, because *both* the hydrogen atoms *and* the lone pairs are omitted. Keep in mind the following:

- A charge on a carbon atom takes the place of **one** hydrogen atom.
- The charge determines the number of lone pairs. Negatively charged carbon atoms have **one** lone pair and positively charged carbon atoms have none.




Skeletal structures often leave out lone pairs on heteroatoms, but *don't forget about them.*


Use the formal charge on an atom to determine the number of lone pairs. For example, a neutral O atom with two bonds needs two additional lone pairs, and a positively charged O atom with three bonds needs only one lone pair.



Problem 1.27 Draw in all hydrogens and lone pairs on the charged carbons in each ion.



Problem 1.28 Draw a skeletal structure for the molecules in parts (a) and (b), and a condensed structure for the molecules in parts (c) and (d).




1.8 Hybridization

What orbitals do the first- and second-row atoms use to form bonds? Let's begin with hydrogen and then examine the orbitals used for bonding by atoms in the second row.

1.8A Hydrogen

Recall from Section 1.2 that two hydrogen atoms share each of their electrons to form H_2 . Thus, the $1s$ orbital on one H overlaps with the $1s$ orbital on the other H to form a bond that concentrates electron density between the two nuclei. This type of bond, called a σ (sigma) **bond**, is cylindrically symmetrical because the electrons forming the bond are distributed symmetrically about an imaginary line connecting the two nuclei.




- A σ bond concentrates electron density on the axis that joins two nuclei. All single bonds are σ bonds.

1.8B Bonding in Methane


To account for the bonding patterns observed in more complex molecules, we must take a closer look at the $2s$ and $2p$ orbitals of atoms of the second row. Let's illustrate this with methane, CH_4 .

Carbon has two core electrons, plus **four valence electrons**. To fill atomic orbitals in the most stable arrangement, electrons are placed in the orbitals of lowest energy. For carbon, this places two in the $2s$ orbital and one each in two $2p$ orbitals.




- This lowest energy arrangement of electrons for an atom is called its **ground state**.

In this description, **carbon should form only two bonds** because it has only two unpaired valence electrons, and CH_2 should be a stable molecule. In reality, however, CH_2 is a highly reactive species that cannot be isolated under typical laboratory conditions. In CH_2 , carbon would not have an octet of electrons.



There is a second possibility. Promotion of an electron from a $2s$ to a vacant $2p$ orbital would form **four** unpaired electrons for bonding. This process requires energy because it moves an electron to a higher energy orbital. This higher energy electron configuration is called an electronically **excited state**.




This description is still not adequate. Carbon would form two different types of bonds: three with $2p$ orbitals and one with a $2s$ orbital. **But experimental evidence points to carbon forming four identical bonds in methane.**

To solve this dilemma, chemists have proposed that atoms like carbon do not use pure s and pure p orbitals in forming bonds. Instead, atoms use a set of new orbitals called **hybrid orbitals**. The mathematical process by which these orbitals are formed is called **hybridization**.

- **Hybridization** is the combination of two or more atomic orbitals to form the same number of hybrid orbitals, each having the same shape and energy.

Hybridization of one $2s$ orbital and three $2p$ orbitals for carbon forms four hybrid orbitals, each with one electron. These new hybrid orbitals are intermediate in energy between the $2s$ and $2p$ orbitals.

Forming four sp^3 hybrid orbitals for carbon




- These hybrid orbitals are called sp^3 hybrids because they are formed from *one* s orbital and *three* p orbitals.



What do these new hybrid orbitals look like? Mixing a spherical $2s$ orbital and three dumbbell-shaped $2p$ orbitals together produces four orbitals having one large lobe and one small lobe, oriented toward the corners of a tetrahedron. Each large lobe concentrates electron density in the bonding direction between two nuclei. This makes bonds formed from hybrid orbitals **stronger** than bonds formed from pure p orbitals.

Shape and orientation of sp^3 hybrid orbitals




The four hybrid orbitals form four equivalent bonds. We can now explain the observed bonding in CH_4 .

- Each bond in CH_4 is formed by overlap of an sp^3 hybrid orbital of carbon with a $1s$ orbital of hydrogen. These four bonds point to the corners of a tetrahedron.

All four C–H bonds in methane are **σ bonds**, because the electron density is concentrated on the axis joining C and H. An orbital picture of the bonding in CH_4 is given in Figure 1.8.

Figure 1.8
Bonding in CH_4 using sp^3 hybrid orbitals



Problem 1.29 What orbitals are used to form each of the C–C and C–H bonds in $\text{CH}_3\text{CH}_2\text{CH}_3$ (propane)? How many σ bonds are present in this molecule?

1.8C Other Hybridization Patterns— sp and sp^2 Hybrid Orbitals

Forming sp^3 hybrid orbitals is just one way that $2s$ and $2p$ orbitals can hybridize. In fact, three common modes of hybridization are seen in organic molecules. The number of orbitals is always conserved in hybridization; that is, a **given number of atomic orbitals hybridize to form an equivalent number of hybrid orbitals**.

- One $2s$ orbital and *three* $2p$ orbitals form *four* sp^3 hybrid orbitals.
- One $2s$ orbital and *two* $2p$ orbitals form *three* sp^2 hybrid orbitals.
- One $2s$ orbital and *one* $2p$ orbital form *two* sp hybrid orbitals.

We have already seen pictorially how four sp^3 hybrid orbitals are formed from one $2s$ and three $2p$ orbitals. Figures 1.9 and 1.10 illustrate the same process for sp and sp^2 hybrids. Each sp and sp^2 hybrid orbital has one large and one small lobe, much like an sp^3 hybrid orbital. Note, however, that both sp^2 and sp hybridization **leave one and two $2p$ orbitals unhybridized**, respectively, on each atom.

The **superscripts** for hybrid orbitals correspond to the **number of atomic orbitals** used to form them. The number “1” is understood.

For example: $sp^3 = s^1p^3$


one $2s$ + three $2p$ orbitals used to make each hybrid orbital

To determine the hybridization of an atom in a molecule, we count groups around the atom, just as we did in determining geometry. **The number of groups (atoms and nonbonded electron pairs) corresponds to the number of atomic orbitals that must be hybridized to form the hybrid orbitals.**

number of groups around an atom	number of orbitals used	type of hybrid orbital
2	2	two sp hybrid orbitals
3	3	three sp^2 hybrid orbitals
4	4	four sp^3 hybrid orbitals


Let's illustrate this phenomenon with BeH_2 , BF_3 , NH_3 , and H_2O . We already determined the geometry in these molecules by counting groups in Section 1.6.

Figure 1.9
Forming two sp hybrid orbitals



- Forming **two sp hybrid orbitals** uses **one $2s$ and one $2p$ orbital**, leaving **two $2p$ orbitals unhybridized**.


Figure 1.10
Forming three sp^2 hybrid orbitals



- Forming **three sp^2 hybrid orbitals** uses **one $2s$ and two $2p$ orbitals**, leaving **one $2p$ orbital unhybridized**.

In BeH_2 , the **Be atom is sp hybridized** because it is surrounded by two groups (two H atoms). Each Be–H bond is formed by overlap of an sp hybrid orbital from Be and a $1s$ orbital from H. The sp hybrid orbitals are oriented 180° away from each other.

In BF_3 , the **B atom is sp^2 hybridized** because it is surrounded by three groups (three F atoms). Each B–F bond is formed by overlap of an sp^2 hybrid orbital from B and a $2p$ orbital from F. The sp^2 hybrid orbitals all lie in a plane, and are oriented 120° apart. The B atom also has a vacant unhybridized $2p$ orbital. This orbital is located **above and below the plane** of the BF_3 molecule.




The N atom in NH_3 and the O atom in H_2O are both surrounded by four groups, making them sp^3 hybridized. Each N–H and O–H bond in these molecules is formed by overlap of an sp^3 hybrid orbital with a $1s$ orbital from H. The lone pairs of electrons on N and O also occupy sp^3 hybrid orbitals, as shown in Figure 1.11.

Sample Problem 1.9

What orbitals are used to form each bond in methanol, CH_3OH ?

Solution

To solve this problem, draw a valid Lewis structure and count groups around each atom. Then, use the rule to determine hybridization: two groups = sp , three groups = sp^2 , and four groups = sp^3 .




- All C–H bonds are formed from $\text{C}_{sp^3} - \text{H}_{1s}$.
- The C–O bond is formed from $\text{C}_{sp^3} - \text{O}_{sp^3}$.
- The O–H bond is formed from $\text{O}_{sp^3} - \text{H}_{1s}$.

Problem 1.30

What orbitals are used to form each bond in the following molecules?


- a. CH_3BeH b. $(\text{CH}_3)_3\text{B}$ c. CH_3OCH_3

Figure 1.11
Hybrid orbitals of NH_3 and H_2O



1.9 Ethane, Ethylene, and Acetylene

Let's now use the principles of hybridization to determine the type of bonds in **ethane**, **ethylene**, and **acetylene**.



ethane



ethylene




acetylene

1.9A Ethane— CH_3CH_3


According to the Lewis structure for **ethane**, CH_3CH_3 , each carbon atom is singly bonded to four other atoms. As a result:

- Each carbon is tetrahedral.
- Each carbon is sp^3 hybridized.




Ethane is a constituent of natural gas.

All of the bonds in ethane are σ bonds. The C–H bonds are formed from the overlap of one of the three sp^3 hybrid orbitals on each carbon atom with the $1s$ orbital on hydrogen. The C–C bond is formed from the overlap of an sp^3 hybrid orbital on each carbon atom.



Finally, a model of ethane shows that **rotation can occur around the central C–C σ bond**. Note how the relative position of the H atoms on the adjacent CH_3 groups changes from one representation to another. This process is discussed in greater detail in Chapter 4.




1.9B Ethylene— C_2H_4

Based on the Lewis structure of **ethylene**, $\text{CH}_2=\text{CH}_2$, each carbon atom is singly bonded to two H atoms and doubly bonded to the other C atom, so each C is surrounded by three groups. As a result:




Ethylene is an important starting material in the preparation of the plastic polyethylene.

- Each carbon is trigonal planar (Section 1.6B).
- Each carbon is sp^2 hybridized.




What orbitals are used to form the two bonds of the C–C double bond? Recall from Section 1.8 that sp^2 hybrid orbitals are formed from **one 2s and two 2p orbitals**, leaving one **2p orbital unhybridized**. Because carbon has four valence electrons, **each of these orbitals has one electron** that can be used to form a bond.

Forming an sp^2 hybridized carbon atom




Each C–H bond results from the end-on overlap of an sp^2 hybrid orbital on carbon and the 1s orbital on hydrogen. Similarly, one of the C–C bonds results from the end-on overlap of an sp^2 hybrid orbital on each carbon atom. Each of these bonds is a σ bond.



The second C–C bond results from the side-by-side overlap of the 2p orbitals on each carbon. Side-by-side overlap creates an area of electron density above and below the plane containing the sp^2 hybrid orbitals (that is, the plane containing the six atoms in the σ bonding system).

Overlap of the 2p orbitals forms the second C–C bond.



In this second bond, the electron density is *not* concentrated on the axis joining the two nuclei. This new type of bond is called a π bond. Because the electron density in a π bond is farther from the two nuclei, π bonds are usually weaker and therefore more easily broken than σ bonds.


Thus, a carbon–carbon double bond has two components:

- a σ bond, formed by end-on overlap of two sp^2 hybrid orbitals;
- a π bond, formed by side-by-side overlap of two 2p orbitals.

Figure 1.12 summarizes the bonding observed in ethylene.


Figure 1.12

Summary: The σ and π bonds in ethylene



Unlike the C–C single bond in ethane, rotation about the C–C double bond in ethylene is **restricted**. It can occur only if the π bond first breaks and then re-forms, a process that requires considerable energy.


All double bonds are composed of one σ and one π bond.



1.9C Acetylene— C_2H_2


Based on the Lewis structure of **acetylene**, $HC\equiv CH$, each carbon atom is singly bonded to one hydrogen atom and triply bonded to the other carbon atom, so each carbon atom is surrounded by two groups. As a result:

- Each carbon is linear (Section 1.6B).
- Each carbon is sp hybridized.




Because acetylene produces a very hot flame on burning, it is often used in welding torches. The fire is very bright, too, so it was once used in the lamps worn by spelunkers—people who study and explore caves.

What orbitals are used to form the bonds of the C–C triple bond? Recall from Section 1.8 that sp hybrid orbitals are formed from **one $2s$ and one $2p$ orbital**, leaving **two $2p$ orbitals unhybridized**. Because carbon has four valence electrons, **each of these orbitals has one electron** that can be used to form a bond.




Each C–H bond results from the end-on overlap of an sp hybrid orbital on carbon and the $1s$ orbital on hydrogen. Similarly, one of the C–C bonds results from the end-on overlap of an sp hybrid orbital on each carbon atom. Each of these bonds is a σ bond.



- Each C has two sp hybrid orbitals.
- The C–H bonds and C–C bond are σ bonds.

Each carbon atom also has two **unhybridized $2p$ orbitals** that are perpendicular to each other and to the sp hybrid orbitals. Side-by-side overlap between the two $2p$ orbitals on one carbon with the two $2p$ orbitals on the other carbon creates the second and third bonds of the C–C triple bond. The electron density from one of these two bonds is above and below the axis joining the two nuclei, and the electron density from the second of these two bonds is in front of and behind the axis, so both of these bonds are π bonds.



The side-by-side overlap of two p orbitals always forms a π bond.

All triple bonds are composed of one σ and two π bonds.


Thus, a carbon–carbon triple bond has three components:

- a σ bond, formed by end-on overlap of two sp hybrid orbitals;
- two π bonds, formed by side-by-side overlap of two sets of $2p$ orbitals.

Figure 1.13 summarizes the bonding observed in acetylene. Figure 1.14 summarizes the three possible types of bonding in carbon compounds.

Sample Problem 1.10

Answer each question for the molecule acetone, drawn below:



- Determine the hybridization of the indicated atoms.
- What orbitals are used to form the C–O double bond?
- In what type of orbital does each lone pair reside?

Solution





Figure 1.13


Summary: The σ and π bonds in acetylene

The three σ bonds are labeled.



Overlap of the two sp hybrid orbitals forms the C–C σ bond.




Two π bonds extend out from the axis of the linear molecule.



Overlap of two sets of two $2p$ orbitals forms two C–C π bonds.


Figure 1.14

A summary of covalent bonding seen in carbon compounds


Number of groups bonded to C	Hybridization	Bond angle	Example	Observed bonding
4	sp^3	109.5°	CH_3CH_3 ethane	 <p>one σ bond</p> <p>$\text{C}_{sp^3}-\text{C}_{sp^3}$</p>
3	sp^2	120°	$\text{CH}_2=\text{CH}_2$ ethylene	 <p>one σ bond + one π bond</p> <p>$\text{C}_{sp^2}-\text{C}_{sp^2}$ $\text{C}_{2p}-\text{C}_{2p}$</p>
2	sp	180°	$\text{HC}\equiv\text{CH}$ acetylene	 <p>one σ bond + two π bonds</p> <p>$\text{C}_{sp}-\text{C}_{sp}$ $\text{C}_{2p}-\text{C}_{2p}$ $\text{C}_{2p}-\text{C}_{2p}$</p>

- b. • The σ bond is formed from the end-on overlap of $\text{C}_{sp^2}-\text{O}_{sp^2}$.
 • The π bond is formed from the side-by-side overlap of $\text{C}_{2p}-\text{O}_{2p}$.
- c. The O atom has three sp^2 hybrid orbitals.
 • One is used for the σ bond of the double bond.
 • The remaining two sp^2 hybrids are occupied by the lone pairs.

Problem 1.31 Determine the hybridization around the indicated atoms in the following molecules:



Problem 1.32 Classify each bond in the following molecules as σ or π :




1.10 Bond Length and Bond Strength

Let's now examine the relative bond length and bond strength of the C–C and C–H bonds in ethane, ethylene, and acetylene.

1.10A A Comparison of Carbon–Carbon Bonds

While the SI unit of energy is the **joule** (J), organic chemists often report energy values in **calories** (cal). For this reason, energy values in the tables in this text are reported in joules, followed by the number of calories in parentheses. 1 cal = 4.18 J



- As the number of electrons between two nuclei *increases*, bonds become shorter and stronger.

Note the inverse relationship between bond length and bond strength. The shorter the bond, the closer the electron density is kept to the nucleus, and the harder the bond is to break.

Shorter bonds are stronger bonds.

- Thus, triple bonds are shorter and stronger than double bonds, which are shorter and stronger than single bonds.

Values for bond lengths and bond strengths for CH_3CH_3 , $\text{CH}_2=\text{CH}_2$ and $\text{HC}\equiv\text{CH}$ are listed in Table 1.3. Be careful not to confuse two related but different principles regarding multiple bonds such as C–C double bonds. **Double bonds, consisting of both a σ and a π bond, are strong.** The π component of the double bond, however, is usually much *weaker* than the σ component. This is a particularly important consideration when studying alkenes in Chapter 10.

Table 1.3 Bond Lengths and Bond Strengths for Ethane, Ethylene, and Acetylene

Compound	C–C bond length (pm)	Bond strength kJ/mol (kcal/mol)
$\text{CH}_3\text{—CH}_3$	153	368 (88)
$\text{CH}_2=\text{CH}_2$	134	635 (152)
$\text{HC}\equiv\text{CH}$	121	837 (200)


↑ Increasing bond length ↓ Increasing bond strength

Compound	C–H bond length (pm)	Bond strength kJ/mol (kcal/mol)
$\text{CH}_3\text{CH}_2\text{—H}$	111	410 (98)
$\text{CH}_2=\text{C—H}$ H	110	435 (104)
$\text{HC}\equiv\text{C—H}$	109	523 (125)

↑ Increasing bond length ↓ Increasing bond strength

1.10B A Comparison of Carbon–Hydrogen Bonds

The length and strength of a C–H bond vary slightly depending on the hybridization of the carbon atom.



To understand why this is so, we must look at the atomic orbitals used to form each type of hybrid orbital. A single $2s$ orbital is always used, but the number of $2p$ orbitals varies with the type of hybridization. A quantity called **percent s -character** indicates the fraction of a hybrid orbital due to the $2s$ orbital used to form it.

$$sp \text{ hybrid} \quad \frac{\text{one } 2s \text{ orbital}}{\text{two hybrid orbitals}} = 50\% \text{ } s\text{-character}$$

$$sp^2 \text{ hybrid} \quad \frac{\text{one } 2s \text{ orbital}}{\text{three hybrid orbitals}} = 33\% \text{ } s\text{-character}$$


$$sp^3 \text{ hybrid} \quad \frac{\text{one } 2s \text{ orbital}}{\text{four hybrid orbitals}} = 25\% \text{ } s\text{-character}$$

Why should the percent s -character of a hybrid orbital affect the length of a C–H bond? A $2s$ orbital keeps electron density closer to a nucleus compared to a $2p$ orbital. As the **percent**


s-character increases, a hybrid orbital holds its electrons closer to the nucleus, and the **bond becomes shorter and stronger**.

Increased percent s-character \rightarrow Increased bond strength \rightarrow Decreased bond length

Problem 1.33 Rank the indicated bonds in each compound in order of (a) increasing bond strength and (b) increasing bond length.



Problem 1.34 Which of the indicated bonds in each pair of compounds is shorter?



1.11 Electronegativity and Bond Polarity

Electronegativity is a measure of an atom's attraction for electrons in a bond. Thus, electronegativity indicates how much a particular atom “wants” electrons. The following trends in electronegativity are observed in the periodic table:

- Electronegativity *increases* across a row of the periodic table as the nuclear charge increases (excluding the noble gases).
- Electronegativity *decreases* down a column of the periodic table as the atomic radius increases, pushing the valence electrons farther from the nucleus.

As a result, the *most* electronegative elements are located at the **upper right-hand corner** of the periodic table, and the *least* electronegative elements in the **lower left-hand corner**. A scale has been established to represent electronegativity values arbitrarily, from 0 to 4, as shown in Figure 1.15.

Electronegativity values are relative, so they can be used for comparison purposes only. When comparing two different elements, one is **more electronegative** than the other if it attracts elec-

Figure 1.15

Electronegativity values for some common elements

Increasing electronegativity \rightarrow

1A								
H								
2.2	2A		3A	4A	5A	6A	7A	
Li	Be	//	B	C	N	O	F	
1.0	1.6	//	1.8	2.5	3.0	3.4	4.0	
Na	Mg		Si	P	S	Cl		
0.9	1.3		1.9	2.2	2.6	3.2		
K							Br	
0.8							3.0	
							I	
							2.7	


Increasing electronegativity \uparrow

tron density toward itself. One is less electronegative—**more electropositive**—if it gives up electron density to the other element.


Problem 1.35 Rank the following atoms in order of increasing electronegativity. Label the most electronegative and most electropositive atom in each group.

- a. Se, O, S b. P, Na, Cl c. Cl, S, F d. O, P, N

Electronegativity values are used as a guideline to indicate whether the electrons in a bond are **equally shared** or **unequally shared** between two atoms. For example, whenever two identical atoms are bonded together, each atom attracts the electrons in the bond to the same extent. The electrons are equally shared, and the **bond is nonpolar**. Thus, a **carbon–carbon bond is nonpolar**. The same is true whenever two different atoms having similar electronegativities are bonded together. **C–H bonds are considered to be nonpolar**, because the electronegativity difference between C (2.5) and H (2.2) is small.



Bonding between atoms of different electronegativity values results in the **unequal sharing** of electrons. For example, in a C–O bond, the electrons are pulled away from C (2.5) toward O (3.4), the element of higher electronegativity. **The bond is polar**, or **polar covalent**. The bond is said to have a **dipole**; that is, a **separation of charge**.



The direction of polarity in a bond is often indicated by an arrow, with the head of the arrow pointing toward the more electronegative element. The tail of the arrow, with a perpendicular line drawn through it, is positioned at the less electronegative element. Alternatively, the symbols δ^+ and δ^- indicate this unequal sharing of electron density.

- δ^+ means an atom is electron deficient (has a partial positive charge).
- δ^- means the atom is electron rich (has a partial negative charge).


Problem 1.36 Show the direction of the dipole in each bond. Label the atoms with δ^+ and δ^- .

- a. H–F b. $\begin{array}{c} | \\ -\text{B}-\text{C}- \\ | \end{array}$ c. $\begin{array}{c} | \\ -\text{C}-\text{Li} \\ | \end{array}$ d. $\begin{array}{c} | \\ -\text{C}-\text{Cl} \\ | \end{array}$

Students often wonder how large an electronegativity difference must be to consider a bond polar. That's hard to say. We will set an arbitrary value for this difference and use it as an *approximation*. **Usually, a polar bond will be one in which the electronegativity difference between two atoms is ≥ 0.5 units.**

The distribution of electron density in a molecule can be shown using an **electrostatic potential map**. These maps are color coded to illustrate areas of high and low electron density. Electron-rich

Figure 1.16
Electrostatic potential
plot of CH_3Cl



regions are indicated in red, and electron-deficient sites are indicated in blue. Regions of intermediate electron density are shown in orange, yellow, and green.

For example, an electrostatic potential map of CH_3Cl clearly indicates the polar nature of the C–Cl bond (Figure 1.16). The more electronegative Cl atom pulls electron density toward it, making it electron rich. This is indicated by the red around the Cl in the plot. The carbon is electron deficient, and this is shown with blue.

When comparing two maps, the comparison is useful only if they are plotted *using the same scale* of color gradation. For this reason, whenever we compare two plots in this text, they will be drawn side by side using the same scale. It will be difficult to compare two plots in different parts of the book, because the scale may be different. Despite this limitation, an electrostatic potential plot is a useful tool for visually evaluating the distribution of electron density in a molecule, and with care, comparing the electron density in two different molecules.

1.12 Polarity of Molecules


A **polar molecule** has either one polar bond, or two or more bond dipoles that reinforce. A **nonpolar molecule** has either no polar bonds, or two or more bond dipoles that cancel.

Whenever C or H is bonded to N, O, and all halogens, the bond is **polar**. Thus, the C–I bond is considered polar even though the electronegativity difference between C and I is small. Remember, electronegativity is just an approximation.

Thus far, we have been concerned with the polarity of one bond. To determine whether a molecule has a net dipole, use the following two-step procedure:


- [1] Use electronegativity differences to **identify all of the polar bonds and the directions of the bond dipoles**.
- [2] **Determine the geometry** around individual atoms by counting groups, and decide if individual dipoles **cancel** or **reinforce each other in space**.

The two molecules H_2O and CO_2 illustrate different outcomes of this process. In H_2O , each O–H bond is polar because the electronegativity difference between O (3.4) and H (2.2) is large. Since H_2O is a **bent** molecule, the two dipoles reinforce (both point *up*). Thus, H_2O has a **net dipole, making it a polar molecule**. CO_2 also has polar C–O bonds because the electronegativity difference between O (3.4) and C (2.5) is large. However, CO_2 is a **linear** molecule, so the two dipoles, which are equal and opposite in direction, **cancel**. Thus, CO_2 is a **nonpolar molecule with no net dipole**.



The net dipole bisects the H–O–H bond angle.
The two individual dipoles reinforce.

H_2O is a polar molecule.



no net dipole
The two dipoles cancel.

CO_2 is a nonpolar molecule.

Electrostatic potential plots for H_2O and CO_2 appear in Figure 1.17. Additional examples of polar and nonpolar molecules are given in Figure 1.18.

Problem 1.37

Indicate which of the following molecules is polar because it possesses a net dipole. Show the direction of the net dipole if one exists.




Figure 1.17

Electrostatic potential plots for H_2O and CO_2 

- The electron-rich (red) region is concentrated on the electronegative O atom. Both H atoms are electron deficient (blue-green).
- Both electronegative O atoms are electron rich (red) and the central C atom is electron deficient (blue).


Figure 1.18

More examples of polar and nonpolar molecules



1.13 L-Dopa—A Representative Organic Molecule

The principles learned in this chapter apply to all organic molecules regardless of size or complexity. For example, we now know a great deal about the structure of **L-dopa**, a drug used to treat Parkinson's disease described in the chapter opener.




For example, you should be able to do all of the following:

- Convert this skeletal structure of L-dopa to a Lewis structure.
- Determine the hybridization and geometry around every atom.
- Label all polar and nonpolar bonds.
- Compare bond length and bond strength for certain bonds.

Some of these concepts are illustrated in the Lewis structure for L-dopa. As we continue our study of organic chemistry you will see that these fundamental properties about a molecule determine its physical properties and its behavior in chemical reactions.

The C—O and O—H bonds are polar.



Lewis structure of L-dopa

Problem 1.38

Provide the following information about L-dopa. In all cases, label different sites than those identified in the structure shown.

- Label two polar and two nonpolar bonds.
- Label all sp^3 hybridized C atoms.
- Label all H atoms that bear a partial positive charge (δ^+).
- Draw another resonance structure.

KEY CONCEPTS


Structure and Bonding

Important Facts

- **The general rule of bonding:** Atoms “strive” to attain a complete outer shell of valence electrons (Section 1.2). H “wants” two electrons. Second-row elements “want” eight electrons.
- **Formal charge** is the difference between the number of valence electrons of an atom and the number of electrons it “owns” (Section 1.3C). See Sample Problem 1.4 for a stepwise example.
- **Curved arrow notation** shows the movement of an electron pair. The tail of the arrow always begins at an electron pair, either in a bond or a lone pair. The head points to where the electron pair “moves” (Section 1.5).
- **Electrostatic potential plots** are color-coded maps of electron density, indicating electron-rich and electron-deficient regions (Section 1.11).

The Importance of Lewis Structures (Sections 1.3, 1.4)

A properly drawn Lewis structure shows the number of bonds and lone pairs present around each atom in a molecule. In a valid Lewis structure, each H has two electrons, and each second-row element has no more than eight. This is the first step needed to determine many properties of a molecule.




Resonance (Section 1.5)

The basic principles:

- Resonance exists when a compound cannot be represented by a single Lewis structure.
- Resonance structures differ in the position of only nonbonded electrons and π bonds, not atoms.
- The resonance hybrid is the only accurate representation for a resonance-stabilized compound. A hybrid represents the compound better than any single resonance structure because electron density is delocalized.

The difference between resonance structures and isomers:

- Two **isomers** differ in the arrangement of *both* atoms and electrons.
- **Resonance structures** differ *only* in the arrangement of electrons.



Geometry and Hybridization


The number of groups around an atom determines both its geometry (Section 1.6) and hybridization (Section 1.8).

Number of groups	Geometry	Bond angle ($^\circ$)	Hybridization
2	linear	180	sp
3	trigonal planar	120	sp^2
4	tetrahedral	109.5	sp^3


- 1.45 Draw an acceptable Lewis structure from each condensed structure, such that all atoms have zero formal charge.
- diethyl ether, $(\text{CH}_3\text{CH}_2)_2\text{O}$, the first general anesthetic used in medical procedures
 - acrylonitrile, CH_2CHCN , starting material used to manufacture synthetic Orlon fibers
 - dihydroxyacetone, $(\text{HOCH}_2)_2\text{CO}$, an ingredient in sunless tanning products
 - acetic anhydride, $(\text{CH}_3\text{CO})_2\text{O}$, a reagent used to synthesize aspirin

Isomers and Resonance Structures


- 1.46 Draw all possible isomers for each molecular formula.
- $\text{C}_3\text{H}_7\text{Cl}$ (two isomers)
 - $\text{C}_2\text{H}_4\text{O}$ (three isomers)
 - $\text{C}_3\text{H}_9\text{N}$ (four isomers)
- 1.47 Draw Lewis structures for the nine isomers having molecular formula $\text{C}_3\text{H}_6\text{O}$, with all atoms having a zero formal charge.
- 1.48 With reference to compound **A** drawn below, label each compound as an isomer, a resonance structure, or neither.




- 1.49 With reference to species **B**, label each species as an isomer, a resonance structure, or neither.




- 1.50 How are the molecules or ions in each pair related? Classify them as resonance structures, isomers, or neither.




- 1.51 Add curved arrows to show how the first resonance structure can be converted into the second.



- 1.52 Follow the curved arrows to draw a second resonance structure for each species.




- 1.53 Draw a second resonance structure for each ion.




- 1.54 For each ion in Problem 1.53 draw the resonance hybrid.


- 1.55 Draw all reasonable resonance structures for each species.




1.56 Draw four additional resonance structures for the following cation. Then draw the resonance hybrid.



1.57 Rank the resonance structures in each group in order of increasing contribution to the resonance hybrid. Label the resonance structure that contributes the most as **3** and the resonance structure that contributes the least as **1**. Label the intermediate contributor as **2**.




1.58 Consider the compounds and ions with curved arrows drawn below. When the curved arrows give a second valid resonance structure, draw the resonance structure. When the curved arrows generate an invalid Lewis structure, explain why the structure is unacceptable.




Geometry

1.59 Predict all bond angles in each compound.




1.60 Predict the geometry around each indicated atom.




1.61 Draw the structure of halothane, CF_3CHClBr , in three dimensions, using solid lines, wedges, and dashes to illustrate the position of atoms. Halothane is a safe and widely used general anesthetic.

Drawing Organic Molecules




1.62 How many hydrogens are present around each carbon atom in the following molecules?





1.63 Draw in all the carbon and hydrogen atoms in each molecule.






1.64 Convert each molecule into a skeletal structure.







limonene
(oil of lemon)







1.65 Convert the following condensed formulas into Lewis structures.







1.66 Draw in all the hydrogen atoms and nonbonded electron pairs in each ion.



1.67 Draw in all H atoms and lone pairs in each ion.










1.68 Each of the following condensed or skeletal structures is an incorrect representation of a molecule or ion. Explain what is wrong in each structure.







Hybridization


1.69 Predict the hybridization and geometry around each indicated atom.




1.70 What orbitals are used to form each indicated bond? For multiple bonds, indicate the orbitals used in individual bonds.



1.71 Citric acid and zingerone are two naturally occurring compounds. What orbitals are used to form each indicated bond?





citric acid
(responsible for the tartness
of citrus fruits)



zingerone
(responsible for the pungent
taste of ginger)

1.72 Ketene, $\text{CH}_2=\text{C}=\text{O}$, is an unusual organic molecule that has a single carbon atom doubly bonded to two different atoms. Determine the hybridization of both C atoms and the O in ketene. Then, draw a diagram showing what orbitals are used to form each bond (similar to Figures 1.12 and 1.13).





1.73 Consider the unstable cation and anion drawn below.






- What is the hybridization of each carbon atom in these ions?
- What orbitals are used to form both bonds in each carbon-carbon double bond?

Bond Length and Strength


1.74 Rank the following bonds in order of *increasing* bond length.



1.75 Indicate the longer labeled bond in each compound.





1.76 Answer the following questions about compound **A**.



- Label the shortest C-C single bond.
- Label the longest C-C single bond.
- Considering all the bonds, label the shortest C-C bond.
- Label the weakest C-C bond.
- Label the strongest C-H bond.
- Explain why bond (1) and bond (2) are different in length, even though they are both C-C single bonds.

1.77 A σ bond formed from two sp^2 hybridized C atoms is stronger than a σ bond formed from two sp^3 hybridized C atoms. Explain.

1.78 Two useful organic compounds that contain Cl atoms are drawn below. Vinyl chloride is the starting material used to prepare poly(vinyl chloride), a plastic used in insulation, pipes, and bottles. Chloroethane (ethyl chloride) is a local anesthetic. Why is the C-Cl bond of vinyl chloride stronger than the C-Cl bond in chloroethane?






Bond Polarity

1.79 Answer each question with a brief explanation and an example to illustrate your answer.




- Can a compound be nonpolar if it contains one polar bond?
- Can a compound be nonpolar if it contains two or more polar bonds?
- Can a compound be polar if it contains no polar bonds?

1.80 Use the symbols δ^+ and δ^- to indicate the polarity of the labeled bonds.



1.81 Label the polar bonds in each molecule. Indicate the direction of the net dipole (if there is one).

- CHBr_3
- $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$
- CBr_4




General Questions

1.82 Answer the following questions about acetonitrile ($\text{CH}_3\text{C}\equiv\text{N}$).

- Determine the hybridization of both C atoms and the N atom.
- Label all bonds as σ or π .
- In what type of orbital does the lone pair on N reside?
- Label all bonds as polar or nonpolar.

1.83 Benzene is the simplest member of a whole class of compounds called aromatic hydrocarbons.




- How is each carbon atom hybridized?
- What is the geometry around each carbon atom? What is the overall geometry of the benzene ring?
- Draw a diagram showing the orbitals used to join the carbon atoms of the ring.
- Follow the indicated curved arrow notation to draw a second resonance structure.




- Benzene and other aromatic hydrocarbons are shown in Chapter 17 to be very stable. Offer an explanation.

1.84 The principles of this chapter can be applied to organic molecules of any size. Answer the following questions about amoxicillin, an antibiotic from the penicillin family.




- Predict the hybridization and geometry around each indicated atom.
- Label five polar bonds using the symbols δ^+ and δ^- .
- Draw a skeletal structure.
- How many π bonds does amoxicillin have? Label them.
- Find a C–H bond containing a carbon atom having a hybrid orbital with 33% s-character.

1.85



- What is the hybridization of each N atom in nicotine?
- What is the geometry around each N atom?
- In what type of orbital does the lone pair on each N atom reside?
- Draw a constitutional isomer of nicotine.
- Draw a resonance structure of nicotine.



1.86 Stalevo is the trade name for a medication used for Parkinson's disease, which contains both L-dopa and entacapone.



- Draw a Lewis structure for entacapone.
- Which C–C bond in entacapone is the longest?
- Which C–C single bond is the shortest?
- Which C–N bond is the longest?
- Which C–N bond is the shortest?
- Use curved arrows to draw a resonance structure that is an equal contributor to the resonance hybrid.
- Use curved arrows to draw a resonance structure that is a minor contributor to the resonance hybrid.


1.87 CH_3^+ and CH_3^- are two highly reactive carbon species.

- What is the predicted hybridization and geometry around each carbon atom?
- Two electrostatic potential plots are drawn for these species. Which ion corresponds to which diagram and why?




Challenge Problems

- 1.88** When two carbons having different hybridization are bonded together, the C–C bond contains a slight dipole. In a $C_{sp^2}-C_{sp^3}$ bond, what is the direction of the dipole? Which carbon is considered more electronegative?
- 1.89** Draw all possible isomers having molecular formula C_4H_8 that contain one π bond.
- 1.90** Use the principles of resonance theory to explain why carbocation **A** is more stable than carbocation **B**.



- 1.91** The curved arrow notation introduced in Section 1.5 is a powerful method used by organic chemists to show the movement of electrons not only in resonance structures, but also in chemical reactions. Since each curved arrow shows the movement of two electrons, following the curved arrows illustrates what bonds are broken and formed in a reaction. Consider the following three-step process. (a) Add curved arrows in Step [1] to show the movement of electrons. (b) Use the curved arrows drawn in Step [2] to identify the structure of **X**. **X** is converted in Step [3] to phenol and HCl.



2

Acids and Bases

- 2.1 Brønsted–Lowry acids and bases
- 2.2 Reactions of Brønsted–Lowry acids and bases
- 2.3 Acid strength and pK_a
- 2.4 Predicting the outcome of acid–base reactions
- 2.5 Factors that determine acid strength
- 2.6 Common acids and bases
- 2.7 Aspirin
- 2.8 Lewis acids and bases



Aspirin is one of the most widely used over-the-counter drugs. Whether you purchase Anacin, Bufferin, Bayer, or a generic, the active ingredient is the same—**acetylsalicylic acid**. Aspirin, a synthetic compound that does not occur in nature, was first marketed to the general public in 1899, and is now used regularly by over 100 million people throughout the world. Like many drugs, aspirin undergoes a proton transfer reaction after ingestion. In Chapter 2, we learn about acidity and the role of acid–base reactions in aspirin’s chemistry.

Chemical terms such as *anion* and *cation* may be unfamiliar to most nonscientists, but *acid* has found a place in everyday language. Commercials advertise the latest remedy for the heartburn caused by excess stomach *acid*. The nightly news may report the latest environmental impact of *acid* rain. Wine lovers know that wine sours because its alcohol has turned to *acid*. *Acid* comes from the Latin word *acidus*, meaning sour, because when tasting compounds was a routine method of identification, these compounds were sour.

In Chapter 2, we will concentrate on two definitions of acids and bases: the **Brønsted–Lowry** definition, which describes acids as *proton donors* and bases as *proton acceptors*, and the **Lewis** definition, which describes acids as *electron pair acceptors* and bases as *electron pair donors*.

2.1 Brønsted–Lowry Acids and Bases

The general words “acid” and “base” usually mean a *Brønsted–Lowry* acid and *Brønsted–Lowry* base.

H^+ = proton

H–A = general Brønsted–Lowry acid.

B: = general Brønsted–Lowry base.

The Brønsted–Lowry definition describes acidity in terms of protons: positively charged **hydrogen ions, H^+** .


- A Brønsted–Lowry acid is a *proton donor*.
- A Brønsted–Lowry base is a *proton acceptor*.

A Brønsted–Lowry acid must contain a hydrogen atom. This definition of an acid is often familiar to students, because many inorganic acids in general chemistry are Brønsted–Lowry acids. The symbol **H–A** is used for a general Brønsted–Lowry acid.

A Brønsted–Lowry base must be able to form a bond to a proton. Because a proton has no electrons, a base must contain an “available” **electron pair** that can be easily donated to form a new bond. These include **lone pairs** or electron pairs in **π bonds**. The symbol **B:** is used for a general Brønsted–Lowry base.

Examples of Brønsted–Lowry acids and bases are given in Figure 2.1.

Charged species such as ^-OH and $^-NH_2$ are used as **salts**, with cations such as Li^+ , Na^+ , or K^+ to balance the negative charge. These cations are called **counterions** or **spectator ions**, and their **identity is usually inconsequential**. For this reason, the counterion is often omitted.




Compounds like H_2O and CH_3OH that contain both hydrogen atoms and lone pairs may be either an acid or a base, depending on the particular reaction. These fundamental principles are true no matter how complex the compound. For example, the addictive pain reliever **morphine** is a

Figure 2.1


Examples of Brønsted–Lowry acids and bases

Brønsted–Lowry acids [H–A]

Inorganic




Organic




- All Brønsted–Lowry acids contain a proton.
- The net charge may be zero, (+), or (–).

Brønsted–Lowry bases [B:]


Inorganic



Organic




- All Brønsted–Lowry bases contain a lone pair of electrons or a π bond.
- The net charge may be zero or (–).



Morphine is obtained from the opium poppy.

Brønsted–Lowry acid because it contains many hydrogen atoms. It is also a Brønsted–Lowry base because it has lone pairs on O and N, and four π bonds.



Problem 2.1

- Which compounds are Brønsted–Lowry acids: HBr, NH_3 , CCl_4 ?
- Which compounds are Brønsted–Lowry bases: CH_3CH_3 , $(\text{CH}_3)_3\text{CO}^-$, $\text{HC}\equiv\text{CH}$?
- Classify each compound as an acid, a base, or both: $\text{CH}_3\text{CH}_2\text{OH}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CO}_2\text{CH}_3$.


2.2 Reactions of Brønsted–Lowry Acids and Bases

A Brønsted–Lowry acid–base reaction results in transfer of a proton from an acid to a base. These acid–base reactions, also called *proton transfer reactions*, are fundamental to the study of organic chemistry.

Consider, for example, the reaction of the acid $\text{H}-\text{A}$ with the base $:\text{B}$. In an acid–base reaction, one bond is broken and one is formed.

- The electron pair of the base B : forms a new bond to the proton of the acid.
- The acid $\text{H}-\text{A}$ loses a proton, leaving the electron pair in the $\text{H}-\text{A}$ bond on A .

Recall from Section 1.5 that a curved arrow shows the movement of an **electron pair**. The tail of the arrow always begins at an electron pair and the head points to where that electron pair “moves.”



This “movement” of electrons in reactions can be illustrated using curved arrow notation. Because **two electron pairs** are involved in this reaction, **two curved arrows** are needed. Two products are formed.


- Loss of a proton from an acid forms its *conjugate base*.
- Gain of a proton by a base forms its *conjugate acid*.

Keep in mind two other facts about this general reaction:

- The **net charge must be the same** on both sides of any equation. In this example, the net charge on each side is zero. Individual charges can be calculated using formal charges.
- A **double reaction arrow** is used between starting materials and products to indicate that the reaction can proceed in the forward and reverse directions. These are **equilibrium arrows**.


Two examples of proton transfer reactions are drawn here with curved arrow notation.

A double reaction arrow indicates equilibrium.



equilibrium arrows

Remove H^+ from an acid to form its conjugate base.
Add H^+ to a base to form its conjugate acid.



- Brønsted–Lowry acid–base reactions always result in the transfer of a proton from an acid to a base.

The ability to identify and draw a conjugate acid or base from a given starting material is illustrated in Sample Problems 2.1 and 2.2.


Sample Problem 2.1

- What is the conjugate acid of CH_3O^- ?
- What is the conjugate base of NH_3 ?

Solution


- Add H^+ to CH_3O^- to form its conjugate acid.

Add a proton to a lone pair.



- Remove H^+ from NH_3 to form its conjugate base.

Leave the electron pair on N.



Problem 2.2


- Draw the conjugate acid of each base: NH_3 , Cl^- , $(\text{CH}_3)_2\text{C}=\text{O}$.
- Draw the conjugate base of each acid: HBr , HSO_4^- , CH_3OH .

Problem 2.3

The presence of a π bond also makes a compound a base. With this in mind, draw the conjugate acid of ethylene, $\text{CH}_2=\text{CH}_2$. What is ethylene's conjugate base?


Sample Problem 2.2

Label the acid and base, and the conjugate acid and base in the following reaction. Use curved arrow notation to show the movement of electron pairs.




Solution

CH_3^- is the base because it accepts a proton, forming its conjugate acid, CH_4 . H_2O is the acid because it donates a proton, forming its conjugate base, OH^- . Two curved arrows are needed. One shows that the lone pair on CH_3^- bonds to a proton of H_2O , and the second shows that the electron pair in the $\text{O}-\text{H}$ bond remains on O.



Problem 2.4 Label the acid and base, and the conjugate acid and base in the following reactions. Use curved arrows to show the movement of electron pairs.




In all proton transfer reactions, the **electron-rich base** donates an electron pair to the acid, which usually has a polar H–A bond. Thus, the H of the acid bears a partial positive charge, making it **electron deficient**. This is the first example of a general pattern of reactivity.

- Electron-rich species react with electron-deficient ones.

Given two starting materials, how do you know which is the acid and which is the base in a proton transfer reaction? The following generalizations will help to decide this in many reactions:

- [1] Common acids and bases introduced in general chemistry will often be used in the same way in organic reactions. HCl and H₂SO₄ are strong acids, and [−]OH is a strong base.
- [2] When only one starting material contains a hydrogen, it must be the acid. If only one starting material has a lone pair or a π bond, it must be the base.
- [3] A starting material with a net positive charge is usually the acid. A starting material with a negative charge is usually the base.

Problem 2.5 Draw the products of each proton transfer reaction.



Problem 2.6 Draw the products formed from the acid–base reaction of HCl with each compound.




2.3 Acid Strength and pK_a

Acid strength is the tendency of an acid to donate a proton.

- The more readily a compound donates a proton, the stronger the acid.

Acidity is measured by an equilibrium constant. When a Brønsted–Lowry acid H–A is dissolved in water, an acid–base reaction occurs, and an equilibrium constant K_{eq} can be written for the reaction.



Equilibrium constant

$$K_{\text{eq}} = \frac{[\text{products}]}{[\text{starting materials}]} = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{H-A}][\text{H}_2\text{O}]}$$

Because the concentration of the solvent H₂O is essentially constant, the equation can be rearranged and a new equilibrium constant, called the **acidity constant**, K_{a} , can be defined.

$$\text{Acidity constant} = K_{\text{a}} = [\text{H}_2\text{O}]K_{\text{eq}} = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{H-A}]}$$

How is the magnitude of K_{a} related to acid strength?


- The *stronger the acid*, the further the equilibrium lies to the right, and the *larger the K_a* .

For most organic compounds, K_a is small, typically 10^{-5} to 10^{-50} . This contrasts with the K_a values for many inorganic acids, which range from 10^0 to 10^{10} . Because dealing with exponents can be cumbersome, it is often more convenient to use pK_a values instead of K_a values.

Definition: $pK_a = -\log K_a$


How does pK_a relate to acid strength?

Recall that a **log** is an **exponent**; for example, $\log 10^{-5} = -5$.



- The *smaller the pK_a* , the *stronger the acid*.

Problem 2.7 Which compound in each pair is the stronger acid?



Problem 2.8 Use a calculator when necessary to answer the following questions.

- What is the pK_a for each K_a : 10^{-10} , 10^{-21} , and 5.2×10^{-6} ?
- What is the K_a for each pK_a : 7, 11, and 3.2?

Understanding acid strength is the first step to understanding base strength (and vice versa), because **an inverse relationship exists between acidity and basicity**.

- A **strong acid** readily donates a proton, forming a **weak conjugate base**.
- A **strong base** readily accepts a proton, forming a **weak conjugate acid**.

Table 2.1 is a brief list of pK_a values for some common compounds, ranked in order of **increasing pK_a** and therefore **decreasing acidity**. Because strong acids form weak conjugate bases, this list also ranks their conjugate bases, in order of **increasing basicity**. For example, CH_4 is the

Table 2.1 Selected pK_a Values

	Acid	p <i>K</i> _a	Conjugate base
↑ Increasing acidity of the acid	H–Cl	–7	Cl [–]
	CH ₃ COO–H	4.8	CH ₃ COO [–]
	HO–H	15.7	HO [–]
	CH ₃ CH ₂ O–H	16	CH ₃ CH ₂ O [–]
	HC≡CH	25	HC≡C [–]
	H–H	35	H [–]
	H ₂ N–H	38	H ₂ N [–]
	CH ₂ =CH ₂	44	CH ₂ =C [–] H
	CH ₃ –H	50	CH ₃ [–]
			↓ Increasing basicity of the conjugate base

weakest acid in the list, because it has the highest pK_a (50). Its conjugate base, CH_3^- , is therefore the strongest conjugate base. An extensive pK_a table is located in Appendix A.

Comparing pK_a values thus provides two useful bits of information: the **relative acidity of two acids**, and the **relative basicity of their conjugate bases**, as shown in Sample Problem 2.3.


Sample Problem 2.3

Rank the following compounds in order of increasing acidity, and then rank their conjugate bases in order of increasing basicity.




Solution

The pK_a values in Table 2.1 allow us to rank these compounds in order of increasing acidity: **the lower the pK_a , the stronger the acid**.



Because strong acids form weak conjugate bases, the **basicity of conjugate bases increases with increasing pK_a** of their acids.



Problem 2.9

Rank the conjugate bases of each group of acids in order of increasing basicity.

- a. NH_3 , H_2O , CH_4 b. $\text{CH}_2=\text{CH}_2$, $\text{HC}\equiv\text{CH}$, CH_4

Problem 2.10

Consider two acids: HCO_2H (formic acid, $pK_a = 3.8$) and pivalic acid [$(\text{CH}_3)_3\text{CCO}_2\text{H}$, $pK_a = 5.0$]. (a) Which acid has the larger K_a ? (b) Which acid is the stronger acid? (c) Which acid forms the stronger conjugate base? (d) When each acid is dissolved in water, for which acid does the equilibrium lie further to the right?


The pK_a values in Table 2.1 span a large range (-7 to 50). The **pK_a scale is logarithmic**, so a small difference in pK_a translates into a large numerical difference. For example, the difference between the pK_a of NH_3 (38) and $\text{CH}_2=\text{CH}_2$ (44) is six pK_a units. This means that NH_3 is 10^6 or **one million times more acidic** than $\text{CH}_2=\text{CH}_2$.

Although Table 2.1 is abbreviated, it gives pK_a values of many common compounds seen in organic chemistry. It is also a useful tool for *estimating* the pK_a of a compound similar though not identical to one in the table.

Suppose you are asked to estimate the pK_a of the $\text{N}-\text{H}$ bond of CH_3NH_2 . Although CH_3NH_2 is not listed in the table, we have enough information to *approximate* its pK_a . Because the pK_a of the $\text{N}-\text{H}$ bond of NH_3 is 38 , we can estimate the pK_a of the $\text{N}-\text{H}$ bond of CH_3NH_2 to be 38 . Its actual pK_a is 40 , so this is a good first approximation.

Problem 2.11

Estimate the pK_a of each of the indicated bonds.



2.4 Predicting the Outcome of Acid–Base Reactions

In a proton transfer reaction, the **stronger acid reacts with the stronger base** to form the weaker acid and the weaker base.

A proton transfer reaction represents an equilibrium. Because an acid donates a proton to a base, thus forming a conjugate acid and conjugate base, there are always two acids and two bases in the reaction mixture. Which pair of acids and bases is favored at equilibrium? **The position of the equilibrium depends on the relative strengths of the acids and bases.**

- Equilibrium always favors formation of the weaker acid and base.

Because a strong acid readily donates a proton and a strong base readily accepts one, these two species react to form a weaker conjugate acid and base that do not donate or accept a proton as readily. Comparing pK_a values allows us to determine the position of equilibrium, as illustrated in Sample Problem 2.4.

Sample Problem 2.4

Determine the direction of equilibrium when acetylene ($\text{HC}\equiv\text{CH}$) reacts with NH_2^- in a proton transfer reaction.


Solution

Follow three steps to determine the position of equilibrium:


Step [1] Identify the acid and base in the starting materials.

- Assume NH_2^- is the base because it bears a net negative charge. That makes $\text{HC}\equiv\text{CH}$ the acid.

Step [2] Draw the products of proton transfer and identify the conjugate acid and base in the products.




Step [3] Compare the pK_a values of the acid and the conjugate acid. Equilibrium favors formation of the weaker acid with the higher pK_a .



- Because the pK_a of the starting acid (25) is **lower** than the pK_a of the conjugate acid (38), $\text{HC}\equiv\text{CH}$ is a **stronger** acid and equilibrium favors the products.

Problem 2.12

Draw the products of each reaction and determine the direction of equilibrium.



How can we know if a particular base is strong enough to deprotonate a given acid, so that the equilibrium lies to the right? The pK_a table readily gives us this information.


Compare any two entries in Table 2.1, such as ethanol ($\text{CH}_3\text{CH}_2\text{OH}$; $pK_a = 16$) and acetylene ($\text{HC}\equiv\text{CH}$; $pK_a = 25$), and their conjugate bases, ethoxide ($\text{CH}_3\text{CH}_2\text{O}^-$) and acetylide ($\text{HC}\equiv\text{C}^-$). Ethanol is a stronger acid than acetylene, so acetylide is a stronger base than ethoxide.

Because Table 2.1 is arranged from low to high pK_a , **an acid can be deprotonated by the conjugate base of any acid below it in the table.**

	acid	pK_a	conjugate base
stronger acid →	$\text{CH}_3\text{CH}_2\ddot{\text{O}}-\text{H}$ ethanol	16	$\text{CH}_3\text{CH}_2\ddot{\text{O}}^-$ ethoxide
	$\text{H}-\text{C}\equiv\text{C}-\text{H}$ acetylene	25	$\text{H}-\text{C}\equiv\text{C}^-$ ← stronger base


Two proton transfer reactions are possible.

- [1] Reaction of acetylene with ethoxide forms acetylide and ethanol. Because the stronger acid is the product of the reaction, *equilibrium favors the starting materials*. The base ethoxide is *not strong enough to deprotonate acetylene*.



Equilibrium favors the starting materials.

- [2] Reaction of ethanol with acetylide forms ethoxide and acetylene. Because the weaker acid is the product of the reaction, *equilibrium favors the products*. Thus, the base acetylide is *strong enough to deprotonate ethanol*.



Equilibrium favors the products.

In the second reaction, **ethanol is deprotonated by acetylide, the conjugate base of an acid weaker than itself**. This is a specific example of a general fact.

- An acid can be deprotonated by the conjugate base of any acid having a *higher* pK_a .

Problem 2.13 Answer the following questions by referring to Table 2.1:

- Which of the following bases is strong enough to deprotonate CH_3COOH : H^- , $\text{HC}\equiv\text{C}^-$, and Cl^- ?
- List four bases that are strong enough to deprotonate $\text{HC}\equiv\text{CH}$.

Problem 2.14 Using the data in Appendix A, determine which of the following bases is strong enough to deprotonate acetonitrile (CH_3CN) so that equilibrium favors the products: (a) NaH ; (b) Na_2CO_3 ; (c) NaOH ; (d) NaNH_2 ; (e) NaHCO_3 .

2.5 Factors That Determine Acid Strength

We have already learned in Section 2.3 that a tremendous difference in acidity exists among compounds. HCl ($pK_a < 0$) is an extremely strong acid. Water ($pK_a = 15.7$) is moderate in acidity, and CH_4 ($pK_a = 50$) is an extremely weak acid. How are these differences explained? There is one general rule.

- Anything that stabilizes a conjugate base A^- makes the starting acid $\text{H}-\text{A}$ more acidic.

For now we will concentrate on how structural differences between molecules can profoundly affect acidity. In Chapter 6, we will learn how to relate the stability of a species to its relative potential energy.

Four factors affect the acidity of $\text{H}-\text{A}$:

- Element effects**
- Inductive effects**
- Resonance effects**
- Hybridization effects**

No matter which factor is discussed, the same procedure is always followed. To compare the acidity of any two acids:

- Always draw the conjugate bases.
- Determine which conjugate base is more stable.
- The *more stable* the conjugate base, the *more acidic* the acid.


2.5A Element Effects—Trends in the Periodic Table

The most important factor determining the acidity of $\text{H}-\text{A}$ is the location of A in the periodic table.

Comparing Elements in the Same Row of the Periodic Table


Examine acidity trends **across a row** of the periodic table by comparing CH_4 and H_2O , two compounds having H atoms bonded to a second-row element. We know from Table 2.1 that **H_2O has a much lower pK_a and therefore is much more acidic than CH_4** , but why is this the case?

To answer this question, first draw both conjugate bases and then determine which is more stable. Each conjugate base has a net negative charge, but the negative charge in OH^- is on oxygen and in CH_3^- it is on carbon.



Because the oxygen atom is much **more electronegative** than carbon, oxygen more readily accepts a negative charge, making OH^- much more stable than CH_3^- . **H_2O is a stronger acid than CH_4 because OH^- is a more stable conjugate base than CH_3^- .** This is a specific example of a general trend.

- Across a row of the periodic table, the acidity of $\text{H}-\text{A}$ increases as the electronegativity of A increases.



The enormity of this effect is evident by noting the approximate pK_a values for these bonds. A **$\text{C}-\text{H}$ bond is approximately 10^{47} times less acidic than $\text{H}-\text{F}$.**

Comparing Elements Down a Column of the Periodic Table


Now examine acidity trends down a column of the periodic table by comparing $\text{H}-\text{F}$ and $\text{H}-\text{Br}$. Once again, first draw both conjugate bases and then determine which is more stable. In this case, removal of a proton forms F^- and Br^- .



There are two important differences between F^- and Br^- —electronegativity and size. In this case, **size is more important than electronegativity**. The size of an atom or ion increases down a column of the periodic table, so Br^- is much larger than F^- , and this stabilizes the negative charge.


- Positive or negative charge is stabilized when it is spread over a larger volume.

Because Br^- is larger than F^- , Br^- is more stable than F^- , and $\text{H}-\text{Br}$ is a stronger acid than $\text{H}-\text{F}$.



This again is a specific example of a general trend.

- Down a column of the periodic table, the acidity of $\text{H}-\text{A}$ increases as the size of A increases.



Because of carbon's position in the periodic table (in the second row and to the left of O, N, and the halogens), **C-H bonds are usually the least acidic bonds in a molecule.**

This is *opposite* to what would be expected on the basis of electronegativity differences between F and Br, because F is more electronegative than Br. **Size and not electronegativity determines acidity down a column.** Combining both trends together:

- The acidity of $\text{H}-\text{A}$ increases both left-to-right across a row and down a column of the periodic table.

Sample Problem 2.5

Without reference to a $\text{p}K_{\text{a}}$ table, decide which compound in each pair is the stronger acid:

- a. H_2O or HF b. H_2S or H_2O

Solution

- a. H_2O and $\text{H}-\text{F}$ both have H atoms bonded to a second-row element. Because the acidity of $\text{H}-\text{A}$ increases across a row of the periodic table, the $\text{H}-\text{F}$ bond is more acidic than the $\text{H}-\text{O}$ bond. **HF is a stronger acid than H_2O .**
- b. H_2O and H_2S both have H atoms bonded to elements in the same column. Because the acidity of $\text{H}-\text{A}$ increases down a column of the periodic table, the $\text{H}-\text{S}$ bond is more acidic than the $\text{H}-\text{O}$ bond. **H_2S is a stronger acid than H_2O .**

Problem 2.15

Without reference to a $\text{p}K_{\text{a}}$ table, decide which compound in each pair is the stronger acid.

- a. NH_3 or H_2O b. HBr or HCl c. H_2S or HBr

Problem 2.16


Which compound in each pair of isomers is the stronger acid?

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$ or $(\text{CH}_3)_3\text{N}$ b. $\text{CH}_3\text{CH}_2\text{OCH}_3$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$

When discussing acidity, the most acidic proton in a compound is the one removed first by a base. Although four factors determine the overall acidity of a particular hydrogen atom, **element effects—the identity of A—is the single most important factor in determining the acidity of the $\text{H}-\text{A}$ bond.**

To decide which hydrogen is most acidic, **first determine what element each hydrogen is bonded to and then decide its acidity based on periodic trends.** For example, $\text{CH}_3\text{NHCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$


contains only C–H and N–H bonds. Since the acidity of H–A increases across a row of the periodic table, the single H on N is the most acidic H in this compound.



Problem 2.17 Which hydrogen in each molecule is most acidic?

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ b. $\text{HOCH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ c. $(\text{CH}_3)_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}_2$

Problem 2.18 Which hydrogen in pseudoephedrine, the nasal decongestant in the commercial medication Sudafed, is most acidic?




pseudoephedrine

2.5B Inductive Effects




Because the pseudoephedrine (Problem 2.18) in Sudafed can be readily converted to the illegal, addictive drug methamphetamine, products that contain pseudoephedrine are now stocked behind the pharmacy counter so their sale can be more closely monitored. Sudafed PE is a related product that contains a decongestant less easily converted to methamphetamine.


A second factor affecting the acidity of H–A is the presence of electronegative atoms. To illustrate this phenomenon, compare ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) and 2,2,2-trifluoroethanol ($\text{CF}_3\text{CH}_2\text{OH}$), two different compounds containing O–H bonds. The $\text{p}K_a$ table in Appendix A indicates that $\text{CF}_3\text{CH}_2\text{OH}$ is a stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$. Because we are comparing the acidity of the O–H bond in both compounds, what causes the difference?



Once again first draw both conjugate bases and then determine which is more stable. Both bases have a negative charge on an electronegative oxygen, but the second anion has three additional electronegative fluorine atoms. These fluorine atoms withdraw electron density from the carbon to which they are bonded, making it electron deficient. Furthermore, this electron-deficient carbon pulls electron density through σ bonds from the negatively charged oxygen atom, stabilizing the negative charge. This is called an **inductive effect**.



No additional electronegative atoms stabilize the conjugate base.



CF_3 withdraws electron density, stabilizing the conjugate base.


- An *inductive effect* is the pull of electron density through σ bonds caused by electronegativity differences of atoms.

In this case, the electron density is pulled away from the negative charge through σ bonds by the very electronegative fluorine atoms, and so it is called an **electron-withdrawing inductive effect**. Thus, the three electronegative fluorine atoms stabilize the negatively charged conjugate base $\text{CF}_3\text{CH}_2\text{O}^-$, making $\text{CF}_3\text{CH}_2\text{OH}$ a stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$. We have learned two important principles from this discussion:


- More electronegative atoms stabilize regions of high electron density by an electron-withdrawing inductive effect.
- The acidity of H–A increases with the presence of electron-withdrawing groups in A.

Figure 2.2

Electrostatic potential plots of $\text{CH}_3\text{CH}_2\text{O}^-$ and $\text{CF}_3\text{CH}_2\text{O}^-$



The dark red of the O atom indicates a region of high electron density.



The O atom is yellow, indicating it is less electron rich.

Inductive effects result because an electronegative atom stabilizes the negative charge of the conjugate base. **The more electronegative the atom and the closer it is to the site of the negative charge, the greater the effect.** This effect is discussed in greater detail in Chapter 19.

Electrostatic potential plots in Figure 2.2 compare the electron density around the oxygen atoms in these conjugate bases. The darker red around the O atom of $\text{CH}_3\text{CH}_2\text{O}^-$ indicates a higher concentration of electron density compared to the O atom of $\text{CF}_3\text{CH}_2\text{O}^-$.

Problem 2.19


Which compound in each pair is the stronger acid?

- a. ClCH_2COOH or FCH_2COOH c. CH_3COOH or $\text{O}_2\text{NCH}_2\text{COOH}$
 b. $\text{Cl}_2\text{CHCH}_2\text{OH}$ or $\text{Cl}_2\text{CHCH}_2\text{CH}_2\text{OH}$

Problem 2.20

Glycolic acid, $\text{HOCH}_2\text{CO}_2\text{H}$, is the simplest member of a group of compounds called α -hydroxy acids, ingredients in skin care products that have an OH group on the carbon adjacent to a CO_2H group. Would you expect $\text{HOCH}_2\text{CO}_2\text{H}$ to be a stronger or weaker acid than acetic acid, $\text{CH}_3\text{CO}_2\text{H}$?


2.5C Resonance Effects




α -Hydroxy acids (Problem 2.20) are used in skin care products that purportedly smooth fine lines and improve skin texture by reacting with the outer layer of skin cells, causing them to loosen and flake off.

Recall that resonance structures are two Lewis structures having the same placement of atoms but a different arrangement of electrons.


A third factor that determines acidity is resonance. Recall from Section 1.5 that resonance occurs whenever two or more different Lewis structures can be drawn for the same arrangement of atoms. To illustrate this phenomenon, compare ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) and acetic acid (CH_3COOH), two different compounds containing O–H bonds. Based on Table 2.1, CH_3COOH is a stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$:



Draw the conjugate bases of these acids to illustrate the importance of resonance. For ethoxide ($\text{CH}_3\text{CH}_2\text{O}^-$), the conjugate base of ethanol, only one Lewis structure can be drawn. The negative charge of this conjugate base is *localized* on the O atom.



With acetate (CH_3COO^-), however, two resonance structures can be drawn.



Resonance delocalization often produces a larger effect on pK_a than the inductive effects discussed in Section 2.5B. Resonance makes CH_3COOH ($pK_a = 4.8$) a much stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$ ($pK_a = 16$), while the inductive effects due to three electronegative F atoms make $\text{CF}_3\text{CH}_2\text{OH}$ ($pK_a = 12.4$) a somewhat stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$.

The difference in these two resonance structures is the **position of a π bond** and a **lone pair**. Although each resonance structure of acetate implies that the negative charge is localized on an O atom, in actuality, charge is **delocalized** over both O atoms. **Delocalization of electron density stabilizes acetate, making it a weaker base.**

Remember that neither resonance form adequately represents acetate. The true structure is a **hybrid** of both structures. In the hybrid, the electron pairs drawn in different locations in individual resonance structures are **delocalized**. With acetate, a dashed line is used to show that each C–O bond has partial double bond character. The symbol δ^- (partial negative) indicates that the charge is delocalized on both O atoms in the hybrid.

Thus, **resonance delocalization makes CH_3COO^- more stable than $\text{CH}_3\text{CH}_2\text{O}^-$, so CH_3COOH is a stronger acid than $\text{CH}_3\text{CH}_2\text{OH}$.** This is another example of a general rule.

- The acidity of H–A increases when the conjugate base A^- is resonance stabilized.

Electrostatic potential plots of $\text{CH}_3\text{CH}_2\text{O}^-$ and CH_3COO^- in Figure 2.3 indicate that the negative charge is concentrated on a single O in $\text{CH}_3\text{CH}_2\text{O}^-$, but delocalized over the O atoms in CH_3COO^- .


Problem 2.21 The C–H bond in acetone, $(\text{CH}_3)_2\text{C}=\text{O}$, has a pK_a of 19.2. Draw two resonance structures for its conjugate base. Then, explain why acetone is much more acidic than propane, $\text{CH}_3\text{CH}_2\text{CH}_3$ ($pK_a = 50$).

Problem 2.22 Acetonitrile (CH_3CN) has a pK_a of 25, making it more acidic than many other compounds having only C–H bonds. Draw Lewis structures for acetonitrile and its conjugate base. Use resonance structures to account for the acidity of acetonitrile.


2.5D Hybridization Effects

The final factor affecting the acidity of H–A is the hybridization of A. To illustrate this phenomenon, compare ethane (CH_3CH_3), ethylene ($\text{CH}_2=\text{CH}_2$), and acetylene ($\text{HC}\equiv\text{CH}$), three different compounds containing C–H bonds. Appendix A indicates that there is a considerable difference in the pK_a values of these compounds.


Figure 2.3
Electrostatic potential plots of $\text{CH}_3\text{CH}_2\text{O}^-$ and CH_3COO^-




The negative charge is concentrated on the single oxygen atom, making this anion *less stable*.



The negative charge is delocalized over both oxygen atoms, making this anion *more stable*.



The conjugate bases formed by removing a proton from ethane, ethylene, and acetylene are each carbanions—**species with a negative charge on carbon**.



Note, however, that the hybridization of the carbon bearing the negative charge is different in each anion, so the lone pair of electrons occupies an orbital with a different percent *s*-character in each case. A higher percent *s*-character means a hybrid orbital has a larger fraction of the lower energy *s* orbital.

- The *higher* the percent *s*-character of the hybrid orbital, the **more stable** the conjugate base.

Thus, **acidity increases from CH_3CH_3 to $\text{CH}_2=\text{CH}_2$ to $\text{HC}\equiv\text{CH}$ as the negative charge of the conjugate base is stabilized by increasing percent *s*-character**. Once again this is a specific example of a general trend.

- The acidity of $\text{H}-\text{A}$ increases as the percent *s*-character of the A^- increases.

Electrostatic potential plots of these carbanions appear in Figure 2.4.

Problem 2.23

For each pair of compounds: [1] Which indicated H is more acidic? [2] Draw the conjugate base of each acid. [3] Which conjugate base is stronger?





Figure 2.4
Electrostatic potential plots
of three carbanions



- As the lone pair of electrons is pulled closer to the nucleus, the negatively charged carbon appears less intensely red.

Figure 2.5

Summary of the factors that determine acidity

Factor	Example		
1. Element effects: The acidity of H–A increases both left-to-right across a row and down a column of the periodic table.			
2. Inductive effects: The acidity of H–A increases with the presence of electron-withdrawing groups in A.	$\text{CH}_3\text{CH}_2\text{O}-\text{H}$	$\text{CF}_3\text{CH}_2\text{O}-\text{H}$	more acidic
3. Resonance effects: The acidity of H–A increases when the conjugate base A^- is resonance stabilized.	$\text{CH}_3\text{CH}_2\text{O}-\text{H}$	$\text{CH}_3\text{COO}-\text{H}$	more acidic
4. Hybridization effects: The acidity of H–A increases as the percent s-character of A^- increases.	CH_3CH_3	$\text{CH}_2=\text{CH}_2$	$\text{H}-\text{C}\equiv\text{C}-\text{H}$

2.5E Summary of Factors Determining Acid Strength

The ability to recognize the most acidic site in a molecule will be important throughout the study of organic chemistry. All the factors that determine acidity are therefore summarized in Figure 2.5. The following two-step procedure shows how these four factors can be used to determine the relative acidity of protons.

HOW TO Determine the Relative Acidity of Protons

Step [1] Identify the atoms bonded to hydrogen, and use periodic trends to assign relative acidity.

- The most common H–A bonds in organic compounds are C–H, N–H, and O–H. Because acidity increases left-to-right across a row, the relative acidity of these bonds is C–H < N–H < O–H. Therefore, H atoms bonded to C atoms are usually *less acidic* than H atoms bonded to any heteroatom.


Step [2] If the two H atoms in question are bonded to the same element, draw the conjugate bases and look for other points of difference. Ask three questions:

- Do electron-withdrawing groups stabilize the conjugate base?
- Is the conjugate base resonance stabilized?
- How is the conjugate base hybridized?


Sample Problem 2.6 shows how to apply this procedure to actual compounds.

Sample Problem 2.6

Rank the following compounds in order of increasing acidity of their most acidic hydrogen atom.



A



B




C

Solution

[1] Compounds **A**, **B**, and **C** contain C–H, N–H, and O–H bonds. Because acidity increases left-to-right across a row of the periodic table, the O–H bonds are most acidic. Compound **C** is thus the least acidic because it has *no* O–H bonds.


[2] The only difference between compounds **A** and **B** is the presence of an electronegative Cl in **A**. The Cl atom stabilizes the conjugate base of **A**, making it more acidic than **B**. Thus,



Problem 2.24 Rank the compounds in each group in order of increasing acidity.

- $\text{CH}_3\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{OH}$, $\text{CH}_3\text{CH}_2\text{NH}_2$
- BrCH_2COOH , $\text{CH}_3\text{CH}_2\text{COOH}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
- $\text{CH}_3\text{CH}_2\text{NH}_2$, $(\text{CH}_3)_3\text{N}$, $\text{CH}_3\text{CH}_2\text{OH}$


Problem 2.25 Which proton in each of the following drugs is most acidic? THC is the active component in marijuana, ketoprofen is an anti-inflammatory agent, and propranolol is an antihypertensive agent—that is, it lowers blood pressure.



2.6 Common Acids and Bases

Many strong or moderately strong acids and bases are used as reagents in organic reactions.


2.6A Common Acids



Sulfuric acid is the most widely produced industrial chemical. It is also formed when sulfur oxides, emitted into the atmosphere by burning fossil fuels high in sulfur content, dissolve in water. This makes rainwater acidic, forming acid rain.

Several organic reactions are carried out in the presence of strong inorganic acids, most commonly HCl and H_2SO_4 . These strong acids, with $\text{p}K_a$ values ≤ 0 , should be familiar from previous chemistry courses.

Two organic acids are also commonly used, namely **acetic acid** and ***p*-toluenesulfonic acid** (usually abbreviated as **TsOH**). Although acetic acid has a higher $\text{p}K_a$ than the inorganic acids, making it a weaker acid, it is more acidic than most organic compounds. *p*-Toluenesulfonic acid is similar in acidity to the strong inorganic acids. Because it is a solid, small quantities can be easily weighed on a balance and then added to a reaction mixture.



2.6B Common Bases

Common strong bases used in organic reactions are more varied in structure. Three common kinds of negatively charged bases include:



- Negatively charged oxygen bases: OH^- (hydroxide) and its organic derivatives
- Negatively charged nitrogen bases: NH_2^- (amide) and its organic derivatives
- Hydride (H^-)

Figure 2.6 gives examples of these strong bases. Each negatively charged base is used as a salt with a spectator ion (usually Li^+ , Na^+ , or K^+) that serves to balance charge.

Figure 2.6

Some common negatively charged bases


The modern history of aspirin (Section 2.7) dates back to 1763 when Reverend Edmund Stone reported on the analgesic effect of chewing on the bark of the willow tree. Willow bark is now known to contain *salicin*, which is structurally related to aspirin.




- Strong bases have weak conjugate acids with high $\text{p}K_a$ values, usually > 12 .

Strong bases have a net negative charge, but not all negatively charged species are strong bases. For example, none of the halides, F^- , Cl^- , Br^- , or I^- , is a strong base. These anions have very strong conjugate acids and have little affinity for donating their electron pairs to a proton.

Carbanions, negatively charged carbon atoms discussed in Section 2.5D, are especially strong bases. Perhaps the most common example is **butyllithium**. Butyllithium and related compounds are discussed in greater detail in Chapter 20.



Two other weaker organic bases are **triethylamine** and **pyridine**. These compounds have a lone pair on nitrogen, making them basic, but they are considerably weaker than the amide bases because they are neutral, not negatively charged.




Problem 2.26

Draw the products formed when 2-propanol [$(\text{CH}_3)_2\text{CHOH}$], the main ingredient in rubbing alcohol, is treated with each acid or base: (a) NaH ; (b) H_2SO_4 ; (c) $\text{Li}^+ \text{ } ^-\text{N}[\text{CH}(\text{CH}_3)_2]_2$; (d) $\text{CH}_3\text{CO}_2\text{H}$.

2.7 Aspirin

Aspirin, or acetylsalicylic acid, is the most well known member of a group of compounds called **salicylates**. Although aspirin was first used in medicine for its analgesic (pain-relieving), anti-pyretic (fever-reducing), and anti-inflammatory properties, today it is commonly used as an anti-platelet agent in the treatment and prevention of heart attacks and strokes. **Aspirin is a synthetic compound**; it does not occur in nature, though some related salicylates are found in willow bark and meadowsweet blossoms.


Like many drugs, aspirin is capable of undergoing a proton transfer reaction. Its most acidic proton is the H bonded to O, and in the presence of base, this H is readily removed.



Why is this acid–base reaction important? After ingestion, aspirin first travels into the stomach and then the intestines. In the acidic environment of the stomach, aspirin remains in its neutral form, but in the basic environment of the small intestine, aspirin is deprotonated to form its conjugate base, an ion.

Whether aspirin is present as its acid or its conjugate base is very important in determining whether it can permeate a cell. **To be active, aspirin must cross a cell membrane, and to do so, it must be neutral, not ionic.** This means that aspirin crosses a cell membrane and is absorbed by the body in its neutral form in the stomach. Aspirin's mechanism of action is discussed in greater detail in Chapter 19.

Problem 2.27



amphetamine

Compounds like amphetamine that contain nitrogen atoms are protonated by the HCl in the gastric juices of the stomach, and the resulting salt is then deprotonated in the basic environment of the intestines to regenerate the neutral form. Write proton transfer reactions for both of these processes. Where is amphetamine likely to be absorbed by the body?


2.8 Lewis Acids and Bases

The Lewis definition of acids and bases is more general than the Brønsted–Lowry definition.

- A Lewis acid is an *electron pair acceptor*.
- A Lewis base is an *electron pair donor*.


Lewis bases are structurally the same as Brønsted–Lowry bases. Both have an **available electron pair**—a lone pair or an electron pair in a π bond. A Brønsted–Lowry base always donates this electron pair to a proton, but a Lewis base donates this electron pair to anything that is electron deficient.

All Brønsted–Lowry bases are Lewis bases.



A Lewis acid must be able to accept an electron pair, but there are many ways for this to occur. **All Brønsted–Lowry acids are also Lewis acids, but the reverse is not necessarily true.** Any species that is electron deficient and capable of accepting an electron pair is also a Lewis acid.

Common examples of Lewis acids (which are not Brønsted–Lowry acids) include BF_3 and AlCl_3 . These compounds contain elements in group 3A of the periodic table that can accept an electron pair because they do not have filled valence shells of electrons.



Problem 2.28

Which compounds are Lewis bases?

- a. NH_3 b. $\text{CH}_3\text{CH}_2\text{CH}_3$ c. H^- d. $\text{H}-\text{C}\equiv\text{C}-\text{H}$

Problem 2.29

Which compounds are Lewis acids?


- a. BBr_3 b. $\text{CH}_3\text{CH}_2\text{OH}$ c. $(\text{CH}_3)_3\text{C}^+$ d. Br^-

Any reaction in which one species donates an electron pair to another species is a Lewis acid–base reaction.

In a Lewis acid–base reaction, a Lewis base donates an electron pair to a Lewis acid. Most of the reactions in organic chemistry involving movement of electron pairs can be classified as Lewis acid–base reactions. Lewis acid–base reactions illustrate a general pattern of reactivity in organic chemistry.

- Electron-rich species react with electron-poor species.

In the simplest Lewis acid–base reaction one bond is formed and no bonds are broken. This is illustrated with the reaction of BF_3 with H_2O . BF_3 has only six electrons around B, so it is the electron-deficient Lewis acid. H_2O has two lone pairs on O, so it is the electron-rich Lewis base.




H_2O donates an electron pair to BF_3 to form one new bond. The electron pair in the new B–O bond comes from the oxygen atom, and a single product is formed. Both B and O bear formal charges in the product, but the overall product is neutral.

Nucleophile = nucleus loving.
Electrophile = electron loving.

- A Lewis acid is also called an *electrophile*.
- When a Lewis base reacts with an electrophile other than a proton, the Lewis base is also called a *nucleophile*.



In this Lewis acid–base reaction, **BF_3 is the electrophile** and **H_2O is the nucleophile**.

Two other examples are drawn. In each reaction the **electron pair is not removed from the Lewis base**; instead, the electron pair is donated to an atom of the Lewis acid, and one new covalent bond is formed.




Problem 2.30

For each reaction, label the Lewis acid and base. Use curved arrow notation to show the movement of electron pairs.



Problem 2.31 Draw the products of each reaction, and label the nucleophile and electrophile.




Problem 2.32 Draw the product formed when $(\text{CH}_3\text{CH}_2)_3\text{N:}$, a Lewis base, reacts with each Lewis acid: (a) $\text{B}(\text{CH}_3)_3$; (b) $(\text{CH}_3)_3\text{C}^+$; (c) AlCl_3 .

In some Lewis acid–base reactions, one bond is formed and one bond is broken. To draw the products of these reactions, keep the following steps in mind.


- [1] Always identify the Lewis acid and base first.
- [2] Draw a curved arrow from the electron pair of the base to the electron-deficient atom of the acid.
- [3] Count electron pairs and break a bond when needed to keep the correct number of valence electrons.

Recall from Chapter 1 that a positively charged carbon atom is called a **carbocation**.




In the reaction of cyclohexene with HCl, the new bond to H could form at **either carbon of the double bond**, because the same carbocation results.

For example, draw the Lewis acid–base reaction between cyclohexene and H–Cl. The Brønsted–Lowry acid HCl is also a Lewis acid, and cyclohexene, having a π bond, is the Lewis base.



To draw the product of this reaction, the electron pair in the π bond of the Lewis base forms a new bond to the proton of the Lewis acid, forming a carbocation. The H–Cl bond must break, giving its two electrons to Cl, forming Cl^- . Because two electron pairs are involved, two curved arrows are needed.



The Lewis acid–base reaction of cyclohexene with HCl is a specific example of a fundamental reaction of compounds containing C–C double bonds, as discussed in Chapter 10.

Problem 2.33 Label the Lewis acid and base. Use curved arrow notation to show the movement of electron pairs.



KEY CONCEPTS


Acids and Bases

A Comparison of Brønsted–Lowry and Lewis Acids and Bases


Type	Definition	Structural feature	Examples
Brønsted–Lowry acid (2.1)	proton donor	a proton	HCl, H ₂ SO ₄ , H ₂ O, CH ₃ COOH, TsOH
Brønsted–Lowry base (2.1)	proton acceptor	a lone pair <i>or</i> a π bond	OH^- , OCH_3^- , H^- , NH_2^- , NH_3 , $\text{CH}_2=\text{CH}_2$
Lewis acid (2.8)	electron pair acceptor	a proton, <i>or</i> an unfilled valence shell, <i>or</i> a partial (+) charge	BF_3 , AlCl_3 , HCl, CH ₃ COOH, H ₂ O
Lewis base (2.8)	electron pair donor	a lone pair <i>or</i> a π bond	OH^- , OCH_3^- , H^- , NH_2^- , NH_3 , $\text{CH}_2=\text{CH}_2$

Acid–Base Reactions

[1] A Brønsted–Lowry acid donates a proton to a Brønsted–Lowry base (2.2).



[2] A Lewis base donates an electron pair to a Lewis acid (2.8).




- Electron-rich species react with electron-poor ones.
- Nucleophiles react with electrophiles.

Important Facts

- Definition: $\text{p}K_{\text{a}} = -\log K_{\text{a}}$. The lower the $\text{p}K_{\text{a}}$, the stronger the acid (2.3).
- The stronger the acid, the weaker the conjugate base (2.3).
- In proton transfer reactions, equilibrium favors the weaker acid and weaker base (2.4).
- An acid can be deprotonated by the conjugate base of an acid having a higher $\text{p}K_{\text{a}}$ (2.4).

Periodic Trends in Acidity and Basicity (2.5A)



Factors That Determine Acidity (2.5)

- | | |
|---|---|
| [1] Element effects (2.5A) | The acidity of H–A increases both left-to-right across a row and down a column of the periodic table. |
| [2] Inductive effects (2.5B) | The acidity of H–A increases with the presence of electron-withdrawing groups in A. |
| [3] Resonance effects (2.5C) | The acidity of H–A increases when the conjugate base A^- is resonance stabilized. |
| [4] Hybridization effects (2.5D) | The acidity of H–A increases as the percent s-character of the A^- increases. |

PROBLEMS

Brønsted–Lowry Acids and Bases

2.34 What is the conjugate acid of each base?

- a. H_2O b. NH_2^- c. HCO_3^- d. $\text{CH}_3\text{CH}_2\text{NHCH}_3$ e. CH_3OCH_3 f. CH_3COO^-

2.35 What is the conjugate base of each acid?





- a. HCN b. HCO_3^- c. $(\text{CH}_3)_2\text{NH}_2^+$ d. $\text{HC}\equiv\text{CH}$ e. $\text{CH}_3\text{CH}_2\text{COOH}$ f. $\text{CH}_3\text{SO}_3\text{H}$

Reactions of Brønsted–Lowry Acids and Bases




2.36 Draw the products formed from the acid–base reaction of H_2SO_4 with each compound.

- a. b. c. d.




2.37 Draw the products formed from the acid–base reaction of KOH with each compound.




2.38 Draw the products of each proton transfer reaction. Label the acid and base in the starting materials, and the conjugate acid and base in the products.





2.39 Draw the products of each acid–base reaction.





2.40 Amphetamine is a powerful stimulant of the central nervous system. Draw the products formed from the acid–base reaction of amphetamine with each reagent: (a) HCl; (b) NaH.



2.41 Draw the products of each acid–base reaction.






2.42 Fenfluramine and phentermine are two components of fen–phen, an appetite suppressant withdrawn from the market in 1997 after it was shown to damage the heart valves in some patients. What products are formed when fenfluramine and phentermine are each treated with acetic acid ($\text{CH}_3\text{CO}_2\text{H}$)?










$\text{p}K_a$, K_a , and the Direction of Equilibrium

2.43 What is K_a for each compound? Use a calculator when necessary.








2.44 What is the $\text{p}K_a$ for each compound? Use a calculator when necessary.






- 2.45** Which bases in Table 2.1 are strong enough to deprotonate each compound?
 a. H_2O b. NH_3 c. CH_4
- 2.46** Which of the following bases are strong enough to deprotonate $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$ ($\text{p}K_{\text{a}} = 25$), so that equilibrium favors the products: (a) H_2O ; (b) NaOH ; (c) NaNH_2 ; (d) NH_3 ; (e) NaH ; (f) CH_3Li ?
- 2.47** Which compounds can be deprotonated by OH^- , so that equilibrium favors the products? Refer to the $\text{p}K_{\text{a}}$ table in Appendix A.
 a. HCOOH b. H_2S c.  d. CH_3NH_2
- 2.48** Draw the products of each reaction. Use the $\text{p}K_{\text{a}}$ table in Appendix A to decide if the equilibrium favors the starting materials or products.
- a.  + $\text{O}^-\text{CH}_2\text{CH}_3 \rightleftharpoons$
- b.  + $\text{NaCl} \rightleftharpoons$
- c. $(\text{CH}_3)_3\text{COH} + \text{H}_2\text{SO}_4 \rightleftharpoons$
- d.  + $\text{NaHCO}_3 \rightleftharpoons$
- e. $\text{H}-\text{C}\equiv\text{C}-\text{H} + \text{Li}^+\text{O}^-\text{CH}_2\text{CH}_3 \rightleftharpoons$
- f. $\text{CH}_3\text{NH}_2 + \text{H}_2\text{SO}_4 \rightleftharpoons$

Relative Acid Strength


- 2.49** Rank the following compounds in order of increasing acidity.
 a. NH_3 , H_2O , HF
 b. HBr , HCl , HF
 c. H_2O , H_3O^+ , HO^-
 d. NH_3 , H_2O , H_2S
 e. CH_3OH , CH_3NH_2 , CH_3CH_3
 f. HCl , H_2O , H_2S
 g. $\text{CH}_3\text{CH}_2\text{CH}_3$, $\text{ClCH}_2\text{CH}_2\text{OH}$, $\text{CH}_3\text{CH}_2\text{OH}$
 h. $\text{HC}\equiv\text{CCH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CH}=\text{CHCH}_3$
- 2.50** Rank the following ions in order of increasing basicity.
 a. $\text{CH}_3\text{C}\bar{\text{H}}_2$, CH_3O^- , CH_3NH^-
 b. CH_3^- , HO^- , Br^-
 c. CH_3COO^- , $\text{CH}_3\text{CH}_2\text{O}^-$, $\text{ClCH}_2\text{COO}^-$
 d. , , 
- 2.51** The $\text{p}K_{\text{a}}$'s of the two ammonium cations drawn below are 8.33 and 11.1. Which $\text{p}K_{\text{a}}$ corresponds to which cation? Explain your choice.



- 2.52** Explain why the $\text{p}K_{\text{a}}$ of propiolic acid ($\text{HC}\equiv\text{CCO}_2\text{H}$, $\text{p}K_{\text{a}} = 1.8$) is significantly lower than the $\text{p}K_{\text{a}}$ of propanoic acid ($\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$, $\text{p}K_{\text{a}} = 4.9$).
- 2.53** The $\text{p}K_{\text{a}}$ of three different C–H bonds is given below.




- a. For each compound, draw the conjugate base, including all possible resonance structures.
 b. Explain the observed trend in $\text{p}K_{\text{a}}$.
- 2.54** a. What is the conjugate acid of **A**?
 b. What is the conjugate base of **A**?



2.55 Many drugs are Brønsted–Lowry acids or bases.


- What is the most acidic proton in the analgesic ibuprofen? Draw the conjugate base.
- What is the most basic electron pair in cocaine? Draw the conjugate acid.



2.56 The pK_a of nitromethane (CH_3NO_2) is 10, making its C–H bond more acidic than most C–H bonds. Explain.


2.57 Dimethyl ether (CH_3OCH_3) and ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) are isomers, but CH_3OCH_3 has a pK_a of 40 and $\text{CH}_3\text{CH}_2\text{OH}$ has a pK_a of 16. Why are these pK_a values so different?

2.58 Atenolol is a β (beta) blocker, a drug used to treat high blood pressure. Which of the indicated N–H bonds is more acidic? Explain your reasoning.



2.59 Ethyl butanoate, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$, is one of the many organic compounds isolated from mangoes. Which hydrogen is most readily removed when ethyl butanoate is treated with base? Propose a reason for your choice, and using the data in Appendix A, estimate its pK_a .

2.60 Use the principles in Section 2.5 to label the most acidic hydrogen in each drug. Explain your choice.



2.61 Label the three most acidic hydrogen atoms in lactic acid, $\text{CH}_3\text{CH}(\text{OH})\text{CO}_2\text{H}$, and rank them in order of decreasing acidity. Explain your reasoning.

Lewis Acids and Bases

2.62 Classify each compound as a Lewis base, a Brønsted–Lowry base, both, or neither.




2.63 Classify each species as a Lewis acid, a Brønsted–Lowry acid, both, or neither.




Lewis Acid–Base Reactions

2.64 Label the Lewis acid and Lewis base in each reaction. Use curved arrows to show the movement of electron pairs.




2.65 Draw the products of each Lewis acid–base reaction. Label the electrophile and nucleophile.



2.66 Draw the product formed when the Lewis acid $(\text{CH}_3\text{CH}_2)_3\text{C}^+$ reacts with each Lewis base: (a) H_2O ; (b) CH_3OH ; (c) $(\text{CH}_3)_2\text{O}$; (d) NH_3 ; (e) $(\text{CH}_3)_2\text{NH}$.

General


2.67 Classify each reaction as either a proton transfer reaction, or a reaction of a nucleophile with an electrophile. Use curved arrows to show how the electron pairs move.



2.68 Hydroxide (OH^-) can react as a Brønsted–Lowry base (and remove a proton), or a Lewis base (and attack a carbon atom).

(a) What organic product is formed when OH^- reacts with the carbocation $(\text{CH}_3)_3\text{C}^+$ as a Brønsted–Lowry base? (b) What organic product is formed when OH^- reacts with $(\text{CH}_3)_3\text{C}^+$ as a Lewis base?

2.69 Answer the following questions about esmolol, a drug used to treat high blood pressure sold under the trade name Brevibloc.




- Label the two most acidic hydrogen atoms in esmolol, and explain which H is more acidic.
- What products are formed when esmolol is treated with NaH ?
- What products are formed when esmolol is treated with HCl ?
- Label all sp^2 hybridized C atoms.
- Label the only trigonal pyramidal atom.
- Label all C's that bear a δ^+ charge.


Challenge Problems

2.70 Molecules like acetamide (CH_3CONH_2) can be protonated on either their O or N atoms when treated with a strong acid like HCl . Which site is more readily protonated and why?


2.71 Two pK_a values are reported for malonic acid, a compound with two COOH groups. Explain why one pK_a is lower and one pK_a is higher than the pK_a of acetic acid (CH_3COOH , $pK_a = 4.8$).




2.72 Amino acids such as glycine are the building blocks of large molecules called proteins that give structure to muscle, tendon, hair, and nails.



- Explain why glycine does not actually exist in the form with all atoms uncharged, but actually exists as a salt called a zwitterion.
 - What product is formed when glycine is treated with concentrated HCl?
 - What product is formed when glycine is treated with NaOH?
- 2.73** Write a stepwise reaction sequence using proton transfer reactions to show how the following reaction occurs. (Hint: As a first step, use OH^- to remove a proton from the CH_2 group between the $\text{C}=\text{O}$ and $\text{C}=\text{C}$.)




2.74 Which H atom in vitamin C (ascorbic acid) is most acidic?



Introduction to Organic Molecules and Functional Groups

3

- 3.1 Functional groups
- 3.2 An overview of functional groups
- 3.3 Intermolecular forces
- 3.4 Physical properties
- 3.5 Application: Vitamins
- 3.6 Application of solubility: Soap
- 3.7 Application: The cell membrane
- 3.8 Functional groups and reactivity
- 3.9 Biomolecules



Vitamin C, or **ascorbic acid**, is important in the formation of collagen, a protein that holds together the connective tissues of skin, muscle, and blood vessels. Although citrus fruits (oranges, grapefruit, and lemons) are well known sources of vitamin C, guava, kiwi, and rose hips contain vitamin C, too. A deficiency of vitamin C causes scurvy, a common disease of sailors in the 1600s who had no access to fresh fruits on long voyages. In Chapter 3, we learn why some vitamins like vitamin A can be stored in the fat cells in the body, whereas others like vitamin C are excreted in urine.

Having learned some basic concepts about structure, bonding, and acid–base chemistry in Chapters 1 and 2, we will now concentrate on organic molecules.

- What are the characteristic features of an organic compound?
- What determines the properties of an organic compound?

After these questions are answered, we can understand some important phenomena. For example, why do we store some vitamins in the body and readily excrete others? How does soap clean away dirt? We will also use the properties of organic molecules to explain some basic biological phenomena, such as the structure of cell membranes and the transport of species across these membranes.

3.1 Functional Groups

What are the characteristic features of an organic compound? Most organic molecules have C–C and C–H σ bonds. These bonds are strong, nonpolar, and not readily broken. Organic molecules may have the following structural features as well:


- **Heteroatoms**—atoms other than carbon or hydrogen. Common heteroatoms are nitrogen, oxygen, sulfur, phosphorus, and the halogens.
- **π Bonds**. The most common π bonds occur in C–C and C–O double bonds.

These structural features distinguish one organic molecule from another. They determine a molecule's geometry, physical properties, and reactivity, and comprise what is called a **functional group**.


- A **functional group** is an atom or a group of atoms with characteristic chemical and physical properties. It is the **reactive part** of the molecule.

Why do heteroatoms and π bonds confer reactivity on a particular molecule?

- Heteroatoms have lone pairs and create electron-deficient sites on carbon.
- π Bonds are easily broken in chemical reactions. A π bond makes a molecule a base and a nucleophile.




Don't think, though, that the C–C and C–H σ bonds are unimportant. They form the **carbon backbone** or **skeleton** to which the functional groups are bonded. A functional group usually behaves the same whether it is bonded to a carbon skeleton having as few as two or as many as 20 carbons. For this reason, we often abbreviate the carbon and hydrogen portion of the molecule by a capital letter **R**, and draw the **R** bonded to a particular functional group.



Ethane, for example, has only C–C and C–H σ bonds, so it has *no* functional group. Ethane has no polar bonds, no lone pairs, and no π bonds, so it has **no reactive sites**. Because of this, ethane and molecules like it are very unreactive.

Ethanol, on the other hand, has two carbons and five hydrogens in its carbon backbone, as well as an OH group, a functional group called a **hydroxy** group. Ethanol has lone pairs and polar bonds that make it reactive with a variety of reagents, including the acids and bases discussed in Chapter 2.

The hydroxy group makes the properties of ethanol very different from the properties of ethane. Moreover, any organic molecule containing a hydroxy group has properties similar to ethanol.



Most organic compounds can be grouped into a relatively small number of categories, based on the structure of their functional group. Ethane, for example, is an **alkane**, whereas ethanol is a simple **alcohol**.

Problem 3.1 What reaction occurs when $\text{CH}_3\text{CH}_2\text{OH}$ is treated with (a) H_2SO_4 ? (b) NaH ? What happens when CH_3CH_3 is treated with these same reagents?

3.2 An Overview of Functional Groups

We can subdivide the most common functional groups into three types. A more complete list of functional groups is presented on the inside front cover.

- **Hydrocarbons**
- **Compounds containing a C–Z σ bond** where Z = an electronegative element
- **Compounds containing a C=O group**

3.2A Hydrocarbons


To review the structure and bonding of the simple aliphatic hydrocarbons, return to Section 1.9.

The word *aliphatic* is derived from the Greek word *aleiphas* meaning *fat*. Aliphatic compounds have physical properties similar to fats.

Hydrocarbons are compounds made up of only the elements carbon and hydrogen. They may be **aliphatic** or **aromatic**.





- [1] **Aliphatic hydrocarbons.** Aliphatic hydrocarbons can be divided into three subgroups.
- **Alkanes** have only C–C σ bonds and no functional group. Ethane, CH_3CH_3 , is a simple alkane.
 - **Alkenes** have a C–C double bond as a functional group. Ethylene, $\text{CH}_2=\text{CH}_2$, is a simple alkene.
 - **Alkynes** have a C–C triple bond as a functional group. Acetylene, $\text{HC}\equiv\text{CH}$, is a simple alkyne.
- [2] **Aromatic hydrocarbons.** This class of hydrocarbons was so named because many of the earliest known aromatic compounds had strong, characteristic odors.

The simplest aromatic hydrocarbon is **benzene**. The six-membered ring and three π bonds of benzene comprise a *single* functional group. Benzene is a component of the **BTX** mixture (**B** for benzene) added to gasoline to boost octane ratings.




When a benzene ring is bonded to another group, it is called a **phenyl group**. In phenylcyclohexane, for example, a phenyl group is bonded to the six-membered cyclohexane ring. Table 3.1 summarizes the four different types of hydrocarbons.

Table 3.1 Hydrocarbons

Type of compound	General structure	Example	Functional group
Alkane	R-H	CH ₃ CH ₃	—
Alkene			double bond
Alkyne	—C≡C—	H-C≡C-H	triple bond
Aromatic compound			phenyl group


Polyethylene is a synthetic plastic first produced in the 1930s, and initially used as insulating material for radar during World War II. It is now a plastic used in milk containers, sandwich bags, and plastic wrapping. Over 100 billion pounds of polyethylene are manufactured each year.

Alkanes, which have no functional groups, are notoriously unreactive except under very drastic conditions. For example, **polyethylene** is a synthetic plastic and high molecular weight alkane, consisting of chains of $-\text{CH}_2-$ groups bonded together, hundreds or even thousands of atoms long. Because it is an alkane with no reactive sites, it is a very stable compound that does not readily degrade and thus persists for years in landfills.



3.2B Compounds Containing C-Z σ Bonds

Several types of functional groups that contain C-Z σ bonds are listed in Table 3.2. The electronegative heteroatom Z creates a polar bond, making carbon electron deficient. The lone pairs on Z are available for reaction with protons and other electrophiles, especially when Z = N or O.



Several simple compounds in this category are widely used. As an example, chloroethane ($\text{CH}_3\text{CH}_2\text{Cl}$, commonly called ethyl chloride) is an alkyl halide used as a local anesthetic. Chloroethane quickly evaporates when sprayed on a wound, causing a cooling sensation that numbs the site of an injury.

Molecules containing these functional groups may be simple or very complex. Diethyl ether, the first common general anesthetic, is a simple ether because it contains a single O atom, depicted in red, bonded to two C atoms. Hemibrevetoxin B, on the other hand, contains four ether groups, in addition to other functional groups.



Chloroethane, $\text{CH}_3\text{CH}_2\text{Cl}$, is a local anesthetic.

Hemibrevetoxin B is a neurotoxin produced by algal blooms referred to as “red tides,” because of the color often seen in shallow ocean waters when these algae proliferate.





Table 3.2 Compounds Containing C–Z σ Bonds

Type of compound	General structure	Example	3-D structure	Functional group
Alkyl halide	$R-\ddot{X}:$ (X = F, Cl, Br, I)	$CH_3-\ddot{Br}:$		-X halo group
Alcohol	$R-\ddot{O}H$	$CH_3-\ddot{O}H$		-OH hydroxy group
Ether	$R-\ddot{O}-R$	$CH_3-\ddot{O}-CH_3$		-OR alkoxy group
Amine	$R-\ddot{N}H_2$ or $R_2\ddot{N}H$ or $R_3\ddot{N}$	$CH_3-\ddot{N}H_2$		-NH2 amino group
Thiol	$R-\ddot{S}H$	$CH_3-\ddot{S}H$		-SH mercapto group
Sulfide	$R-\ddot{S}-R$	$CH_3-\ddot{S}-CH_3$		-SR alkylthio group

3.2C Compounds Containing a C=O Group

Many different types of functional groups possess a C–O double bond (a **carbonyl group**), as shown in Table 3.3. The polar C–O bond makes the carbonyl carbon an electrophile, while the lone pairs on O allow it to react as a nucleophile and base. The carbonyl group also contains a π bond that is more easily broken than a C–O σ bond.

Reactive features of a carbonyl group



Atenolol and donepezil are examples of useful drugs that contain a variety of functional groups. **Atenolol is a β blocker**, a group of drugs used to treat hypertension. Donepezil, sold under the trade name Aricept, is used to treat mild to moderate dementia associated with Alzheimer's disease.























Table 3.3 Compounds Containing a C=O Group


Type of compound	General structure	Example	3-D structure	Functional group
Aldehyde				C=O carbonyl group
Ketone				C=O carbonyl group
Carboxylic acid				-COOH carboxy group
Ester				-COOR
Amide				-CONH ₂ , -CONHR, or -CONR ₂
Acid chloride				-COCl

The importance of a functional group cannot be overstated. A functional group determines all the following properties of a molecule:

- bonding and shape
- type and strength of intermolecular forces
- physical properties
- nomenclature
- chemical reactivity

Problem 3.2

Oseltamivir (trade name Tamiflu), the most effective antiviral drug against avian influenza currently available, can be prepared in 10 steps from shikimic acid. Identify the functional groups in Tamiflu and shikimic acid.



- Problem 3.3** Draw the structure of a compound fitting each description:
- a. An aldehyde with molecular formula C_4H_8O c. A carboxylic acid with molecular formula $C_4H_8O_2$
 b. A ketone with molecular formula C_4H_8O d. An ester with molecular formula $C_4H_8O_2$

- Problem 3.4** Draw structures that fit each description and name the functional group in each molecule:
 (a) two constitutional isomers with molecular formula $C_5H_{10}O$ that contain different functional groups;
 (b) two constitutional isomers with molecular formula $C_6H_{10}O$ that contain the same functional group.


3.3 Intermolecular Forces

Intermolecular forces are the interactions that exist *between* molecules. A functional group determines the type and strength of these interactions.

3.3A Ionic Compounds

Intermolecular forces are also referred to as **noncovalent interactions** or **nonbonded interactions**.

Ionic compounds contain oppositely charged particles held together by **extremely strong electrostatic interactions**. These ionic interactions are much stronger than the intermolecular forces present between covalent molecules, so it takes a great deal of energy to separate oppositely charged ions from each other.



3.3B Covalent Compounds

Covalent compounds are composed of discrete molecules. The nature of the forces between the molecules depends on the functional group present. There are three different types of interactions, presented here in order of *increasing strength*:

- van der Waals forces
- dipole-dipole interactions
- hydrogen bonding

van der Waals Forces


van der Waals forces, also called **London forces**, are very weak interactions caused by the **momentary changes in electron density in a molecule**. van der Waals forces are the only attractive forces present in nonpolar compounds.

For example, although a nonpolar CH_4 molecule has no net dipole, at any one instant its electron density may not be completely symmetrical, creating a *temporary dipole*. This can induce a temporary dipole in another CH_4 molecule, with the partial positive and negative charges arranged close to each other. **The weak interaction of these temporary dipoles constitutes van der Waals forces.** All compounds exhibit van der Waals forces.



Although any single van der Waals interaction is weak, a large number of van der Waals interactions creates a strong force. For example, geckos stick to walls and ceilings by van der Waals interactions of the surfaces with the 500,000 tiny hairs on each foot.

van der Waals interaction between two CH_4 molecules



Unsymmetrical electron density creates a temporary dipole.

The surface area of a molecule determines the strength of the van der Waals interactions. **The larger the surface area, the larger the attractive force between two molecules, and the stronger the intermolecular forces.** Long, sausage-shaped molecules such as $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ (pentane) have stronger van der Waals interactions than compact spherical ones like $\text{C}(\text{CH}_3)_4$ (neopentane), as shown in Figure 3.1.

Another factor affecting the strength of van der Waals forces is **polarizability**.

- **Polarizability** is a measure of how the electron cloud around an atom responds to changes in its electronic environment.


Larger atoms like iodine, which have more loosely held valence electrons, are more polarizable than smaller atoms like fluorine, which have more tightly held electrons. Because larger atoms have more easily induced dipoles, compounds containing them possess stronger intermolecular interactions.

Thus, two F_2 molecules have little force of attraction between them, because the electrons are held very tightly and temporary dipoles are difficult to induce. On the other hand, two I_2 molecules exhibit a much stronger force of attraction because the electrons are held much more loosely and temporary dipoles are easily induced.


Figure 3.1

Surface area and van der Waals forces


A long, cylindrical molecule




larger surface area
stronger van der Waals interactions




A compact, spherical molecule




smaller surface area
weaker van der Waals interactions






Dipole–Dipole Interactions

Dipole–dipole interactions are the attractive forces between the permanent dipoles of two polar molecules. In acetone, $(\text{CH}_3)_2\text{C}=\text{O}$, for example, the dipoles in adjacent molecules align so that the partial positive and partial negative charges are in close proximity. These attractive forces caused by permanent dipoles are much stronger than weak van der Waals forces.



Hydrogen Bonding

Hydrogen bonding typically occurs when a hydrogen atom bonded to O, N, or F, is electrostatically attracted to a lone pair of electrons on an O, N, or F atom in another molecule. Thus, H_2O molecules can hydrogen bond to each other. When they do, an H atom covalently bonded to O in one water molecule is attracted to a lone pair of electrons on the O in another water molecule. Hydrogen bonds are the strongest of the three types of intermolecular forces, though they are still much weaker than any covalent bond.



Hydrogen bonding helps determine the three-dimensional shape of large biomolecules such as carbohydrates and proteins. See Chapters 27 and 28 for details.


Sample Problem 3.1 illustrates how to determine the relative strength of intermolecular forces for a group of compounds. Table 3.4 summarizes the four types of interactions that affect the properties of all compounds.

Table 3.4 Summary of Types of Intermolecular Forces


Type of force	Relative strength	Exhibited by	Example
van der Waals	weak	all molecules	CH ₃ CH ₂ CH ₂ CH ₂ CH ₃ CH ₃ CH ₂ CH ₂ CHO CH ₃ CH ₂ CH ₂ CH ₂ OH
dipole–dipole	moderate	molecules with a net dipole	CH ₃ CH ₂ CH ₂ CHO CH ₃ CH ₂ CH ₂ CH ₂ OH
hydrogen bonding	strong	molecules with an O–H, N–H, or H–F bond	CH ₃ CH ₂ CH ₂ CH ₂ OH
ion–ion	very strong	ionic compounds	NaCl, LiF

Sample Problem 3.1



Rank the following compounds in order of increasing strength of intermolecular forces: CH₃CH₂CH₂CH₂CH₃ (pentane), CH₃CH₂CH₂CH₂OH (1-butanol), and CH₃CH₂CH₂CHO (butanal).

Solution

- Pentane has only nonpolar C–C and C–H bonds, so its molecules are held together by only **van der Waals** forces.
- 1-Butanol is a polar bent molecule, so it can have **dipole–dipole** interactions in addition to **van der Waals** forces. Because it has an O–H bond, 1-butanol molecules are held together by intermolecular **hydrogen bonds** as well.
- Butanal has a trigonal planar carbon with a polar C=O bond, so it exhibits **dipole–dipole** interactions in addition to **van der Waals** forces. There is *no* H atom bonded to O, so two butanal molecules *cannot* hydrogen bond to each other.



Problem 3.5 What types of intermolecular forces are present in each compound?

- | | | |
|--|--|---|
| a.  | c. (CH ₃ CH ₂) ₃ N | e. CH ₃ CH ₂ CH ₂ COOH |
| b.  | d. CH ₂ =CHCl | f. CH ₃ –C≡C–CH ₃ |

3.4 Physical Properties


The strength of a compound's intermolecular forces determines many of its physical properties, including its boiling point, melting point, and solubility.

3.4A Boiling Point (bp)


The boiling point of a compound is the temperature at which a liquid is converted to a gas. In boiling, energy is needed to overcome the attractive forces in the more ordered liquid state.

- The stronger the intermolecular forces, the higher the boiling point.

Because **ionic compounds** are held together by extremely strong interactions, they have **very high boiling points**. The boiling point of NaCl, for example, is 1413 °C. **With covalent molecules, the boiling point depends on the identity of the functional group.** For compounds of approximately the same molecular weight:




Recall from Sample Problem 3.1, for example, that the relative strength of the intermolecular forces increases from pentane to butanal to 1-butanol. The boiling points of these compounds increase in the same order.



Because surface area and polarizability affect the strength of intermolecular forces, they also affect the boiling point. For two compounds with similar functional groups:


- The larger the surface area, the higher the boiling point.
- The more polarizable the atoms, the higher the boiling point.

Examples of each phenomenon are illustrated in Figure 3.2. In comparing two ketones that differ in size, 3-pentanone has a higher boiling point than acetone because it has a greater molecular weight and larger surface area. In comparing two alkyl halides having the same number of carbon atoms, CH_3I has a higher boiling point than CH_3F because I is more polarizable than F.




- The stronger the intermolecular forces, the higher the melting point.
- Given the same functional group, the more symmetrical the compound, the higher the melting point.


Because **ionic compounds** are held together by extremely strong interactions, they have **very high melting points**. For example, the melting point of NaCl is 801 °C. With covalent molecules, the melting point once again depends on the identity of the functional group. For compounds of approximately the same molecular weight:




The trend in the melting points of pentane, butanal, and 1-butanol parallels the trend observed in their boiling points.



Symmetry also plays a role in determining the melting points of compounds having the same functional group and similar molecular weights, but very different shapes. A compact symmetrical molecule like neopentane packs well into a crystalline lattice whereas isopentane, which has a CH_3 group dangling from a four-carbon chain, does not. Thus, neopentane has a much higher melting point.



Problem 3.8 Predict which compound in each pair has the higher melting point.




Problem 3.9 Consider acetic acid ($\text{CH}_3\text{CO}_2\text{H}$) and its conjugate base, sodium acetate ($\text{CH}_3\text{CO}_2\text{Na}$). (a) What intermolecular forces are present in each compound? (b) Explain why the melting point of sodium acetate (324 °C) is considerably higher than the melting point of acetic acid (17 °C).

Problem 3.10 Why do you suppose that symmetry affects the melting point of a compound but not its boiling point?

3.4C Solubility

Solubility is the extent to which a compound, called the *solute*, dissolves in a liquid, called the *solvent*. In dissolving a compound, the energy needed to break up the interactions between the molecules or ions of the solute comes from new interactions between the solute and the solvent.

Quantitatively, a compound may be considered soluble when 3 g of solute dissolves in 100 mL of solvent.



Compounds dissolve in solvents having similar kinds of intermolecular forces.

- “Like dissolves like.”
- Polar compounds dissolve in polar solvents. Nonpolar or weakly polar compounds dissolve in nonpolar or weakly polar solvents.

Water and **organic liquids** are two different kinds of solvents. Water is very polar because it is capable of hydrogen bonding with a solute. Many organic solvents are either nonpolar, like carbon tetrachloride (CCl_4) and hexane [$\text{CH}_3(\text{CH}_2)_4\text{CH}_3$], or weakly polar like diethyl ether ($\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$).

Ionic compounds are held together by strong electrostatic forces, so they need very polar solvents to dissolve. **Most ionic compounds are soluble in water, but are insoluble in organic solvents.** To dissolve an ionic compound, the strong ion-ion interactions must be replaced by many weaker **ion-dipole interactions**, as illustrated in Figure 3.4.

Most organic compounds are soluble in organic solvents (remember, *like dissolves like*). **An organic compound is water soluble only if it contains one polar functional group capable of hydrogen bonding with the solvent for every five C atoms it contains.** In other words, a water-soluble organic compound has an O- or N-containing functional group that solubilizes its nonpolar carbon backbone.

Compare, for example, the solubility of butane and acetone in H_2O and CCl_4 .





Figure 3.4


Dissolving an ionic compound in H_2O



- When an ionic solid is dissolved in H_2O , the ion-ion interactions are replaced by ion-dipole interactions. Though these forces are weaker, there are so many of them that they compensate for the stronger ionic bonds.

Because butane and acetone are both organic compounds having a C–C and C–H backbone, they are soluble in the organic solvent CCl_4 . Butane, a nonpolar molecule, is insoluble in the polar solvent H_2O . Acetone, however, is H_2O soluble because it contains only three C atoms and its O atom can hydrogen bond with one H atom of H_2O . In fact, acetone is so soluble in water that acetone and water are **miscible**—they form solutions in all proportions with each other.


$(\text{CH}_3)_2\text{C}=\text{O}$ molecules cannot hydrogen bond to each other because they have no OH group. However, $(\text{CH}_3)_2\text{C}=\text{O}$ can hydrogen bond to H_2O because its O atom can hydrogen bond to one of the H atoms of H_2O .



Hydrogen bonding makes the small polar molecule acetone H_2O soluble.

For an organic compound with one functional group, **a compound is water soluble only if it has \leq five C atoms and contains an O or N atom.**

The size of an organic molecule with a polar functional group determines its water solubility. A low molecular weight alcohol like **ethanol is water soluble** because it has a small carbon skeleton (\leq five C atoms) compared to the size of its polar OH group. Cholesterol, on the other hand, has 27 carbon atoms and only one OH group. Its carbon skeleton is too large for the OH group to solubilize by hydrogen bonding, so **cholesterol is insoluble in water**.





Hydrophobic = afraid of H_2O ;
hydrophilic = H_2O loving.

- The nonpolar part of a molecule that is not attracted to H_2O is said to be *hydrophobic*.
- The polar part of a molecule that can hydrogen bond to H_2O is said to be *hydrophilic*.

In cholesterol, for example, the **hydroxy group is hydrophilic**, whereas the **carbon skeleton is hydrophobic**.

MTBE (*tert*-butyl methyl ether) and 4,4'-dichlorobiphenyl (a polychlorinated biphenyl, abbreviated as PCB) demonstrate that solubility properties can help determine the fate of organic compounds in the environment.



Using **MTBE** as a high-octane additive in unleaded gasoline has had a negative environmental impact. Although MTBE is not toxic or carcinogenic, it has a distinctive, nauseating odor, and **it is water soluble**. Small amounts of MTBE have contaminated the drinking water in several communities, making it unfit for consumption. For this reason, the use of MTBE as a gasoline additive has steadily declined in the United States since 1999.

4,4'-Dichlorobiphenyl is a polychlorinated biphenyl (**PCB**), a compound that contains two benzene rings joined by a C–C bond, and substituted by one or more chlorine atoms on each ring. PCBs

Figure 3.5
Solubility summary



Type of compound	Solubility in H ₂ O	Solubility in organic solvents (such as CCl ₄)
• Ionic NaCl	soluble	insoluble
• Covalent CH ₃ CH ₂ CH ₂ CH ₃	insoluble (no N or O atom to hydrogen bond to H ₂ O)	soluble
CH ₃ CH ₂ CH ₂ OH	soluble (≤ 5 C's and an O atom for hydrogen bonding to H ₂ O)	soluble
CH ₃ (CH ₂) ₁₀ OH	insoluble (> 5 C's; too large to be soluble even though it has an O atom for hydrogen bonding to H ₂ O)	soluble

have been used as plasticizers in polystyrene coffee cups and coolants in transformers. They have been released into the environment during production, use, storage, and disposal, making them one of the most widespread organic pollutants. **PCBs are insoluble in H₂O, but very soluble in organic media**, so they are soluble in fatty tissue, including that found in all types of fish and birds around the world. Although PCBs are not acutely toxic, frequently ingesting large quantities of fish contaminated with PCBs has been shown to retard growth and memory retention in children.

Solubility properties of some representative compounds are summarized in Figure 3.5.

Sample Problem 3.3

Which compounds are water soluble?



Solution

A has five C atoms and a polar C–Cl bond, but it is incapable of hydrogen bonding with H₂O, making it H₂O insoluble. **B** has five C atoms and a polar OH group able to hydrogen bond to H₂O, making it H₂O soluble.



Problem 3.11

Which compounds are water soluble?

- a. CH₃CH₂OCH₂CH₃ b. CH₃CH₂CH₂CH₂CH₃ c. (CH₃CH₂CH₂CH₂)₃N



Sample Problem 3.4

- a. Which of the following molecules can hydrogen bond to another molecule like itself?
b. Which of the following molecules can hydrogen bond with water?




Solution

- a. Compounds **A** and **B** have polar C–O and C–N bonds but **A** does not have an O–H bond, so it cannot hydrogen bond to another **A** molecule. **B**, on the other hand, has an N–H bond, so two **B** molecules can hydrogen bond with each other.
b. Because **A** has an electronegative O atom and **B** has an electronegative N atom, both **A** and **B** can hydrogen bond with a hydrogen atom in water.



Problem 3.12 Label the hydrophobic and hydrophilic portions of each molecule:



3.5 Application: Vitamins

Vitamins are organic compounds needed in small amounts for normal cell function. Our bodies cannot synthesize these compounds, so they must be obtained in the diet. Most vitamins are identified by a letter, such as A, C, D, E, and K. There are several different B vitamins, though, so a subscript is added to distinguish them: for example, B₁, B₂, and B₁₂.

Whether a vitamin is **fat soluble** (it dissolves in organic media) or **water soluble** can be determined by applying the solubility principles discussed in Section 3.4C. Vitamins A and C illustrate the differences between fat-soluble and water-soluble vitamins.


3.5A Vitamin A

The name **vitamin** was first used in 1912 by the Polish chemist Casimir Funk, who called them *vitamines*, because he thought that they all contained an *amine* functional group. Later the word was shortened to vitamin, because some are amines but others, like vitamins A and C, are not.



Vitamin A is synthesized from β -carotene, the orange pigment in carrots.


Vitamin A, or retinol, is an essential component of the vision receptors in the eyes. It also helps to maintain the health of mucous membranes and the skin, so many anti-aging creams contain vitamin A. A deficiency of this vitamin leads to a loss of night vision.



Vitamin A contains 20 carbons and a single OH group, making it **water insoluble**. Because it is organic, it is **soluble in any organic medium**. To understand the consequences of these solubility characteristics, we must learn about the chemical environment of the body.

About 70% of the body is composed of water. Fluids such as blood, gastric juices in the stomach, and urine are largely water with dissolved ions such as Na⁺ and K⁺. Vitamin A is insoluble in these fluids. There are also fat cells composed of organic compounds having C–C and C–H bonds. Vitamin A is soluble in this organic environment, and thus it is readily stored in these fat cells, particularly in the liver.

Vitamin A may be obtained directly from the diet. In addition, β -carotene, the orange pigment found in many plants including carrots, is readily converted to vitamin A in our bodies.




Eating too many carrots does not result in an excess of stored vitamin A. If you consume more β -carotene than you need, your body stores this precursor until it needs more vitamin A. Some β -carotene reaches the surface tissues of the skin and eyes, giving them an orange color. This phenomenon may look odd, but it is harmless and reversible. When stored β -carotene is converted to vitamin A and is no longer in excess, these tissues will return to their normal hue.

3.5B Vitamin C





Vitamin C is obtained by eating citrus fruits and a wide variety of other fruits and vegetables. Individuals can also obtain the recommended daily dose of vitamin C by taking tablets that contain vitamin C prepared in the laboratory. Both the “natural” vitamin C in oranges and the “synthetic” vitamin C in vitamin supplements are identical.



Vitamin C has six carbon atoms, each bonded to an oxygen atom that is capable of hydrogen bonding, making it **water soluble**. Vitamin C thus dissolves in urine. Although it has been acclaimed as a deterrent for all kinds of diseases, from the common cold to cancer, the consequences of taking large amounts of vitamin C are not really known, because any excess of the minimum daily requirement is excreted in the urine.

Problem 3.13

Predict the water solubility of each vitamin.



3.6 Application of Solubility: Soap

Soap has been used by humankind for some 2000 years. Historical records describe its manufacture in the first century and document the presence of a soap factory in Pompeii. Before this time clothes were cleaned by rubbing them on rocks in water, or by forming soapy lathers from the roots, bark, and leaves of certain plants. These plants produced natural materials called *saponins*, which act in much the same way as modern soaps.

Soap molecules have two distinct parts:

- a hydrophilic portion composed of ions called the *polar head*
- a hydrophobic carbon chain of nonpolar C–C and C–H bonds, called the *nonpolar tail*





Figure 3.6
Dissolving soap in water




- When soap is dissolved in H_2O , the molecules form micelles with the nonpolar tails in the interior and the polar heads on the surface. The polar heads are solvated by ion-dipole interactions with H_2O molecules.

Dissolving soap in water forms *micelles*, spherical droplets having the ionic heads on the surface and the nonpolar tails packed together in the interior, as shown in Figure 3.6. In this arrangement, the ionic heads are solvated by the polar solvent water, thus solubilizing the nonpolar, “greasy” hydrocarbon portion of the soap.

How does soap dissolve grease and oil? Water alone cannot dissolve dirt, which is composed largely of nonpolar hydrocarbons. When soap is mixed with water, however, the nonpolar hydrocarbon tails dissolve the dirt in the interior of the micelle. The polar head of the soap remains on the surface of the micelle to interact with water. The nonpolar tails of the soap molecules are so well sealed off from the water by the polar head groups that the micelles are water soluble, allowing them to separate from the fibers of our clothes and be washed down the drain with water. In this way, soaps do a seemingly impossible task: they remove nonpolar hydrocarbon material from skin and clothes, by solubilizing it in the polar solvent water.


Cross-section of a soap micelle with a grease particle dissolved in the interior



Problem 3.14 Which of the following structures represent soaps? Explain your answers.

- a. $\text{CH}_3\text{CO}_2^-\text{Na}^+$ b. $\text{CH}_3(\text{CH}_2)_{14}\text{CO}_2^-\text{Na}^+$ c. $\text{CH}_3(\text{CH}_2)_{12}\text{COOH}$ d. $\text{CH}_3(\text{CH}_2)_9\text{CO}_2^-\text{Na}^+$

Problem 3.15 Today, synthetic detergents like the compound drawn here, not soaps, are used to clean clothes. Explain how this detergent cleans away dirt.




3.7 Application: The Cell Membrane

The cell membrane is a beautifully complex example of how the principles of organic chemistry come into play in a biological system.

3.7A Structure of the Cell Membrane

The basic unit of living organisms is the **cell**. The cytoplasm is the aqueous medium inside the cell, separated from water outside the cell by the **cell membrane**. The cell membrane serves two apparently contradictory functions. It acts as a barrier to the passage of ions, water, and other molecules into and out of the cell, and it is also selectively permeable, letting nutrients in and waste out.

A major component of the cell membrane is a group of organic molecules called **phospholipids**. Like soap, they contain a hydrophilic ionic portion, and a hydrophobic hydrocarbon portion, in this case two long carbon chains composed of C–C and C–H bonds. **Phospholipids thus contain a polar head and two nonpolar tails.**



When phospholipids are mixed with water, they assemble in an arrangement called a **lipid bilayer**, with the ionic heads oriented on the outside and the nonpolar tails on the inside. The polar heads electrostatically interact with the polar solvent H_2O , while the nonpolar tails are held in close proximity by numerous van der Waals interactions. This is schematically illustrated in Figure 3.7.


Cell membranes are composed of these lipid bilayers. The charged heads of the phospholipids are oriented toward the aqueous interior and exterior of the cell. The nonpolar tails form the hydrophobic interior of the membrane, thus serving as an insoluble barrier that protects the cell from the outside.

3.7B Transport Across a Cell Membrane

How does a polar molecule or ion in the water outside a cell pass through the nonpolar interior of the cell membrane and enter the cell? Some nonpolar molecules like O_2 are small enough to enter and exit the cell by diffusion. Polar molecules and ions, on the other hand, may be too large or too polar to diffuse efficiently. Some ions are transported across the membrane with the help of molecules called **ionophores**.

Ionophores are organic molecules that complex cations. They have a hydrophobic exterior that makes them soluble in the nonpolar interior of the cell membrane, and a central cavity with


Figure 3.7
The cell membrane



- Phospholipids contain an ionic or polar head, and two long nonpolar hydrocarbon tails. In an aqueous environment, phospholipids form a lipid bilayer, with the polar heads oriented toward the aqueous exterior and the nonpolar tails forming a hydrophobic interior. Cell membranes are composed largely of this lipid bilayer.

several oxygen atoms whose lone pairs complex with a given ion. The size of the cavity determines the identity of the cation with which the ionophore complexes. Two naturally occurring antibiotics that act as ionophores are **nonactin** and **valinomycin**.

Naturally occurring antibiotic ionophores




Each molecule contains a large central cavity to hold a cation.

Several synthetic ionophores have also been prepared, including one group called **crown ethers**. **Crown ethers** are cyclic ethers containing several oxygen atoms that bind specific cations depending on the size of their cavity. Crown ethers are named according to the general format


Figure 3.8

Transport of ions across a cell membrane



- By binding an ion on one side of a lipid bilayer (where the concentration of the ion is high) and releasing it on the other side of the bilayer (where the concentration of the ion is low), an ionophore transports an ion across a cell membrane.

x-crown-y, where x is the total number of atoms in the ring and y is the number of oxygen atoms. For example, 18-crown-6 contains 18 atoms in the ring, including 6 O atoms. This crown ether binds potassium ions. Sodium ions are too small to form a tight complex with the O atoms, and larger cations do not fit in the cavity.



How does an ionophore transfer an ion across a membrane? The ionophore binds the ion on one side of the membrane in its polar interior. It can then move across the membrane because its hydrophobic exterior interacts with the hydrophobic tails of the phospholipid. The ionophore then releases the ion on the other side of the membrane. This ion-transfer role is essential for normal cell function. This process is illustrated in Figure 3.8.

In this manner, antibiotic ionophores like nonactin transport ions across a cell membrane of bacteria. This disrupts the normal ionic balance in the cell, thus interfering with cell function and causing the bacteria to die.

Problem 3.16 Nonactin and valinomycin each contain only two different types of functional groups. What two functional groups are present in nonactin? In valinomycin?

3.8 Functional Groups and Reactivity


Much of Chapter 3 has been devoted to how a functional group determines the strength of intermolecular forces and, consequently, the physical properties of molecules. A functional group also determines reactivity. What type of reaction does a particular kind of organic compound undergo? Begin by recalling two fundamental concepts.

- Functional groups create reactive sites in molecules.
- Electron-rich sites react with electron-poor sites.


All functional groups contain a heteroatom, a π bond, or both, and these features make electron-deficient (or electrophilic) sites and electron-rich (or nucleophilic) sites in a mol-

ecule. Molecules react at these sites. To predict reactivity, first locate the functional group and then determine the resulting electron-rich or electron-deficient sites it creates. Keep three guidelines in mind.


- An electronegative heteroatom like N, O, or X makes a carbon atom *electrophilic*.



- A lone pair on a heteroatom makes it basic and nucleophilic.




- π Bonds create *nucleophilic* sites and are more easily broken than σ bonds.



Problem 3.17

Label the electrophilic and nucleophilic sites in each molecule.



By identifying the nucleophilic and electrophilic sites in a compound you can begin to understand how it will react. In general, electron-rich sites react with electron-deficient sites:


$:\text{Nu}^-$ = a nucleophile;
 E^+ = an electrophile.

- An electron-deficient carbon atom reacts with a nucleophile, symbolized as $:\text{Nu}^-$.
- An electron-rich carbon reacts with an electrophile, symbolized as E^+ .


At this point we don't know enough organic chemistry to draw the products of many reactions with confidence. We do know enough, however, to begin to predict if two compounds might react together based solely on electron density arguments, and at what atoms that reaction is most likely to occur.

For example, alkenes contain an electron-rich C–C double bond and so they react with electrophiles, E^+ . On the other hand, alkyl halides possess an electrophilic carbon atom, so they react with electron-rich nucleophiles.

Alkenes react with electrophiles.




Alkyl halides react with nucleophiles.




For now, you don't need to worry about the products of these reactions. At this point you should only be able to find reactive sites in molecules and begin to understand why a reaction might occur at these sites. After you learn more about the structure of organic molecules in Chapters 4 and 5, we will begin a detailed discussion of organic reactions in Chapter 6.

Problem 3.18 Considering only electron density, state whether the following reactions will occur:




3.9 Biomolecules


Biomolecules are organic compounds found in biological systems. Many are relatively small, with molecular weights of less than 1000 g/mol. There are four main families of these small molecules—simple sugars, amino acids, lipids, and nucleotides. Many simple biomolecules are used to synthesize larger compounds that have important cellular functions.




glucose
a simple sugar



oleic acid
a fatty acid




alanine
an amino acid



deoxyadenosine 5'-monophosphate
a nucleotide

Figure 3.9

The double helix of DNA



- DNA, which is contained in the chromosomes of the nucleus of a cell, stores all of the genetic information in an organism. DNA consists of two long strands of polynucleotides held together by hydrogen bonding.


Simple sugars such as glucose combine to form the complex carbohydrates starch and cellulose, as described in Chapter 27. Alanine is an amino acid used to synthesize proteins, the subject of Chapter 28. Fatty acids such as oleic acid react with alcohols to form triacylglycerols, the most prevalent lipids, first mentioned in Chapter 10, and discussed in more detail in Chapters 22 and 29. While these biomolecules all contain more than one functional group, their properties and reactions are explained by the principles of basic organic chemistry.

Finally, deoxyadenosine 5'-monophosphate is a nucleotide that combines with thousands of other nucleotides to form DNA, deoxyribonucleic acid, the high molecular weight polynucleotide that stores the genetic information of an organism. DNA consists of two polynucleotide chains that wind together in a double helix. Figure 3.9 illustrates the importance of hydrogen bonding in the structure of DNA. The two polynucleotide chains are held together by an extensive network of hydrogen bonds in which the N–H groups on one chain intermolecularly hydrogen bond to an oxygen or nitrogen atom on the adjacent chain.










KEY CONCEPTS

Introduction to Organic Molecules and Functional Groups

Types of Intermolecular Forces (3.3)

	Type of force	Cause
 Increasing strength	van der Waals	Caused by the interaction of temporary dipoles <ul style="list-style-type: none"> • Larger surface area, stronger forces • Larger, more polarizable atoms, stronger forces
	dipole–dipole hydrogen bonding	Caused by the interaction of permanent dipoles Caused by the electrostatic interaction of a H atom in an O–H, N–H, or H–F bond with the lone pair of another N, O, or F atom
	ion–ion	Caused by the charge attraction of two ions

Physical Properties

Property	Observation														
Boiling point (3.4A)	<ul style="list-style-type: none"> • For compounds of comparable molecular weight, the stronger the intermolecular forces the higher the bp. <table style="margin-left: auto; margin-right: auto; text-align: center;"> <tr> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW bp = 36 °C</td> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD bp = 76 °C</td> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB bp = 118 °C</td> </tr> <tr> <td colspan="3" style="text-align: center;">  Increasing strength of intermolecular forces Increasing boiling point </td> </tr> </table> • For compounds with similar functional groups, the larger the surface area, the higher the bp. <table style="margin-left: auto; margin-right: auto; text-align: center;"> <tr> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 0 °C</td> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 36 °C</td> </tr> <tr> <td colspan="2" style="text-align: center;">  Increasing surface area Increasing boiling point </td> </tr> </table> • For compounds with similar functional groups, the more polarizable the atoms, the higher the bp. <table style="margin-left: auto; margin-right: auto; text-align: center;"> <tr> <td>CH_3F bp = –78 °C</td> <td>CH_3I bp = 42 °C</td> </tr> <tr> <td colspan="2" style="text-align: center;">  Increasing polarizability Increasing boiling point </td> </tr> </table> 	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW bp = 36 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD bp = 76 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB bp = 118 °C	 Increasing strength of intermolecular forces Increasing boiling point			$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 0 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 36 °C	 Increasing surface area Increasing boiling point		CH_3F bp = –78 °C	CH_3I bp = 42 °C	 Increasing polarizability Increasing boiling point	
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW bp = 36 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD bp = 76 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB bp = 118 °C													
 Increasing strength of intermolecular forces Increasing boiling point															
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 0 °C	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 36 °C														
 Increasing surface area Increasing boiling point															
CH_3F bp = –78 °C	CH_3I bp = 42 °C														
 Increasing polarizability Increasing boiling point															

Melting point (3.4B)	<ul style="list-style-type: none"> For compounds of comparable molecular weight, the stronger the intermolecular forces the higher the mp. <table style="width: 100%; text-align: center; margin: 10px 0;"> <tr> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW mp = $-130\text{ }^\circ\text{C}$</td> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD mp = $-96\text{ }^\circ\text{C}$</td> <td>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB mp = $-90\text{ }^\circ\text{C}$</td> </tr> </table> <p style="text-align: center; margin: 10px 0;"> Increasing strength of intermolecular forces Increasing melting point </p> <ul style="list-style-type: none"> For compounds with similar functional groups, the more symmetrical the compound, the higher the mp. <table style="width: 100%; text-align: center; margin: 10px 0;"> <tr> <td>$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ mp = $-160\text{ }^\circ\text{C}$</td> <td>$(\text{CH}_3)_4\text{C}$ mp = $-17\text{ }^\circ\text{C}$</td> </tr> </table> <p style="text-align: center; margin: 10px 0;"> Increasing symmetry Increasing melting point </p> 	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW mp = $-130\text{ }^\circ\text{C}$	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD mp = $-96\text{ }^\circ\text{C}$	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB mp = $-90\text{ }^\circ\text{C}$	$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ mp = $-160\text{ }^\circ\text{C}$	$(\text{CH}_3)_4\text{C}$ mp = $-17\text{ }^\circ\text{C}$
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW mp = $-130\text{ }^\circ\text{C}$	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD mp = $-96\text{ }^\circ\text{C}$	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB mp = $-90\text{ }^\circ\text{C}$				
$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ mp = $-160\text{ }^\circ\text{C}$	$(\text{CH}_3)_4\text{C}$ mp = $-17\text{ }^\circ\text{C}$					
Solubility (3.4C)	<p>Types of water-soluble compounds:</p> <ul style="list-style-type: none"> Ionic compounds Organic compounds having ≤ 5 C's, and an O or N atom for hydrogen bonding (for a compound with one functional group). <p>Types of compounds soluble in organic solvents:</p> <ul style="list-style-type: none"> Organic compounds regardless of size or functional group. 					

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding







Reactivity (3.8)

- Nucleophiles react with electrophiles.**
- Electronegative heteroatoms create electrophilic carbon atoms, which tend to react with nucleophiles.
- Lone pairs and π bonds are nucleophilic sites that tend to react with electrophiles.

PROBLEMS


Functional Groups

3.19 Identify the functional groups in each molecule.



3.20 Draw the seven constitutional isomers having molecular formula $\text{C}_4\text{H}_{10}\text{O}$. Identify the functional group in each isomer.


3.21 Identify each functional group located in the following rings. Which structure represents a lactone—a cyclic ester—and which represents a lactam—a cyclic amide?




3.22 Draw seven constitutional isomers with molecular formula $C_3H_6O_2$ that contain a carbonyl group. Identify the functional group(s) in each isomer.

Intermolecular Forces


3.23 What types of intermolecular forces are exhibited by each compound?




3.24 Rank the following compounds in order of increasing strength of intermolecular forces:



3.25 Carboxylic acids ($RCOOH$) can exist as dimers in some situations, with two molecules held together by two intermolecular hydrogen bonds. Show how two molecules of acetic acid, the carboxylic acid present in vinegar, can hydrogen bond to each other.





3.26 Intramolecular forces of attraction are often important in holding large molecules together. For example, some proteins fold into compact shapes, held together by attractive forces between nearby functional groups. A schematic of a folded protein is drawn here, with the protein backbone indicated by a blue-green ribbon, and various appendages drawn dangling from the chain. What types of intramolecular forces occur at each labeled site (A–F)?



Physical Properties


3.27 Rank the compounds in each group in order of increasing boiling point.

- a. $\text{CH}_3(\text{CH}_2)_4\text{I}$, $\text{CH}_3(\text{CH}_2)_5\text{I}$, $\text{CH}_3(\text{CH}_2)_6\text{I}$
 b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$, $(\text{CH}_3)_3\text{N}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$
 c. $(\text{CH}_3)_3\text{COC}(\text{CH}_3)_3$, $\text{CH}_3(\text{CH}_2)_3\text{O}(\text{CH}_2)_3\text{CH}_3$, $\text{CH}_3(\text{CH}_2)_7\text{OH}$




3.28 Explain why $\text{CH}_3\text{CH}_2\text{NHCH}_3$ has a higher boiling point than $(\text{CH}_3)_3\text{N}$, even though they have the same molecular weight.

3.29 Menthone and menthol are both isolated from mint. Explain why menthol is a solid at room temperature but menthone is a liquid.



3.30 Rank the compounds in each group in order of increasing melting point.


- a. $(\text{CH}_3)_2\text{CHOH}$, $(\text{CH}_3)_3\text{CH}$, $(\text{CH}_3)_2\text{C}=\text{O}$
 b. CH_3F , CH_3Cl , CH_3I



3.31 Explain the observed trend in the melting points for four isomers of molecular formula C_7H_{16} .


	mp (°C)
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$	-119
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$	-118
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	-91
$(\text{CH}_3)_2\text{C}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$	-25

3.32 Explain why benzene has a lower boiling point but much higher melting point than toluene.



3.33 Rank the compounds in each group in order of increasing water solubility.

- a. $(\text{CH}_3)_3\text{CH}$, $\text{CH}_3\text{OCH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$




3.34 Which of the following molecules can hydrogen bond to another molecule of itself? Which can hydrogen bond with water?

- a. CBr_3CH_3 c. CH_3OCH_3 e. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CONH}_2$ g. CH_3SOCH_3
 b. CH_3NH_2 d. $(\text{CH}_3\text{CH}_2)_3\text{N}$ f. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ h. $\text{CH}_3\text{CH}_2\text{COOCH}_3$

3.35 Explain why diethyl ether ($\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$) and 1-butanol ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$) have similar solubility properties in water, but 1-butanol has a much higher boiling point.

3.36 Predict the water solubility of each of the following organic molecules:




3.37 A mixture of four alkanes (**A–D**) was distilled and four fractions (1–4, in order of increasing boiling point) were collected. Which alkane was the principal component in each fraction?




Applications


3.38 Predict the solubility of each of the following vitamins in water and in organic solvents:





3.39 Avobenzene and dioxybenzone are two commercial sunscreens. Using the principles of solubility, predict which sunscreen is more readily washed off when an individual goes swimming. Explain your choice.





3.40 Poly(ethylene glycol) (PEG) and poly(vinyl chloride) (PVC) are examples of polymers, large organic molecules composed of repeating smaller units covalently bonded together. Polymers have very different properties depending (in part) on their functional groups. Discuss the water solubility of each polymer and suggest why PEG is used in shampoos while PVC is used to make garden hoses and pipes. Synthetic polymers are discussed in detail in Chapters 15 and 30.





- 3.41 THC is the active component in marijuana, and ethanol is the alcohol in alcoholic beverages. Explain why drug screenings are able to detect the presence of THC but not ethanol weeks after these substances have been introduced into the body.



- 3.42 Cocaine is a widely abused, addictive drug. Cocaine is usually obtained as its hydrochloride salt (cocaine hydrochloride) but can be converted to crack (the neutral molecule) by treatment with base. Which of the two compounds here has a higher boiling point? Which is more soluble in water? How does the relative solubility explain why crack is usually smoked but cocaine hydrochloride is injected directly into the bloodstream?




- 3.43 Unlike soap, which is ionic, some liquid laundry detergents are neutral molecules. Explain how each of the following molecules behaves like soap and cleans away dirt.



- 3.44 Most mayonnaise recipes call for oil (nonpolar, mostly hydrocarbon), vinegar (H_2O and CH_3COOH), and egg yolk. The last ingredient is a source of phospholipids that act as emulsifying agents. Explain.






- 3.45 Many drugs are sold as their hydrochloride salts ($\text{R}_2\text{NH}_2^+ \text{Cl}^-$), formed by reaction of an amine (R_2NH) with HCl .




- Draw the product (a hydrochloride salt) formed by reaction of acebutolol with HCl . Acebutolol is a β blocker used to treat high blood pressure.
- Discuss the solubility of acebutolol and its hydrochloride salt in water.
- Offer a reason as to why the drug is marketed as a hydrochloride salt rather than a neutral amine.

Reactivity of Organic Molecules

- 3.46 Label the electrophilic and nucleophilic sites in each molecule.




3.47 By using only electron density arguments, determine whether the following reactions will occur:



Cell Membrane

3.48 The composition of a cell membrane is not uniform for all types of cells. Some cell membranes are more rigid than others. Rigidity is determined by a variety of factors, one of which is the structure of the carbon chains in the phospholipids that comprise the membrane. One example of a phospholipid was drawn in Section 3.7A, and another, having C–C double bonds in its carbon chains, is drawn here. Which phospholipid would be present in the more rigid cell membrane and why?




a phospholipid with π bonds in the long hydrocarbon chains

General Question


3.49 Vancomycin is an especially useful antibiotic for treating infections in cancer patients on chemotherapy and renal patients on dialysis. Unlike mammalian cells, bacterial cells are surrounded by a fairly rigid cell wall, which is crucial to the bacterium's survival. Vancomycin kills bacteria by interfering with their cell wall synthesis.

- How many amide functional groups are present in vancomycin?
- Which OH groups are bonded to sp^3 hybridized carbon atoms and which are bonded to sp^2 hybridized carbons?
- Would you expect vancomycin to be water soluble? Explain.
- Which proton is the most acidic?
- Label three *different* functional groups capable of hydrogen bonding.





Challenge Problems

3.50 Explain why **A** is less water soluble than **B**, even though both compounds have the same functional groups.



3.51 Recall from Section 1.9B that there is restricted rotation around carbon-carbon double bonds. Maleic acid and fumaric acid are two isomers with vastly different physical properties and pK_a values for loss of both protons. Explain why each of these differences occurs.



mp (°C)	130	286
solubility (g/L) in H ₂ O at 25 °C	788	7
pK_{a1}	1.9	3.0
pK_{a2}	6.5	4.5

Alkanes

4

- 4.1 Alkanes—An introduction
- 4.2 Cycloalkanes
- 4.3 An introduction to nomenclature
- 4.4 Naming alkanes
- 4.5 Naming cycloalkanes
- 4.6 Common names
- 4.7 Fossil fuels
- 4.8 Physical properties of alkanes
- 4.9 Conformations of acyclic alkanes—Ethane
- 4.10 Conformations of butane
- 4.11 An introduction to cycloalkanes
- 4.12 Cyclohexane
- 4.13 Substituted cycloalkanes
- 4.14 Oxidation of alkanes
- 4.15 Lipids—Part 1



Alkanes, the simplest hydrocarbons, are found in all shapes and sizes and occur widely in nature. They are the major constituents of petroleum, a complex mixture of compounds that includes hydrocarbons such as **hexane** and **decane**. Crude petroleum spilled into the sea from a ruptured oil tanker creates an insoluble oil slick on the surface. Petroleum is refined to produce gasoline, diesel fuel, home heating oil, and a myriad of other useful compounds. In Chapter 4, we learn about the properties of alkanes, how to name them (nomenclature), and oxidation—one of their important reactions.

In Chapter 4, we apply the principles of bonding, shape, and reactivity discussed in Chapters 1–3 to our first family of organic compounds, the **alkanes**. Because alkanes have no functional group, they are much less reactive than other organic compounds, and for this reason, much of Chapter 4 is devoted to learning how to name and draw them, as well as to understanding what happens when rotation occurs about their carbon–carbon single bonds.

Studying alkanes also provides an opportunity to learn about **lipids**, a group of biomolecules similar to alkanes, in that they are composed mainly of nonpolar carbon–carbon and carbon–hydrogen σ bonds. Section 4.15 serves as a brief introduction only, so we will return to lipids in Chapters 10 and 29.

4.1 Alkanes—An Introduction




Secretion of **undecane** by a cockroach causes other members of the species to aggregate. Undecane is a **pheromone**, a chemical substance used for communication in an animal species, most commonly an insect population. Minute amounts elicit an activity such as mating, aggregation, or defense.

Recall from Section 3.2 that **alkanes are aliphatic hydrocarbons having only C–C and C–H σ bonds**. Because their carbon atoms can be joined together in chains or rings, they can be categorized as acyclic or cyclic.

- **Acyclic alkanes** have the molecular formula C_nH_{2n+2} (where n = an integer) and contain only linear and branched chains of carbon atoms. They are also called **saturated hydrocarbons** because they have the maximum number of hydrogen atoms per carbon.
- **Cycloalkanes** contain carbons joined in one or more rings. Because their general formula is C_nH_{2n} , they have two fewer H atoms than an acyclic alkane with the same number of carbons.

Undecane and cyclohexane are two naturally occurring alkanes.




4.1A Acyclic Alkanes Having One to Five C Atoms




Cyclohexane is one component of the mango, the most widely consumed fruit in the world.

Structures for the two simplest acyclic alkanes were given in Chapter 1. **Methane**, CH_4 , has a single carbon atom, and **ethane**, CH_3CH_3 , has two. All C atoms in an alkane are surrounded by four groups, making them sp^3 hybridized and **tetrahedral**, and all bond angles are 109.5° .



The three-carbon alkane $CH_3CH_2CH_3$, called **propane**, has molecular formula C_3H_8 . Each carbon in the three-dimensional drawing has two bonds in the plane (solid lines), one bond in front (on a wedge), and one bond behind the plane (on a dashed line).

To draw the structure of an alkane, join the carbon atoms together with single bonds, and add enough H atoms to make each C tetravalent.



Problem 4.1

Three components of the sex pheromone of the female sand bee (*Ophrys sphegodes*) are saturated hydrocarbons containing 23, 25, and 27 carbon atoms. How many H atoms does each of these alkanes contain? Interestingly, the early spider orchid emits a similar hydrocarbon mixture to attract male sand bees to pollinate its flowers.

The molecular formulas for methane, ethane, and propane fit into the general molecular formula for an alkane, $\text{C}_n\text{H}_{2n+2}$:

- Methane = $\text{CH}_4 = \text{C}_1\text{H}_{2(1)+2}$
- Ethane = $\text{C}_2\text{H}_6 = \text{C}_2\text{H}_{2(2)+2}$
- Propane = $\text{C}_3\text{H}_8 = \text{C}_3\text{H}_{2(3)+2}$


The three-dimensional representations and the ball-and-stick models for these alkanes indicate the tetrahedral geometry around each carbon atom. In contrast, **the Lewis structures are not meant to imply any three-dimensional arrangement.** Moreover, in propane and higher molecular weight alkanes, the carbon skeleton can be drawn in a variety of different ways and still represent the same molecule.



For example, the three carbons of propane can be drawn in a horizontal row or with a bend. **These representations are equivalent.** If you follow the carbon chain from one end to the other, you move across the *same* three carbon atoms in both representations.

- The bends in a carbon chain don't matter when it comes to identifying different compounds.

There are two different ways to arrange four carbons, giving two compounds with molecular formula C_4H_{10} , named **butane** and **isobutane**.



Constitutional isomers like butane and isobutane belong to the same family of compounds: they are both **alkanes**. In contrast, constitutional isomers like $\text{CH}_3\text{CH}_2\text{OH}$ and CH_3OCH_3 have different functional groups and belong to different families: $\text{CH}_3\text{CH}_2\text{OH}$ is an **alcohol** and CH_3OCH_3 is an **ether**.


Butane and isobutane are **isomers**, two different compounds with the same molecular formula (Section 1.4A). They belong to one of the two major classes of isomers called **constitutional** or **structural isomers**. The two isomers discussed in Section 1.4A, CH_3OCH_3 and $\text{CH}_3\text{CH}_2\text{OH}$, are also constitutional isomers. We will learn about the second major class of isomers, called **stereoisomers**, in Section 4.13B.

- **Constitutional isomers** differ in the way the atoms are connected to each other.


Butane, which has four carbons in a row, is a **straight-chain** or **normal alkane** (an *n*-alkane). Isobutane, on the other hand, is a **branched-chain alkane**.

With alkanes having more than four carbons, the names of the straight-chain isomers are systematic and derive from Greek roots: *pentane* for five C atoms, *hexane* for six, and so on. There are three constitutional isomers for the five-carbon alkane, each having molecular formula C_5H_{12} : **pentane**, **isopentane** (or 2-methylbutane), and **neopentane** (or 2,2-dimethylpropane).

Three constitutional isomers having molecular formula C_5H_{12}




Problem 4.2 Which of the following is *not* another representation for isopentane?



Carbon atoms in alkanes and other organic compounds are classified by the number of other carbons directly bonded to them.


- A *primary carbon* (1° carbon) is bonded to *one* other C atom.
- A *secondary carbon* (2° carbon) is bonded to *two* other C atoms.
- A *tertiary carbon* (3° carbon) is bonded to *three* other C atoms.
- A *quaternary carbon* (4° carbon) is bonded to *four* other C atoms.




Hydrogen atoms are classified as **primary** (1°), **secondary** (2°), or **tertiary** (3°) depending on the **type of carbon atom** to which they are bonded.

- A *primary hydrogen* (1° H) is on a C bonded to one other C atom.
- A *secondary hydrogen* (2° H) is on a C bonded to two other C atoms.
- A *tertiary hydrogen* (3° H) is on a C bonded to three other C atoms.

Classification of hydrogen atoms




Example




Sample Problem 4.1

(a) Classify the designated carbon atoms as 1° , 2° , 3° , or 4° . (b) Classify the designated hydrogen atoms as 1° , 2° , or 3° .




Solution




Problem 4.3

(a) Classify the carbon atoms in each compound as 1° , 2° , 3° , or 4° . (b) Classify the hydrogen atoms in each compound as 1° , 2° , or 3° .



Problem 4.4

Classifying a carbon atom by the number of carbons to which it is bonded can also be done in more complex molecules that contain heteroatoms. Classify each tetrahedral carbon atom in the female sex hormone estrone as 1° , 2° , 3° , or 4° .



4.1B Acyclic Alkanes Having More Than Five C Atoms

The maximum number of possible constitutional isomers increases dramatically as the number of carbon atoms in the alkane increases, as shown in Table 4.1. For example, there are 75 possible isomers for an alkane having 10 carbon atoms, but 366,319 possible isomers for one having 20 carbons.

Each entry in Table 4.1 is formed from the preceding entry by adding a CH_2 group. A CH_2 group is called a *methylene group*. A group of compounds that differ by only a CH_2 group is called a *homologous series*. The names of all alkanes end in the suffix *-ane*, and the syllable preceding the suffix identifies the number of carbon atoms in the chain.

The suffix *-ane* identifies a molecule as an *alkane*.

Problem 4.5

Draw the five constitutional isomers having molecular formula C_6H_{14} .




Table 4.1 Summary: Straight-Chain Alkanes

Number of C atoms	Molecular formula	Name (<i>n</i> -alkane)	Number of constitutional isomers
1	CH ₄	methane	—
2	C ₂ H ₆	ethane	—
3	C ₃ H ₈	propane	—
4	C ₄ H ₁₀	butane	2
5	C ₅ H ₁₂	pentane	3
6	C ₆ H ₁₄	hexane	5
7	C ₇ H ₁₆	heptane	9
8	C ₈ H ₁₈	octane	18
9	C ₉ H ₂₀	nonane	35
10	C ₁₀ H ₂₂	decane	75
20	C ₂₀ H ₄₂	eicosane	366,319

Problem 4.6 Draw all constitutional isomers having molecular formula C₈H₁₈ that contain seven carbons in the longest chain and a single CH₃ group bonded to the chain.

Problem 4.7 Draw the structure of an alkane with molecular formula C₇H₁₆ that contains (a) one 4° carbon; (b) only 1° and 2° carbons; (c) 1°, 2°, and 3° hydrogens.



Problem 4.8 Considering compounds **A–C**, which two structures represent the same compound?

**A****B****C**

4.2 Cycloalkanes

Cycloalkanes have molecular formula C_nH_{2n} and contain carbon atoms arranged in a ring. Think of a cycloalkane as being formed by removing two H atoms from the end carbons of a chain, and then bonding the two carbons together. Simple cycloalkanes are named by adding the prefix **cyclo-** to the name of the acyclic alkane having the same number of carbons.

Cycloalkanes having three to six carbon atoms are shown in the accompanying figure. They are most often drawn in skeletal representations.

cyclopropane
C₃H₆cyclobutane
C₄H₈cyclopentane
C₅H₁₀cyclohexane
C₆H₁₂

Problem 4.9 Draw the five constitutional isomers that have molecular formula C_5H_{10} and contain one ring.

4.3 An Introduction to Nomenclature




Garlic has been a valued commodity throughout history. It has been used in Chinese herbal medicine for more than 4000 years, as a form of currency in Siberia, and as a repellent for witches by the Saxons. Today it is used as a dietary supplement because of its reported health benefits. **Allicin**, the molecule responsible for garlic's odor, is a rather unstable molecule that is not stored in the garlic bulb, but rather is produced by the action of enzymes when the bulb is crushed or bruised.


How are organic compounds named? Long ago, the name of a compound was often based on the plant or animal source from which it was obtained. For example, the name for **formic acid**, a caustic compound isolated from certain ants, comes from the Latin word *formica*, meaning *ant*; and **allicin**, the pungent principle of garlic, is derived from the botanical name for garlic, *Allium sativum*. Other compounds were named by their discoverer for more personal reasons. Adolf von Baeyer supposedly named barbituric acid after a woman named Barbara, although speculation continues on Barbara's identity—a lover, a Munich waitress, or even St. Barbara.



formic acid
(obtained from certain ants)



allicin
(odor of garlic)



barbituric acid
(named for Barbara whom?)


With the isolation and preparation of thousands of new organic compounds it became clear that each organic compound must have an unambiguous name, derived from a set of easily remembered rules. A systematic method of naming compounds was developed by the *International Union of Pure and Applied Chemistry*. It is referred to as the **IUPAC system of nomenclature**; how it can be used to name alkanes is explained in Sections 4.4 and 4.5.

The IUPAC system of nomenclature has been regularly revised since it was first adopted in 1892. Revisions in 1979 and 1993 and recent extensive recommendations in 2004 have given chemists a variety of acceptable names for compounds. Many changes are minor. For example, the 1979 nomenclature rules assign the name 1-butene to $CH_2=CHCH_2CH_3$, while the 1993 rules assign the name but-1-ene; that is, only the position of the number differs. In this text, the most generally used IUPAC conventions will be given, and often a margin note will be added to mention the differences between past and recent recommendations.

Naming organic compounds has become big business for drug companies. The IUPAC name of an organic compound can be long and complex, and may be comprehensible only to a chemist. As a result, most drugs have three names:

- **Systematic:** The systematic name follows the accepted rules of nomenclature and indicates the compound's chemical structure; this is the IUPAC name.
- **Generic:** The generic name is the official, internationally approved name for the drug.
- **Trade:** The trade name for a drug is assigned by the company that manufactures it. Trade names are often "catchy" and easy to remember. Companies hope that the public will continue to purchase a drug with an easily recalled trade name long after a cheaper generic version becomes available.


In the world of over-the-counter anti-inflammatory agents, the compound a chemist calls 2-[4-(2-methylpropyl)phenyl]propanoic acid has the generic name ibuprofen. It is marketed under a variety of trade names including Motrin and Advil.




Systematic name: 2-[4-(2-methylpropyl)phenyl]propanoic acid
Generic name: ibuprofen
Trade name: Motrin or Advil

The prefix *sec-* is short for *secondary*. A *sec*-butyl group is formed by removal of a **2° H**. The prefix *tert-* is short for *tertiary*. A *tert*-butyl group is formed by removal of a **3° H**.

From **butane**:



From **isobutane**:



Abbreviations are sometimes used for certain common alkyl groups.

Alkyl group (abbreviation)

- methyl (Me)
- ethyl (Et)
- butyl (Bu)
- *tert*-butyl (*t*-Bu)

The names isopropyl, *sec*-butyl, isobutyl, and *tert*-butyl are recognized as acceptable substituent names in both the 1979 and 1993 revisions of IUPAC nomenclature. A general method to name these substituents, as well as alkyl groups that contain five or more carbon atoms, is described in Appendix B.


4.4B Naming an Acyclic Alkane

Four steps are needed to name an alkane. In the following examples, only the C atoms of the carbon skeleton are drawn. **Remember each C has enough H atoms to make it tetravalent.**


HOW TO Name an Alkane Using the IUPAC System

Step [1] Find the parent carbon chain and add the suffix.

- Find the longest continuous carbon chain, and name the molecule by using the parent name for that number of carbons, given in Table 4.1. To the name of the parent, add the suffix **-ane** for an alkane. Each functional group has its own characteristic suffix.



- Finding the longest chain is just a matter of trial and error. Place your pencil on one end of the chain, go to the other end without picking it up, and count carbons. Repeat this procedure until you have found the chain with the largest number of carbons.
- It does not matter if the chain is *straight* or has *bends*. All of the following representations are equivalent.




—Continued

HOW TO, continued . . .

- If there are two chains of equal length, pick the chain with more substituents. In the following example, two different chains in the same alkane contain 7 C's, but the compound on the left has two alkyl groups attached to its long chain, whereas the compound to the right has only one.


Example: Having two different longest chains of the same length



7 atoms in the longest chain
2 substituents

more substituents

Correct





7 atoms in the longest chain
only 1 substituent

fewer substituents

Incorrect



Step [2] Number the atoms in the carbon chain.

- Number the longest chain to give the *first* substituent the lower number.





- If the first substituent is the same distance from both ends, number the chain to give the *second* substituent the lower number. Always look for the first point of difference in numbering from each end of the longest chain.

Example: Giving a lower number to the second substituent



- When numbering a carbon chain results in the same numbers from either end of the chain, assign the lower number *alphabetically* to the first substituent.


Example: Two different groups equidistant from the ends



HOW TO, continued . . .

Step [3] Name and number the substituents.

- Name the substituents as alkyl groups, and use the numbers from Step 2 to designate their location.




- Every carbon belongs to *either* the longest chain or a substituent, but *not both*.
- Each substituent needs its *own* number.
- If two or more identical substituents are bonded to the longest chain, use prefixes to indicate how many: **di-** for two groups, **tri-** for three groups, **tetra-** for four groups, and so forth. The preceding molecule has two methyl substituents, and so its name contains the prefix **di-** before the word methyl → *dimethyl*.


Step [4] Combine substituent names and numbers + parent + suffix.

- Precede the name of the parent by the names of the substituents.
- Alphabetize the names of the substituents, ignoring all prefixes except *iso*, as in isopropyl and isobutyl.
- Precede the name of each substituent by the number that indicates its location. There must be **one number for each substituent**.
- Separate numbers by commas and separate numbers from letters by hyphens. The name of an alkane is a single word, with no spaces after hyphens or commas.

[1] Identify all the pieces of a compound, using Steps 1–3.




[2] Then, put the pieces of the name together.




Several additional examples of alkane nomenclature are given in Figure 4.1.

Figure 4.1
Examples of alkane
nomenclature




2,3-dimethylpentane

[Number to give the 1st methyl group the lower number.]




4-ethyl-5-methyloctane

[Assign the lower number to the 1st substituent alphabetically: the **e** of **ethyl** before the **m** of **methyl**.]



4-ethyl-3,4-dimethyloctane

[Alphabetize the **e** of **ethyl** before the **m** of **methyl**.]




2,3,5-trimethyl-4-propylheptane

[Pick the long chain with more substituents.]

- The carbon atoms of each long chain are **drawn in red**.

Sample Problem 4.2 Give the IUPAC name for the following compound.




Solution

To help identify which carbons belong to the longest chain and which are substituents, always draw a box around the atoms of the long chain. Every other carbon atom then becomes a substituent that needs its own name as an alkyl group.


[H's on C's are omitted in the answer, for clarity.]

Step 1: Name the parent.




9 C's in the longest chain ----> **nonane**

Step 3: Name and number the substituents.



Step 2: Number the chain.







Step 4: Combine the parts.






- Alphabetize: the **b** of **butyl** before the **m** of **methyl**

Answer: 5-tert-butyl-3-methylnonane

Problem 4.10 Give the IUPAC name for each compound.



Problem 4.11 Give the IUPAC name for each compound.



You must also know how to derive a structure from a given name. Sample Problem 4.3 illustrates a stepwise method.

Sample Problem 4.3 Give the structure corresponding to the following IUPAC name: 6-isopropyl-3,3,7-trimethyldecane.

Solution

Follow three steps to derive a structure from a name.


Step [1] Identify the parent name and functional group found at the *end* of the name.



HOW TO, continued . . .

- For rings with more than one substituent, begin numbering at one substituent and proceed around the ring clockwise or counterclockwise to give the second substituent the lower number.


numbering clockwise



CH₃ groups at C1 and C3
The 2nd substituent has a lower number.

Correct: 1,3-dimethylcyclohexane

numbering counterclockwise




CH₃ groups at C1 and C5

Incorrect: 1,5-dimethylcyclohexane

- With two different substituents, number the ring to assign the lower number to the substituents *alphabetically*.

Begin numbering at the ethyl group.




- ethyl group at C1
- methyl group at C3

earlier letter → lower number

Correct: 1-ethyl-3-methylcyclohexane

Begin numbering at the methyl group.



- methyl group at C1
- ethyl group at C3

Incorrect: 3-ethyl-1-methylcyclohexane

When an alkane is composed of both a ring and a long chain, what determines whether a compound is named as an acyclic alkane or a cycloalkane? If the number of carbons in the ring is greater than or equal to the number of carbons in the longest chain, the compound is named as a **cycloalkane**, as shown in Figure 4.2. Several examples of cycloalkane nomenclature are given in Figure 4.3.

Problem 4.14 Give the IUPAC name for each compound.










Figure 4.2

Two contrasting examples—
Naming compounds containing both a ring and a long chain of carbon atoms

More carbons in the ring

4 C's in the chain —
a **butyl** group




6 C's in the ring—**cyclohexane**

Name as a **cyclohexane** with a substituent.

Answer: butylcyclohexane

More carbons in the chain

6 C's in the chain —
a **hexane**




4 C's in the ring—a **cyclobutyl** group

Name as a **hexane** with a substituent.


Answer: 1-cyclobutylhexane

Figure 4.3
Examples of cycloalkane
nomenclature




ethylcyclobutane

[No number is needed
with only one substituent.]




1-sec-butyl-3-methylcyclohexane

[Assign the lower number to the 1st substituent
alphabetically: the **b** of butyl before the **m** of methyl.]



1,2-dimethylcyclohexane

[Number to give the 2nd CH₃ group
the lower number: 1,2- not 1,6-.]



1,2,4-triethylcyclopentane

[Number to give the 2nd CH₃CH₂ group the
lower number: 1,2,4- not 1,3,4- or 1,3,5-.]


Problem 4.15 Give the structure corresponding to each IUPAC name.

- 1,2-dimethylcyclobutane
- 1,1,2-trimethylcyclopropane
- 4-ethyl-1,2-dimethylcyclohexane
- 1-sec-butyl-3-isopropylcyclopentane
- 1,1,2,3,4-pentamethylcycloheptane


Problem 4.16 Give the IUPAC names for the five cyclic isomers of molecular formula C₅H₁₀ in Problem 4.9.

4.6 Common Names

Some organic compounds are identified using **common names** that do not follow the IUPAC system of nomenclature. Many of these names were given to molecules long ago, before the IUPAC system was adopted. These names are still widely used. For example, isopentane, an older name for one of the C₅H₁₂ isomers, is still allowed by IUPAC rules, although it can also be named 2-methylbutane. We will follow the IUPAC system except in cases in which a common name is widely accepted.



isopentane or 2-methylbutane




dodecahedrane

In the past several years organic chemists have attempted to synthesize some unusual cycloalkanes not found in nature. **Dodecahedrane**, a beautifully symmetrical compound composed of 12 five-membered rings, is one such molecule. It was first prepared at The Ohio State University in 1982. The IUPAC name for dodecahedrane is undecacyclo[9.9.9.0^{2,9}.0^{3,7}.0^{4,20}.0^{3,15}.0^{6,16}.0^{8,15}.0^{10,14}.0^{12,19}.0^{13,17}]icicosane, a name so complex that few trained organic chemists would be able to identify its structure.

Because these systematic names are so unwieldy, organic chemists often assign a name to a polycyclic compound that is more descriptive of its shape and structure. Dodecahedrane is named because its 12 five-membered rings resemble a dodecahedron. Figure 4.4 shows the names and structures of several other cycloalkanes whose names were inspired by the shape of their carbon skeletons. All the names end in the suffix **-ane**, indicating that they refer to alkanes.

Figure 4.4

Common names for some polycyclic alkanes



For a more comprehensive list of unusual polycyclic alkanes (including windowpane, davidane, catenane, propellane, and many others), see *Organic Chemistry: The Name Game* by Alex Nickon and Ernest Silversmith, Pergamon Press, 1987.

4.7 Fossil Fuels

Natural gas is odorless. The smell observed in a gas leak is due to minute amounts of a sulfur additive such as methanethiol, CH_3SH , which provides an odor for easy detection.

Methane is formed and used in a variety of ways. The CH_4 released from decaying vegetable matter in New York City's main landfill is used for heating homes. CH_4 generators in China convert cow manure into energy in rural farming towns.


Many alkanes occur in nature, primarily in natural gas and petroleum. Both of these fossil fuels serve as energy sources, formed from the degradation of organic material long ago.

Natural gas is composed largely of **methane** (60% to 80% depending on its source), with lesser amounts of ethane, propane, and butane. These organic compounds burn in the presence of oxygen, releasing energy for cooking and heating.

Petroleum is a complex mixture of compounds, most of which are hydrocarbons containing 1–40 carbon atoms. Distilling crude petroleum, a process called **refining**, separates it into usable fractions that differ in boiling point (Figure 4.5). Most products of petroleum refining provide fuel for home heating, automobiles, diesel engines, and airplanes. Each fuel type has a different composition of hydrocarbons:

- **gasoline:** C_5H_{12} – $\text{C}_{12}\text{H}_{26}$
- **kerosene:** $\text{C}_{12}\text{H}_{26}$ – $\text{C}_{16}\text{H}_{34}$
- **diesel fuel:** $\text{C}_{15}\text{H}_{32}$ – $\text{C}_{18}\text{H}_{38}$

Petroleum provides more than fuel. About 3% of crude oil is used to make plastics and other synthetic compounds including drugs, fabrics, dyes, and pesticides. These products are responsible for many of the comforts we now take for granted in industrialized countries. Imagine what life would be like without air conditioning, refrigeration, anesthetics, and pain relievers, all products of the petroleum industry. Consider college students living without CDs and spandex!




products made from petroleum

Figure 4.5

Refining crude petroleum into usable fuel and other petroleum products



a. **An oil refinery.** At an oil refinery, crude petroleum is separated into fractions of similar boiling point by the process of **distillation**.



b. **Schematic of a refinery tower.** As crude petroleum is heated, the lower boiling, more volatile components distill first, followed by fractions of progressively higher boiling point.

Energy from petroleum is *nonrenewable*, and the remaining known oil reserves are limited. Given our dependence on petroleum, not only for fuel, but also for the many necessities of modern society, it becomes clear that we must both conserve what we have and find alternate energy sources.

4.8 Physical Properties of Alkanes


Alkanes contain only nonpolar C–C and C–H bonds, and as a result they exhibit only **weak van der Waals forces**. Table 4.2 summarizes how these intermolecular forces affect the physical properties of alkanes.

The gasoline industry exploits the dependence of boiling point and melting point on alkane size by seasonally changing the composition of gasoline in locations where it gets very hot in the summer and very cold in the winter. Gasoline is refined to contain a larger fraction of higher boiling hydrocarbons in warmer weather, so it evaporates less readily. In colder weather, it is refined to contain more lower boiling hydrocarbons, so it freezes less readily.

The mutual insolubility of nonpolar oil and very polar water leads to the common expression, “Oil and water don’t mix.”

Because nonpolar alkanes are not water soluble, crude petroleum that leaks into the sea from an oil tanker creates an insoluble oil slick on the surface. The insoluble hydrocarbon oil poses a special threat to birds whose feathers are coated with natural nonpolar oils for insulation. Because these hydrophobic oils dissolve in the crude petroleum, birds lose their layer of natural protection and many die.

Problem 4.17 Arrange the following compounds in order of increasing boiling point.



4.9 Conformations of Acyclic Alkanes—Ethane


Let’s now take a closer look at the three-dimensional structure of alkanes. The three-dimensional structure of molecules is called **stereochemistry**. In Chapter 4 we examine the effect of rotation around single bonds. In Chapter 5, we will learn about other aspects of stereochemistry.

Recall from Section 1.9A that **rotation occurs around carbon–carbon σ bonds**. Thus, the two CH_3 groups of ethane rotate, allowing the hydrogens on one carbon to adopt different orientations relative to the hydrogens on the other carbon. These arrangements are called **conformations**.

Table 4.2 Physical Properties of Alkanes

Property	Observation
Boiling point	<ul style="list-style-type: none"> Alkanes have low bp's compared to more polar compounds of comparable size. <div style="text-align: center;"> <p> $\text{CH}_3\text{CH}_2\text{CH}_3$ CH_3CHO $\text{CH}_3\text{CH}_2\text{OH}$ VDW VDW, DD VDW, DD, HB MW = 44 MW = 44 MW = 46 bp = -42 °C bp = 21 °C bp = 79 °C </p> <p>low bp →</p> <p>Increasing strength of intermolecular forces Increasing boiling point</p> </div>
	<ul style="list-style-type: none"> Bp increases as the number of carbons increases because of increased surface area. <div style="text-align: center;"> <p> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 0 °C bp = 36 °C bp = 69 °C </p> <p>Increasing surface area Increasing boiling point</p> </div>
	<ul style="list-style-type: none"> The bp of isomers decreases with branching because of decreased surface area. <div style="text-align: center;"> <p> $(\text{CH}_3)_4\text{C}$ $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ bp = 10 °C bp = 30 °C bp = 36 °C </p> <p>Increasing surface area Increasing boiling point</p> </div>
Melting point	<ul style="list-style-type: none"> Alkanes have low mp's compared to more polar compounds of comparable size. <div style="text-align: center;"> <p> $\text{CH}_3\text{CH}_2\text{CH}_3$ CH_3CHO VDW VDW, DD mp = -190 °C mp = -121 °C </p> <p>low mp →</p> <p>Increasing strength of intermolecular forces Increasing melting point</p> </div>
	<ul style="list-style-type: none"> Mp increases as the number of carbons increases because of increased surface area. <div style="text-align: center;"> <p> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ mp = -138 °C mp = -95 °C </p> <p>Increasing surface area Increasing melting point</p> </div>
	<ul style="list-style-type: none"> Mp increases with increased symmetry. <div style="text-align: center;"> <p> $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)_2$ $(\text{CH}_3)_4\text{C}$ mp = -160 °C mp = -17 °C </p> <p>Increasing symmetry Increasing melting point</p> </div>
Solubility	<ul style="list-style-type: none"> Alkanes are soluble in organic solvents. Alkanes are insoluble in water.


Key: bp = boiling point; mp = melting point; VDW = van der Waals; DD = dipole-dipole; HB = hydrogen bonding; MW = molecular weight




- *Conformations* are different arrangements of atoms that are interconverted by rotation about single bonds.

Names are given to two different arrangements.

- In the *eclipsed conformation*, the C–H bonds on one carbon are directly aligned with the C–H bonds on the adjacent carbon.
- In the *staggered conformation*, the C–H bonds on one carbon bisect the H–C–H bond angle on the adjacent carbon.




Rotating the atoms on one carbon by 60° converts an eclipsed conformation into a staggered conformation, and vice versa. These conformations are often viewed **end-on**—that is, looking directly down the carbon–carbon bond. The angle that separates a bond on one atom from a bond on an adjacent atom is called a **dihedral angle**. For ethane in the staggered conformation, the dihedral angle for the C–H bonds is 60° . For eclipsed ethane, it is 0° .




End-on representations for conformations are commonly drawn using a convention called a **Newman projection**. A Newman projection is a graphic that shows the three groups bonded to each carbon atom in a particular C–C bond, as well as the dihedral angle that separates them.

HOW TO Draw a Newman Projection

Step [1] Look directly down the C–C bond (end-on), and draw a circle with a dot in the center to represent the carbons of the C–C bond.



Step [2] Draw in the bonds.



Step [3] Add the atoms on each bond.




Figure 4.6 illustrates the Newman projections for both the staggered and eclipsed **conformations** for ethane.

Follow this procedure for any C–C bond. With a Newman projection, **always consider one C–C bond only and draw the atoms bonded to the carbon atoms, not the carbon atoms in the bond itself**. Newman projections for the staggered and eclipsed conformations of propane are drawn in Figure 4.7.

Problem 4.18 Draw the staggered and eclipsed conformations that result from rotation around the C–C bond in $\text{CH}_3\text{--CH}_2\text{Br}$.

The staggered and eclipsed conformations of ethane interconvert at room temperature, but **each conformation is not equally stable**.

- The staggered conformations are more stable (lower in energy) than the eclipsed conformations.

Figure 4.6
Newman projections for the staggered and eclipsed conformations of ethane






Figure 4.7

Newman projections for the staggered and eclipsed conformations of propane





Electron–electron repulsion between the bonds in the eclipsed conformation increases its energy compared to the staggered conformation, where the bonding electrons are farther apart.



In a Newman projection it doesn't matter which C you pick to be in the front or the back. All of the Newman projections shown here represent the staggered conformation of propane.

These C–H bonds are farther apart.

These C–H bonds are closer together.




The difference in energy between the staggered and eclipsed conformations is 12 kJ/mol (2.9 kcal/mol), a small enough difference that the rotation is still very rapid at room temperature, and the conformations cannot be separated. Because three eclipsed C–H bonds increase the energy of a conformation by 12 kJ/mol, **each eclipsed C–H bond results in an increase in energy of 4.0 kJ/mol (1.0 kcal/mol)**. The energy difference between the staggered and eclipsed conformations is called **torsional energy**. Thus, eclipsing introduces **torsional strain** into a molecule.

- **Torsional strain** is an increase in energy caused by eclipsing interactions.

The graph in Figure 4.8 shows how the potential energy of ethane changes with dihedral angle as one CH₃ group rotates relative to the other. **The staggered conformation is the most stable**

Figure 4.8

Graph: Energy versus dihedral angle for ethane



- Note the position of the labeled H atom after each 60° rotation. All three staggered conformations are identical (except for the position of the label), and the same is true for all three eclipsed conformations.

Each H,H eclipsing interaction contributes 4.0 kJ/mol of destabilization to the eclipsed conformation.

Strain results in an **increase in energy**. Torsional strain is the first of three types of strain discussed in this text. The other two are steric strain (Section 4.10) and angle strain (Section 4.11).

arrangement, so it is at an energy minimum. As the C–H bonds on one carbon are rotated relative to the C–H bonds on the other carbon, the energy increases as the C–H bonds get closer until a **maximum is reached after 60° rotation to the eclipsed conformation.** As rotation continues, the energy decreases until after 60° rotation, when the staggered conformation is reached once again. At any given moment, all ethane molecules do not exist in the more stable staggered conformation; rather, a higher percentage of molecules is present in the more stable staggered conformation than any other possible arrangement.

- An energy minimum and maximum occur every 60° as the conformation changes from staggered to eclipsed. Conformations that are neither staggered nor eclipsed are intermediate in energy.

Problem 4.19


Draw an energy versus rotation diagram similar to Figure 4.8 for rotation around a C–C bond in propane.

Problem 4.20

The torsional energy in propane is 14 kJ/mol (3.4 kcal/mol). Because each H,H eclipsing interaction is worth 4.0 kJ/mol (1.0 kcal/mol) of destabilization, how much is one H,CH₃ eclipsing interaction worth in destabilization? (See Section 4.10 for an alternate way to arrive at this value.)

4.10 Conformations of Butane

Butane and higher molecular weight alkanes have several carbon–carbon bonds, all capable of rotation.




To analyze the different conformations that result from rotation about the C2–C3 bond, begin arbitrarily with one—for example, the staggered conformation that places two CH₃ groups 180° from each other—then,

- Rotate one carbon atom in 60° increments either clockwise or counterclockwise, while keeping the other carbon fixed. Continue until you return to the original conformation.

It takes six 60° rotations to return to the original conformation.

Figure 4.9 illustrates the six possible conformations that result from this process.

Figure 4.9
Six different conformations of butane



Although each 60° bond rotation converts a staggered conformation into an eclipsed conformation (or vice versa), neither all the staggered conformations nor all the eclipsed conformations are the same. For example, the dihedral angle between the methyl groups in staggered conformations **3** and **5** are both 60° , whereas it is 180° in staggered conformation **1**.


- A staggered conformation with two larger groups 180° from each other is called *anti*.
- A staggered conformation with two larger groups 60° from each other is called *gauche*.

Similarly, the methyl groups in conformations **2** and **6** both eclipse hydrogen atoms, whereas they eclipse each other in conformation **4**.

The staggered conformations (**1**, **3**, and **5**) are lower in energy than the eclipsed conformations (**2**, **4**, and **6**), but how do the energies of the individual staggered and eclipsed conformations compare to each other? The relative energies of the individual staggered conformations (or the individual eclipsed conformations) depend on their **steric strain**.

- **Steric strain** is an increase in energy resulting when atoms are forced too close to one another.


The methyl groups are farther apart in the anti conformation (**1**) than in the gauche conformations (**3** and **5**), so amongst the staggered conformations, **1** is lower in energy (more stable) than **3** and **5**. In fact, the anti conformation is 3.8 kJ/mol (0.9 kcal/mol) lower in energy than either gauche conformation because of the steric strain that results from the proximity of the methyl groups in **3** and **5**.



- Gauche conformations are generally higher in energy than anti conformations because of steric strain.

Steric strain also affects the relative energies of eclipsed conformations. Conformation **4** is higher in energy than **2** or **6**, because the two larger CH₃ groups are forced close to each other, introducing considerable steric strain.

Steric strain caused by two eclipsed CH₃ groups




To graph energy versus dihedral angle, keep in mind two considerations:

- Staggered conformations are at energy minima and eclipsed conformations are at energy maxima.
- Unfavorable steric interactions increase energy.

Figure 4.10

Graph: Energy versus dihedral angle for butane



- Staggered conformations **1**, **3**, and **5** are at energy minima.
- Anti conformation **1** is lower in energy than gauche conformations **3** and **5**, which possess steric strain.
- Eclipsed conformations **2**, **4**, and **6** are at energy maxima.
- Eclipsed conformation **4**, which has additional steric strain due to two eclipsed CH₃ groups, is highest in energy.

For butane, this means that anti conformation **1** is lowest in energy, and conformation **4** with two eclipsed CH₃ groups is the highest in energy. The relative energy of other conformations is depicted in the energy versus rotation diagram for butane in Figure 4.10.

We can now use the values in Figure 4.10 to estimate the destabilization caused by other eclipsed groups. For example, conformation **4** is 19 kJ/mol less stable than the anti conformation **1**. Conformation **4** possesses two H,H eclipsing interactions, worth 4.0 kJ/mol each in destabilization (Section 4.9), and one CH₃,CH₃ eclipsing interaction. Thus, the **CH₃,CH₃ interaction** is worth $19 - 2(4.0) = 11$ kJ/mol of destabilization.

Similarly, conformation **2** is 16 kJ/mol less stable than the anti conformation **1**, and possesses one H,H eclipsing interaction (worth 4.0 kJ/mol of destabilization), and two H,CH₃ interactions. Thus **each H,CH₃ interaction** is worth $1/2(16 - 4.0) = 6.0$ kJ/mol of destabilization. These values are summarized in Table 4.3.


- The energy difference between the lowest and highest energy conformations is called the *barrier to rotation*.

Table 4.3 Summary: Torsional and Steric Strain Energies in Acyclic Alkanes


Type of interaction	Energy increase	
	kJ/mol	kcal/mol
H,H eclipsing	4.0	1.0
H,CH ₃ eclipsing	6.0	1.4
CH ₃ ,CH ₃ eclipsing	11	2.6
gauche CH ₃ groups	3.8	0.9

We can use these same principles to determine conformations and relative energies for any acyclic alkane. Because the lowest energy conformation has all bonds staggered and all large groups anti, alkanes are often drawn in zigzag skeletal structures to indicate this.

A zigzag arrangement keeps all carbons **staggered** and **anti**.



Problem 4.21



- Draw the three staggered and three eclipsed conformations that result from rotation around the designated bond using Newman projections.
- Label the most stable and least stable conformation.


Problem 4.22

Consider rotation around the carbon-carbon bond in 1,2-dichloroethane (ClCH₂CH₂Cl).

- Using Newman projections, draw all of the staggered and eclipsed conformations that result from rotation around this bond.
- Graph energy versus dihedral angle for rotation around this bond.

Problem 4.23

Calculate the destabilization present in each eclipsed conformation.




4.11 An Introduction to Cycloalkanes

Besides torsional strain and steric strain, the conformations of cycloalkanes are also affected by **angle strain**.

- Angle strain** is an increase in energy when tetrahedral bond angles deviate from the optimum angle of 109.5°.


Originally cycloalkanes were thought to be flat rings, with the bond angles between carbon atoms determined by the size of the ring. For example, a flat cyclopropane ring would have 60° internal bond angles, a flat cyclobutane ring would have 90° angles, and large flat rings would have very large angles. It was assumed that rings with bond angles so different from the tetrahedral bond angle would be very strained and highly reactive. This is called the Baeyer strain theory.



It turns out, though, that **cycloalkanes with more than three C atoms in the ring are not flat molecules**. They are puckered to **reduce strain**, both angle strain and torsional strain. The three-dimensional structures of some simple cycloalkanes are shown in Figure 4.11. Three- and four-membered rings still possess considerable angle strain, but puckering reduces the internal bond angles in larger rings, thus reducing angle strain.

Figure 4.11


Three-dimensional structure of some cycloalkanes



Keep in mind the three different types of strain in organic molecules:

- **Torsional strain:** strain caused by eclipsing interactions.
- **Steric strain:** strain produced when atoms are forced too close to each other.
- **Angle strain:** strain produced when bond angles deviate from 109.5° (for sp^3 hybridized atoms).

Many polycyclic hydrocarbons are of interest to chemists. For example, **dodecahedrane**, containing 12 five-membered rings bonded together, is one member of a family of three hydrocarbons that contain several rings of one size joined together. The two other members of this family are **tetrahedrane**, consisting of four three-membered rings, and **cubane**, consisting of six four-membered rings. These compounds are the simplest regular polyhedra whose structures resemble three of the highly symmetrical Platonic solids: the tetrahedron, the cube, and the dodecahedron.



How stable are these compounds? Tetrahedrane (with internal 60° bond angles) is so strained that all attempts to prepare it have been thus far unsuccessful. Although cubane is also highly strained because of its 90° bond angles, it was first synthesized in 1964 and is a stable molecule at room temperature. Finally, dodecahedrane is very stable because it has bond angles very close to the tetrahedral bond angle (108° versus 109.5°). Its synthesis eluded chemists for years not because of its strain or inherent instability, but because of the enormous challenge of joining 12 five-membered rings together to form a sphere.

4.12 Cyclohexane


Let's now examine in detail the conformation of **cyclohexane**, the most common ring size in naturally occurring compounds.

4.12A The Chair Conformation



Visualizing the chair. If the cyclohexane chair conformation is tipped downward, we can more easily view it as a chair with a back, seat, and foot support.


A planar cyclohexane ring would experience angle strain, because the internal bond angle between the carbon atoms would be 120° , and torsional strain, because all of the hydrogens on adjacent carbon atoms would be eclipsed.



In reality, cyclohexane adopts a puckered conformation, called the **chair** form, which is more stable than any other possible conformation.


Figure 4.12

A three-dimensional model of the chair form of cyclohexane with all H atoms drawn




- Cyclohexane has **six axial H's** and **six equatorial H's**.

The carbon skeleton of chair cyclohexane



The chair conformation is so stable because it eliminates angle strain (all C–C–C bond angles are **109.5°**) and torsional strain (all hydrogens on adjacent carbon atoms are **staggered**, not eclipsed).




- In cyclohexane, three C atoms pucker up and three C atoms pucker down, alternating around the ring. These C atoms are called *up C's* and *down C's*.


Each carbon in cyclohexane has two different kinds of hydrogens.

- *Axial* hydrogens are located above and below the ring (along a perpendicular axis).
- *Equatorial* hydrogens are located in the plane of the ring (around the equator).

3 up C's and 3 down C's



Two kinds of H's




- **Axial** bonds are oriented **above** and **below**.
- **Equatorial** bonds are oriented around the **equator**.

Each cyclohexane carbon atom has one axial and one equatorial hydrogen.

A three-dimensional representation of the chair form is shown in Figure 4.12. Before continuing, we must first learn how to draw the chair form of cyclohexane.

HOW TO Draw the Chair Form of Cyclohexane

Step [1] Draw the carbon skeleton.



- Draw three parts of the chair: **a wedge**, **a set of parallel lines**, and **another wedge**.
- Then, join them together.
- The bottom 3 C's come out of the page, and for this reason, bonds to them are often highlighted in bold.

—Continued

HOW TO, continued . . .

Step [2] Label the *up* C's and *down* C's on the ring.




● = *up* C ○ = *down* C

- There are 3 *up* C's and 3 *down* C's, and they alternate around the ring.

Step [3] Draw in the axial H atoms.

3 axial H's **above** the ring




- On an *up* C the axial H is *up*.
- On a *down* C the axial H is *down*.

3 axial H's **below** the ring




Step [4] Draw in the equatorial H atoms.


- The axial H is **down** on a *down* C, so the equatorial H must be up.
- The axial H is **up** on an *up* C, so the equatorial H must be down.



All equatorial H's drawn in.




All H's drawn in.



Axial H's are drawn in blue.

Problem 4.24 Classify the ring carbons as *up* C's or *down* C's. Identify the bonds highlighted in bold as axial or equatorial.



4.12B Ring-Flipping

Like acyclic alkanes, **cyclohexane does not remain in a single conformation**. The bonds twist and bend, resulting in new arrangements, but the movement is more restricted. One important conformational change involves **ring-flipping**, which can be viewed as a two-step process.





Figure 4.13
Ring-flipping interconverts axial and equatorial hydrogens in cyclohexane



- A **down carbon flips up**. This forms a new conformation of cyclohexane called a **boat**. The boat form has two carbons oriented above a plane containing the other four carbons.
- The boat form can flip in two possible ways. The original carbon (labeled with an open circle) can flip down, re-forming the initial conformation; or the **second up carbon** (labeled with a solid circle) **can flip down**. This forms a **second chair conformation**.

Because of ring-flipping, the **up carbons become down carbons and the down carbons become up carbons**. Thus, cyclohexane exists as two different chair conformations of equal stability, which rapidly interconvert at room temperature.

The process of ring-flipping also affects the orientation of cyclohexane's hydrogen atoms.

- Axial and equatorial H atoms are interconverted during a ring flip. Axial H atoms become equatorial H atoms, and equatorial H atoms become axial H atoms (Figure 4.13).

The chair forms of cyclohexane are 30 kJ/mol more stable than the boat forms. The boat conformation is destabilized by torsional strain because the hydrogens on the four carbon atoms in the plane are eclipsed. Additionally, there is steric strain because two hydrogens at either end of the boat—the **flagpole hydrogens**—are forced close to each other, as shown in Figure 4.14.

4.13 Substituted Cycloalkanes


What happens when one hydrogen on cyclohexane is replaced by a larger substituent? Is there a difference in the stability of the two cyclohexane conformations? To answer these questions, remember one rule.

- The equatorial position has more room than the axial position, so larger substituents are more stable in the equatorial position.

4.13A Cyclohexane with One Substituent

There are two possible chair conformations of a monosubstituted cyclohexane, such as methylcyclohexane.

Figure 4.14
Two views of the boat conformation of cyclohexane




The boat form of cyclohexane is less stable than the chair forms for two reasons.

- Eclipsing interactions between H's cause torsional strain.
- The proximity of the flagpole H's causes steric strain.


HOW TO Draw the Two Conformations for a Substituted Cyclohexane

Step [1] Draw one chair form and add the substituents.

- Arbitrarily pick a ring carbon, classify it as an *up* or *down* carbon, and draw the bonds. Each C has one axial and one equatorial bond.
- Add the substituents, in this case H and CH₃, arbitrarily placing one axial and one equatorial. In this example, the CH₃ group is drawn equatorial.
- This forms one of the two possible chair conformations, labeled Conformation 1.




Step [2] Ring-flip the cyclohexane ring.



- Convert *up* C's to *down* C's and vice versa. The chosen *up* C now puckers down.

Step [3] Add the substituents to the second conformation.

- Draw axial and equatorial bonds. On a *down* C the axial bond is *down*.
- Ring-flipping converts axial bonds to equatorial bonds, and vice versa. The equatorial methyl becomes axial.
- This forms the other possible chair conformation, labeled Conformation 2.




Although the CH₃ group flips from equatorial to axial, it starts on a down bond, and stays on a down bond. It never flips from below the ring to above the ring.

- A substituent always stays on the same side of the ring—either below or above—during the process of ring-flipping.

The two conformations of methylcyclohexane are different, so they are not equally stable. In fact, Conformation 1, which places the larger methyl group in the roomier equatorial position, is considerably more stable than Conformation 2, which places it axial.

Each carbon atom has one *up* and one *down* bond. An *up* bond can be either axial or equatorial, depending on the carbon to which it is attached. On an *up* C, the axial bond is *up*, but on a *down* C, the equatorial bond is *up*.

The equatorial bond is *up*. The axial bond is *up*.



The larger CH₃ group is equatorial.





Figure 4.15

Three-dimensional representations for the two conformations of methylcyclohexane




Why is a substituted cyclohexane ring more stable with a larger group in the equatorial position? Figure 4.15 shows that with an equatorial CH₃ group, steric interactions with nearby groups are minimized. An axial CH₃ group, however, is close to two other axial H atoms, creating two destabilizing steric interactions called **1,3-diaxial interactions**. Each unfavorable H,CH₃ interaction destabilizes the conformation by 3.8 kJ/mol, so Conformation 2 is 7.6 kJ/mol less stable than Conformation 1.

- Larger axial substituents create unfavorable 1,3-diaxial interactions, destabilizing a cyclohexane conformation.

The larger the substituent on the six-membered ring, the higher the percentage of the conformation containing the equatorial substituent at equilibrium. In fact, with a very large substituent like *tert*-butyl [(CH₃)₃C–], essentially none of the conformation containing an axial *tert*-butyl group is present at room temperature, so **the ring is essentially anchored in a single conformation having an equatorial *tert*-butyl group**. This is illustrated in Figure 4.16.

Problem 4.25

Draw a second chair conformation for each cyclohexane. Then decide which conformation is present in higher concentration at equilibrium.

**Problem 4.26**


When an ethyl group (CH₃CH₂–) is bonded to a cyclohexane ring, 96% of the molecules possess an equatorial CH₃CH₂– group at equilibrium. When an ethynyl group (HC≡C–) is bonded to a cyclohexane ring, only 67% of the molecules possess an equatorial HC≡C– group at equilibrium. Suggest a reason for this difference.

4.13B A Disubstituted Cycloalkane

Rotation around the C–C bonds in the ring of a cycloalkane is restricted, so **a group on one side of the ring can never rotate to the other side of the ring**. As a result, there are two different 1,2-dimethylcyclopentanes—one having two CH₃ groups on the **same side** of the ring and one having them on **opposite sides** of the ring.

Figure 4.16

The two conformations of *tert*-butylcyclohexane




Wedges indicate bonds in front of the plane of the ring and dashes indicate bonds behind.

For a review of this convention, see Section 1.6B. In this text, dashes are drawn equal in length, as recommended in the latest IUPAC guidelines. If a ring carbon is bonded to a CH_3 group in **front** of the ring (on a wedge), it is *assumed* that the other atom bonded to this carbon is hydrogen, located **behind** the ring (on a dash).

Cis and **trans** isomers are named by adding the prefixes *cis* and *trans* to the name of the cycloalkane. Thus, **A** is *cis*-1,2-dimethylcyclopentane, and **B** is *trans*-1,2-dimethylcyclopentane.

A disubstituted cycloalkane: 1,2-dimethylcyclopentane

These two compounds cannot be interconverted.




A and **B** are **isomers**, because they are different compounds with the same molecular formula, but they represent the second major class of isomers called **stereoisomers**.

- **Stereoisomers** are isomers that differ *only* in the way the atoms are oriented in space.

The prefixes **cis** and **trans** are used to distinguish these stereoisomers.

- The **cis isomer** has two groups on the *same side* of the ring.
- The **trans isomer** has two groups on *opposite sides* of the ring.

Cis- and *trans*-1,2-dimethylcyclopentane can also be drawn as if the plane of the ring goes through the plane of the page. Each carbon in the ring then has one bond that points above the ring and one that points below.




Problem 4.27 Draw the structure for each compound using wedges and dashes.

a. *cis*-1,2-dimethylcyclopropane b. *trans*-1-ethyl-2-methylcyclopentane

Problem 4.28 For *cis*-1,3-diethylcyclobutane, draw (a) a stereoisomer; (b) a constitutional isomer.

4.13C A Disubstituted Cyclohexane

A disubstituted cyclohexane like 1,4-dimethylcyclohexane also has *cis* and *trans* stereoisomers. In addition, each of these stereoisomers has two possible chair conformations.




All disubstituted cycloalkanes with two groups bonded to *different* atoms have *cis* and *trans* isomers.

To draw both conformations for each stereoisomer, follow the procedure in Section 4.13A for a monosubstituted cyclohexane, keeping in mind that two substituents must now be added to the ring.


HOW TO Draw Two Conformations for a Disubstituted Cyclohexane

Step [1] Draw one chair form and add the substituents.


- For *trans*-1,4-dimethylcyclohexane, arbitrarily pick two C's located 1,4- to each other, classify them as *up* or *down* C's, and draw in the substituents.
- The *trans* isomer must have one group *above* the ring (on an *up* bond) and one group *below* the ring (on a *down* bond). The substituents can be either axial or equatorial, as long as one is up and one is down. The easiest *trans* isomer to visualize has two axial CH₃ groups. This arrangement is said to be **diaxial**.
- This forms one of the two possible chair conformations, labeled Conformation 1.



Step [2] Ring-flip the cyclohexane ring.




Step [3] Add the substituents to the second conformation.



- Ring-flipping converts axial bonds to equatorial bonds, and vice versa. The diaxial CH₃ groups become diequatorial. This *trans* conformation is less obvious to visualize. It is still *trans*, because one CH₃ group is above the ring (on an *up* bond), and one is below (on a *down* bond).


Conformations **1** and **2** are not equally stable. Because **Conformation 2** has both larger CH₃ groups in the roomier equatorial position, it is lower in energy.



The *cis* isomer of 1,4-dimethylcyclohexane also has two conformations, as shown in Figure 4.17. Because each conformation has one CH₃ group axial and one equatorial, they are **identical in energy**. At room temperature, therefore, the two conformations exist in a 50:50 mixture at equilibrium.

Figure 4.17

The two conformations of *cis*-1,4-dimethylcyclohexane



- A *cis* isomer has two groups on the same side of the ring, either both *up* or both *down*. In this example, Conformations **1** and **2** have two CH_3 groups drawn up.
- Both conformations have one CH_3 group axial and one equatorial, making them equally stable.

The relative stability of the two conformations of any disubstituted cyclohexane can be analyzed using this procedure.


- A *cis* isomer has two substituents on the same side, either both on *up* bonds or both on *down* bonds.
- A *trans* isomer has two substituents on opposite sides, one *up* and one *down*.
- Whether substituents are axial or equatorial depends on the relative location of the two substituents (on carbons 1,2-, 1,3-, or 1,4-).

Sample Problem 4.4

Draw both chair conformations for *trans*-1,3-dimethylcyclohexane.


Solution

Step [1] Draw one chair form and add substituents.



- Pick two C's 1,3- to each other.
- The *trans* isomer has two groups on **opposite sides**. In Conformation **1**, this means that one CH_3 is equatorial (on an *up* bond), and one group is axial (on a *down* bond).




Steps [2–3] Ring-flip and add substituents.



- The two *down* C's flip up.
- The axial CH_3 flips equatorial (still a *down* bond) and the equatorial CH_3 flips axial (still an *up* bond). Conformation **2** is *trans* because the two CH_3 's are still on opposite sides.
- **Conformations 1 and 2 are equally stable** because each has one CH_3 equatorial and one axial.

Problem 4.29

Label each compound as *cis* or *trans*. Then draw the second chair conformation.



- Problem 4.30** Consider 1,2-dimethylcyclohexane.
- Draw structures for the cis and trans isomers using a hexagon for the six-membered ring.
 - Draw the two possible chair conformations for the cis isomer. Which conformation, if either, is more stable?
 - Draw the two possible chair conformations for the trans isomer. Which conformation, if either, is more stable?
 - Which isomer, cis or trans, is more stable and why?

- Problem 4.31** Draw a chair conformation of cyclohexane with one CH_3CH_2 group and one CH_3 group that fits each description:
- A 1,1-disubstituted cyclohexane with an axial CH_3CH_2 group
 - A cis-1,2-disubstituted cyclohexane with an axial CH_3 group
 - A trans-1,3-disubstituted cyclohexane with an equatorial CH_3 group
 - A trans-1,4-disubstituted cyclohexane with an equatorial CH_3CH_2 group

4.14 Oxidation of Alkanes

In Chapter 3 we learned that a functional group contains a heteroatom or π bond and constitutes **the reactive part of a molecule**. Alkanes are the only family of organic molecules that have no functional group, and therefore, **alkanes undergo few reactions**. In fact, alkanes are inert to reaction unless forcing conditions are used.

In Chapter 4, we consider only one reaction of alkanes—**combustion**. Combustion is an **oxidation–reduction** reaction.

4.14A Oxidation and Reduction Reactions

Compounds that contain many C–H bonds and few C–Z bonds are said to be in a **reduced state**, whereas those that contain few C–H bonds and more C–Z bonds are in a **more oxidized state**. CH_4 is thus highly reduced, while CO_2 is highly oxidized.

Because Z is more electronegative than C, replacing C–H bonds with C–Z bonds decreases the electron density around C. Loss of electron density = oxidation.

- **Oxidation** is the *loss* of electrons.
- **Reduction** is the *gain* of electrons.

Oxidation and reduction are opposite processes. As in acid–base reactions, there are always two components in these reactions. **One component is oxidized and one is reduced.**


To determine if an organic compound undergoes oxidation or reduction, we concentrate on the carbon atoms of the starting material and product, and **compare the relative number of C–H and C–Z bonds**, where Z = an element *more electronegative* than carbon (usually O, N, or X). Oxidation and reduction are then defined in two complementary ways.

- **Oxidation** results in an *increase* in the number of C–Z bonds; *or*
- **Oxidation** results in a *decrease* in the number of C–H bonds.
- **Reduction** results in a *decrease* in the number of C–Z bonds; *or*
- **Reduction** results in an *increase* in the number of C–H bonds.

Figure 4.18 illustrates the oxidation of CH_4 by replacing C–H bonds with C–O bonds (from left to right). The symbol **[O]** indicates oxidation. Because reduction is the reverse of oxidation, the molecules in Figure 4.18 are progressively reduced moving from right to left, from CO_2 to CH_4 . The symbol **[H]** indicates reduction.


Figure 4.18

The oxidation and reduction of a carbon compound



Sample Problem 4.5

Determine whether the organic compound is oxidized or reduced in each transformation.




Solution

- a. The conversion of ethanol to acetic acid is an **oxidation** because the number of C–O bonds increases: $\text{CH}_3\text{CH}_2\text{OH}$ has one C–O bond and CH_3COOH has three C–O bonds.
- b. The conversion of ethylene to ethane is a **reduction** because the number of C–H bonds increases: ethane has two more C–H bonds than ethylene.

Problem 4.32


Classify each transformation as an oxidation, reduction, or neither.



4.14B Combustion of Alkanes

When an organic compound is *oxidized* by a reagent, the reagent itself is *reduced*. Similarly, when an organic compound is *reduced* by a reagent, the reagent is *oxidized*. **Organic chemists identify a reaction as an oxidation or reduction by what happens to the *organic* component of the reaction.**


Alkanes undergo **combustion**—that is, **they burn in the presence of oxygen to form carbon dioxide and water**. This is a practical example of oxidation. Every C–H and C–C bond in the starting material is converted to a C–O bond in the product. The reactions drawn show the combustion of two different alkanes. Note that the products, $\text{CO}_2 + \text{H}_2\text{O}$, are the same, regardless of the identity of the starting material. Combustion of alkanes in the form of natural gas, gasoline, or heating oil releases energy for heating homes, powering vehicles, and cooking food.



Combustion requires a spark or a flame to initiate the reaction. Gasoline, therefore, which is composed largely of alkanes, can be safely handled and stored in the air, but the presence of a spark or match causes immediate and violent combustion.

Figure 4.19

The changing concentration of CO₂ in the atmosphere since 1958



The increasing level of atmospheric CO₂ is clearly evident on the graph. Two data points are recorded each year. The sawtooth nature of the graph is due to seasonal variation of CO₂ level with the seasonal variation in photosynthesis. (Data recorded at Mauna Loa, Hawaii)



Driving an automobile 10,000 miles at 25 miles per gallon releases ~10,000 lb of CO₂ into the atmosphere.

The combustion of alkanes and other hydrocarbons obtained from fossil fuels adds a tremendous amount of CO₂ to the atmosphere each year. Quantitatively, data show a 22% increase in the atmospheric concentration of CO₂ in the last 49 years (from 315 parts per million in 1958 to 384 parts per million in 2007; Figure 4.19). Although the composition of the atmosphere has changed over the lifetime of the earth, this may be the first time that the actions of humankind have altered that composition significantly and so quickly.

An increased CO₂ concentration in the atmosphere may have long-range and far-reaching effects. CO₂ absorbs thermal energy that normally radiates from the earth's surface, and redirects it back to the surface. Higher levels of CO₂ may therefore contribute to an increase in the average temperature of the earth's atmosphere. This **global warming**, as it has been called, has many consequences—the melting of polar ice caps, the rise in sea level, and drastic global climate changes to name a few. How great a role CO₂ plays in this process is hotly debated.

Problem 4.33

Draw the products of each combustion reaction.

**4.15 Lipids—Part 1**

Lipids are biomolecules whose properties resemble those of alkanes and other hydrocarbons. They are unlike any other class of biomolecules, though, because they are defined by a **physical property**, not by the presence of a particular functional group.


- Lipids are biomolecules that are soluble in organic solvents and insoluble in water.

Lipids that contain carbon-carbon double bonds are discussed in Section 10.6.

Lipids have varied sizes and shapes, and a diverse number of functional groups. Fat-soluble vitamins like vitamin A and the phospholipids that comprise cell membranes are two examples of lipids that were presented in Sections 3.5 and 3.7. Other examples are shown in Figure 4.20. One unifying feature accounts for their solubility.

Figure 4.20

Three representative lipid molecules



- Lipids are composed of many nonpolar C–H and C–C bonds, and have few polar functional groups.

Waxes are lipids having two long alkyl chains joined by a single oxygen-containing functional group. Because of their many C–C and C–H bonds, **waxes are hydrophobic**. They form a protective coating on the feathers of birds to make them water repellent, and on leaves to prevent water evaporation. Bees secrete $\text{CH}_3(\text{CH}_2)_{14}\text{COO}(\text{CH}_2)_{29}\text{CH}_3$, a wax that forms the honeycomb in which they lay eggs.

$\text{PGF}_{2\alpha}$ belongs to a class of lipids called **prostaglandins**. Prostaglandins contain many C–C and C–H bonds and a single COOH group (a **carboxy group**). Prostaglandins possess a wide range of biological activities. They control inflammation, affect blood-platelet aggregation, and stimulate uterine contractions. Nonsteroidal anti-inflammatory drugs such as ibuprofen operate by blocking the synthesis of prostaglandins, as discussed in Sections 19.6 and 29.6.


Cholesterol is a member of the steroid family, a group of lipids having four rings joined together. Because it has just one polar OH group, cholesterol is insoluble in the aqueous medium of the blood. It is synthesized in the liver and transported to other cells bound to water-soluble organic molecules. Elevated cholesterol levels can lead to coronary artery disease.

Cholesterol is a vital component of the cell membrane. Its hydrophobic carbon chain is embedded in the interior of the lipid bilayer, and its hydrophilic hydroxy group is oriented toward the aqueous exterior (Figure 4.21). Because its tetracyclic carbon skeleton is quite rigid compared to the long floppy side chains of a phospholipid, cholesterol stiffens the cell membrane somewhat, giving it more strength.

More details concerning cholesterol's structure and properties are presented in Section 29.8.

Figure 4.21



Cholesterol embedded in a lipid bilayer of a cell membrane



- The nonpolar hydrocarbon skeleton of cholesterol is embedded in the nonpolar interior of the cell membrane. Its rigid carbon skeleton stiffens the fluid lipid bilayer, giving it strength.
- Cholesterol's polar OH group is oriented toward the aqueous media inside and outside the cell.

Lipids have a high energy content, meaning that much energy is released on their metabolism. Because lipids are composed mainly of C–C and C–H bonds, they are oxidized with the release of energy, just like alkanes are. In fact, lipids are the most efficient biomolecules for the storage of energy. **The combustion of alkanes provides heat for our homes, and the metabolism of lipids provides energy for our bodies.**

Problem 4.34 Which of the following compounds can be classified as lipids?



Problem 4.35 Explain why beeswax is insoluble in H_2O , slightly soluble in ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), and soluble in chloroform (CHCl_3).

KEY CONCEPTS

Alkanes


General Facts About Alkanes (4.1–4.3)

- Alkanes are composed of **tetrahedral, sp^3** hybridized C atoms.
- There are two types of alkanes: acyclic alkanes having molecular formula $\text{C}_n\text{H}_{2n+2}$, and cycloalkanes having molecular formula C_nH_{2n} .
- Alkanes have only **nonpolar C–C and C–H bonds** and no functional group, so they undergo few reactions.
- Alkanes are named with the suffix **-ane**.

Classifying C Atoms and H Atoms (4.1A)


- Carbon atoms are classified by the number of carbon atoms bonded to them; a **1° carbon is bonded to one other carbon**, and so forth.
- Hydrogen atoms are classified by the type of carbon atom to which they are bonded; a **1° H is bonded to a 1° carbon**, and so forth.

Names of Alkyl Groups (4.4A)



Conformations in Acyclic Alkanes (4.9, 4.10)

- Alkane conformations can be classified as **eclipsed**, **staggered**, **anti**, or **gauche** depending on the relative orientation of the groups on adjacent carbons.




- A staggered conformation is **lower in energy** than an eclipsed conformation.
- An anti conformation is **lower in energy** than a gauche conformation.

Types of Strain

- Torsional strain**—an increase in energy caused by eclipsing interactions (4.9).
- Steric strain**—an increase in energy when atoms are forced too close to each other (4.10).
- Angle strain**—an increase in energy when tetrahedral bond angles deviate from 109.5° (4.11).


Two Types of Isomers

- Constitutional isomers**—isomers that differ in the way the atoms are connected to each other (4.1A).
- Stereoisomers**—isomers that differ *only* in the way the atoms are oriented in space (4.13B).



Conformations in Cyclohexane (4.12, 4.13)

- Cyclohexane exists as **two chair conformations** in rapid equilibrium at room temperature.
- Each carbon atom on a cyclohexane ring has **one axial** and **one equatorial hydrogen**. Ring-flipping converts axial H's to equatorial H's, and vice versa.



- In substituted cyclohexanes, groups larger than hydrogen are more stable in the **roomier equatorial position**.
- Disubstituted cyclohexanes with substituents on different atoms exist as two possible stereoisomers.
 - The **cis** isomer has two groups on the **same side** of the ring, either both up or both down.
 - The **trans** isomer has two groups on **opposite sides** of the ring, one up and one down.


Oxidation–Reduction Reactions (4.14)

- Oxidation** results in an **increase in the number of C–Z bonds** or a **decrease in the number of C–H bonds**.
- Reduction** results in a **decrease in the number of C–Z bonds** or an **increase in the number of C–H bonds**.

PROBLEMS

Classifying Carbons and Hydrogens


4.36 For each alkane: (a) classify each carbon atom as 1°, 2°, 3°, or 4°; (b) classify each hydrogen atom as 1°, 2°, or 3°.



4.37 Draw the structure of an alkane that:

- Contains only 1° and 4° carbons.
- Contains only 2° carbons.
- Contains only 1° and 2° hydrogens.
- Contains only 1° and 3° hydrogens.

4.38 Like ginkgolide B, the cover molecule described in the Prologue, bilobalide is also isolated from *Ginkgo biloba* extracts. Classify each sp^3 hybridized carbon atom in bilobalide as 1°, 2°, 3°, or 4°.















Constitutional Isomers

4.39 Draw the structure of all compounds that fit the following descriptions.

- Five constitutional isomers having the molecular formula C_4H_8 .
- Nine constitutional isomers having the molecular formula C_7H_{16} .
- Twelve constitutional isomers having the molecular formula C_6H_{12} and containing one ring.

IUPAC Nomenclature

4.40 Give the IUPAC name for each compound.



4.41 Give the structure and IUPAC name for each of the nine isomers having molecular formula C_9H_{20} that contains seven carbons in the longest chain and two methyl groups as substituents.


4.42 Draw the structure corresponding to each IUPAC name.

- 3-ethyl-2-methylhexane
- sec-butylcyclopentane
- 4-isopropyl-2,4,5-trimethylheptane
- cyclobutylcycloheptane
- 3-ethyl-1,1-dimethylcyclohexane
- 4-butyl-1,1-diethylcyclooctane
- 6-isopropyl-2,3-dimethylnonane
- 2,2,6,6,7-pentamethyloctane
- cis-1-ethyl-3-methylcyclopentane
- trans-1-tert-butyl-4-ethylcyclohexane

4.43 Each of the following IUPAC names is incorrect. Explain why it is incorrect and give the correct IUPAC name.

- | | |
|--------------------------------|-------------------------------------|
| a. 2,2-dimethyl-4-ethylheptane | e. 1-ethyl-2,6-dimethylcycloheptane |
| b. 5-ethyl-2-methylhexane | f. 5,5,6-trimethyloctane |
| c. 2-methyl-2-isopropylheptane | g. 3-butyl-2,2-dimethylhexane |
| d. 1,5-dimethylcyclohexane | h. 1,3-dimethylbutane |

4.44 Give the IUPAC name for each compound.



Physical Properties


4.45 Rank each group of alkanes in order of increasing boiling point. Explain your choice of order.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_3\text{CH}_2\text{CH}_3$
 b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $\text{CH}_3(\text{CH}_2)_4\text{CH}_3$, $(\text{CH}_3)_2\text{CHCH}(\text{CH}_3)_2$

4.46 The melting points and boiling points of two isomeric alkanes are as follows: $\text{CH}_3(\text{CH}_2)_6\text{CH}_3$, mp = -57°C and bp = 126°C ; $(\text{CH}_3)_3\text{CC}(\text{CH}_3)_3$, mp = 102°C and bp = 106°C . (a) Explain why one isomer has a lower melting point but higher boiling point. (b) Explain why there is a small difference in the boiling points of the two compounds, but a huge difference in their melting points.

Conformation of Acyclic Alkanes


4.47 Which conformation in each pair is *higher* in energy? Calculate the energy difference between the two conformations using the values given in Table 4.3.




4.48 Considering rotation around the indicated bond in each compound, draw Newman projections for the most stable and least stable conformations.




4.49 Convert each three-dimensional model to a Newman projection around the indicated bond.




4.50 Convert each structure to a Newman projection around the indicated bond.





4.51 (a) Using Newman projections, draw all staggered and eclipsed conformations that result from rotation around the indicated bond in each molecule; (b) draw a graph of energy versus dihedral angle for rotation around this bond.



4.52 Label the sites of torsional and steric strain in each conformation.



4.53 Calculate the barrier to rotation for each designated bond.






4.54 The eclipsed conformation of $\text{CH}_3\text{CH}_2\text{Cl}$ is 15 kJ/mol less stable than the staggered conformation. How much is the H,Cl eclipsing interaction worth in destabilization?

4.55 (a) Draw the anti and gauche conformations for ethylene glycol ($\text{HOCH}_2\text{CH}_2\text{OH}$). (b) Ethylene glycol is unusual in that the gauche conformation is more stable than the anti conformation. Offer an explanation.

Conformations and Stereoisomers in Cycloalkanes

4.56 For each compound drawn below:

- Label each OH, Br, and CH_3 group as axial or equatorial.
- Classify each conformation as cis or trans.
- Translate each structure into a representation with a hexagon for the six-membered ring, and wedges and dashes for groups above and below the ring.
- Draw the second possible chair conformation for each compound.






4.57 Draw the two possible chair conformations for *cis*-1,3-dimethylcyclohexane. Which conformation, if either, is more stable?

4.58 For each disubstituted cyclohexane, indicate the axial/equatorial position of the substituents in the following table. The first entry has been completed for you.

Disubstituted cyclohexane	Axial/equatorial substituent location	
	Conformation 1	Conformation 2
a. 1,2- <i>cis</i> disubstituted	Axial/equatorial	Equatorial/axial
b. 1,2- <i>trans</i> disubstituted		
c. 1,3- <i>cis</i> disubstituted		
d. 1,3- <i>trans</i> disubstituted		
e. 1,4- <i>cis</i> disubstituted		
f. 1,4- <i>trans</i> disubstituted		

4.59 For each compound drawn below:


- Draw representations for the *cis* and *trans* isomers using a hexagon for the six-membered ring, and wedges and dashes for substituents.
- Draw the two possible chair conformations for the *cis* isomer. Which conformation, if either, is more stable?
- Draw the two possible chair conformations for the *trans* isomer. Which conformation, if either, is more stable?
- Which isomer, *cis* or *trans*, is more stable and why?



4.60 Which isomer in each pair of compounds is *lower* in energy?

- cis*- or *trans*-1,2-diethylcyclohexane
- cis*- or *trans*-1-ethyl-3-isopropylcyclohexane


4.61 Which of the given 1,3,5-trimethylcyclohexane isomers is more stable? Explain your choice.



4.62 Convert each of the following structures into its more stable chair form. One structure represents menthol and one represents isomenthol. Menthol, the more stable isomer, is used in lip balms and mouthwash. Which structure corresponds to menthol?




4.63 Glucose is a simple sugar with five substituents bonded to a six-membered ring.



- Using a chair representation, draw the most stable arrangement of these substituents on the six-membered ring.
- Convert this representation into one that uses a hexagon with wedges and dashes.


4.64 Galactose is a simple sugar formed when lactose, a carbohydrate in milk, is hydrolyzed. Individuals with galactosemia, a rare inherited disorder, lack an enzyme needed to metabolize galactose, and must avoid cow's milk and all products derived from cow's milk. Galactose is a stereoisomer of glucose (Problem 4.63).




- Draw both chair forms of galactose and label the more stable conformation.
- Which simple sugar, galactose or glucose, is more stable? Explain.
- Draw a constitutional isomer of galactose.
- Draw a stereoisomer of galactose that is different from glucose.

Constitutional Isomers and Stereoisomers


4.65 Classify each pair of compounds as constitutional isomers, stereoisomers, identical molecules, or not isomers of each other.



4.66 Classify each pair of compounds as constitutional isomers or identical molecules.






4.67 Draw a constitutional isomer and a stereoisomer for each compound.





4.68 Draw the three constitutional isomers having molecular formula C_7H_{14} that contain a five-membered ring and two methyl groups as substituents. For each constitutional isomer that can have cis and trans isomers, draw the two stereoisomers.

Oxidation and Reduction


4.69 Classify each reaction as oxidation, reduction, or neither.



4.70 Draw the products of combustion of each alkane.




4.71 Hydrocarbons like benzene are metabolized in the body to arene oxides, which rearrange to form phenols. This is an example of a general process in the body, in which an unwanted compound (benzene) is converted to a more water-soluble derivative called a *metabolite*, so that it can be excreted more readily from the body.




- Classify each of these reactions as oxidation, reduction, or neither.
- Explain why phenol is more water soluble than benzene. This means that phenol dissolves in urine, which is largely water, to a greater extent than benzene.

Lipids

4.72 Which of the following compounds are lipids?




- 4.73 Cholic acid, a compound called a **bile acid**, is converted to a **bile salt** in the body. Bile salts have properties similar to soaps, and they help transport lipids through aqueous solutions. Explain why this is so.




- 4.74 Mineral oil, a mixture of high molecular weight alkanes, is sometimes used as a laxative. Why are individuals who use mineral oil for this purpose advised to avoid taking it at the same time they consume foods rich in fat-soluble vitamins such as vitamin A?

Challenge Problems

- 4.75 Although penicillin G has two amide functional groups, one is much more reactive than the other. Which amide is more reactive and why?



- 4.76 Haloethanes ($\text{CH}_3\text{CH}_2\text{X}$, $\text{X} = \text{Cl}, \text{Br}, \text{I}$) have similar barriers to rotation (13.4–15.5 kJ/mol) despite the fact that the size of the halogen increases, $\text{Cl} \rightarrow \text{Br} \rightarrow \text{I}$. Offer an explanation.
- 4.77 When two six-membered rings share a C–C bond, this bicyclic system is called a **decalin**. There are two possible arrangements: *trans*-decalin having two hydrogen atoms at the ring fusion on opposite sides of the rings, and *cis*-decalin having the two hydrogens at the ring fusion on the same side.



- Draw *trans*- and *cis*-decalin using the chair form for the cyclohexane rings.
 - The *trans* isomer is more stable. Explain why.
- 4.78 Read Appendix B on naming branched alkyl substituents, and draw all possible alkyl groups having the formula C_5H_{11} . Give the IUPAC names for the eight compounds of molecular formula $\text{C}_{10}\text{H}_{20}$ that contain a cyclopentane ring with each of these alkyl groups as a substituent.

Stereochemistry

5

- 5.1 Starch and cellulose
- 5.2 The two major classes of isomers
- 5.3 Looking glass chemistry—Chiral and achiral molecules
- 5.4 Stereogenic centers
- 5.5 Stereogenic centers in cyclic compounds
- 5.6 Labeling stereogenic centers with *R* or *S*
- 5.7 Diastereomers
- 5.8 Meso compounds
- 5.9 *R* and *S* assignments in compounds with two or more stereogenic centers
- 5.10 Disubstituted cycloalkanes
- 5.11 Isomers—A summary
- 5.12 Physical properties of stereoisomers
- 5.13 Chemical properties of enantiomers



(S)-Naproxen is the active ingredient in the widely used pain relievers Naprosyn and Aleve. The three-dimensional orientation of two atoms at a single carbon in naproxen determines its therapeutic properties. Changing the position of these two atoms converts this anti-inflammatory agent into a liver toxin. In Chapter 5, we learn more about stereochemistry and how small structural differences can have a large effect on the properties of a molecule.

Are you left-handed or right-handed? If you're right-handed, you've probably spent little time thinking about your hand preference. If you're left-handed, though, you probably learned at an early age that many objects—like scissors and baseball gloves—“fit” for righties, but are “backwards” for lefties. **Hands, like many objects in the world around us, are mirror images that are *not* identical.**

In Chapter 5 we examine the “handedness” of molecules, and ask, “How important is the three-dimensional shape of a molecule?”


5.1 Starch and Cellulose

Recall from Chapter 4 that **stereochemistry is the three-dimensional structure of a molecule.** How important is stereochemistry? Two biomolecules—starch and cellulose—illustrate how apparently minute differences in structure can result in vastly different properties.


Starch and cellulose are two polymers that belong to the family of biomolecules called **carbohydrates** (Figure 5.1). A **polymer is a large molecule composed of repeating smaller units—called monomers—that are covalently bonded together.**

Starch is the main carbohydrate in the seeds and roots of plants. When we humans ingest wheat, rice, or potatoes, for example, we consume starch, which is then hydrolyzed to the simple sugar **glucose**, one of the compounds our bodies use for energy. **Cellulose**, nature's most abundant organic material, gives rigidity to tree trunks and plant stems. Wood, cotton, and flax are composed largely of cellulose. Complete hydrolysis of cellulose also forms glucose, but unlike starch, humans cannot metabolize cellulose to glucose. In other words, we can digest starch but not cellulose.

Cellulose and starch are both composed of the same repeating unit—a six-membered ring containing an oxygen atom and three OH groups—joined by an oxygen atom. They differ in the position of the O atom joining the rings together.




- In cellulose, the O atom joins two rings using two equatorial bonds.
- In starch, the O atom joins two rings using one equatorial and one axial bond.



Starch and cellulose are **isomers** because they are different compounds with the same molecular formula ($C_6H_{10}O_5$)_n. They are **stereoisomers** because only the three-dimensional arrangement of atoms is different.

How the six-membered rings are joined together has an enormous effect on the shape and properties of these carbohydrate molecules. Cellulose is composed of long chains held together by intermolecular hydrogen bonds, thus forming sheets that stack in an extensive three-dimensional

Figure 5.1
Starch and
cellulose—Two
common
carbohydrates




network. The axial–equatorial ring junction in starch creates chains that fold into a helix (Figure 5.2). Moreover, the human digestive system contains the enzyme necessary to hydrolyze starch by cleaving its axial C–O bond, but not an enzyme to hydrolyze the equatorial C–O bond in cellulose.

Thus, an **apparently minor difference in the three-dimensional arrangement of atoms confers very different properties on starch and cellulose.**

Figure 5.2

Three-dimensional structure of cellulose and starch



Problem 5.1 Cellulose is water insoluble, despite its many OH groups. Considering its three-dimensional structure, why do you think this is so?

5.2 The Two Major Classes of Isomers

Because an understanding of isomers is integral to the discussion of stereochemistry, let's begin with an overview of isomers.

- Isomers are different compounds with the same molecular formula.

There are two major classes of isomers: **constitutional isomers** and **stereoisomers**. *Constitutional (or structural) isomers differ in the way the atoms are connected to each other.* Constitutional isomers have:

- different IUPAC names;
- the same or different functional groups;
- different physical properties, so they are separable by physical techniques such as distillation; and
- different chemical properties. They behave differently or give different products in chemical reactions.


Stereoisomers differ only in the way atoms are oriented in space. Stereoisomers have identical IUPAC names (except for a prefix like *cis* or *trans*). Because they differ only in the three-dimensional arrangement of atoms, stereoisomers always have the same functional group(s).

A particular three-dimensional arrangement is called a **configuration**. Thus, stereoisomers differ in configuration. The *cis* and *trans* isomers in Section 4.13B and the biomolecules starch and cellulose in Section 5.1 are two examples of stereoisomers.

Figure 5.3 illustrates examples of both types of isomers. Most of Chapter 5 relates to the types and properties of stereoisomers.


Figure 5.3

A comparison of constitutional isomers and stereoisomers



Problem 5.2

Classify each pair of compounds as constitutional isomers or stereoisomers.




5.3 Looking Glass Chemistry—Chiral and Achiral Molecules



The dominance of right-handedness over left-handedness occurs in all races and cultures. Despite this fact, even identical twins can exhibit differences in hand preference. Pictured are Matthew (right-handed) and Zachary (left-handed), identical twin sons of the author.


Everything has a mirror image. What's important in chemistry is **whether a molecule is identical to or different from its mirror image.**

Some molecules are like hands. **Left and right hands are mirror images of each other, but they are not identical.** If you try to mentally place one hand inside the other hand, you can never superimpose either all the fingers, or the tops and palms. To *superimpose* an object on its mirror image means to align *all* parts of the object with its mirror image. With molecules, this means aligning all atoms and all bonds.



- A molecule (or object) that is *not* superimposable on its mirror image is said to be *chiral*.

Other molecules are like socks. **Two socks from a pair are mirror images that are superimposable.** One sock can fit inside another, aligning toes and heels, and tops and bottoms. A sock and its mirror image are *identical*.



- A molecule (or object) that is superimposable on its mirror image is said to be *achiral*.

Let's determine whether three molecules— H_2O , CH_2BrCl , and CHBrClF —are superimposable on their mirror images; that is, are H_2O , CH_2BrCl , and CHBrClF **chiral** or **achiral**?


To test chirality:

- Draw the molecule in three dimensions.
- Draw its mirror image.
- Try to align all bonds and atoms. To superimpose a molecule and its mirror image you can perform any rotation but **you cannot break bonds**.


Following this procedure, H_2O and CH_2BrCl are both **achiral** molecules because each molecule is superimposable on its mirror image.

The adjective **chiral** comes from the Greek *cheir*, meaning “hand.” Left and right hands are **chiral**: they are mirror images that do not superimpose on each other.

Few beginning students of organic chemistry can readily visualize whether a compound and its mirror image are superimposable by looking at drawings on a two-dimensional page. Molecular models can help a great deal in this process.



With CHBrClF , the result is different. The molecule (labeled **A**) and its mirror image (labeled **B**) are not superimposable. No matter how you rotate **A** and **B**, all the atoms never align. **CHBrClF is thus a chiral molecule**, and **A** and **B** are different compounds.



A and **B** are **stereoisomers** because they are isomers differing only in the three-dimensional arrangement of substituents. These stereoisomers are called **enantiomers**.

- *Enantiomers* are mirror images that are not superimposable.

CHBrClF contains a carbon atom bonded to four different groups. **A carbon atom bonded to four different groups is called a tetrahedral stereogenic center**. Most chiral molecules contain one or more stereogenic centers.

The general term *stereogenic center* refers to any site in a molecule at which the interchange of two groups forms a stereoisomer. **A carbon atom with four different groups is a tetrahedral**

Naming a carbon atom with four different groups is a topic that currently has no firm agreement among organic chemists. The IUPAC recommends the term *chirality center*, but the term has not gained wide acceptance among organic chemists since it was first suggested in 1996. Other terms in common use are chiral center, chiral carbon, asymmetric carbon, stereocenter, and stereogenic center, the term used in this text.

stereogenic center, because the interchange of two groups converts one enantiomer into another. We will learn about another type of stereogenic center in Section 8.2B.

We have now learned two related but different concepts, and it is necessary to distinguish between them.


- A molecule that is not superimposable on its mirror image is a *chiral molecule*.
- A carbon atom bonded to four different groups is a *stereogenic center*.

Molecules can contain zero, one, or more stereogenic centers.

- **With no stereogenic centers, a molecule generally is not chiral.** H_2O and CH_2BrCl have no stereogenic centers and are *achiral* molecules. (There are a few exceptions to this generalization, as we will learn in Section 17.5.)
- **With one tetrahedral stereogenic center, a molecule is always chiral.** CHBrClF is a *chiral* molecule containing *one* stereogenic center.
- **With two or more stereogenic centers, a molecule may or may not be chiral**, as we will learn in Section 5.8.

Problem 5.3

Draw the mirror image of each compound. Label each molecule as chiral or achiral.



When trying to distinguish between chiral and achiral compounds, keep in mind the following:

- A *plane of symmetry* is a mirror plane that cuts a molecule in half, so that one half of the molecule is a reflection of the other half.
- Achiral molecules usually contain a plane of symmetry but chiral molecules do not.

The achiral molecule CH_2BrCl has a plane of symmetry, but the chiral molecule CHBrClF does not.





Figure 5.4 summarizes the main facts about chirality we have learned thus far.

Figure 5.4
Summary: The basic principles of chirality

- Everything has a mirror image. The fundamental question is whether a molecule and its mirror image are superimposable.
- If a molecule and its mirror image are *not* superimposable, the molecule and its mirror image are *chiral*.
- The terms *stereogenic center* and *chiral molecule* are related but distinct. In general, a chiral molecule must have one or more stereogenic centers.
- The presence of a *plane of symmetry* makes a molecule achiral.

Problem 5.4 Draw in a plane of symmetry for each molecule.




Problem 5.5



When a right-handed shell is held in the right hand with the thumb pointing towards the wider end, the opening is on the right side.

A molecule is achiral if it has a plane of symmetry in *any* conformation. The given conformation of 2,3-dibromobutane does not have a plane of symmetry, but rotation around the C2–C3 bond forms a conformation that does have a plane of symmetry. Draw this conformation.




Stereochemistry may seem esoteric, but chirality pervades our very existence. On a molecular level, many biomolecules fundamental to life are chiral. On a macroscopic level, many naturally occurring objects possess handedness. Examples include chiral helical seashells shaped like right-handed screws, and plants such as honeysuckle that wind in a chiral left-handed helix. The human body is chiral, and hands, feet, and ears are not superimposable.

5.4 Stereogenic Centers


A necessary skill in the study of stereochemistry is the ability to locate and draw tetrahedral stereogenic centers.

5.4A Stereogenic Centers on Carbon Atoms That Are Not Part of a Ring

Recall from Section 5.3 that any carbon atom bonded to four different groups is a tetrahedral stereogenic center. To locate a stereogenic center, examine each *tetrahedral* carbon atom in a molecule, and look at the four *groups*—not the four *atoms*—bonded to it. CBrClFI has one stereogenic center because its central carbon atom is bonded to four different elements. 3-Bromohexane also has one stereogenic center because one carbon is bonded to H, Br, CH₂CH₃, and CH₂CH₂CH₃. We consider all atoms in a group as a *whole unit*, not just the atom directly bonded to the carbon in question.






Ephedrine is isolated from ma huang, an herb used to treat respiratory ailments in traditional Chinese medicine. Once a popular drug to promote weight loss and enhance athletic performance, ephedrine has now been linked to episodes of sudden death, heart attack, and stroke.





Always omit from consideration all C atoms that can't be tetrahedral stereogenic centers. These include:

- CH₂ and CH₃ groups (more than one H bonded to C)
- any *sp* or *sp*² hybridized C (less than four groups around C)

Larger organic molecules can have two, three, or even hundreds of stereogenic centers. **Pro-poxyphene** and **ephedrine** each contain two stereogenic centers, and **fructose**, a simple carbohydrate, has three.

**Sample Problem 5.1**



Locate the stereogenic center in each drug. Albuterol is a bronchodilator—that is, it widens airways—so it is used to treat asthma. Chloramphenicol is an antibiotic used extensively in developing countries because of its low cost.



Heteroatoms surrounded by four different groups are also stereogenic centers. Stereogenic N atoms are discussed in Chapter 25.

Solution

Omit all CH_2 and CH_3 groups and all doubly bonded (sp^2 hybridized) C's. In albuterol, one C has three CH_3 groups bonded to it, so it can be eliminated as well. When a molecule is drawn as a skeletal structure, draw in H atoms on tetrahedral C's to more clearly see the groups. This leaves one C in albuterol and two C's in chloramphenicol surrounded by four different groups, making them stereogenic centers.

**Problem 5.6**

Locate any stereogenic center in the given molecules. (Some compounds contain no stereogenic centers.)

- a. $\text{CH}_3\text{CH}_2\text{CH}(\text{Cl})\text{CH}_2\text{CH}_3$ d. $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
 b. $(\text{CH}_3)_3\text{CH}$ e. $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$
 c. $\text{CH}_3\text{CH}(\text{OH})\text{CH}=\text{CH}_2$ f. $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$


Problem 5.7

Locate the stereogenic centers in each molecule. Compounds may have one or more stereogenic centers.


- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ c.
- b. $(\text{CH}_3)_2\text{CHCH}_2\text{CH}(\text{NH}_2)\text{COOH}$ d.

Problem 5.8

The facts in Section 5.4A can be used to locate stereogenic centers in any molecule, no matter how complicated. Always look for carbons surrounded by four different groups. With this in mind, locate the four stereogenic centers in aliskiren, a drug introduced in 2007 for the treatment of hypertension.




5.4B Drawing a Pair of Enantiomers



- Any molecule with one tetrahedral stereogenic center is a chiral compound and exists as a pair of enantiomers.


2-Butanol, for example, has one stereogenic center. To draw both enantiomers, use the typical convention for depicting a tetrahedron: **place two bonds in the plane, one in front of the plane on a wedge, and one behind the plane on a dash.** Then, to form the first enantiomer **A**, arbitrarily place the four groups—H, OH, CH₃, and CH₂CH₃—on any bond to the stereogenic center.



Then, draw a mirror plane and arrange the substituents in the mirror image so that they are a reflection of the groups in the first molecule, forming **B**. No matter how **A** and **B** are rotated, it is impossible to align all of their atoms. Because **A** and **B** are mirror images and not superimposable, **A** and **B** are a pair of **enantiomers**. Two other pairs of enantiomers are drawn in Figure 5.5.

Problem 5.9

Locate the stereogenic center in each compound and draw both enantiomers.




Problem 5.10

The smallest chiral molecule ever prepared in the laboratory has one stereogenic center substituted by the three isotopes of hydrogen [hydrogen (H), deuterium (D), and tritium (T)] and a methyl group, forming CH_3CHDT (*Journal of the American Chemical Society*, **1997**, 119, 1818–1827). Draw the structure for the lowest molecular weight alkane (general molecular formula $\text{C}_n\text{H}_{2n+2}$, having only C and H and no isotopes) that contains a stereogenic center.

5.5 Stereogenic Centers in Cyclic Compounds

Stereogenic centers may also occur at carbon atoms that are part of a ring. To find stereogenic centers on ring carbons always draw the rings as flat polygons, and look for tetrahedral carbons that are bonded to four different groups, as usual. Each ring carbon is bonded to two other atoms in the ring, as well as two substituents attached to the ring. When the two substituents on the ring are *different*, we must compare the ring atoms equidistant from the atom in question.

Figure 5.5
Three-dimensional representations for pairs of enantiomers



In drawing a tetrahedron using solid lines, wedges, and dashes, always draw the two solid lines first; then draw the wedge and the dash on the *opposite side* of the solid lines.


If you draw the two solid lines down...

then add the wedge and dash above.



If you draw the two solid lines on the left...

then add the wedge and dash to the right.





Two enantiomers are *different* compounds. To convert one enantiomer to another you must **switch the position of two atoms**. This amounts to breaking bonds.

Although it is a potent teratogen (a substance that causes fetal abnormalities), thalidomide exhibits several beneficial effects. It is now prescribed under strict control for the treatment of Hansen's disease (leprosy) and certain forms of cancer.

Does methylcyclopentane have a stereogenic center? All of the carbon atoms are bonded to two or three hydrogen atoms except for C1, the ring carbon bonded to the methyl group. Next, compare the ring atoms and bonds on both sides equidistant from C1, and continue until a point of difference is reached, or until both sides meet, either at an atom or in the middle of a bond. In this case, there is no point of difference on either side, so C1 is bonded to identical alkyl groups that happen to be part of a ring. **C1, therefore, is not a stereogenic center.**



Is C1 a stereogenic center?




NO, C1 is **not** a stereogenic center.

With 3-methylcyclohexene, the result is different. All carbon atoms are bonded to two or three hydrogen atoms or are sp^2 hybridized except for C3, the ring carbon bonded to the methyl group. In this case, the atoms equidistant from C3 are different, so C3 is bonded to *different* alkyl groups in the ring. **C3 is therefore bonded to four different groups, making it a stereogenic center.**

Is C3 a stereogenic center?




Because 3-methylcyclohexene has one tetrahedral stereogenic center it is a chiral compound and exists as a pair of enantiomers.



Many biologically active compounds contain one or more stereogenic centers on ring carbons. For example, **thalidomide**, which contains one such stereogenic center, was used as a popular sedative and anti-nausea drug for pregnant women in Europe and Great Britain from 1959–1962.

Two enantiomers of thalidomide




Unfortunately thalidomide was sold as a mixture of its two enantiomers, and each of these stereoisomers has a different biological activity. This is a property not uncommon in chiral drugs, as we will see in Section 5.13. Although one enantiomer had the desired therapeutic effect, the other enantiomer was responsible for thousands of catastrophic birth defects in children born to women who took the drug during pregnancy. Thalidomide was never approved for use in the United States



Initial studies with taxol were carried out with material isolated from the bark of the Pacific yew tree, but stripping the bark killed these magnificent trees. Taxol can now be synthesized in four steps from a compound isolated from the needles of the common English yew tree.


due to the diligence of Frances Oldham Kelsey, a medical reviewing officer for the Food and Drug Administration, who insisted that the safety data on thalidomide were inadequate.

Sucrose and **taxol** are two useful molecules with several stereogenic centers at ring carbons. Identify the stereogenic centers in these more complicated compounds in exactly the same way, looking at one carbon at a time. **Sucrose**, with nine stereogenic centers on two rings, is the carbohydrate used as table sugar. **Taxol**, with 11 stereogenic centers, is an anticancer agent active against ovarian, breast, and some lung tumors.




Problem 5.11

Locate the stereogenic centers in each compound. A molecule may have zero, one, or more stereogenic centers. Gabapentin [part (d)] is used clinically to treat seizures and certain types of chronic pain. Gabapentin enacarbil [part (e)] is a related compound that is three times more potent.



Problem 5.12

Locate the stereogenic centers in each compound.



5.6 Labeling Stereogenic Centers with *R* or *S*


Naming enantiomers with the prefixes *R* or *S* is called the Cahn–Ingold–Prelog system after the three chemists who devised it.

Because enantiomers are two different compounds, we need to distinguish them by name. This is done by adding the prefix *R* or *S* to the IUPAC name of the enantiomer. To designate an enantiomer as *R* or *S*, first **assign a priority** (1, 2, 3, or 4) to each group bonded to the stereogenic center, and then use these priorities to label one enantiomer *R* and one *S*.

Rules Needed to Assign Priority

Rule 1 Assign priorities (1, 2, 3, or 4) to the atoms directly bonded to the stereogenic center in order of decreasing atomic number. The atom of *highest* atomic number gets the *highest* priority (1).


- In CHBrClF, priorities are assigned as follows: Br (1, highest) → Cl (2) → F (3) → H (4, lowest). In many molecules the lowest priority group will be H.




Rule 2 If two atoms on a stereogenic center are the *same*, assign priority based on the atomic number of the atoms bonded to these atoms. *One* atom of higher atomic number determines a higher priority.

- With 2-butanol, the O atom gets highest priority (1) and H gets lowest priority (4) using Rule 1. To assign priority (either 2 or 3) to the two C atoms, look at what atoms (other than the stereogenic center) are bonded to each C.

Following Rule 1:



Adding Rule 2:



- The order of priority of groups in 2-butanol is: -OH (1), -CH₂CH₃ (2), -CH₃ (3), and -H (4).
- If priority still cannot be assigned, continue along a chain until a point of difference is reached.


Rule 3 If two isotopes are bonded to the stereogenic center, assign priorities in order of decreasing mass number.

- In comparing the three isotopes of hydrogen, the order of priorities is:

	Mass number	Priority
T (tritium)	3 (1 proton + 2 neutrons)	1
D (deuterium)	2 (1 proton + 1 neutron)	2
H (hydrogen)	1 (1 proton)	3

Rule 4 To assign a priority to an atom that is part of a multiple bond, treat a multiply bonded atom as an equivalent number of singly bonded atoms.

- For example, the C of a C=O is considered to be bonded to two O atoms.



- Other common multiple bonds are drawn below.




Figure 5.6

Examples of assigning priorities to stereogenic centers




Figure 5.6 gives examples of priorities assigned to stereogenic centers.

Problem 5.13

Which group in each pair is assigned the *higher* priority?

- a. $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$ c. $-\text{H}$, $-\text{D}$ e. $-\text{CH}_2\text{CH}_2\text{Cl}$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$
 b. $-\text{I}$, $-\text{Br}$ d. $-\text{CH}_2\text{Br}$, $-\text{CH}_2\text{CH}_2\text{Br}$ f. $-\text{CH}_2\text{OH}$, $-\text{CHO}$

Problem 5.14

Rank the following groups in order of *decreasing* priority.


- a. $-\text{COOH}$, $-\text{H}$, $-\text{NH}_2$, $-\text{OH}$ c. $-\text{CH}_2\text{CH}_3$, $-\text{CH}_3$, $-\text{H}$, $-\text{CH}(\text{CH}_3)_2$
 b. $-\text{H}$, $-\text{CH}_3$, $-\text{Cl}$, $-\text{CH}_2\text{Cl}$ d. $-\text{CH}=\text{CH}_2$, $-\text{CH}_3$, $-\text{C}\equiv\text{CH}$, $-\text{H}$

R is derived from the Latin word *rectus* meaning “right” and *S* is from the Latin word *sinister* meaning “left.”

Once priorities are assigned to the four groups around a stereogenic center, we can use three steps to designate the center as either *R* or *S*.

HOW TO Assign *R* or *S* to a Stereogenic Center


Example Label each enantiomer as *R* or *S*.



two enantiomers of 2-butanol

Step [1] Assign priorities from 1 to 4 to each group bonded to the stereogenic center.


- The priorities for the four groups around the stereogenic center in 2-butanol were given in Rule 2, on page 171.



HOW TO, continued . . .


Step [2] Orient the molecule with the lowest priority group (4) *back* (on a *dash*), and visualize the relative positions of the remaining three groups (priorities 1, 2, and 3).

- For each enantiomer of 2-butanol, look toward the lowest priority group, drawn behind the plane, down the C–H bond.





Step [3] Trace a circle from priority group 1 → 2 → 3.

- If tracing the circle goes in the **clockwise** direction—to the right from the noon position—the isomer is named **R**.
- If tracing the circle goes in the **counterclockwise** direction—to the left from the noon position—the isomer is named **S**.




- The letters *R* or *S* precede the IUPAC name of the molecule. For the enantiomers of 2-butanol:




Sample Problem 5.2 Label the following compound as *R* or *S*.

Solution


[1] Assign priorities.



[2] Look down the C–H bond, toward the lowest priority group (H).




[3] Trace a circle, 1 → 2 → 3.


counterclockwise

 Answer: **S** isomer


How do you assign *R* or *S* to a molecule when the lowest priority group is not oriented toward the back, on a dashed line? You could rotate and flip the molecule until the lowest priority group is in the back, as shown in Figure 5.7; then follow the stepwise procedure for assigning the configuration. Or, if manipulating and visualizing molecules in three dimensions is difficult for you, try the procedure suggested in Sample Problem 5.3.

Sample Problem 5.3 Label the following compound as *R* or *S*.

Solution


In this problem, the lowest priority group (H) is oriented in **front** of, not behind, the page. To assign *R* or *S* in this case:

- **Switch** the position of the lowest priority group (H) with the group located behind the page (–CH₂CH₃).
- Determine *R* or *S* in the usual manner.
- **Reverse the answer.** Because we switched the position of two groups on the stereogenic center to begin with, and there are only two possibilities, the answer is **opposite** to the correct answer.


[1] Assign priorities.



[2] Switch groups 4 and 3.




[3] Trace a circle, 1 → 2 → 3, and reverse the answer.


 Answer: **R** isomer





counterclockwise

 It looks like an *S* isomer, but we must reverse the answer because we switched groups 3 and 4, *S* → *R*.


Figure 5.7
Examples: Orienting the lowest
priority group in back



Problem 5.15 Label each compound as *R* or *S*.



Problem 5.16 Draw both enantiomers of clopidogrel (trade name Plavix), a drug given to prevent the formation of blood clots in persons who have a history of stroke or coronary artery disease. Plavix is sold as a single enantiomer with the *S* configuration. Which enantiomer is Plavix?



5.7 Diastereomers


We have now seen many examples of compounds containing one tetrahedral stereogenic center. The situation is more complex for compounds with two stereogenic centers, because more stereoisomers are possible. Moreover, a molecule with two stereogenic centers *may or may not be chiral*.

- For n stereogenic centers, the maximum number of stereoisomers is 2^n .
- When $n = 1$, $2^1 = 2$. With one stereogenic center there are always two stereoisomers and they are **enantiomers**.
- When $n = 2$, $2^2 = 4$. With two stereogenic centers, the maximum number of stereoisomers is four, although sometimes there are *fewer* than four.


Problem 5.17 What is the maximum number of stereoisomers possible for a compound with: (a) three stereogenic centers; (b) eight stereogenic centers?

Let's illustrate a stepwise procedure for finding all possible stereoisomers using 2,3-dibromopentane.

In testing to see if one compound is superimposable on another, rotate atoms and flip the entire molecule, but **do not break any bonds**.




Add substituents around stereogenic centers with the bonds **eclipsed**, for easier visualization.




Don't forget, however, that the staggered arrangement is more stable.

HOW TO Find and Draw All Possible Stereoisomers for a Compound with Two Stereogenic Centers

Step [1] Draw one stereoisomer by arbitrarily arranging substituents around the stereogenic centers. Then draw its mirror image.



- Arbitrarily add the H, Br, CH₃, and CH₂CH₃ groups to the stereogenic centers, forming **A**. Then draw the mirror image **B** so that substituents in **B** are a reflection of the substituents in **A**.
- Determine whether **A** and **B** are superimposable by flipping or rotating one molecule to see if all the atoms align.
- If you have drawn the compound and the mirror image in the described manner, you only have to do two operations to see if the atoms align. Place **B** directly on top of **A** (either in your mind or use models); and, rotate **B** 180° and place it on top of **A** to see if the atoms align.




- In this case, the atoms of **A** and **B** do not align, making **A** and **B** nonsuperimposable mirror images—**enantiomers**. **A** and **B** are two of the four possible stereoisomers for 2,3-dibromopentane.

Step [2] Draw a third possible stereoisomer by switching the positions of any two groups on *one* stereogenic center only. Then draw its mirror image.

- Switching the positions of H and Br (or any two groups) on one stereogenic center of either **A** or **B** forms a new stereoisomer (labeled **C** in this example), which is different from both **A** and **B**. Then draw the mirror image of **C**, labeled **D**. **C** and **D** are nonsuperimposable mirror images—**enantiomers**. We have now drawn four stereoisomers for 2,3-dibromopentane, the maximum number possible.

HOW TO, continued . . .



There are only two types of stereoisomers: **Enantiomers** are stereoisomers that are mirror images. **Diastereomers** are stereoisomers that are not mirror images.


There are four stereoisomers for 2,3-dibromopentane: enantiomers **A** and **B**, and enantiomers **C** and **D**. What is the relationship between two stereoisomers like **A** and **C**? **A** and **C** represent the second broad class of stereoisomers, called **diastereomers**. **Diastereomers** are stereoisomers that are not mirror images of each other. **A** and **B** are diastereomers of **C** and **D**, and vice versa. Figure 5.8 summarizes the relationships between the stereoisomers of 2,3-dibromopentane.

Problem 5.18

Label the two stereogenic centers in each compound and draw all possible stereoisomers: (a) $\text{CH}_3\text{CH}_2\text{CH}(\text{Cl})\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$; (b) $\text{CH}_3\text{CH}(\text{Br})\text{CH}_2\text{CH}(\text{Cl})\text{CH}_3$.

5.8 Meso Compounds


Whereas 2,3-dibromopentane has two stereogenic centers and the maximum of four stereoisomers, **2,3-dibromobutane** has two stereogenic centers but fewer than the maximum number of stereoisomers.



With two stereogenic centers, the **maximum** number of stereoisomers = **4**.


To find and draw all the stereoisomers of 2,3-dibromobutane, follow the same stepwise procedure outlined in Section 5.7. Arbitrarily add the H, Br, and CH_3 groups to the stereogenic centers,

Figure 5.8
Summary: The four stereoisomers of 2,3-dibromopentane




- Pairs of enantiomers: **A** and **B**; **C** and **D**.
- Pairs of diastereomers: **A** and **C**; **A** and **D**; **B** and **C**; **B** and **D**.

forming one stereoisomer **A**, and then draw its mirror image **B**. **A** and **B** are nonsuperimposable mirror images—**enantiomers**.




To find the other two stereoisomers (if they exist), switch the position of two groups on *one* stereogenic center of *one* enantiomer only. In this case, switching the positions of H and Br on one stereogenic center of **A** forms **C**, which is different from both **A** and **B** and is thus a new stereoisomer.



However, the mirror image of **C**, labeled **D**, is superimposable on **C**, so **C** and **D** are identical. Thus, **C** is achiral, even though it has two stereogenic centers. **C** is a **meso compound**.

- A *meso compound* is an achiral compound that contains tetrahedral stereogenic centers.

C contains a **plane of symmetry**. **Meso compounds generally have a plane of symmetry**, so they possess two identical halves.




Because one stereoisomer of 2,3-dibromobutane is superimposable on its mirror image, there are only three stereoisomers and not four, as summarized in Figure 5.9.

Problem 5.19


Draw all the possible stereoisomers for each compound and label pairs of enantiomers and diastereomers: (a) $\text{CH}_3\text{CH}(\text{OH})\text{CH}(\text{OH})\text{CH}_3$; (b) $\text{CH}_3\text{CH}(\text{OH})\text{CH}(\text{Cl})\text{CH}_3$.

Figure 5.9
Summary: The three stereoisomers of 2,3-dibromobutane




- Pair of enantiomers: **A** and **B**.
- Pairs of diastereomers: **A** and **C**; **B** and **C**.

Problem 5.20 Which compounds are meso compounds?




Problem 5.21 Draw a meso compound for each of the following molecules.



5.9 R and S Assignments in Compounds with Two or More Stereogenic Centers

When a compound has more than one stereogenic center, the *R* or *S* configuration must be assigned to each of them. In the stereoisomer of 2,3-dibromopentane drawn here, C2 has the *S* configuration and C3 has the *R*, so the complete name of the compound is (2*S*,3*R*)-2,3-dibromopentane.



Complete name:
(2*S*,3*R*)-2,3-dibromopentane

one stereoisomer of 2,3-dibromopentane



Sorbitol (Problem 5.24) occurs naturally in some berries and fruits. It is used as a substitute sweetener in sugar-free—that is, sucrose-free—candy and gum.

R,S configurations can be used to determine whether two compounds are identical, enantiomers, or diastereomers.

- Identical compounds have the *same* *R,S* designations at every tetrahedral stereogenic center.
- Enantiomers have *exactly opposite* *R,S* designations.
- Diastereomers have the *same* *R,S* designation for at least one stereogenic center and the *opposite* for at least one of the other stereogenic centers.


For example, if a compound has two stereogenic centers, both with the *R* configuration, then its enantiomer is *S,S* and the diastereomers are either *R,S* or *S,R*.

Problem 5.22 If the two stereogenic centers of a compound are *R,S* in configuration, what are the *R,S* assignments for its enantiomer and two diastereomers?


- The trans isomer **B** is *not* superimposable on its mirror image, labeled **C**, making **B** and **C** different compounds. Thus, **B** and **C** are **enantiomers**.

Because one stereoisomer of 1,3-dibromocyclopentane is superimposable on its mirror image, there are only three stereoisomers, not four. **A** is an achiral meso compound and **B** and **C** are a pair of chiral enantiomers. **A** and **B** are diastereomers, as are **A** and **C**.

Problem 5.25 Which of the following cyclic molecules are *meso* compounds?



Problem 5.26 Draw all possible stereoisomers for each compound. Label pairs of enantiomers and diastereomers.



5.11 Isomers—A Summary

Before moving on to other aspects of stereochemistry, take the time to review Figures 5.10 and 5.11. Keep in mind the following facts, and use Figure 5.10 to summarize the types of isomers.

- There are two major classes of isomers: constitutional isomers and stereoisomers.
- There are only two kinds of stereoisomers: enantiomers and diastereomers.

Then, to determine the relationship between two nonidentical molecules, refer to the flowchart in Figure 5.11.

Problem 5.27 State how each pair of compounds is related. Are they enantiomers, diastereomers, constitutional isomers, or identical?




Figure 5.10
Summary—Types of isomers





Figure 5.11

Determining the relationship between two nonidentical molecules



5.12 Physical Properties of Stereoisomers


Recall from Section 5.2 that constitutional isomers have different physical and chemical properties. How, then, do the physical and chemical properties of enantiomers compare?

- The chemical and physical properties of two enantiomers are *identical* except in their interaction with *chiral* substances.

5.12A Optical Activity


Two enantiomers have identical physical properties—melting point, boiling point, solubility—except for how they interact with plane-polarized light.

What is plane-polarized light? Ordinary light consists of electromagnetic waves that oscillate in all planes perpendicular to the direction in which the light travels. Passing light through a polarizer allows light in only one plane to come through. This is **plane-polarized light** (or simply **polarized light**), and it has an electric vector that oscillates in a single plane.




A **polarimeter** is an instrument that allows plane-polarized light to travel through a sample tube containing an organic compound. After the light exits the sample tube, an analyzer slit is rotated to determine the direction of the plane of the polarized light exiting the sample tube. There are two possible results.

With **achiral compounds**, the light exits the sample tube *unchanged*, and the plane of the polarized light is in the same position it was before entering the sample tube. A **compound that does not change the plane of polarized light is said to be optically inactive**.




With **chiral compounds**, the plane of the polarized light is rotated through an angle α . The angle α , measured in degrees ($^\circ$), is called the **observed rotation**. A compound that rotates the plane of polarized light is said to be **optically active**.



For example, the achiral compound CH_2BrCl is optically inactive, whereas a single enantiomer of CHBrClF , a chiral compound, is optically active.


The rotation of polarized light can be in the **clockwise** or **counterclockwise** direction.

- If the rotation is *clockwise* (to the right from the noon position), the compound is called *dextrorotatory*. The rotation is labeled *d* or (+).
- If the rotation is *counterclockwise* (to the left from noon), the compound is called *levorotatory*. The rotation is labeled *l* or (–).



No relationship exists between the *R* and *S* prefixes that designate configuration and the (+) and (–) designations indicating optical rotation. For example, the *S* enantiomer of lactic acid is dextrorotatory (+), whereas the *S* enantiomer of glyceraldehyde is levorotatory (–).

How does the rotation of two enantiomers compare?



- Two enantiomers rotate plane-polarized light to an equal extent but in the opposite direction.

Thus, if enantiomer **A** rotates polarized light $+5^\circ$, then the same concentration of enantiomer **B** rotates it -5° .

5.12B Racemic Mixtures

What is the observed rotation of an equal amount of two enantiomers? Because **two enantiomers rotate plane-polarized light to an equal extent but in opposite directions, the rotations cancel**, and no rotation is observed.

- An equal amount of two enantiomers is called a *racemic mixture* or a *racemate*. A racemic mixture is optically inactive.

Table 5.1 The Physical Properties of Enantiomers A and B Compared

Property	A alone	B alone	Racemic A + B
Melting point	identical to B	identical to A	may be different from A and B
Boiling point	identical to B	identical to A	may be different from A and B
Optical rotation	equal in magnitude but opposite in sign to B	equal in magnitude but opposite in sign to A	0°

Besides optical rotation, other physical properties of a racemate are not readily predicted. The melting point and boiling point of a racemic mixture are not necessarily the same as either pure enantiomer, and this fact is not easily explained. The physical properties of two enantiomers and their racemic mixture are summarized in Table 5.1.

5.12C Specific Rotation

The observed rotation depends on the number of chiral molecules that interact with polarized light. This in turn depends on the concentration of the sample and the length of the sample tube. To standardize optical rotation data, the quantity **specific rotation** ($[\alpha]$) is defined using a specific sample tube length (usually 1 dm), concentration, temperature (25 °C), and wavelength (589 nm, the D line emitted by a sodium lamp).

$$\text{specific rotation} = [\alpha] = \frac{\alpha}{l \times c}$$


α = observed rotation (°)
 l = length of sample tube (dm)
 c = concentration (g/mL)

$$\left[\begin{array}{l} \text{dm} = \text{decimeter} \\ 1 \text{ dm} = 10 \text{ cm} \end{array} \right]$$

Specific rotations are physical constants just like melting points or boiling points, and are reported in chemical reference books for a wide variety of compounds.

Problem 5.28

The amino acid (S)-alanine has the physical characteristics listed under the structure.



(S)-alanine
 $[\alpha] = +8.5$
 $\text{mp} = 297 \text{ }^\circ\text{C}$

- What is the melting point of (R)-alanine?
- How does the melting point of a racemic mixture of (R)- and (S)-alanine compare to the melting point of (S)-alanine?
- What is the specific rotation of (R)-alanine, recorded under the same conditions as the reported rotation of (S)-alanine?
- What is the optical rotation of a racemic mixture of (R)- and (S)-alanine?
- Label each of the following as optically active or inactive: a solution of pure (S)-alanine; an equal mixture of (R)- and (S)-alanine; a solution that contains 75% (S)- and 25% (R)-alanine.

Problem 5.29

A natural product was isolated in the laboratory, and its observed rotation was +10° when measured in a 1 dm sample tube containing 1.0 g of compound in 10 mL of water. What is the specific rotation of this compound?

5.12D Enantiomeric Excess

Sometimes in the laboratory we have neither a pure enantiomer nor a racemic mixture, but rather a mixture of two enantiomers in which one enantiomer is present in excess of the other. The **enantiomeric excess** (*ee*), also called the **optical purity**, tells how much more there is of one enantiomer.

- Enantiomeric excess = $ee = \% \text{ of one enantiomer} - \% \text{ of the other enantiomer}$.

Enantiomeric excess tells how much one enantiomer is present in excess of the racemic mixture. For example, if a mixture contains 75% of one enantiomer and 25% of the other, the enantiomeric excess is $75\% - 25\% = 50\%$. There is a 50% excess of one enantiomer over the racemic mixture.

Problem 5.30 What is the *ee* for each of the following mixtures of enantiomers **A** and **B**?
 a. 95% **A** and 5% **B** b. 85% **A** and 15% **B**

Knowing the *ee* of a mixture makes it possible to calculate the amount of each enantiomer present, as shown in Sample Problem 5.4.

Sample Problem 5.4 If the enantiomeric excess is 95%, how much of each enantiomer is present?

Solution

Label the two enantiomers **A** and **B** and assume that **A** is in excess. A 95% *ee* means that the solution contains an excess of 95% of **A**, and 5% of the racemic mixture of **A** and **B**. Because a racemic mixture is an equal amount of both enantiomers, it has 2.5% of **A** and 2.5% of **B**.

- Total amount of **A** = 95% + 2.5% = 97.5%
- Total amount of **B** = 2.5% (or 100% – 97.5%)

Problem 5.31 For the given *ee* values, calculate the percentage of each enantiomer present.
 a. 90% *ee* b. 99% *ee* c. 60% *ee*

The enantiomeric excess can also be calculated if two quantities are known—the specific rotation $[\alpha]$ of a mixture and the specific rotation $[\alpha]$ of a pure enantiomer.

$$ee = \frac{[\alpha] \text{ mixture}}{[\alpha] \text{ pure enantiomer}} \times 100\%$$


Sample Problem 5.5 Pure cholesterol has a specific rotation of -32 . A sample of cholesterol prepared in the lab had a specific rotation of -16 . What is the enantiomeric excess of this sample of cholesterol?

Solution

Calculate the *ee* of the mixture using the given formula.

$$ee = \frac{[\alpha] \text{ mixture}}{[\alpha] \text{ pure enantiomer}} \times 100\% = \frac{-16}{-32} \times 100\% = 50\% ee$$

Problem 5.32 Pure MSG, a common flavor enhancer, exhibits a specific rotation of $+24$. (a) Calculate the *ee* of a solution whose $[\alpha]$ is $+10$. (b) If the *ee* of a solution of MSG is 80%, what is $[\alpha]$ for this solution?



MSG
monosodium glutamate


Problem 5.33 (S)-Lactic acid has a specific rotation of $+3.8$. (a) If the *ee* of a solution of lactic acid is 60%, what is $[\alpha]$ for this solution? (b) How much of the dextrorotatory and levorotatory isomers does the solution contain?

5.12E The Physical Properties of Diastereomers

Diastereomers are not mirror images of each other, and as such, **their physical properties are different, including optical rotation**. Figure 5.12 compares the physical properties of the three stereoisomers of tartaric acid, consisting of a meso compound that is a diastereomer of a pair of enantiomers.

Figure 5.12

The physical properties of the three stereoisomers of tartaric acid




Property	A	B	C	A + B (1:1)
melting point (°C)	171	171	146	206
solubility (g/100 mL H ₂ O)	139	139	125	139
[α]	+13	-13	0	0
<i>R,S</i> designation	<i>R,R</i>	<i>S,S</i>	<i>R,S</i>	—
<i>d,l</i> designation	<i>d</i>	<i>l</i>	none	<i>d,l</i>

- The physical properties of **A** and **B** differ from their diastereomer **C**.
- The physical properties of a racemic mixture of **A** and **B** (last column) can also differ from either enantiomer and diastereomer **C**.
- **C** is an achiral meso compound, so it is optically inactive; [α] = 0.

Whether the physical properties of a set of compounds are the same or different has practical applications in the lab. Physical properties characterize a compound's physical state, and two compounds can usually be separated only if their physical properties are different.

- Because two enantiomers have identical physical properties, they cannot be separated by common physical techniques like distillation.
- Diastereomers and constitutional isomers have different physical properties, and therefore they can be separated by common physical techniques.

Problem 5.34 Compare the physical properties of the three stereoisomers of 1,3-dimethylcyclopentane.



- How do the boiling points of **A** and **B** compare? What about **A** and **C**?
- Characterize a solution of each of the following as optically active or optically inactive: pure **A**; pure **B**; pure **C**; an equal mixture of **A** and **B**; an equal mixture of **A** and **C**.
- A reaction forms a 1:1:1 mixture of **A**, **B**, and **C**. If this mixture is distilled, how many fractions would be obtained? Which fractions would be optically active and which would be optically inactive?

5.13 Chemical Properties of Enantiomers

When two enantiomers react with an achiral reagent, they react at the same rate, but when they react with a chiral, non-racemic reagent, they react at different rates.

- Two enantiomers have exactly the same chemical properties except for their reaction with chiral, non-racemic reagents.

For an everyday analogy, consider what happens when you are handed an achiral object like a pen and a chiral object like a right-handed glove. Your left and right hands are enantiomers, but they can both hold the achiral pen in the same way. With the glove, however, only your right hand can fit inside it, not your left.


We will examine specific reactions of chiral molecules with both chiral and achiral reagents later in this text. Here, we examine two more general applications.

5.13A Chiral Drugs


A living organism is a sea of chiral molecules. Many drugs are chiral, and often they must interact with a chiral receptor or a chiral enzyme to be effective. One enantiomer of a drug may effectively treat a disease whereas its mirror image may be ineffective. Alternatively, one enantiomer may trigger one biochemical response and its mirror image may elicit a totally different response.

For example, the drugs ibuprofen and fluoxetine each contain one stereogenic center, and thus exist as a pair of enantiomers, only one of which exhibits biological activity. (**S**)-**Ibuprofen** is the active component of the anti-inflammatory agents Motrin and Advil, and (**R**)-**fluoxetine** is the active component in the antidepressant Prozac.

Although (*R*)-ibuprofen shows no anti-inflammatory activity itself, it is slowly converted to the *S* enantiomer in vivo.




(**S**)-**ibuprofen**
anti-inflammatory agent



(**R**)-**fluoxetine**
antidepressant

The *S* enantiomer of **naproxen**, the molecule that introduced Chapter 5, is an active anti-inflammatory agent, but the *R* enantiomer is a harmful liver toxin. Changing the orientation of two substituents to form a mirror image can thus alter biological activity to produce an undesirable side effect in the other enantiomer.



(**S**)-**naproxen**
anti-inflammatory agent

(**R**)-**naproxen**
liver toxin

For more examples of two enantiomers that exhibit very different biochemical properties, see *Journal of Chemical Education*, **1996**, 73, 481–484.

If a chiral drug could be sold as a single active enantiomer, it should be possible to use smaller doses with fewer side effects. Many chiral drugs continue to be sold as racemic mixtures, however, because it is more difficult and therefore more costly to obtain a single enantiomer. An enantiomer is not easily separated from a racemic mixture because the two enantiomers have the same physical properties. In Chapter 12 we will study a reaction that can form a single active enantiomer, an important development in making chiral drugs more readily available.

Recent rulings by the Food and Drug Administration have encouraged the development of so-called **racemic switches**, the patenting and marketing of a single enantiomer that was originally sold as a racemic mixture. To obtain a new patent on a single enantiomer, however, a company must show evidence that it provides significant benefit over the racemate.


5.13B Enantiomers and the Sense of Smell

Research suggests that the odor of a particular molecule is determined more by its shape than by the presence of a particular functional group. For example, hexachloroethane (Cl_3CCCl_3) and cyclooctane have no obvious structural similarities, but they both have a camphor-like odor, a fact attributed to their similar spherical shape. Each molecule binds to spherically shaped olfactory receptors present on the nerve endings in the nasal passage, resulting in similar odors (Figure 5.13).

Because enantiomers interact with chiral smell receptors, some enantiomers have different odors. There are a few well-characterized examples of this phenomenon in nature. For example,


Figure 5.13

The shape of molecules and the sense of smell



Cyclooctane and other molecules similar in shape bind to a particular olfactory receptor on the nerve cells that lie at the top of the nasal passage. Binding results in a nerve impulse that travels to the brain, which interprets impulses from particular receptors as specific odors.

(*S*)-carvone is responsible for the odor of caraway, whereas (*R*)-carvone is responsible for the odor of spearmint.



These examples demonstrate that understanding the three-dimensional structure of a molecule is very important in organic chemistry.

KEY CONCEPTS


Stereochemistry

Isomers Are Different Compounds with the Same Molecular Formula (5.2, 5.11).

- [1] **Constitutional isomers**—isomers that differ in the way the atoms are connected to each other. They have:
 - different IUPAC names
 - the same or different functional groups
 - different physical and chemical properties
- [2] **Stereoisomers**—isomers that differ only in the way atoms are oriented in space. They have the same functional group and the same IUPAC name except for prefixes such as *cis*, *trans*, *R*, and *S*.
 - **Enantiomers**—stereoisomers that are nonsuperimposable mirror images of each other (5.4).
 - **Diastereomers**—stereoisomers that are not mirror images of each other (5.7).

Some Basic Principles

- When a compound and its mirror image are **superimposable**, they are **identical achiral compounds**. When a compound has a plane of symmetry in one conformation, the compound is achiral (5.3).
- When a compound and its mirror image are **not superimposable**, they are **different chiral compounds** called **enantiomers**. A chiral compound has no plane of symmetry in any conformation (5.3).
- A **tetrahedral stereogenic center** is a carbon atom bonded to four different groups (5.4, 5.5).
- For n **stereogenic centers**, the maximum number of stereoisomers is 2^n (5.7).



Optical Activity Is the Ability of a Compound to Rotate Plane-Polarized Light (5.12).

- An optically active solution contains a chiral compound.
- An optically inactive solution contains one of the following:
 - an achiral compound with no stereogenic centers
 - a meso compound—an achiral compound with two or more stereogenic centers
 - a racemic mixture—an equal amount of two enantiomers

The Prefixes *R* and *S* Compared with *d* and *l*

The prefixes *R* and *S* are labels used in nomenclature. Rules on assigning *R,S* are found in Section 5.6.

- An enantiomer has every stereogenic center opposite in configuration. If a compound with two stereogenic centers has the *R,R* configuration, its enantiomer has the *S,S* configuration.
- A diastereomer of this same compound has either the *R,S* or *S,R* configuration; one stereogenic center has the same configuration and one is opposite.

The prefixes *d* (or +) and *l* (or -) tell the direction a compound rotates plane-polarized light (5.12).

- Dextrorotatory (*d* or +) compounds rotate polarized light clockwise.
- Levorotatory (*l* or -) compounds rotate polarized light counterclockwise.
- There is no relation between whether a compound is *R* or *S* and whether it is *d* or *l*.

The Physical Properties of Isomers Compared (5.12)

Type of isomer	Physical properties
Constitutional isomers	Different
Enantiomers	Identical except for the direction polarized light is rotated
Diastereomers	Different
Racemic mixture	Possibly different from either enantiomer

Equations

- Specific rotation (5.12C):

$$\text{specific rotation} = [\alpha] = \frac{\alpha}{l \times c}$$

α = observed rotation ($^{\circ}$)
 l = length of sample tube (dm)
 c = concentration (g/mL)

$$\left[\begin{array}{l} \text{dm = decimeter} \\ 1 \text{ dm} = 10 \text{ cm} \end{array} \right]$$


- Enantiomeric excess (5.12D):

$$\begin{aligned} ee &= \% \text{ of one enantiomer} - \% \text{ of the other enantiomer} \\ &= \frac{[\alpha] \text{ mixture}}{[\alpha] \text{ pure enantiomer}} \times 100\% \end{aligned}$$

PROBLEMS


Constitutional Isomers versus Stereoisomers

5.35 Label each pair of compounds as constitutional isomers, stereoisomers, or not isomers of each other.




Mirror Images and Chirality


5.36 Draw the mirror image of each compound, and label the compound as chiral or achiral.



5.37 Determine if each compound is identical to or an enantiomer of **A**.




5.38 Indicate a plane of symmetry for each molecule that contains one. Some molecules require rotation around a carbon-carbon bond to see the plane of symmetry.





Finding and Drawing Stereogenic Centers

5.39 Locate the stereogenic center(s) in each compound. A molecule may have zero, one, or more stereogenic centers.



5.40 Draw the eight constitutional isomers having molecular formula $\text{C}_5\text{H}_{11}\text{Cl}$. Label any stereogenic centers.







5.41 Draw both enantiomers for each biologically active compound.








5.42 Draw the lowest molecular weight chiral compound that contains only C, H, and O and fits each description: (a) an acyclic alcohol; (b) a ketone; (c) a cyclic ether.

Nomenclature









5.43 Which group in each pair is assigned the higher priority in *R,S* nomenclature?




5.44 Rank the following groups in order of decreasing priority.






5.45 Label each stereogenic center as *R* or *S*.



5.46 Draw the structure for each compound.






5.47 Give the IUPAC name for each compound, including the *R,S* designation for each stereogenic center.




5.48 Draw the two enantiomers for the amino acid leucine, HOOCCH(NH₂)CH₂CH(CH₃)₂, and label each enantiomer as *R* or *S*. Only the *S* isomer exists in nature, and it has a bitter taste. Its enantiomer, however, is sweet.


5.49 Label the stereogenic center(s) in each drug as *R* or *S*. L-Dopa is used to treat Parkinson's disease (Chapter 1). Ketamine is an anesthetic. Enalapril belongs to a class of drugs called ACE inhibitors, which are used to lower blood pressure.



5.50 Methylphenidate (trade name: Ritalin) is prescribed for attention deficit hyperactivity disorder (ADHD). Ritalin is a mixture of *R,R* and *S,S* isomers, even though only the *R,R* isomer is active in treating ADHD. (The single *R,R* enantiomer, called dexmethylphenidate, is now sold under the trade name Focalin.) Draw the structure of the *R,R* and *S,S* isomers of methylphenidate.



- 5.51** The shrub ma huang (Section 5.4A) contains two biologically active stereoisomers—ephedrine and pseudoephedrine—with two stereogenic centers as shown in the given structure. Ephedrine is one component of a once popular combination drug used by body builders to increase energy and alertness, while pseudoephedrine is a nasal decongestant.






isolated from ma huang


- Draw the structure of naturally occurring (–)-ephedrine, which has the $1R,2S$ configuration.
- Draw the structure of naturally occurring (+)-pseudoephedrine, which has the $1S,2S$ configuration.
- How are ephedrine and pseudoephedrine related?
- Draw all other stereoisomers of ephedrine and pseudoephedrine and give the R,S designation for all stereogenic centers.
- How is each compound drawn in part (d) related to ephedrine?

Compounds with More Than One Stereogenic Center

- 5.52** Locate the stereogenic centers in each drug.







- 5.53** What is the maximum number of stereoisomers possible for each compound?






- 5.54** Draw all possible stereoisomers for each compound. Label pairs of enantiomers and diastereomers. Label any meso compound.



- 5.55** Draw the enantiomer and a diastereomer for each compound.








- 5.56** Draw all possible stereoisomers for each cycloalkane. Label pairs of enantiomers and diastereomers. Label any meso compound.



- 5.57** Draw all possible constitutional and stereoisomers for a compound of molecular formula C_6H_{12} having a cyclobutane ring and two methyl groups as substituents. Label each compound as chiral or achiral.


- 5.58** Explain each statement by referring to compounds **A–E**.




- A** has a mirror image but no enantiomer.
- B** has an enantiomer and no diastereomer.
- C** has both an enantiomer and a diastereomer.
- D** has a diastereomer but no enantiomer.
- E** has a diastereomer but no enantiomer.

Comparing Compounds: Enantiomers, Diastereomers, and Constitutional Isomers


5.59 How is each compound related to the simple sugar D-erythrose? Is it an enantiomer, diastereomer, or identical?




5.60 Consider Newman projections (A–D) for four-carbon carbohydrates. How is each pair of compounds related: (a) A and B; (b) A and C; (c) A and D; (d) C and D? Choose from identical molecules, enantiomers, or diastereomers.



5.61 How is compound A related to compounds B–E? Choose from enantiomers, diastereomers, constitutional isomers, or identical molecules.




5.62 How are the compounds in each pair related to each other? Are they identical, enantiomers, diastereomers, constitutional isomers, or not isomers of each other?




Physical Properties of Isomers

5.63 Drawn are four isomeric dimethylcyclopropanes.




- How are the compounds in each pair related (enantiomers, diastereomers, constitutional isomers): **A and B**; **A and C**; **B and C**; **C and D**?
- Label each compound as chiral or achiral.
- Which compounds, alone, would be optically active?
- Which compounds have a plane of symmetry?
- How do the boiling points of the compounds in each pair compare: **A and B**; **B and C**; **C and D**?
- Which of the compounds are meso compounds?
- Would an equal mixture of compounds **C** and **D** be optically active? What about an equal mixture of **B** and **C**?

5.64 The $[\alpha]$ of pure quinine, an antimalarial drug, is -165 .



- Calculate the ee of a solution with the following $[\alpha]$ values: -50 , -83 , and -120 .
- For each ee, calculate the percent of each enantiomer present.
- What is $[\alpha]$ for the enantiomer of quinine?
- If a solution contains 80% quinine and 20% of its enantiomer, what is the ee of the solution?
- What is $[\alpha]$ for the solution described in part (d)?


5.65 Amygdalin, a compound isolated from the pits of apricots, peaches, and wild cherries, is sometimes called laetrile. Although it has no known therapeutic value, amygdalin has been used as an unsanctioned anticancer drug both within and outside of the United States. One hydrolysis product formed from amygdalin is mandelic acid, used in treating common skin problems caused by photo-aging and acne.



- How many stereogenic centers are present in amygdalin? What is the maximum number of stereoisomers possible?
- Draw both enantiomers of mandelic acid and label each stereogenic center as *R* or *S*.
- Pure (*R*)-mandelic acid has a specific rotation of -154 . If a sample contains 60% of the *R* isomer and 40% of its enantiomer, what is $[\alpha]$ of this solution?
- Calculate the ee of a solution of mandelic acid having $[\alpha] = +50$. What is the percentage of each enantiomer present?


General Problems

5.66 Artemisinin and mefloquine are widely used antimalarial drugs.



- Locate the stereogenic centers in both drugs.
- Label each stereogenic center in mefloquine as *R* or *S*.
- What is the maximum number of stereoisomers possible for artemisinin?
- How are the N atoms in mefloquine hybridized?
- Can two molecules of artemisinin intermolecularly hydrogen bond to each other?
- What product is formed when mefloquine is treated with HCl?




- 5.67 Saquinavir (trade name Invirase) belongs to a class of drugs called protease inhibitors, which are used to treat HIV (human immunodeficiency virus).




- Locate all stereogenic centers in saquinavir, and label each stereogenic center as *R* or *S*.
 - Draw the enantiomer of saquinavir.
 - Draw a diastereomer of saquinavir.
 - Draw a constitutional isomer that contains at least one different functional group.
- 5.68 Salicin is an analgesic isolated from willow bark.
-
- salicin
- Convert the given skeletal structure to a representation that shows the more stable chair form of the six-membered ring.
 - Draw a diastereomer of salicin at C1 and label each substituent on the six-membered ring as axial or equatorial.
 - Draw the enantiomer of salicin.

Challenge Problems


- 5.69 A limited number of chiral compounds having no stereogenic centers exist. For example, although **A** is achiral, constitutional isomer **B** is chiral. Make models and explain this observation. Compounds containing two double bonds that share a single carbon atom are called *allenes*. Locate the allene in the antibiotic mycomycin and decide whether mycomycin is chiral or achiral.



- 5.70 a. Locate all the tetrahedral stereogenic centers in discodermolide, a natural product isolated from the Caribbean marine sponge *Discodermia dissoluta*. Discodermolide is a potent tumor inhibitor, and shows promise as a drug for treating colon, ovarian, and breast cancers.
- b. Certain carbon-carbon double bonds can also be stereogenic centers. With reference to the definition in Section 5.3, explain how this can occur, and then locate the three additional stereogenic centers in discodermolide.
- c. Considering all stereogenic centers, what is the maximum number of stereoisomers possible for discodermolide?



- 5.71 An acid-base reaction of (*R*)-*sec*-butylamine with a racemic mixture of 2-phenylpropanoic acid forms two products having different melting points and somewhat different solubilities. Draw the structure of these two products. Assign *R* and *S* to any stereogenic centers in the products. How are the two products related? Choose from enantiomers, diastereomers, constitutional isomers, or not isomers.



6

Understanding Organic Reactions

- 6.1 Writing equations for organic reactions
- 6.2 Kinds of organic reactions
- 6.3 Bond breaking and bond making
- 6.4 Bond dissociation energy
- 6.5 Thermodynamics
- 6.6 Enthalpy and entropy
- 6.7 Energy diagrams
- 6.8 Energy diagram for a two-step reaction mechanism
- 6.9 Kinetics
- 6.10 Catalysts
- 6.11 Enzymes



Isooctane, a component of petroleum, and **glucose**, a simple sugar formed from starch during digestion, are very different organic molecules that share a common feature. On oxidation, both compounds release a great deal of energy. Isooctane is burned in gasoline to power automobiles, and glucose is metabolized in the body to provide energy for exercise. In Chapter 6, we learn about these energy changes that accompany chemical reactions.

Why do certain reactions occur when two compounds are mixed together whereas others do not? To answer this question we must learn how and why organic compounds react.

Reactions are at the heart of organic chemistry. An understanding of chemical processes has made possible the conversion of natural substances into new compounds with different, and sometimes superior, properties. Aspirin, ibuprofen, nylon, and polyethylene are all products of chemical reactions between substances derived from petroleum.

Reactions are difficult to learn when each reaction is considered a unique and isolated event. *Avoid this tendency.* Virtually all chemical reactions are woven together by a few basic themes. After we learn the general principles, specific reactions then fit neatly into a general pattern.

In our study of organic reactions we will begin with the functional groups, looking for electron-rich and electron-deficient sites, and bonds that might be broken easily. These reactive sites give us a clue as to the general type of reaction a particular class of compound undergoes. Finally, we will learn about how a reaction occurs. Does it occur in one step or in a series of steps? Understanding the details of an organic reaction allows us to determine when it might be used in preparing interesting and useful organic compounds.

6.1 Writing Equations for Organic Reactions


Often the solvent and temperature of a reaction are omitted from chemical equations, to further focus attention on the main substances involved in the reaction.

Solvent. Most organic reactions take place in a **liquid solvent**. Solvents solubilize key reaction components and serve as heat reservoirs to maintain a given temperature. Chapter 7 presents the two major types of reaction solvents and how they affect substitution reactions.

Like other reactions, equations for organic reactions are usually drawn with a single reaction arrow (\rightarrow) between the starting material and product, but other conventions make these equations look different from those encountered in general chemistry.


The **reagent**, the chemical substance with which an organic compound reacts, is sometimes drawn on the left side of the equation with the other reactants. At other times, the reagent is drawn above the reaction arrow itself, to focus attention on the organic starting material by itself on the left side. The solvent and temperature of a reaction may be added above or below the arrow. **The symbols “ $h\nu$ ” and “ Δ ” are used for reactions that require light or heat, respectively.** Figure 6.1 presents an organic reaction in different ways.

When two sequential reactions are carried out without drawing any intermediate compound, the steps are usually numbered above or below the reaction arrow. This convention signifies that the first step occurs *before* the second, and the reagents are added *in sequence*, not at the same time.



In this equation only the organic product is drawn on the right side of the arrow. Although the reagent CH_3MgBr contains both Mg and Br, these elements do not appear in the organic product, and they are often omitted on the product side of the equation. These elements have not disappeared. They are part of an inorganic by-product (HOMgBr in this case), and are often of little interest to an organic chemist.


Figure 6.1
Different ways of writing organic reactions



6.2C Addition Reactions



- **Addition** is a reaction in which elements are added to a starting material.

A general addition reaction




In an addition reaction, new groups X and Y are added to a starting material. A π bond is broken and two σ bonds are formed.

Examples







A summary of the general types of organic reactions is given in Appendix G.

Addition and elimination reactions are exactly opposite. A π bond is *formed* in elimination reactions, whereas a π bond is *broken* in addition reactions.



Problem 6.1 Classify each transformation as substitution, elimination, or addition.



Problem 6.2 Many classes of organic compounds undergo one or two characteristic types of reaction. For example, reaction of ethylene, $\text{CH}_2=\text{CH}_2$, with HCl forms $\text{CH}_3\text{CH}_2\text{Cl}$, and reaction with Br_2 forms $\text{BrCH}_2\text{CH}_2\text{Br}$. If these reactions are observed in all alkenes, what is the general type of reaction that alkenes undergo?

6.3 Bond Breaking and Bond Making


Having now learned how to write and identify some common kinds of organic reactions, we can turn to a discussion of **reaction mechanism**.

- A *reaction mechanism* is a detailed description of how bonds are broken and formed as a starting material is converted to a product.

A reaction mechanism describes the relative order and rate of bond cleavage and formation. It explains all the known facts about a reaction and accounts for all products formed, and it is subject to modification or refinement as new details are discovered.

A reaction can occur either in one step or in a series of steps.

- A **one-step reaction** is called a *concerted reaction*. No matter how many bonds are broken or formed, a starting material is converted *directly* to a product.




- A **stepwise reaction** involves more than one step. A starting material is first converted to an unstable intermediate, called a **reactive intermediate**, which then goes on to form the product.




6.3A Bond Cleavage

Bonds are broken and formed in all chemical reactions. No matter how many steps there are in the reaction, however, there are only two ways to break (cleave) a bond: the electrons in the bond can be divided **equally** or **unequally** between the two atoms of the bond.

- Breaking a bond by **equally dividing the electrons** between the two atoms in the bond is called **homolysis** or **homolytic cleavage**.



- Breaking a bond by **unequally dividing the electrons** between the two atoms in the bond is called **heterolysis** or **heterolytic cleavage**. Heterolysis of a bond between **A** and **B** can give either **A** or **B** the two electrons in the bond. When **A** and **B** have different electronegativities, the *electrons normally end up on the more electronegative atom*.



Homolysis and heterolysis require energy. Both processes generate reactive intermediates, but the products are different in each case.

- Homolysis generates uncharged reactive intermediates with *unpaired* electrons.
- Heterolysis generates *charged* intermediates.


Each of these reactive intermediates has a very short lifetime, and each reacts quickly to form a stable organic product.

6.3B Radicals, Carbocations, and Carbanions

The curved arrow notation first discussed in Section 1.5 works fine for heterolytic bond cleavage because it illustrates the movement of an **electron pair**. For homolytic cleavage, however, one electron moves to one atom in the bond and one electron moves to the other, so a different kind of curved arrow is needed.


- To illustrate the movement of a single electron, use a half-headed curved arrow, sometimes called a *fishhook*.

Homolysis



Two **half-headed** curved arrows are needed for two **single** electrons.

Heterolysis



One **full-headed** curved arrow is needed for one electron **pair**.



A **full-headed curved arrow** () shows the movement of an electron *pair*. A **half-headed curved arrow** () shows the movement of a *single* electron.

Figure 6.2 illustrates homolysis and two different heterolysis reactions for a carbon compound using curved arrows. Three different reactive intermediates are formed.

Homolysis of the C–Z bond generates two uncharged products with unpaired electrons.

- A reactive intermediate with a single unpaired electron is called a *radical*.


Most radicals are highly unstable because they contain an atom that does not have an octet of electrons. Radicals typically have **no charge**. They are **intermediates in a group of reactions called radical reactions**, which are discussed in detail in Chapter 15.

Heterolysis of the C–Z bond can generate a **carbocation** or a **carbanion**.


- Giving two electrons to Z and none to carbon generates a positively charged carbon intermediate called a *carbocation*.
- Giving two electrons to C and none to Z generates a negatively charged carbon species called a *carbanion*.

Both carbocations and carbanions are unstable reactive intermediates: A carbocation contains a carbon atom surrounded by only six electrons. A carbanion has a negative charge on carbon, which is not a very electronegative atom. **Carbocations (electrophiles)** and **carbanions (nucleophiles)** can be intermediates in *polar reactions*—reactions in which a nucleophile reacts with an electrophile.

Figure 6.2
Three reactive intermediates resulting from homolysis and heterolysis of a C–Z bond





Thus, homolysis and heterolysis generate radicals, carbocations, and carbanions, the three most common reactive intermediates in organic chemistry.



- Radicals and carbocations are electrophiles because they contain an electron-deficient carbon.
- Carbanions are nucleophiles because they contain a carbon with a lone pair.

Problem 6.3


By taking into account electronegativity differences, draw the products formed by heterolysis of the carbon–heteroatom bond in each molecule. Classify the organic reactive intermediate as a carbocation or a carbanion.




6.3C Bond Formation

Like bond cleavage, bond formation occurs in two different ways. Two radicals can each donate **one electron** to form a two-electron bond. Alternatively, two ions with unlike charges can come together, with the negatively charged ion donating **both electrons** to form the resulting two-electron bond. **Bond formation always releases energy.**

Forming a bond from two radicals



Forming a bond from two ions



6.3D All Kinds of Arrows

Table 6.1 summarizes the many kinds of arrows used in describing organic reactions. Curved arrows are especially important because they explicitly show what electrons are involved in a reaction, how these electrons move in forming and breaking bonds, and if a reaction proceeds via a radical or polar pathway.


A more complete summary of the arrows used in organic chemistry is given in the table Common Abbreviations, Arrows, and Symbols, located on the inside back cover.

Table 6.1 A Summary of Arrow Types in Chemical Reactions

Arrow	Name	Use
\longrightarrow	Reaction arrow	Drawn between the starting materials and products in an equation (6.1)
\rightleftharpoons	Double reaction arrows (equilibrium arrows)	Drawn between the starting materials and products in an equilibrium equation (2.2)
\longleftrightarrow	Double-headed arrow	Drawn between resonance structures (1.5)
	Full-headed curved arrow	Shows movement of an electron pair (1.5, 2.2)
	Half-headed curved arrow (fishhook)	Shows movement of a single electron (6.3)


Sample Problem 6.1

Use full-headed or half-headed curved arrows to show the movement of electrons in each equation.




Solution

a. In this reaction, the C–O bond is broken heterolytically. Because only one electron pair is involved, **one full-headed curved arrow** is needed.






b. This reaction involves radicals, so half-headed curved arrows are needed to show the movement of single electrons. One new two-electron bond is formed between H and Cl, and an unpaired electron is left on C. Because a total of three electrons are involved, **three half-headed curved arrows are needed**.



Problem 6.4


Use curved arrows to show the movement of electrons in each equation.



6.4 Bond Dissociation Energy

Bond breaking can be quantified using the bond dissociation energy.

- The *bond dissociation energy* is the energy needed to homolytically cleave a covalent bond.




The energy absorbed or released in any reaction, symbolized by ΔH° , is called the **enthalpy change** or **heat of reaction**.

- When ΔH° is positive (+), energy is absorbed and the reaction is *endothermic*.
- When ΔH° is negative (–), energy is released and the reaction is *exothermic*.

The superscript ($^\circ$) means that values are determined under standard conditions (pure compounds in their most stable state at 25 °C and 1 atm pressure).

A bond dissociation energy is the ΔH° for a specific kind of reaction—the homolysis of a covalent bond to form two radicals. Because bond breaking requires energy, **bond dissociation energies are always positive numbers**, and homolysis is always **endothermic**. Conversely, **bond formation always releases energy**, so this reaction is always **exothermic**. The H–H bond

requires +435 kJ/mol to cleave and releases -435 kJ/mol when formed. Table 6.2 contains a representative list of bond dissociation energies for many common bonds.



Additional bond dissociation energies for C-C multiple bonds are given in Table 1.3.

A table of bond dissociation energies also appears in Appendix C.

Comparing bond dissociation energies is equivalent to comparing **bond strength**.

- The **stronger** the bond, the **higher** its bond dissociation energy.


For example, the H-H bond is stronger than the Cl-Cl bond because its bond dissociation energy is higher [Table 6.2: 435 kJ/mol (H₂) versus 242 kJ/mol (Cl₂)]. The data in Table 6.2 demonstrate that **bond dissociation energies decrease down a column of the periodic table as the valence electrons used in bonding are farther from the nucleus**. Bond dissociation energies for a group of methyl-halogen bonds exemplify this trend.

Table 6.2 Bond Dissociation Energies for Some Common Bonds [A-B → A• + •B]


Bond	ΔH° kJ/mol	(kcal/mol)	Bond	ΔH° kJ/mol	(kcal/mol)
H-Z bonds			R-X bonds		
H-F	569	(136)	CH ₃ -F	456	(109)
H-Cl	431	(103)	CH ₃ -Cl	351	(84)
H-Br	368	(88)	CH ₃ -Br	293	(70)
H-I	297	(71)	CH ₃ -I	234	(56)
H-OH	498	(119)	CH ₃ CH ₂ -F	448	(107)
Z-Z bonds			CH ₃ CH ₂ -Cl	339	(81)
H-H	435	(104)	CH ₃ CH ₂ -Br	285	(68)
F-F	159	(38)	CH ₃ CH ₂ -I	222	(53)
Cl-Cl	242	(58)	(CH ₃) ₂ CH-F	444	(106)
Br-Br	192	(46)	(CH ₃) ₂ CH-Cl	335	(80)
I-I	151	(36)	(CH ₃) ₂ CH-Br	285	(68)
HO-OH	213	(51)	(CH ₃) ₂ CH-I	222	(53)
R-H bonds			(CH ₃) ₃ C-F	444	(106)
CH ₃ -H	435	(104)	(CH ₃) ₃ C-Cl	331	(79)
CH ₃ CH ₂ -H	410	(98)	(CH ₃) ₃ C-Br	272	(65)
CH ₃ CH ₂ CH ₂ -H	410	(98)	(CH ₃) ₃ C-I	209	(50)
(CH ₃) ₂ CH-H	397	(95)	R-OH bonds		
(CH ₃) ₃ C-H	381	(91)	CH ₃ -OH	389	(93)
CH ₂ =CH-H	435	(104)	CH ₃ CH ₂ -OH	393	(94)
HC≡C-H	523	(125)	CH ₃ CH ₂ CH ₂ -OH	385	(92)
CH ₂ =CHCH ₂ -H	364	(87)	(CH ₃) ₂ CH-OH	401	(96)
C ₆ H ₅ -H	460	(110)	(CH ₃) ₃ C-OH	401	(96)
C ₆ H ₅ CH ₂ -H	356	(85)	R-R bonds		
R-R bonds			CH ₃ -CH ₃	368	(88)
CH ₃ -CH ₃	368	(88)	CH ₃ -CH ₂ CH ₃	356	(85)
CH ₃ -CH ₂ CH ₃	356	(85)	CH ₃ -CH=CH ₂	385	(92)
CH ₃ -CH=CH ₂	385	(92)	CH ₃ -C≡CH	489	(117)
CH ₃ -C≡CH	489	(117)			

Because ΔH° is a negative value, this reaction is **exothermic** and energy is released. **The bonds broken in the starting material are weaker than the bonds formed in the product.**

Problem 6.6 Use the values in Table 6.2 to calculate ΔH° for each reaction. Classify each reaction as endothermic or exothermic.



The oxidation of both isooctane and glucose, the two molecules that introduced Chapter 6, forms CO_2 and H_2O .



ΔH° is negative for both oxidations, so both reactions are exothermic. **Both isooctane and glucose release energy on oxidation because the bonds in the products are stronger than the bonds in the reactants.**

Bond dissociation energies have two important limitations. They present *overall* energy changes only. They reveal nothing about the reaction mechanism or how fast a reaction proceeds. Moreover, bond dissociation energies are determined for reactions in the gas phase, whereas most organic reactions are carried out in a liquid solvent where solvation energy contributes to the overall enthalpy of a reaction. As such, bond dissociation energies are imperfect indicators of energy changes in a reaction. Despite these limitations, using bond dissociation energies to calculate ΔH° gives a useful approximation of the energy changes that occur when bonds are broken and formed in a reaction.

Problem 6.7 Calculate ΔH° for each oxidation reaction. Each equation is balanced as written; remember to take into account the coefficients in determining the number of bonds broken or formed.

[ΔH° for $\text{O}_2 = 497 \text{ kJ/mol}$; ΔH° for one $\text{C}=\text{O}$ in $\text{CO}_2 = 535 \text{ kJ/mol}$]



6.5 Thermodynamics

For a reaction to be practical, the equilibrium must favor the products, *and* the reaction rate must be fast enough to form them in a reasonable time. These two conditions depend on the **thermodynamics** and the **kinetics** of a reaction, respectively.


- **Thermodynamics** describes energy and equilibrium. How do the *energies* of the reactants and the products compare? What are the *relative amounts* of reactants and products at equilibrium?
- **Kinetics** describes reaction rates. How *fast* are reactants converted to products?

Reaction kinetics are discussed in Section 6.9.

6.5A Equilibrium Constant and Free Energy Changes

The **equilibrium constant**, K_{eq} , is a mathematical expression that relates the amount of starting material and product at equilibrium. For example, when starting materials **A** and **B** react to form products **C** and **D**, the equilibrium constant is given by the following expression.

K_{eq} was first defined in Section 2.3 for acid–base reactions.



The size of K_{eq} tells about the position of equilibrium; that is, it expresses whether the starting materials or products predominate once equilibrium has been reached.

- When $K_{\text{eq}} > 1$, equilibrium favors the products (**C** and **D**) and the equilibrium lies to the right as the equation is written.
 - When $K_{\text{eq}} < 1$, equilibrium favors the starting materials (**A** and **B**) and the equilibrium lies to the left as the equation is written.
- For a reaction to be useful, the equilibrium must favor the products, and $K_{\text{eq}} > 1$.

What determines whether equilibrium favors the products in a given reaction? **The position of equilibrium is determined by the relative energies of the reactants and products.** The free energy of a molecule, also called its **Gibbs free energy**, is symbolized by G° . The **change in free energy** between reactants and products, symbolized by ΔG° , determines whether the starting materials or products are favored at equilibrium.

- ΔG° is the overall energy difference between reactants and products.

Free energy change	$\Delta G^\circ =$	$G^\circ_{\text{products}}$	$-$	$G^\circ_{\text{reactants}}$
		↑		↑
		free energy of the products		free energy of the reactants

ΔG° is related to the equilibrium constant K_{eq} by the following equation:

$$\Delta G^\circ = -2.303RT \log K_{\text{eq}} \quad \left[\begin{array}{l} R = 8.314 \text{ J}/(\text{K}\cdot\text{mol}), \text{ the gas constant} \\ T = \text{Kelvin temperature (K)} \end{array} \right]$$

K_{eq} depends on the energy difference between reactants and products.

At 25 °C, $2.303 RT = 5.9 \text{ kJ/mol}$; thus, $\Delta G^\circ = -5.9 \log K_{\text{eq}}$.

$K_{\text{eq}} > 1$ when $\Delta G^\circ < 0$, and equilibrium favors the products. $K_{\text{eq}} < 1$ when $\Delta G^\circ > 0$, and equilibrium favors the starting materials.

Using this expression we can determine the relationship between the equilibrium constant and the free energy change between reactants and products.

- When $K_{\text{eq}} > 1$, $\log K_{\text{eq}}$ is positive, making ΔG° **negative**, and energy is released. Thus, **equilibrium favors the products when the energy of the products is lower than the energy of the reactants.**
- When $K_{\text{eq}} < 1$, $\log K_{\text{eq}}$ is negative, making ΔG° **positive**, and energy is absorbed. Thus, **equilibrium favors the reactants when the energy of the products is higher than the energy of the reactants.**

Compounds that are lower in energy have increased stability. Thus, **equilibrium favors the products when they are more stable (lower in energy) than the starting materials of a reaction.** This is summarized in Figure 6.3.

Figure 6.3

Summary of the relationship between ΔG° and K_{eq}




Table 6.3 Representative Values for ΔG° and K_{eq} at 25 °C, for a Reaction $A \rightarrow B$

	ΔG° (kJ/mol)	K_{eq}	Relative amount of A and B at equilibrium	
→	+18	10^{-3}	Essentially all A (99.9%)	↑ Increasing [product] ↓
	+12	10^{-2}	100 times as much A as B	
	+6	10^{-1}	10 times as much A as B	
	0	1	Equal amounts of A and B	
	-6	10^1	10 times as much B as A	
→	-12	10^2	100 times as much B as A	
	-18	10^3	Essentially all B (99.9%)	

A small difference in free energy means a large difference in the amount of A and B at equilibrium.

Because ΔG° depends on the logarithm of K_{eq} , a **small change in energy corresponds to a large difference in the relative amount of starting material and product at equilibrium**. Several values of ΔG° and K_{eq} are given in Table 6.3. For example, a difference in energy of only ~6 kJ/mol means that there is 10 times as much of the more stable species at equilibrium. A difference in energy of ~18 kJ/mol means that there is essentially only one compound, either starting material or product, at equilibrium.

The symbol ~ means approximately.

Problem 6.8

- a. Which K_{eq} corresponds to a negative value of ΔG° , $K_{\text{eq}} = 1000$ or $K_{\text{eq}} = .001$?
 b. Which K_{eq} corresponds to a lower value of ΔG° , $K_{\text{eq}} = 10^{-2}$ or $K_{\text{eq}} = 10^{-5}$?

Problem 6.9

- Given each of the following values, is the starting material or product favored at equilibrium?
 a. $K_{\text{eq}} = 5.5$ b. $\Delta G^\circ = 40$ kJ/mol


Problem 6.10

- Given each of the following values, is the starting material or product lower in energy?
 a. $\Delta G^\circ = 8.0$ kJ/mol b. $K_{\text{eq}} = 10$ c. $\Delta G^\circ = -12$ kJ/mol d. $K_{\text{eq}} = 10^{-3}$

6.5B Energy Changes and Conformational Analysis


These equations can be used for any process with two states in equilibrium. As an example, monosubstituted cyclohexanes exist as two different chair conformations that rapidly interconvert at room temperature, with the conformation having the substituent in the roomier equatorial position favored (Section 4.13). Knowing the energy difference between the two conformations allows us to calculate the amount of each at equilibrium.

For example, the energy difference between the two chair conformations of phenylcyclohexane is -12.1 kJ/mol, as shown in the accompanying equation. Using the values in Table 6.3, this corresponds to an equilibrium constant of ~100, meaning that there is approximately 100 times more B (equatorial phenyl group) than A (axial phenyl group) at equilibrium.



Problem 6.11

The equilibrium constant for the conversion of the axial to the equatorial conformation of methoxycyclohexane is 2.7.



- Given these data, which conformation is present in the larger amount at equilibrium?
- Is ΔG° for this process positive or negative?
- From the values in Table 6.3, approximate the size of ΔG° .

6.6 Enthalpy and Entropy

The **free energy change** (ΔG°) depends on the **enthalpy change** (ΔH°) and the **entropy change** (ΔS°). ΔH° indicates relative bond strength, but what does ΔS° measure?

Entropy (S°) is a measure of the randomness in a system. The more freedom of motion or the more disorder present, the higher the entropy. Gas molecules move more freely than liquid molecules and are higher in entropy. Cyclic molecules have more restricted bond rotation than similar acyclic molecules and are lower in entropy.

The **entropy change** (ΔS°) is the change in the amount of disorder between reactants and products. ΔS° is positive (+) when the products are more disordered than the reactants. ΔS° is negative (–) when the products are less disordered (more ordered) than the reactants.

- Reactions resulting in an increase in entropy are favored.

ΔG° is related to ΔH° and ΔS° by the following equation:


$$\text{Total energy change} \rightarrow \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ \quad [T = \text{Kelvin temperature}]$$

↑ change in bonding energy
 ↑ change in disorder

This equation tells us that the total energy change in a reaction is due to two factors: the change in the **bonding energy** and the change in **disorder**. The change in bonding energy can be calculated from bond dissociation energies (Section 6.4). Entropy changes, on the other hand, are more difficult to access, but they are important in the following two cases:

- When the number of molecules of starting material *differs* from the number of molecules of product in the balanced chemical equation.
- When an acyclic molecule is *cyclized* to a cyclic one, or a cyclic molecule is converted to an acyclic one.

For example, **when a single starting material forms two products**, as in the homolytic cleavage of a bond to form two radicals, **entropy increases** and favors formation of the products. In contrast, **entropy decreases when an acyclic compound forms a ring**, because a ring has fewer degrees of freedom. In this case, therefore, entropy does *not* favor formation of the product.



Entropy is a rather intangible concept that comes up again and again in chemistry courses. One way to remember the relation between entropy and disorder is to consider a handful of chopsticks. Dropped on the floor, they are arranged randomly (a state of high entropy). Placed end-to-end in a straight line, they are arranged intentionally (a state of low entropy). The more disordered, random arrangement is favored and easier to achieve energetically.

In most other reactions that are not carried out at high temperature, the entropy term ($T\Delta S^\circ$) is small compared to the enthalpy term (ΔH°) and it can be neglected. Thus, **we will often approximate the overall free energy change of a reaction by the change in the bonding energy only.** Keep in mind that this is an approximation, but it gives us a starting point from which to decide if the reaction is energetically favorable.

Recall from Section 6.4 that a reaction is endothermic when ΔH° is positive and exothermic when ΔH° is negative. A reaction is **endergonic when ΔG° is positive** and **exergonic when ΔG° is negative.** ΔG° is usually approximated by ΔH° in this text, so the terms endergonic and exergonic are rarely used.

$$\Delta G^\circ \approx \Delta H^\circ$$

- The total energy change is approximated by the change in bonding energy only.

According to this approximation:

- The product is favored in reactions in which ΔH° is a *negative* value; that is, the bonds in the product are *stronger* than the bonds in the starting material.
- The starting material is favored in a reaction in which ΔH° is a *positive* value; that is, the bonds in the starting material are *stronger* than the bonds in the product.

Problem 6.12 Considering each of the following values and neglecting entropy, tell whether the starting material or product is favored at equilibrium: (a) $\Delta H^\circ = 80$ kJ/mol; (b) $\Delta H^\circ = -40$ kJ/mol.


Problem 6.13 For a reaction with $\Delta H^\circ = 40$ kJ/mol, decide which of the following statements is (are) true. Correct any false statement to make it true. (a) The reaction is exothermic; (b) ΔG° for the reaction is positive; (c) K_{eq} is greater than 1; (d) the bonds in the starting materials are stronger than the bonds in the product; and (e) the product is favored at equilibrium.

Problem 6.14 Answer Problem 6.13 for a reaction with $\Delta H^\circ = -20$ kJ/mol.


6.7 Energy Diagrams

An **energy diagram** is a schematic representation of the energy changes that take place as reactants are converted to products. An energy diagram indicates how readily a reaction proceeds, how many steps are involved, and how the energies of the reactants, products, and intermediates compare.

Consider, for example, a concerted reaction between molecule **A–B** with anion **C:[–]** to form products **A:[–]** and **B–C**. If the reaction occurs in a single step, the bond between **A** and **B** is broken *as* the bond between **B** and **C** is formed. Let's assume that the products are lower in energy than the reactants in this hypothetical reaction.



An energy diagram plots **energy on the y axis** versus the progress of reaction, often labeled the **reaction coordinate**, on the **x axis**. As the starting materials **A–B** and **C:[–]** approach one another, their electron clouds feel some repulsion, causing an increase in energy, until a maximum value is reached. This unstable energy maximum is called the **transition state**. In the transition state the bond between **A** and **B** is partially broken, and the bond between **B** and **C** is partially formed. Because it is at the top of an energy “hill,” **a transition state can never be isolated.**



At the transition state, the bond between **A** and **B** can re-form to regenerate starting material, or the bond between **B** and **C** can form to generate product. As the bond forms between **B** and **C** the energy decreases until some stable energy minimum of the products is reached.

- The energy difference between the reactants and products is ΔH° . Because the products are at lower energy than the reactants, this reaction is exothermic and energy is released.
- The energy difference between the transition state and the starting material is called the *energy of activation*, symbolized by E_a .

The *energy of activation* is the minimum amount of energy needed to break bonds in the reactants. It represents an **energy barrier** that must be overcome for a reaction to occur. The size of E_a tells us about the reaction rate.


A slow reaction has a large E_a . A fast reaction has a low E_a .

- The larger the E_a , the greater the amount of energy that is needed to break bonds, and the slower the reaction rate.

How can we draw the structure of the unstable transition state? The structure of the transition state is somewhere in between the structures of the starting material and product. Any bond that is partially broken or formed is drawn with a dashed line. Any atom that gains or loses a charge contains a partial charge in the transition state. Transition states are drawn in brackets, with a superscript double dagger (\ddagger).

In the hypothetical reaction between **A–B** and **C:⁻** to form **A:⁻** and **B–C**, the bond between **A** and **B** is partially broken, and the bond between **B** and **C** is partially formed. Because **A** gains a negative charge and **C** loses a charge in the course of the reaction, each atom bears a partial negative charge in the transition state.

Drawing the structure of a transition state



This bond is partially broken. This bond is partially formed.

Several energy diagrams are drawn in Figure 6.4. For any energy diagram:


- E_a determines the height of the energy barrier.
- ΔH° determines the relative position of the reactants and products.

The two variables, E_a and ΔH° , are **independent** of each other. Two reactions can have identical values for ΔH° but very different E_a values. In Figure 6.5, both reactions have the same

Figure 6.4 Example [1]


Some representative energy diagrams

- Large $E_a \rightarrow$ slow reaction
- (+) $\Delta H^\circ \rightarrow$ endothermic reaction




Example [3]

- Low $E_a \rightarrow$ fast reaction
- (+) $\Delta H^\circ \rightarrow$ endothermic reaction




Example [2]

- Large $E_a \rightarrow$ slow reaction
- (-) $\Delta H^\circ \rightarrow$ exothermic reaction



Example [4]


- Low $E_a \rightarrow$ fast reaction
- (-) $\Delta H^\circ \rightarrow$ exothermic reaction



negative ΔH° favoring the products, but the second reaction has a much higher E_a , so it proceeds more slowly.

Problem 6.15

Draw an energy diagram for a reaction in which the products are higher in energy than the starting materials and E_a is large. Clearly label all of the following on the diagram: the axes, the starting materials, the products, the transition state, ΔH° , and E_a .


Figure 6.5Comparing ΔH° and E_a in two energy diagrams

- Energy diagrams in (a) and (b) both depict exothermic reactions with the same negative value of ΔH° .
- E_a in (a) is lower than E_a in (b), so reaction (a) is faster than reaction (b).

Problem 6.16 Draw the structure for the transition state in each reaction.



Problem 6.17 Compound **A** can be converted to either **B** or **C**. The energy diagrams for both processes are drawn on the graph below.



- Label each reaction as endothermic or exothermic.
- Which reaction is faster?
- Which reaction generates the product lower in energy?
- Which points on the graphs correspond to transition states?
- Label the energy of activation for each reaction.
- Label the ΔH° for each reaction.


6.8 Energy Diagram for a Two-Step Reaction Mechanism

Although the hypothetical reaction in Section 6.7 is concerted, many reactions involve more than one step with formation of a reactive intermediate. Consider the same overall reaction, $\text{A-B} + \text{C}^-$ to form products $\text{A}^- + \text{B-C}$, but in this case begin with the assumption that the reaction occurs by a *stepwise* pathway—that is, bond breaking occurs *before* bond making. Once again, assume that the overall process is exothermic.

Same overall reaction




One possible stepwise mechanism involves heterolysis of the **A-B** bond to form two ions **A⁻** and **B⁺**, followed by reaction of **B⁺** with anion **C⁻** to form product **B-C**, as outlined in the accompanying equations. Species **B⁺** is a **reactive intermediate**. It is formed as a product in Step [1], and then goes on to react with **C⁻** in Step [2].




We must draw an energy diagram for each step, and then combine them in an energy diagram for the overall two-step mechanism. Each step has its own energy barrier, with a transition state at the energy maximum.

Step [1] is endothermic because energy is needed to cleave the **A-B** bond, making ΔH° a positive value and placing the products of Step [1] at higher energy than the starting materials. In the transition state, the **A-B** bond is partially broken.



Step [2] is exothermic because energy is released in forming the **B-C** bond, making ΔH° a negative value and placing the products of Step [2] at lower energy than the starting materials of Step [2]. In the transition state, the **B-C** bond is partially formed.



The overall process is shown in Figure 6.6 as a single energy diagram that combines both steps. Because the reaction has two steps, there are two transition states, each corresponding to an energy barrier. The transition states are separated by an energy minimum, at which the reactive intermediate B^+ is located. Because we made the assumption that the overall two-step process is exothermic, the overall energy difference between the reactants and products, labeled $\Delta H^\circ_{\text{overall}}$, has a negative value, and the final products are at a lower energy than the starting materials.

The energy barrier for Step [1], labeled $E_a[1]$, is higher than the energy barrier for Step [2], labeled $E_a[2]$. This is because bond cleavage (Step [1]) is more difficult (requires more energy) than bond formation (Step [2]). A higher energy transition state for Step [1] makes it the slower step of the mechanism.

- In a multistep mechanism, the step with the highest energy transition state is called the rate-determining step.

In this reaction, the rate-determining step is Step [1].





Figure 6.6
Complete energy diagram for the two-step conversion of $A-B + C:^- \rightarrow A:^- + B-C$

- The transition states are located at energy maxima, while the reactive intermediate B^+ is located at an energy minimum.
- Each step is characterized by its own value of ΔH° and E_a .
- The overall energy difference between starting material and products is called $\Delta H^\circ_{\text{overall}}$. In this example, the products of the two-step sequence are at lower energy than the starting materials.
- Since Step [1] has the higher energy transition state, it is the **rate-determining step**.

Problem 6.18 Consider the following energy diagram.



- How many steps are involved in this reaction?
- Label ΔH° and E_a for each step, and label $\Delta H^\circ_{\text{overall}}$.
- Label each transition state.
- Which point on the graph corresponds to a reactive intermediate?
- Which step is rate-determining?
- Is the overall reaction endothermic or exothermic?

Problem 6.19 Draw an energy diagram for a two-step reaction, $A \rightarrow B \rightarrow C$, where the relative energy of these compounds is $C < A < B$, and the conversion of $B \rightarrow C$ is rate-determining.

6.9 Kinetics

We now turn to a more detailed discussion of **reaction rate**—that is, how fast a particular reaction proceeds. **The study of reaction rates is called kinetics.**

The rate of chemical processes affects many facets of our lives. Aspirin is an effective anti-inflammatory agent because it rapidly inhibits the synthesis of prostaglandins (Section 19.6). Butter turns rancid with time because its lipids are only slowly oxidized by oxygen in the air to undesirable by-products (Section 15.11). DDT (Section 7.4) is a persistent environmental pollutant because it does not react appreciably with water, oxygen, or any other chemical with which it comes into contact. All of these processes occur at different rates, resulting in beneficial or harmful effects.


6.9A Energy of Activation

As we learned in Section 6.7, the energy of activation, E_a , is the energy difference between the reactants and the transition state. It is the **energy barrier** that must be exceeded for reactants to be converted to products.

Some reactions have a very favorable equilibrium constant ($K_{\text{eq}} \gg 1$), but the rate is very slow. The oxidation of alkanes like isooctane to form CO_2 and H_2O is an example of this phenomenon. Without a spark to initiate the reaction, isooctane does not react with O_2 ; and gasoline, which contains isooctane, can be safely handled in the air.



Practically, the effect of temperature on reaction rate is used to an advantage in the kitchen. Food is stored in a cold refrigerator to slow the reactions that cause spoilage. On the other hand, bread is baked in a hot oven to increase the rate of the reactions that occur during baking.



- The *larger* the E_a , the *slower* the reaction.

Concentration and temperature also affect reaction rate.

- The *higher* the concentration, the *faster* the rate. Increasing concentration increases the number of collisions between reacting molecules, which in turn increases the rate.
- The *higher* the temperature, the *faster* the rate. Increasing temperature increases the average kinetic energy of the reacting molecules. Because the kinetic energy of colliding molecules is used for bond cleavage, increasing the average kinetic energy increases the rate.

The E_a values of most organic reactions are 40–150 kJ/mol. When $E_a < 80$ kJ/mol, the reaction occurs readily at or below room temperature. When $E_a > 80$ kJ/mol, higher temperatures are needed. As a rule of thumb, increasing the temperature by 10°C doubles the reaction rate. Thus, reactions in the lab are often heated to increase their rates so they occur in a reasonable amount of time.

Keep in mind that certain **reaction quantities have no effect on reaction rate.**

- ΔG° , ΔH° , and K_{eq} do not determine the rate of a reaction. These quantities indicate the direction of equilibrium and the relative energy of reactants and products.

Problem 6.20 Which value (if any) corresponds to a faster reaction: (a) $E_a = 40$ kJ/mol or $E_a = 4$ kJ/mol; (b) a reaction temperature of 0°C or a reaction temperature of 25°C ; (c) $K_{\text{eq}} = 10$ or $K_{\text{eq}} = 100$; (d) $\Delta H^\circ = -10$ kJ/mol or $\Delta H^\circ = 10$ kJ/mol?


Problem 6.21 Explain why the E_a of an endothermic reaction is at least as large as its ΔH° .

Problem 6.22 For a reaction with $K_{\text{eq}} = 0.8$ and $E_a = 80$ kJ/mol, decide which of the following statements is (are) true. Correct any false statement to make it true. Ignore entropy considerations. (a) The reaction is faster than a reaction with $K_{\text{eq}} = 8$ and $E_a = 80$ kJ/mol. (b) The reaction is faster than a reaction with $K_{\text{eq}} = 0.8$ and $E_a = 40$ kJ/mol. (c) ΔG° for the reaction is a positive value. (d) The starting materials are lower in energy than the products of the reaction. (e) The reaction is exothermic.

6.9B Rate Equations

The rate of a chemical reaction is determined by measuring the decrease in the concentration of the reactants over time, or the increase in the concentration of the products over time. A **rate law** (or **rate equation**) is an equation that shows the relationship between the rate of a reaction and the concentration of the reactants. A rate law is determined *experimentally*, and it depends on the mechanism of the reaction.

A rate law has two important terms: the **rate constant symbolized by k** , and the **concentration of the reactants**. Not all reactant concentrations may appear in the rate equation, as we shall soon see.



A rate constant k is a fundamental characteristic of a reaction. It is a complex mathematical term that takes into account the dependence of a reaction rate on temperature and the energy of activation.


A rate constant k and the energy of activation E_a are inversely related. **A high E_a corresponds to a small k .**

- Fast reactions have large rate constants.
- Slow reactions have small rate constants.

What concentration terms appear in the rate equation? That depends on the mechanism. For the organic reactions we will encounter:

- A rate equation contains concentration terms for *all* reactants involved in a *one-step* mechanism.
- A rate equation contains concentration terms for *only* the reactants involved in the *rate-determining step* in a multistep reaction.


For example, in the one-step reaction of $\text{A}-\text{B} + \text{C}^-$ to form $\text{A}^- + \text{B}-\text{C}$, *both* reactants appear in the transition state of the only step of the mechanism. The **concentration of both reactants affects the reaction rate** and *both* terms appear in the rate equation. This type of reaction involving two reactants is said to be **bimolecular**.



The **order of a rate equation equals the sum of the exponents of the concentration terms** in the rate equation. In the rate equation for the concerted reaction of $\text{A}-\text{B} + \text{C}^-$, there are two concentration terms, each with an exponent of one. Thus, the sum of the exponents is two and the **rate equation is second order** (the reaction follows second-order kinetics).

Because the rate of the reaction depends on the concentration of both reactants, doubling the concentration of *either* $\text{A}-\text{B}$ or C^- doubles the rate of the reaction. Doubling the concentration of *both* $\text{A}-\text{B}$ and C^- increases the reaction rate by a factor of *four*.

The situation is different in the stepwise conversion of $\text{A}-\text{B} + \text{C}^-$ to form $\text{A}^- + \text{B}-\text{C}$. The mechanism shown in Section 6.8 has two steps: a slow step (the **rate-determining step**) in which the $\text{A}-\text{B}$ bond is broken, and a fast step in which the $\text{B}-\text{C}$ bond is formed.



In a multistep mechanism, a reaction can occur no faster than its rate-determining step. **Only the concentrations of the reactants affecting the rate-determining step appear in the rate equation.** In this example, the rate depends on the concentration of $\text{A}-\text{B}$ *only*, because only $\text{A}-\text{B}$ appears in the rate-determining step. A reaction involving only one reactant is said to be **unimolecular**. Because there is only one concentration term (raised to the first power), the **rate equation is first order** (the reaction follows first-order kinetics).

Because the rate of the reaction depends on the concentration of only *one* reactant, doubling the concentration of **A–B** doubles the rate of the reaction, but **doubling the concentration of C:[–] has no effect on the reaction rate.**

This might seem like a puzzling result. If **C:[–]** is involved in the reaction, why doesn't it affect the overall rate of the reaction? Not only can you change the concentration of **C:[–]** and not affect the rate, but you also can replace it by a different anion without affecting the rate. How can this be? **C:[–] is not involved in the slow step of the reaction, so neither its concentration nor its identity affects the reaction rate.**

The following analogy is useful. Let's say three students must make 20 peanut butter and jelly sandwiches for a class field trip. Student **(1)** spreads the peanut butter on the bread. Student **(2)** spreads on the jelly, and student **(3)** cuts the sandwiches in half. Suppose student **(2)** is very slow in spreading the jelly. It doesn't matter how fast students **(1)** and **(3)** are; they can't finish making sandwiches any faster than student **(2)** can add the jelly. Five more students can spread on the peanut butter, or an entirely different individual can replace student **(3)**, and this doesn't speed up the process. How fast the sandwiches are made is determined entirely by the rate-determining step—that is, spreading the jelly.

Rate equations provide very important information about the mechanism of a reaction. Rate laws for new reactions with unknown mechanisms are determined by a set of experiments that measure how a reaction's rate changes with concentration. Then, a mechanism is suggested based on which reactants affect the rate.

Problem 6.23 For each rate equation, what effect does the indicated concentration change have on the overall rate of the reaction?



[1] $\text{rate} = k[\text{CH}_3\text{CH}_2\text{Br}][^-\text{OH}]$

- tripling the concentration of $\text{CH}_3\text{CH}_2\text{Br}$ only
- tripling the concentration of ^-OH only
- tripling the concentration of both $\text{CH}_3\text{CH}_2\text{Br}$ and ^-OH

[2] $\text{rate} = k[(\text{CH}_3)_3\text{COH}]$

- doubling the concentration of $(\text{CH}_3)_3\text{COH}$
- increasing the concentration of $(\text{CH}_3)_3\text{COH}$ by a factor of 10

Problem 6.24 Write a rate equation for each reaction, given the indicated mechanism.



6.10 Catalysts

Some reactions do not occur in a reasonable time unless a **catalyst** is added.

- A **catalyst** is a substance that speeds up the rate of a reaction. A catalyst is recovered unchanged in a reaction, and it does not appear in the product.

Common catalysts in organic reactions are **acids** and **metals**. Two examples are shown in the accompanying equations.





Figure 6.7
The effect of a catalyst
on a reaction






- The catalyst lowers the energy of activation, thus **increasing the rate of the catalyzed reaction**.
- The energy of the reactants and products is the same in both the uncatalyzed and catalyzed reactions, so the **position of equilibrium is unaffected**.

The reaction of acetic acid with ethanol to yield ethyl acetate and water occurs in the presence of an acid catalyst. The acid catalyst is written over or under the arrow to emphasize that it is not part of the starting materials or the products. The details of this reaction are discussed in Chapter 22.

The reaction of ethylene with hydrogen to form ethane occurs only in the presence of a metal catalyst such as palladium, platinum, or nickel. The metal provides a surface that binds both the ethylene and the hydrogen, and in doing so, facilitates the reaction. We return to this mechanism in Chapter 12.

Catalysts accelerate a reaction by lowering the energy of activation (Figure 6.7). They have no effect on the equilibrium constant, so they do not change the amount of reactant and product at equilibrium. Thus, catalysts affect how *quickly* equilibrium is achieved, but not the relative amounts of reactants and products at equilibrium. If a catalyst is somehow used up in one step of a reaction sequence, it must be regenerated in another step. Because only a small amount of a catalyst is needed relative to starting material, it is said to be present in a **catalytic amount**.

Problem 6.25 Identify the catalyst in each equation.



6.11 Enzymes


The catalysts that synthesize and break down biomolecules in living organisms are governed by the same principles as the acids and metals in organic reactions. The catalysts in living organisms, however, are usually protein molecules called **enzymes**.

- **Enzymes** are biochemical catalysts composed of amino acids held together in a very specific three-dimensional shape.

An enzyme contains a region called its **active site**, which binds an organic reactant, called a **substrate**. When bound, this unit is called the **enzyme–substrate complex**, as shown schematically in Figure 6.8 for the enzyme lactase, the enzyme that binds lactose, the principal carbohydrate in milk. Once bound, the organic substrate undergoes a very specific reaction at an enhanced rate. In this example, lactose is converted into two simpler sugars, glucose and galactose. When individuals lack adequate amounts of lactase, they are unable to digest lactose, causing abdominal cramping and diarrhea.

Figure 6.8

Lactase, an example of a biological catalyst



The enzyme lactase binds the carbohydrate lactose ($C_{12}H_{22}O_{11}$) in its active site in Step [1]. Lactose then reacts with water to break a bond and form two simpler sugars, galactose and glucose, in Step [2]. This process is the first step in digesting lactose, the principal carbohydrate in milk.

An enzyme speeds up a biological reaction in a variety of ways. It may hold reactants in the proper conformation to facilitate reaction, or it may provide an acidic site needed for a particular transformation. Once the reaction is completed, the enzyme releases the substrate and it is then able to catalyze another reaction.

KEY CONCEPTS

Understanding Organic Reactions

Writing Equations for Organic Reactions (6.1)

- Use curved arrows to show the movement of electrons. Full-headed arrows are used for electron pairs and half-headed arrows are used for single electrons.
- Reagents can be drawn either on the left side of an equation or over the reaction arrow. Catalysts are drawn over or under the reaction arrow.

Types of Reactions (6.2)

[1] Substitution	$\begin{array}{c} \\ -\text{C}-\text{Z} \\ \end{array} + \text{Y} \longrightarrow \begin{array}{c} \\ -\text{C}-\text{Y} \\ \end{array} + \text{Z} \quad \left[\text{Z} = \text{H or a heteroatom} \right]$ <p style="text-align: center;">Y replaces Z</p>
[2] Elimination	$\begin{array}{c} \quad \\ -\text{C}-\text{C}- \\ \quad \\ \text{X} \quad \text{Y} \end{array} + \text{reagent} \longrightarrow \begin{array}{c} \diagup \quad \diagdown \\ \text{C}=\text{C} \\ \diagdown \quad \diagup \end{array} + \text{X}-\text{Y}$ <p>Two σ bonds are broken. π bond</p>
[3] Addition	$\begin{array}{c} \diagup \quad \diagdown \\ \text{C}=\text{C} \\ \diagdown \quad \diagup \end{array} + \text{X}-\text{Y} \longrightarrow \begin{array}{c} \quad \\ -\text{C}-\text{C}- \\ \quad \\ \text{X} \quad \text{Y} \end{array}$ <p>This π bond is broken. Two σ bonds are formed.</p>

Important Trends


Values compared	Trend
Bond dissociation energy and bond strength Energy and stability E_a and reaction rate E_a and rate constant	The higher the bond dissociation energy, the stronger the bond (6.4). The higher the energy, the less stable the species (6.5A). The larger the energy of activation, the slower the reaction (6.9A). The larger the energy of activation, the smaller the rate constant (6.9B).

Reactive Intermediates (6.3)

- Breaking bonds generates reactive intermediates.
- Homolysis generates radicals with unpaired electrons.
- Heterolysis generates ions.

Reactive intermediate	General structure	Reactive feature	Reactivity
Radical	$\begin{array}{c} \\ -\text{C}\cdot \\ \end{array}$	Unpaired electron	Electrophilic
Carbocation	$\begin{array}{c} \\ -\text{C}^+ \\ \end{array}$	Positive charge; only six electrons around C	Electrophilic
Carbanion	$\begin{array}{c} \\ -\text{C}:^- \\ \end{array}$	Net negative charge; lone electron pair on C	Nucleophilic


Energy Diagrams (6.7, 6.8)



Conditions Favoring Product Formation (6.5, 6.6)

Variable	Value	Meaning
K_{eq}	$K_{\text{eq}} > 1$	More products than reactants are present at equilibrium.
ΔG°	$\Delta G^\circ < 0$	The free energy of the products is lower than the energy of the reactants.
ΔH°	$\Delta H^\circ < 0$	Bonds in the products are stronger than bonds in the reactants.
ΔS°	$\Delta S^\circ > 0$	The products are more disordered than the reactants.

Equations (6.5, 6.6)




Factors Affecting Reaction Rate (6.9)

Factor	Effect
Energy of activation	Larger E_a ----> slower reaction
Concentration	Higher concentration ----> faster reaction
Temperature	Higher temperature ----> faster reaction

PROBLEMS

Types of Reactions

6.26 Classify each transformation as substitution, elimination, or addition.




6.33 Calculate ΔH° for each reaction.

- $\text{CH}_3\text{CH}_3 + \text{Br}_2 \longrightarrow \text{CH}_3\text{CH}_2\text{Br} + \text{HBr}$
- $\text{HO}\cdot + \text{CH}_4 \longrightarrow \cdot\text{CH}_3 + \text{H}_2\text{O}$
- $\text{CH}_3\text{OH} + \text{HBr} \longrightarrow \text{CH}_3\text{Br} + \text{H}_2\text{O}$
- $\text{Br}\cdot + \text{CH}_4 \longrightarrow \text{H}\cdot + \text{CH}_3\text{Br}$

6.34 Explain why the bond dissociation energy for the C–C σ bond in propane is lower than the bond dissociation energy for the C–C σ bond in propene, $\text{CH}_3\text{CH}=\text{CH}_2$.



6.35 Homolysis of the indicated C–H bond in propene forms a resonance-stabilized radical.



- Draw the two possible resonance structures for this radical.
- Use half-headed curved arrows to illustrate how one resonance structure can be converted to the other.
- Draw a structure for the resonance hybrid.

6.36 Because propane ($\text{CH}_3\text{CH}_2\text{CH}_3$) has both 1° and 2° carbon atoms, it has two different types of C–H bonds.

- Draw the carbon radical formed by homolysis of each of these C–H bonds.
- Use the values in Table 6.2 to determine which C–H bond is stronger.
- Explain how this information can also be used to determine the relative stability of the two radicals formed. Which radical formed from propane is more stable?


6.37 Use the bond dissociation energies in Table 1.3 (listed as bond strengths) to estimate the strength of the σ and π components of the double bond in ethylene.

Thermodynamics, ΔG° , ΔH° , ΔS° , and K_{eq}


6.38 Given each value, determine whether the starting material or product is favored at equilibrium.

- | | | |
|---|--|---|
| a. $K_{\text{eq}} = 0.5$ | d. $K_{\text{eq}} = 16$ | g. $\Delta S^\circ = 8 \text{ J/(K}\cdot\text{mol)}$ |
| b. $\Delta G^\circ = -100 \text{ kJ/mol}$ | e. $\Delta G^\circ = 2.0 \text{ kJ/mol}$ | h. $\Delta S^\circ = -8 \text{ J/(K}\cdot\text{mol)}$ |
| c. $\Delta H^\circ = 8.0 \text{ kJ/mol}$ | f. $\Delta H^\circ = 200 \text{ kJ/mol}$ | |


- Which value corresponds to a negative value of ΔG° : $K_{\text{eq}} = 10^{-2}$ or $K_{\text{eq}} = 10^2$?
 - In a unimolecular reaction with five times as much starting material as product at equilibrium, what is the value of K_{eq} ? Is ΔG° positive or negative?
 - Which value corresponds to a larger K_{eq} : $\Delta G^\circ = -8 \text{ kJ/mol}$ or $\Delta G^\circ = 20 \text{ kJ/mol}$?
- 6.40 As we learned in Chapter 4, monosubstituted cyclohexanes exist as an equilibrium mixture of two conformations having either an axial or equatorial substituent.



- When $\text{R} = \text{CH}_3$, which conformation is present in higher concentration?
 - Which R shows the highest percentage of equatorial conformation at equilibrium?
 - Which R shows the highest percentage of axial conformation at equilibrium?
 - For which R is ΔG° most negative?
 - How is the size of R related to the amount of axial and equatorial conformations at equilibrium?
 - Challenge question: Explain why three monosubstituted cycloalkanes [$\text{R} = -\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$] have similar values of K_{eq} , but K_{eq} for *tert*-butylcyclohexane [$\text{R} = -\text{C}(\text{CH}_3)_3$] is much higher.
- 6.41 At 25°C , the energy difference (ΔG°) for the conversion of axial fluorocyclohexane to its equatorial conformation is -1.0 kJ/mol .
- Calculate K_{eq} for this equilibrium.
 - Calculate the percentage of axial and equatorial conformations present at equilibrium.






6.42 For which of the following reactions is ΔS° a positive value?

- a.  $\text{C}_7\text{H}_{16} \longrightarrow \text{C}_6\text{H}_{12} + \text{C}_2\text{H}_6$
- b. $\text{CH}_3\cdot + \text{CH}_3\cdot \longrightarrow \text{CH}_3\text{CH}_3$
- c. $(\text{CH}_3)_2\text{C}(\text{OH})_2 \longrightarrow (\text{CH}_3)_2\text{C}=\text{O} + \text{H}_2\text{O}$
- d. $\text{CH}_3\text{COOCH}_3 + \text{H}_2\text{O} \longrightarrow \text{CH}_3\text{COOH} + \text{CH}_3\text{OH}$

Energy Diagrams and Transition States

6.43 Draw the transition state for each reaction.

- a. 
- b. $\text{BF}_3 + \text{Cl}^- \longrightarrow \text{F}-\text{B}(\text{F})_2-\text{Cl}$
- c. 
- d. 


6.44 Draw an energy diagram for each reaction. Label the axes, the starting material, product, transition state, ΔH° , and E_a .

- a. A concerted, exothermic reaction with a low energy of activation.
- b. A one-step endothermic reaction with a high energy of activation.
- c. A two-step reaction, $\text{A} \rightarrow \text{B} \rightarrow \text{C}$, in which the relative energy of the compounds is $\text{A} < \text{C} < \text{B}$, and the step $\text{A} \rightarrow \text{B}$ is rate-determining.
- d. A concerted reaction with $\Delta H^\circ = -80 \text{ kJ/mol}$ and $E_a = 16 \text{ kJ/mol}$.

6.45 Consider the following reaction: $\text{CH}_4 + \text{Cl}\cdot \rightarrow \cdot\text{CH}_3 + \text{HCl}$.

- a. Use curved arrows to show the movement of electrons.
- b. Calculate ΔH° using the bond dissociation energies in Table 6.2.
- c. Draw an energy diagram assuming that $E_a = 16 \text{ kJ/mol}$.
- d. What is E_a for the reverse reaction ($\cdot\text{CH}_3 + \text{HCl} \rightarrow \text{CH}_4 + \text{Cl}\cdot$)?


6.46 Consider the following energy diagram for the conversion of **A** \rightarrow **G**.



- a. Which points on the graph correspond to transition states?
- b. Which points on the graph correspond to reactive intermediates?
- c. How many steps are present in the reaction mechanism?
- d. Label each step of the mechanism as endothermic or exothermic.
- e. Label the overall reaction as endothermic or exothermic.


6.47 Draw an energy diagram for the Brønsted–Lowry acid–base reaction of $\text{CH}_3\text{CO}_2\text{H}$ with $^-\text{OC}(\text{CH}_3)_3$ to form CH_3CO_2^- and $(\text{CH}_3)_3\text{COH}$. Label the axes, starting materials, products, ΔH° , and E_a . Draw the structure of the transition state.

6.48 Consider the following two-step reaction:



- a. How many bonds are broken and formed in Step [1]? Would you predict ΔH° of Step [1] to be positive or negative?
- b. How many bonds are broken and formed in Step [2]? Would you predict the ΔH° of Step [2] to be positive or negative?
- c. Which step is rate-determining?
- d. Draw the structure for the transition state in both steps of the mechanism.
- e. If $\Delta H^\circ_{\text{overall}}$ is negative for this two-step reaction, draw an energy diagram illustrating all of the information in parts (a)–(d).

6.49 Consider the following energy diagram for the overall reaction: $(\text{CH}_3)_3\text{COH} + \text{HI} \rightarrow (\text{CH}_3)_3\text{CI} + \text{H}_2\text{O}$.



- How many steps are in the reaction mechanism?
- Label the E_a and ΔH° for each step, and the $\Delta H^\circ_{\text{overall}}$ for the reaction.
- Draw the structure of the transition state for each step and indicate its location on the energy diagram.
- Which step is rate-determining? Why?

Kinetics and Rate Laws


6.50 Indicate which factors affect the rate of a reaction.

- | | | |
|---------------------|--------------------|---------------------|
| a. ΔG° | d. temperature | g. k |
| b. ΔH° | e. concentration | h. catalysts |
| c. E_a | f. K_{eq} | i. ΔS° |

6.51 The following is a concerted, bimolecular reaction: $\text{CH}_3\text{Br} + \text{NaCN} \rightarrow \text{CH}_3\text{CN} + \text{NaBr}$.

- What is the rate equation for this reaction?
- What happens to the rate of the reaction if $[\text{CH}_3\text{Br}]$ is doubled?
- What happens to the rate of the reaction if $[\text{NaCN}]$ is halved?
- What happens to the rate of the reaction if $[\text{CH}_3\text{Br}]$ and $[\text{NaCN}]$ are both increased by a factor of five?


6.52 The conversion of acetyl chloride to methyl acetate occurs via the following two-step mechanism:



- Write the rate equation for this reaction, assuming the first step is rate-determining.
 - If the concentration of $^-\text{OCH}_3$ were increased 10 times, what would happen to the rate of the reaction?
 - If the concentrations of both CH_3COCl and $^-\text{OCH}_3$ were increased 10 times, what would happen to the rate of the reaction?
 - Classify the conversion of acetyl chloride to methyl acetate as an addition, elimination, or substitution.
- 6.53 Label each statement as true or false. Correct any false statement to make it true.
- Increasing temperature increases reaction rate.
 - If a reaction is fast, it has a large rate constant.
 - A fast reaction has a large negative ΔG° value.
 - When E_a is large, the rate constant k is also large.
 - Fast reactions have equilibrium constants > 1 .
 - Increasing the concentration of a reactant always increases the rate of a reaction.

General Problems

6.54 The conversion of $(\text{CH}_3)_3\text{CI}$ to $(\text{CH}_3)_2\text{C}=\text{CH}_2$ can occur by either a one-step or a two-step mechanism, as shown in Equations [1] and [2].




- What rate equation would be observed for the mechanism in Equation [1]?
- What rate equation would be observed for the mechanism in Equation [2]?
- What is the order of each rate equation (i.e., first, second, and so forth)?
- How can these rate equations be used to show which mechanism is the right one for this reaction?
- Assume Equation [1] represents an endothermic reaction and draw an energy diagram for the reaction. Label the axes, reactants, products, E_a , and ΔH° . Draw the structure for the transition state.
- Assume Equation [2] represents an endothermic reaction and that the product of the rate-determining step is higher in energy than the reactants or products. Draw an energy diagram for this two-step reaction. Label the axes, reactants and products for each step, and the E_a and ΔH° for each step. Label $\Delta H^\circ_{\text{overall}}$. Draw the structure for both transition states.

Challenge Problems


6.55 Explain why $\text{HC}\equiv\text{CH}$ is more acidic than CH_3CH_3 , even though the C–H bond in $\text{HC}\equiv\text{CH}$ has a higher bond dissociation energy than the C–H bond in CH_3CH_3 .

6.56



- What carbon radical is formed by homolysis of the C–H_a bond in propylbenzene? Draw all reasonable resonance structures for this radical.
- What carbon radical is formed by homolysis of the C–H_b bond in propylbenzene? Draw all reasonable resonance structures for this radical.
- The bond dissociation energy of one of the C–H bonds is considerably less than the bond dissociation energy of the other. Which C–H bond is weaker? Offer an explanation.


6.57 Esterification is the reaction of a carboxylic acid (RCOOH) with an alcohol ($\text{R}'\text{OH}$) to form an ester (RCOOR') with loss of water. Equation [1] is an example of an *intermolecular* esterification reaction. Equation [2] is an example of an *intramolecular* esterification reaction; that is, the carboxylic acid and alcohol are contained in the same starting material, forming a cyclic ester as product. The equilibrium constants for both reactions are given. Explain why K_{eq} is different for these two apparently similar reactions.



6.58 Although K_{eq} of Equation [1] in Problem 6.57 does not greatly favor formation of the product, it is sometimes possible to use Le Châtelier's principle to increase the yield of ethyl acetate. Le Châtelier's principle states that if an equilibrium is disturbed, a system will react to counteract this disturbance. How can Le Châtelier's principle be used to drive the equilibrium to increase the yield of ethyl acetate? Another example of Le Châtelier's principle is given in Section 9.8.

6.59 As we will learn in Section 15.12, many antioxidants—compounds that prevent unwanted radical oxidation reactions from occurring—are phenols, compounds that contain an OH group bonded directly to a benzene ring.

- Explain why homolysis of the O–H bond in phenol requires considerably less energy than homolysis of the O–H bond in ethanol (362 kJ/mol vs. 438 kJ/mol).
- Why is the C–O bond in phenol shorter than the C–O bond in ethanol?



7

Alkyl Halides and Nucleophilic Substitution

- 7.1 Introduction to alkyl halides
- 7.2 Nomenclature
- 7.3 Physical properties
- 7.4 Interesting alkyl halides
- 7.5 The polar carbon–halogen bond
- 7.6 General features of nucleophilic substitution
- 7.7 The leaving group
- 7.8 The nucleophile
- 7.9 Possible mechanisms for nucleophilic substitution
- 7.10 Two mechanisms for nucleophilic substitution
- 7.11 The S_N2 mechanism
- 7.12 Application: Useful S_N2 reactions
- 7.13 The S_N1 mechanism
- 7.14 Carbocation stability
- 7.15 The Hammond postulate
- 7.16 Application: S_N1 reactions, nitrosamines, and cancer
- 7.17 When is the mechanism S_N1 or S_N2 ?
- 7.18 Vinyl halides and aryl halides
- 7.19 Organic synthesis




Adrenaline (or **epinephrine**), a hormone secreted by the adrenal gland, increases blood pressure and heart rate, and dilates lung passages. Individuals often speak of the “rush of adrenaline” when undertaking a particularly strenuous or challenging activity. Adrenaline is made in the body by a simple organic reaction called **nucleophilic substitution**. In Chapter 7 we learn about the mechanism of nucleophilic substitution and how adrenaline is synthesized in organisms.

This is the first of three chapters dealing with an in-depth study of the organic reactions of compounds containing C–Z σ bonds, where Z is an element more electronegative than carbon. In Chapter 7 we learn about **alkyl halides** and one of their characteristic reactions, **nucleophilic substitution**. In Chapter 8, we look at **elimination**, a second general reaction of alkyl halides. We conclude this discussion in Chapter 9 by examining other molecules that also undergo nucleophilic substitution and elimination reactions.

7.1 Introduction to Alkyl Halides

Alkyl halides are organic molecules containing a halogen atom X bonded to an sp^3 hybridized carbon atom. Alkyl halides are classified as **primary** (1°), **secondary** (2°), or **tertiary** (3°) depending on the number of carbons bonded to the carbon with the halogen.

Alkyl halides have the general molecular formula $C_nH_{2n+1}X$, and are formally derived from an alkane by replacing a hydrogen atom with a halogen.



Whether an alkyl halide is 1° , 2° , or 3° is the *most important factor* in determining the course of its chemical reactions. Figure 7.1 illustrates three examples.

Four types of organic halides having the halogen atom in close proximity to a π bond are illustrated in Figure 7.2. **Vinyl halides** have a halogen atom bonded to a carbon–carbon double bond, and **aryl halides** have a halogen atom bonded to a benzene ring. These two types of organic halides with X bonded directly to an sp^2 hybridized carbon atom do *not* undergo the reactions presented in Chapter 7, as discussed in Section 7.18.

Allylic halides and benzylic halides have halogen atoms bonded to sp^3 hybridized carbon atoms and *do* undergo the reactions described in Chapter 7. **Allylic halides** have X bonded to the carbon atom *adjacent* to a carbon–carbon double bond, and **benzylic halides** have X bonded to the carbon atom *adjacent* to a benzene ring. The synthesis of allylic and benzylic halides is discussed in Sections 15.10 and 18.13, respectively.

Problem 7.1 Classify each alkyl halide as 1° , 2° , or 3° .




Figure 7.1

Examples of 1° , 2° , and 3° alkyl halides





Figure 7.2


Four types of organic halides (RX) having X near a π bond



These organic halides are **unreactive** in the reactions discussed in Chapter 7.

These organic halides do participate in the reactions discussed in Chapter 7.

Problem 7.2 Fluticasone is the inhaled anti-inflammatory agent in the nasal spray Flonase and the asthma medication Advair. Classify the circled F atoms in fluticasone as 1°, 2°, or 3°. Why is it impossible to classify the remaining F atom as 1°, 2°, or 3° using the definitions in Section 7.1?



Problem 7.3 Draw the structure of an alkyl bromide with molecular formula $C_6H_{13}Br$ that fits each description: (a) a 1° alkyl bromide with one stereogenic center; (b) a 2° alkyl bromide with two stereogenic centers; (c) an achiral 3° alkyl bromide.

7.2 Nomenclature


The systematic (IUPAC) method for naming alkyl halides follows from the basic rules described in Chapter 4. Common names are also discussed in Section 7.2B, because many low molecular weight alkyl halides are often referred to by their common names.

7.2A IUPAC System


An alkyl halide is named as an alkane with a halogen substituent—that is, as a *halo alkane*. To name a halogen substituent, change the *-ine* ending of the name of the halogen to the suffix *-o* (chlorine \rightarrow chloro).

HOW TO Name an Alkyl Halide Using the IUPAC System

Example Give the IUPAC name of the following alkyl halide:



Step [1] Find the parent carbon chain containing the halogen.




7 C's in the longest chain

7 C's \rightarrow heptane

- Name the parent chain as an *alkane*, with the halogen as a substituent bonded to the longest chain.


Step [2] Apply all other rules of nomenclature.

a. **Number** the chain.



- Begin at the end nearest the first substituent, either alkyl or halogen.

b. **Name and number** the substituents.




c. **Alphabetize:** c for chloro, then m for methyl.

ANSWER: 2-chloro-5-methylheptane

Figure 7.3


Examples: Nomenclature of alkyl halides



7.2B Common Names


Common names for alkyl halides are used only for simple alkyl halides. To assign a common name:

- Name all the carbon atoms of the molecule as a single **alkyl group**.
- Name the halogen bonded to the alkyl group. To name the halogen, change the *-ine* ending of the halogen name to the suffix *-ide*; for example, **bromine** → **bromide**.
- Combine the names of the alkyl group and halide, separating the words with a space.



Other examples of alkyl halide nomenclature are given in Figure 7.3.

Problem 7.4 Give the IUPAC name for each compound.




Problem 7.5 Give the structure corresponding to each name.

- | | |
|---|------------------------------|
| a. 3-chloro-2-methylhexane | d. 1,1,3-tribromocyclohexane |
| b. 4-ethyl-5-iodo-2,2-dimethyloctane | e. propyl chloride |
| c. <i>cis</i> -1,3-dichlorocyclopentane | f. <i>sec</i> -butyl bromide |

7.3 Physical Properties

Alkyl halides are weakly polar molecules. They exhibit **dipole–dipole** interactions because of their polar C–X bond, but because the rest of the molecule contains only C–C and C–H bonds they are incapable of intermolecular hydrogen bonding. How this affects their physical properties is summarized in Table 7.1.




Problem 7.6 Rank the compounds in each group in order of increasing boiling point.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{I}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$
 b. $\text{CH}_3(\text{CH}_2)_4\text{CH}_3$, $\text{CH}_3(\text{CH}_2)_5\text{Br}$, $\text{CH}_3(\text{CH}_2)_5\text{OH}$

Table 7.1 Physical Properties of Alkyl Halides

Property	Observation	
Boiling point and melting point	<ul style="list-style-type: none"> Alkyl halides have higher bp's and mp's than alkanes having the same number of carbons. <div style="text-align: center; margin: 5px 0;"> CH_3CH_3 and $\text{CH}_3\text{CH}_2\text{Br}$ bp = -89°C bp = 39°C </div> Bp's and mp's increase as the size of R increases. <div style="display: flex; justify-content: space-around; align-items: center; margin: 5px 0;"> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{Cl}$ mp = -136°C bp = 12°C </div> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ mp = -123°C bp = 47°C </div> <div style="border: 1px solid black; padding: 2px; margin-left: 10px;"> larger surface area higher mp and bp </div> </div> Bp's and mp's increase as the size of X increases. <div style="display: flex; justify-content: space-around; align-items: center; margin: 5px 0;"> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{Cl}$ mp = -136°C bp = 12°C </div> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{Br}$ mp = -119°C bp = 39°C </div> <div style="border: 1px solid black; padding: 2px; margin-left: 10px;"> more polarizable halogen higher mp and bp </div> </div> 	
	Solubility	<ul style="list-style-type: none"> RX is soluble in organic solvents. RX is insoluble in water.

Problem 7.7 An sp^3 hybridized C–Cl bond is more polar than an sp^2 hybridized C–Cl bond. (a) Explain why this phenomenon arises. (b) Rank the following compounds in order of increasing boiling point.




7.4 Interesting Alkyl Halides

Many simple alkyl halides make excellent solvents because they are not flammable and dissolve a wide variety of organic compounds. Compounds in this category include CHCl_3 (chloroform or trichloromethane) and CCl_4 (carbon tetrachloride or tetrachloromethane). Large quantities of these solvents are produced industrially each year, but like many chlorinated organic compounds, both chloroform and carbon tetrachloride are toxic if inhaled or ingested. Other simple alkyl halides are shown in Figure 7.4.

Synthetic organic halides are also used in insulating materials, plastic wrap, and coatings. Two such compounds are **Teflon** and **poly(vinyl chloride) (PVC)**.






Asparagopsis taxiformis is an edible red seaweed that grows on the edges of reefs in areas of constant water motion. Almost 100 different organic halides have been isolated from this source.



Organic halides constitute a growing list of useful naturally occurring molecules, many produced by marine organisms. Some have irritating odors or an unpleasant taste and are synthesized by organisms for self-defense or feeding deterrents. Examples include $\text{Br}_2\text{C}=\text{CHCHCl}_2$ and $\text{Br}_2\text{C}=\text{CHCHBr}_2$, isolated from the red seaweed *Asparagopsis taxiformis*, known as *limu kohu* (supreme seaweed) in Hawaii. This seaweed has a strong and characteristic odor and flavor, in part probably because of these organic halides.

Figure 7.4


Some simple alkyl halides



- **Chloromethane (CH_3Cl)** is produced by giant kelp and algae and also found in emissions from volcanoes such as Hawaii's Kilauea. Almost all of the atmospheric chloromethane results from these natural sources.
- **Dichloromethane (or methylene chloride, CH_2Cl_2)** is an important solvent, once used to decaffeinate coffee. Coffee is now decaffeinated by using supercritical CO_2 due to concerns over the possible ill effects of trace amounts of residual CH_2Cl_2 in the coffee. Subsequent studies on rats have shown, however, that no cancers occurred when animals ingested the equivalent of over 100,000 cups of decaffeinated coffee per day.
- **Halothane (CF_3CHClBr)** is a safe general anesthetic that has now replaced other organic anesthetics such as CHCl_3 , which causes liver and kidney damage, and $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ (diethyl ether), which is very flammable.

DDT, a nonbiodegradable pesticide, has been labeled both a "miraculous" discovery by Winston Churchill in 1945 and the "elixir of death" by Rachel Carson in her 1962 book *Silent Spring*. DDT use was banned in the United States in 1973, but because of its effectiveness and low cost, it is still widely used to control insect populations in developing countries.

Although the beneficial effects of many organic halides are undisputed, certain synthetic chlorinated organics such as the **chlorofluorocarbons** and the pesticide **DDT** have caused lasting harm to the environment.



Chlorofluorocarbons (CFCs) have the general molecular structure $\text{CF}_x\text{Cl}_{4-x}$. Trichlorofluoromethane [CFCl_3 , CFC 11, or Freon 11 (trade name)] is an example of these easily vaporized compounds, having been extensively used as a refrigerant and an aerosol propellant. CFCs slowly rise to the stratosphere, where sunlight catalyzes their decomposition, a process that contributes to the destruction of the ozone layer, the thin layer of atmosphere that shields the earth's surface from harmful ultraviolet radiation (Section 15.9). Although it is now easy to second-guess the extensive use of CFCs, it is also easy to see why they were used so widely. **CFCs made refrigeration available to the general public.** Would you call your refrigerator a comfort or a necessity?


The story of the insecticide **DDT** (*dichlorodiphenyltrichloroethane*) follows the same theme: DDT is an organic molecule with valuable short-term effects that has caused long-term problems. DDT kills insects that spread diseases such as malaria and typhus, and in controlling insect populations, DDT has saved millions of lives worldwide. DDT is a weakly polar

organic compound that persists in the environment for years. Because DDT is soluble in organic media, it accumulates in fatty tissues. Most adults in the United States have low concentrations of DDT (or a degradation product of DDT) in their bodies. DDT is acutely toxic to many types of marine life (crayfish, sea shrimp, and some fish), but the long-term effect on humans is not known.



Time Magazine, June 30, 1947.

Problem 7.8 Although nonpolar compounds tend to dissolve and remain in fatty tissues, polar substances are more water soluble, and more readily excreted into an environment where they may be degraded by other organisms. Explain why methoxychlor is more biodegradable than DDT.




7.5 The Polar Carbon–Halogen Bond

The properties of alkyl halides dictate their reactivity. The electrostatic potential maps of four simple alkyl halides in Figure 7.5 illustrate that the electronegative halogen X creates a polar C–X bond, making the carbon atom electron deficient. **The chemistry of alkyl halides is determined by this polar C–X bond.**


What kind of reactions do alkyl halides undergo? **The characteristic reactions of alkyl halides are substitution and elimination.** Because alkyl halides contain an electrophilic carbon, they react with electron-rich reagents—Lewis bases (nucleophiles) and Brønsted–Lowry bases.

- Alkyl halides undergo substitution reactions with nucleophiles.



In a substitution reaction of RX, **the halogen X is replaced by an electron-rich nucleophile :Nu[−].** The C–X σ bond is broken and the C–Nu σ bond is formed.

- Alkyl halides undergo elimination reactions with Brønsted–Lowry bases.





In an elimination reaction of RX, **the elements of HX are removed by a Brønsted–Lowry base :B.**

The remainder of Chapter 7 is devoted to a discussion of the substitution reactions of alkyl halides. Elimination reactions are discussed in Chapter 8.

Figure 7.5 Electrostatic potential maps of four halomethanes (CH₃X)


General structure



- The polar C–X bond makes the carbon atom *electron deficient* in each CH₃X molecule.

7.6 General Features of Nucleophilic Substitution


Three components are necessary in any substitution reaction.




- [1] **R**—An alkyl group R containing an sp^3 hybridized carbon bonded to X.
- [2] **X**—An atom X (or a group of atoms) called a **leaving group**, which is able to accept the electron density in the C–X bond. The most common leaving groups are halide anions (X^-), but H_2O (from ROH_2^+) and N_2 (from RN_2^+) are also encountered.
- [3] **:Nu[−]**—A **nucleophile**. Nucleophiles contain a lone pair or a π bond but not necessarily a negative charge.

Because these substitution reactions involve electron-rich nucleophiles, they are called **nucleophilic substitution reactions**. Examples are shown in Equations [1]–[3]. **Nucleophilic substitutions are Lewis acid–base reactions**. The nucleophile donates its electron pair, the alkyl halide (Lewis acid) accepts it, and the C–X bond is heterolytically cleaved. Curved arrow notation can be used to show the movement of electron pairs, as shown in Equation [3].


Examples



Negatively charged nucleophiles like ^-OH and ^-SH are used as **salts** with Li^+ , Na^+ , or K^+ counterions to balance charge. The identity of the cation is usually inconsequential, and therefore it is often omitted from the chemical equation.




When a neutral nucleophile is used, the substitution product bears a positive charge. **Note that all atoms originally bonded to the nucleophile stay bonded to it after substitution occurs.** All three CH_3 groups stay bonded to the N atom in the given example.



The reaction of alkyl halides with NH_3 to form amines (RNH_2) is discussed in Chapter 25.


Furthermore, when the substitution product bears a positive charge and also contains a proton bonded to O or N, the initial substitution product readily loses a proton in a Brønsted–Lowry acid–base reaction, forming a neutral product.




All of these reactions are nucleophilic substitutions and have the same overall result—**replacement of the leaving group by the nucleophile**, regardless of the identity or charge of the nucleophile. To draw any nucleophilic substitution product:

- Find the sp^3 hybridized carbon with the leaving group.
- Identify the nucleophile, the species with a lone pair or π bond.
- Substitute the nucleophile for the leaving group and assign charges (if necessary) to any atom that is involved in bond breaking or bond formation.


Problem 7.9 Identify the nucleophile and leaving group and draw the products of each reaction.



Problem 7.10 Draw the product of nucleophilic substitution with each neutral nucleophile. When the initial substitution product can lose a proton to form a neutral product, draw that product as well.



Problem 7.11 CPC (cetylpyridinium chloride), an antiseptic found in throat lozenges and mouthwash, is synthesized by the following reaction. Draw the structure of CPC.



7.7 The Leaving Group




Cepacol throat lozenges and Crest Pro-Health Mouth Rinse contain the antiseptic CPC, which is prepared by nucleophilic substitution (Problem 7.11).

Nucleophilic substitution is a general reaction of organic compounds. Why, then, are alkyl halides the most common substrates, and halide anions the most common leaving groups? To answer this question, we must understand leaving group ability. **What makes a good leaving group?**


In a nucleophilic substitution reaction of R-X , the C-X bond is heterolytically cleaved, and the leaving group departs with the electron pair in that bond, forming X^- . **The more stable the leaving group X^- , the better able it is to accept an electron pair**, giving rise to the following generalization:

- In comparing two leaving groups, the better leaving group is the weaker base.




For example, H_2O is a better leaving group than ^-OH because H_2O is a weaker base. Moreover, the periodic trends in basicity can now be used to identify **periodic trends in leaving group ability**:

- Left-to-right across a row of the periodic table, basicity *decreases* so leaving group ability *increases*.



- Down a column of the periodic table, basicity *decreases* so leaving group ability *increases*.



All good leaving groups are weak bases with strong conjugate acids having low $\text{p}K_{\text{a}}$ values. Thus, all halide anions except F^- are good leaving groups because their conjugate acids (HCl, HBr, and HI) have low $\text{p}K_{\text{a}}$ values. Tables 7.2 and 7.3 list good and poor leaving groups for nucleophilic substitution reactions, respectively. Nucleophilic substitution does not occur with any of the leaving groups in Table 7.3 because these leaving groups are strong bases.

Table 7.2 Good Leaving Groups for Nucleophilic Substitution

Starting material	Leaving group	Conjugate acid	$\text{p}K_{\text{a}}$
$\text{R}-\text{Cl}$	Cl^-	HCl	-7
$\text{R}-\text{Br}$	Br^-	HBr	-9
$\text{R}-\text{I}$	I^-	HI	-10
$\text{R}-\text{OH}_2^+$	H_2O	H_3O^+	-1.7

↑ These molecules undergo nucleophilic substitution. ↑ good leaving groups

Table 7.3 Poor Leaving Groups for Nucleophilic Substitution

Starting material	Leaving group	Conjugate acid	$\text{p}K_{\text{a}}$
$\text{R}-\text{F}$	F^-	HF	3.2
$\text{R}-\text{OH}$	$\text{}^-\text{OH}$	H_2O	15.7
$\text{R}-\text{NH}_2$	$\text{}^-\text{NH}_2$	NH_3	38
$\text{R}-\text{H}$	H^-	H_2	35
$\text{R}-\text{R}$	R^-	RH	50

↑ These molecules do *not* undergo nucleophilic substitution. ↑ poor leaving groups

Problem 7.12 Which is the better leaving group in each pair?
 a. Cl^- , I^- b. NH_3 , NH_2^- c. H_2O , H_2S


Problem 7.13 Which molecules contain good leaving groups?
 a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ c. $\text{CH}_3\text{CH}_2\text{OH}_2^+$ d. CH_3CH_3

Given a particular nucleophile and leaving group, how can we determine whether the equilibrium will favor products in a nucleophilic substitution? We can often correctly predict the direction of equilibrium by comparing the basicity of the nucleophile and the leaving group.

- Equilibrium favors the products of nucleophilic substitution when the leaving group is a weaker base than the nucleophile.


Sample Problem 7.1 illustrates how to apply this general rule.

Sample Problem 7.1 Will the following substitution reaction favor formation of the products?




Solution

Compare the basicity of the nucleophile (OH^-) and the leaving group (Cl^-) by comparing the pK_a values of their conjugate acids. **The stronger the conjugate acid, the weaker the base, and the better the leaving group.**



Because Cl^- , the leaving group, is a weaker base than OH^- , the nucleophile, the reaction favors the products.

Problem 7.14 Does the equilibrium favor the reactants or products in each substitution reaction?




Problem 7.15 Should it be possible to convert $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ to $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ by a nucleophilic substitution reaction with NaCl ? Explain why or why not.

7.8 The Nucleophile

We use the word *base* to mean *Bronsted-Lowry base* and the word *nucleophile* to mean a *Lewis base* that reacts with electrophiles *other than protons*.

Nucleophiles and bases are structurally similar: both have a lone pair or a π bond. They differ in what they attack.

- Bases attack protons. Nucleophiles attack other electron-deficient atoms (usually carbons).



7.8A Nucleophilicity Versus Basicity

How is **nucleophilicity** (nucleophile strength) related to **basicity**? Although it is generally true that **a strong base is a strong nucleophile**, nucleophile size and steric factors can sometimes change this relationship.

Nucleophilicity parallels basicity in three instances:


[1] **For two nucleophiles with the same nucleophilic atom, the stronger base is the stronger nucleophile.**

- The relative nucleophilicity of OH^- and CH_3COO^- , two oxygen nucleophiles, is determined by comparing the $\text{p}K_{\text{a}}$ values of their conjugate acids (H_2O and CH_3COOH). CH_3COOH ($\text{p}K_{\text{a}} = 4.8$) is a stronger acid than H_2O ($\text{p}K_{\text{a}} = 15.7$), so OH^- is a stronger base and stronger nucleophile than CH_3COO^- .

[2] **A negatively charged nucleophile is always stronger than its conjugate acid.**

- OH^- is a stronger base and stronger nucleophile than H_2O , its conjugate acid.

[3] **Right-to-left across a row of the periodic table, nucleophilicity increases as basicity increases.**



Problem 7.16

Identify the stronger nucleophile in each pair.


- a. NH_3 , NH_2^- b. CH_3^- , HO^- c. CH_3NH_2 , CH_3OH d. CH_3COO^- , $\text{CH}_3\text{CH}_2\text{O}^-$

7.8B Steric Effects and Nucleophilicity

All steric effects arise because two atoms cannot occupy the same space. In Chapter 4, for example, we learned that **steric strain** is an increase in energy when big groups (occupying a large volume) are forced close to each other.

Nucleophilicity does not parallel basicity when **steric hindrance** becomes important. **Steric hindrance** is a decrease in reactivity resulting from the presence of bulky groups at the site of a reaction.

For example, although $\text{p}K_{\text{a}}$ tables indicate that *tert*-butoxide [$(\text{CH}_3)_3\text{CO}^-$] is a stronger base than ethoxide ($\text{CH}_3\text{CH}_2\text{O}^-$), **ethoxide is the stronger nucleophile**. The three CH_3 groups around the O atom of *tert*-butoxide create steric hindrance, making it more difficult for this big, bulky base to attack a tetravalent carbon atom.






Three CH_3 groups sterically hinder the O atom, making it a **weaker nucleophile**.

Steric hindrance decreases nucleophilicity but not basicity. Because bases pull off small, easily accessible protons, they are unaffected by steric hindrance. Nucleophiles, on the other hand, must attack a crowded tetrahedral carbon, so bulky groups decrease reactivity.

Sterically hindered bases that are poor nucleophiles are called nonnucleophilic bases. Potassium *tert*-butoxide [$\text{K}^+\text{OC}(\text{CH}_3)_3$] is a strong, nonnucleophilic base.

Figure 7.6
Examples of polar protic solvents



7.8C Comparing Nucleophiles of Different Size—Solvent Effects


Atoms vary greatly in size down a column of the periodic table, and in this case, **nucleophilicity depends on the solvent used in a substitution reaction**. Although solvent has thus far been ignored, most organic reactions take place in a liquid solvent that dissolves all reactants to some extent. Because substitution reactions involve polar starting materials, polar solvents are used to dissolve them. There are two main kinds of polar solvents—**polar protic solvents and polar aprotic solvents**.

Polar Protic Solvents

In addition to dipole–dipole interactions, **polar protic solvents are capable of intermolecular hydrogen bonding**, because they contain an O–H or N–H bond. The most common polar protic solvents are water and alcohols (ROH), as seen in the examples in Figure 7.6. **Polar protic solvents solvate both cations and anions well**.

- Cations are solvated by ion–dipole interactions.
- Anions are solvated by hydrogen bonding.

For example, if the salt NaBr is used as a source of the nucleophile Br[−] in H₂O, the Na⁺ cations are solvated by ion–dipole interactions with H₂O molecules, and the Br[−] anions are solvated by strong hydrogen bonding interactions.



How do polar protic solvents affect nucleophilicity? **In polar protic solvents, nucleophilicity increases down a column of the periodic table as the size of the anion increases. This is opposite to basicity.** A small electronegative anion like F[−] is very well solvated by hydrogen bonding, effectively shielding it from reaction. On the other hand, a large, less electronegative anion like I[−] does not hold onto solvent molecules as tightly. The solvent does not “hide” a large nucleophile as well, and the nucleophile is much more able to donate its electron pairs in a reaction. Thus, **nucleophilicity increases down a column** even though basicity decreases, giving rise to the following trend in polar protic solvents:







I[−] is a weak base but a strong nucleophile in polar protic solvents.

Down a column of the periodic table



Increasing nucleophilicity in polar protic solvents

Figure 7.7
Examples of polar aprotic solvents




Abbreviations are often used in organic chemistry, instead of a compound's complete name. A list of common abbreviations is given on the inside back cover.

Polar Aprotic Solvents

Polar aprotic solvents also exhibit dipole–dipole interactions, but they have no O–H or N–H bond so they are **incapable of hydrogen bonding**. Examples of polar aprotic solvents are shown in Figure 7.7. **Polar aprotic solvents solvate only cations well.**


- Cations are solvated by ion–dipole interactions.
- Anions are not well solvated because the solvent cannot hydrogen bond to them.

When the salt NaBr is dissolved in acetone, $(\text{CH}_3)_2\text{C}=\text{O}$, the Na^+ cations are solvated by ion–dipole interactions with the acetone molecules, but, with no possibility for hydrogen bonding, the Br^- anions are not well solvated. Often these anions are called **naked anions** because they are not bound by tight interactions with solvent.



How do polar aprotic solvents affect nucleophilicity? Because anions are not well solvated in polar aprotic solvents, there is no need to consider whether solvent molecules more effectively hide one anion than another. Nucleophilicity once again parallels basicity and **the stronger base is the stronger nucleophile**. Because basicity decreases with size down a column, nucleophilicity decreases as well:

Down a column
of the periodic table



← Increasing nucleophilicity
in polar aprotic solvents

Problem 7.17 Classify each solvent as protic or aprotic.
 a. HOCH₂CH₂OH b. CH₃CH₂OCH₂CH₃ c. CH₃COOCH₂CH₃

Problem 7.18 Identify the stronger nucleophile in each pair of anions.
 a. Br⁻ or Cl⁻ in a polar protic solvent c. HS⁻ or F⁻ in a polar protic solvent
 b. HO⁻ or Cl⁻ in a polar aprotic solvent

7.8D Summary

This long discussion of nucleophilicity has brought together many new concepts, such as steric hindrance and solvent effects, both of which we will meet again in our study of organic chemistry. Keep in mind, however, the central relationship between nucleophilicity and basicity in comparing two nucleophiles.

- It is generally true that the *stronger* base is the *stronger* nucleophile.
- In polar protic solvents, however, nucleophilicity increases with increasing size of an anion (opposite to basicity).
- Steric hindrance decreases nucleophilicity without decreasing basicity, making (CH₃)₃CO⁻ a stronger base but a weaker nucleophile than CH₃CH₂O⁻.

Table 7.4 lists some common nucleophiles used in nucleophilic substitution reactions.

Problem 7.19 Rank the nucleophiles in each group in order of increasing nucleophilicity.
 a. ⁻OH, ⁻NH₂, H₂O b. ⁻OH, Br⁻, F⁻ (polar aprotic solvent) c. H₂O, ⁻OH, CH₃COO⁻

Problem 7.20 What nucleophile is needed to convert (CH₃)₂CHCH₂CH₂-Br to each product?
 a. (CH₃)₂CHCH₂CH₂-SH c. (CH₃)₂CHCH₂CH₂-OCOCH₃
 b. (CH₃)₂CHCH₂CH₂-OCH₂CH₃ d. (CH₃)₂CHCH₂CH₂-C≡CH


Table 7.4 Common Nucleophiles in Organic Chemistry

	Negatively charged nucleophiles			Neutral nucleophiles	
Oxygen	⁻ OH	⁻ OR	CH ₃ COO ⁻	H ₂ O	ROH
Nitrogen	N ₃ ⁻			NH ₃	RNH ₂
Carbon	⁻ CN	HC≡C ⁻			
Halogen	Cl ⁻	Br ⁻	I ⁻		
Sulfur	HS ⁻	RS ⁻		H ₂ S	RSH


7.9 Possible Mechanisms for Nucleophilic Substitution

Now that you know something about the general features of nucleophilic substitution, you can begin to understand the mechanism.


Overall reaction




Nucleophilic substitution at an sp^3 hybridized carbon involves two σ bonds: the bond to the leaving group, which is broken, and the bond to the nucleophile, which is formed. To understand the mechanism of this reaction, though, we must know the timing of these two events; that is, **what is the order of bond breaking and bond making?** Do they happen at the same time, or does one event precede the other? There are three possibilities.

[1] Bond breaking and bond making occur at the *same* time.

- If the C–X bond is broken **as** the C–Nu bond is formed, the mechanism has **one step**. As we learned in Section 6.9, the rate of such a bimolecular reaction depends on the concentration of both reactants; that is, the rate equation is **second order**.

[2] Bond breaking occurs *before* bond making.

- If the C–X bond is broken **first** and then the C–Nu bond is formed, the mechanism has **two steps** and a **carbocation** is formed as an intermediate. Because the first step is rate-determining, the rate depends on the concentration of RX only; that is, the rate equation is **first order**.

[3] Bond making occurs *before* bond breaking.


- If the C–Nu bond is formed **first** and then the C–X bond is broken, the mechanism has **two steps**, but this mechanism has an inherent problem. The intermediate generated in the first step has 10 electrons around carbon, violating the octet rule. Because two other mechanistic possibilities do not violate a fundamental rule, this last possibility can be disregarded.

The preceding discussion has generated two possible mechanisms for nucleophilic substitution: **a one-step mechanism in which bond breaking and bond making are simultaneous, and a two-step mechanism in which bond breaking comes before bond making.** In Section 7.10 we look at data for two specific nucleophilic substitution reactions and see if those data fit either of these proposed mechanisms.


7.10 Two Mechanisms for Nucleophilic Substitution

Rate equations for two different reactions give us insight into the possible mechanism for nucleophilic substitution.

Reaction of bromomethane (CH_3Br) with the nucleophile acetate (CH_3COO^-) affords the substitution product methyl acetate with loss of Br^- as the leaving group (Equation [1]). Kinetic data show that the reaction rate depends on the concentration of *both* reactants; that is, the rate equation is **second order**. This suggests a **bimolecular reaction with a one-step mechanism** in which the C–X bond is broken *as* the C–Nu bond is formed.



Equation [2] illustrates a similar nucleophilic substitution reaction with a different alkyl halide, $(\text{CH}_3)_3\text{CBr}$, which also leads to substitution of Br^- by CH_3COO^- . Kinetic data show that this reaction rate depends on the concentration of only *one* reactant, the alkyl halide; that is, the rate equation is **first order**. This suggests a **two-step mechanism in which the rate-determining step involves the alkyl halide only**.



The numbers **1** and **2** in the names $\text{S}_{\text{N}}1$ and $\text{S}_{\text{N}}2$ refer to the kinetic order of the reactions. For example, $\text{S}_{\text{N}}2$ means that the kinetics are **second** order. The number 2 does *not* refer to the number of steps in the mechanism.


How can these two different results be explained? Although these two reactions have the same nucleophile and leaving group, **there must be two different mechanisms** because there are two different rate equations. These equations are specific examples of two well known mechanisms for nucleophilic substitution at an sp^3 hybridized carbon:

- The **$\text{S}_{\text{N}}2$ mechanism (substitution nucleophilic bimolecular)**, illustrated by the reaction in Equation [1].
- The **$\text{S}_{\text{N}}1$ mechanism (substitution nucleophilic unimolecular)**, illustrated by the reaction in Equation [2].

We will now examine the characteristics of the $\text{S}_{\text{N}}2$ and $\text{S}_{\text{N}}1$ mechanisms.

7.11 The $\text{S}_{\text{N}}2$ Mechanism

The reaction of CH_3Br with CH_3COO^- is an example of an **$\text{S}_{\text{N}}2$ reaction**. What are the general features of this mechanism?



7.11A Kinetics

An $\text{S}_{\text{N}}2$ reaction exhibits **second-order kinetics**; that is, the reaction is **bimolecular** and both the alkyl halide and the nucleophile appear in the rate equation.

- $\text{rate} = k[\text{CH}_3\text{Br}][\text{CH}_3\text{COO}^-]$

Changing the concentration of *either* reactant affects the rate. For example, doubling the concentration of *either* the nucleophile or the alkyl halide doubles the rate. Doubling the concentration of *both* reactants increases the rate by a factor of four.

Problem 7.21

What happens to the rate of an $\text{S}_{\text{N}}2$ reaction under each of the following conditions?

- $[\text{RX}]$ is tripled, and $[\text{:Nu}^-]$ stays the same.
- Both $[\text{RX}]$ and $[\text{:Nu}^-]$ are tripled.
- $[\text{RX}]$ is halved, and $[\text{:Nu}^-]$ stays the same.
- $[\text{RX}]$ is halved, and $[\text{:Nu}^-]$ is doubled.


7.11B A One-Step Mechanism

The most straightforward explanation for the observed second-order kinetics is a **concerted reaction**—**bond breaking and bond making occur at the same time**, as shown in Mechanism 7.1.




Mechanism 7.1 The S_N2 Mechanism

One step The C–Br bond breaks as the C–O bond forms.



An energy diagram for the reaction of $\text{CH}_3\text{Br} + \text{CH}_3\text{COO}^-$ is shown in Figure 7.8. The reaction has one step, so there is one energy barrier between reactants and products. Because the equilibrium for this S_N2 reaction favors the products, they are drawn at lower energy than the starting materials.

Problem 7.22 Draw the structure of the transition state in each of the following S_N2 reactions.



Problem 7.23 Draw an energy diagram for the reaction in Problem 7.22a. Label the axes, the starting material, the product, and the transition state. Assume the reaction is exothermic. Label ΔH° and E_a .

7.11C Stereochemistry of the S_N2 Reaction


From what direction does the nucleophile approach the substrate in an S_N2 reaction? There are two possibilities.

- **Frontside attack:** The nucleophile approaches from the *same* side as the leaving group.
- **Backside attack:** The nucleophile approaches from the side *opposite* the leaving group.

The results of frontside and backside attack of a nucleophile are illustrated with $\text{CH}_3\text{CH}(\text{D})\text{Br}$ as substrate and the general nucleophile $:\text{Nu}^-$. This substrate has the leaving group bonded to a stereogenic center, thus allowing us to see the structural difference that results when the nucleophile attacks from two different directions.

Figure 7.8

An energy diagram for the S_N2 reaction:
 $\text{CH}_3\text{Br} + \text{CH}_3\text{COO}^- \rightarrow \text{CH}_3\text{COOCH}_3 + \text{Br}^-$




- In the transition state, the C–Br bond is partially broken, the C–O bond is partially formed, and both the attacking nucleophile and the departing leaving group bear a partial negative charge.

In **frontside attack**, the nucleophile approaches from the **same** side as the leaving group, forming **A**. In this example, the leaving group was drawn on the right, so the nucleophile attacks from the right, and all other groups remain in their original positions. Because the nucleophile and leaving group are in the same position relative to the other three groups on carbon, frontside attack results in **retention of configuration** around the stereogenic center.

Recall from Section 1.1 that D stands for the isotope deuterium (^2H).


Frontside attack



Nu replaces Br on the **same** side.


In **backside attack**, the nucleophile approaches from the **opposite** side to the leaving group, forming **B**. In this example, the leaving group was drawn on the right, so the nucleophile attacks from the left. Because the nucleophile and leaving group are in the opposite position relative to the other three groups on carbon, backside attack results in **inversion of configuration** around the stereogenic center.

Backside attack



Nu replaces Br on the **opposite** side.

The products of frontside and backside attack are *different* compounds. **A** and **B** are stereoisomers that are nonsuperimposable—they are **enantiomers**.



Inversion of configuration in an $\text{S}_{\text{N}}2$ reaction is often called **Walden inversion**, after Latvian chemist Dr. Paul Walden, who first observed this process in 1896.

Which product is formed in an $\text{S}_{\text{N}}2$ reaction? When the stereochemistry of the product is determined, **only B, the product of backside attack, is formed.**

- All $\text{S}_{\text{N}}2$ reactions proceed with backside attack of the nucleophile, resulting in *inversion* of configuration at a stereogenic center.

Backside attack resulting in inversion of configuration occurs in all $\text{S}_{\text{N}}2$ reactions, but we can observe this change only when the leaving group is bonded to a stereogenic center.

One explanation for backside attack is based on an electronic argument. Both the nucleophile and leaving group are electron rich and these like charges repel each other. Backside attack keeps these two groups as far away from each other as possible. In the transition state, the nucleophile and leaving group are 180° away from each other, and the other three groups around carbon occupy a plane, as illustrated in Figure 7.9.

Two additional examples of inversion of configuration in $\text{S}_{\text{N}}2$ reactions are given in Figure 7.10.

Figure 7.9
Stereochemistry of
the S_N2 reaction






Figure 7.10
Two examples of inversion
of configuration in
the S_N2 reaction



- The bond to the nucleophile in the product is always on the **opposite side** relative to the bond to the leaving group in the starting material.


Sample Problem 7.2

Draw the product (including stereochemistry) of the following S_N2 reaction.




Solution

Br⁻ is the leaving group and ⁻CN is the nucleophile. Because S_N2 reactions proceed with **inversion** of configuration and the leaving group is drawn above the ring (on a wedge), the nucleophile must come in from below.



Backside attack converts the starting material, which has two groups **cis** to each other, to a product with two groups **trans** to each other because the nucleophile (⁻CN) attacks from below the plane of the ring.


Problem 7.24 Draw the product of each S_N2 reaction and indicate stereochemistry.



7.11D The Identity of the R Group

How does the rate of an S_N2 reaction change as the alkyl group in the substrate alkyl halide changes from $CH_3 \rightarrow 1^\circ \rightarrow 2^\circ \rightarrow 3^\circ$?

- As the number of R groups on the carbon with the leaving group *increases*, the rate of an S_N2 reaction *decreases*.



- Methyl and 1° alkyl halides undergo S_N2 reactions with ease.
- 2° Alkyl halides react more slowly.
- 3° Alkyl halides *do not* undergo S_N2 reactions.

This order of reactivity can be explained by steric effects. As small H atoms are replaced by larger alkyl groups, **steric hindrance caused by bulky R groups makes nucleophilic attack from the back side more difficult**, slowing the reaction rate. Figure 7.11 illustrates the effect of increasing steric hindrance in a series of alkyl halides.

The effect of steric hindrance on the rate of an S_N2 reaction is reflected in the energy of the transition state, too. Let's compare the reaction of ^-OH with two different alkyl halides, CH_3Br and $(CH_3)_2CHBr$, as shown in Figure 7.12. The transition state of each S_N2 reaction consists of five groups around the central carbon atom—three bonds to either H or R groups and two partial bonds to the leaving group and the nucleophile. **Crowding around the central carbon atom increases as H atoms are successively replaced by R groups**, so the central carbon is much more sterically hindered in the transition state for $(CH_3)_2CHBr$ than for CH_3Br . This increased crowding in the transition state makes it higher in energy (increases E_a), so the rate of the S_N2 reaction decreases.




Figure 7.11
Steric effects in the S_N2 reaction






Figure 7.12

Two energy diagrams depicting the effect of steric hindrance in S_N2 reactions



- CH_3Br is an unhindered alkyl halide. The transition state in the S_N2 reaction is lower in energy, making E_a lower and increasing the reaction rate.

- $(\text{CH}_3)_2\text{CHBr}$ is a sterically hindered alkyl halide. The transition state in the S_N2 reaction is higher in energy, making E_a higher and decreasing the reaction rate.

- Increasing the number of R groups on the carbon with the leaving group *increases crowding in the transition state, decreasing the rate of an S_N2 reaction.*

- The S_N2 reaction is fastest with unhindered halides.

Problem 7.25 Which compound in each pair undergoes a faster S_N2 reaction?



Problem 7.26 Explain why $(\text{CH}_3)_3\text{CCH}_2\text{Br}$, a 1° alkyl halide, undergoes S_N2 reactions very slowly.

Table 7.5 summarizes what we have learned thus far about an S_N2 mechanism.


Table 7.5 Characteristics of the S_N2 Mechanism

Characteristic	Result
Kinetics	<ul style="list-style-type: none"> • Second-order kinetics; rate = $k[\text{RX}][:\text{Nu}^-]$
Mechanism	<ul style="list-style-type: none"> • One step
Stereochemistry	<ul style="list-style-type: none"> • Backside attack of the nucleophile • Inversion of configuration at a stereogenic center
Identity of R	<ul style="list-style-type: none"> • Unhindered halides react fastest. • Rate: $\text{CH}_3\text{X} > \text{RCH}_2\text{X} > \text{R}_2\text{CHX} > \text{R}_3\text{CX}$


Adrenaline (epinephrine), the molecule that opened Chapter 7, is a hormone synthesized in the adrenal glands from noradrenaline (norepinephrine) by nucleophilic substitution using SAM (Figure 7.14). When an individual senses danger or is confronted by stress, the hypothalamus region of the brain signals the adrenal glands to synthesize and release adrenaline, which enters the bloodstream and then stimulates a response in many organs. Stored carbohydrates are metabolized in the liver to form glucose, which is further metabolized to provide an energy boost. Heart rate and blood pressure increase, and lung passages are dilated. These physiological changes prepare an individual for “fight or flight.”

Figure 7.14

Adrenaline synthesis from noradrenaline in response to stress




Problem 7.27 Nicotine, a toxic and addictive component of tobacco, is synthesized from **A** using SAM. Write out the reaction that converts **A** into nicotine.



7.13 The S_N1 Mechanism

The reaction of (CH₃)₃CBr with CH₃COO[−] is an example of the second mechanism for nucleophilic substitution, the **S_N1 mechanism**. What are the general features of this mechanism?



7.13A Kinetics

The S_N1 reaction exhibits **first-order kinetics**.

- rate = $k[(\text{CH}_3)_3\text{CBr}]$

As we learned in Section 7.10, this suggests that the S_N1 mechanism involves **more than one step**, and that the slow step is **unimolecular**, involving *only* the alkyl halide. **The identity and concentration of the nucleophile have no effect on the reaction rate**. For example, doubling the concentration of (CH₃)₃CBr doubles the rate, but doubling the concentration of the nucleophile has no effect.

Problem 7.28 What happens to the rate of an S_N1 reaction under each of the following conditions?

- [RX] is tripled, and [:Nu[−]] stays the same.
- Both [RX] and [:Nu[−]] are tripled.
- [RX] is halved, and [:Nu[−]] stays the same.
- [RX] is halved, and [:Nu[−]] is doubled.

7.13B A Two-Step Mechanism

The most straightforward explanation for the observed first-order kinetics is a **two-step mechanism** in which **bond breaking occurs before bond making**, as shown in Mechanism 7.2.




Mechanism 7.2 The S_N1 Mechanism

Step [1] The C–Br bond is broken.



- **Heterolysis of the C–Br bond** forms an intermediate **carbocation**. This step is rate-determining because it involves only bond cleavage.

Step [2] The C–O bond is formed.



- **Nucleophilic attack of acetate** on the carbocation forms the new C–O bond in the product. This is a **Lewis acid–base reaction**; the nucleophile is the Lewis base and the carbocation is the Lewis acid. Step [2] is *faster* than Step [1] because no bonds are broken and one bond is formed.


The key features of the S_N1 mechanism are:

- The mechanism has two steps.
- Carbocations are formed as reactive intermediates.

An energy diagram for the reaction of $(\text{CH}_3)_3\text{CBr} + \text{CH}_3\text{COO}^-$ is shown in Figure 7.15. Each step has its own energy barrier, with a transition state at each energy maximum. Because the transition state for Step [1] is at higher energy, Step [1] is rate-determining. ΔH° for Step [1] has a positive value because only bond breaking occurs, whereas ΔH° of Step [2] has a negative value because only bond making occurs. The overall reaction is assumed to be exothermic, so the final product is drawn at lower energy than the initial starting material.

7.13C Stereochemistry of the S_N1 Reaction


To understand the stereochemistry of the S_N1 reaction, we must examine the geometry of the carbocation intermediate.




- A carbocation (with three groups around C) is sp^2 hybridized and trigonal planar, and contains a vacant p orbital extending above and below the plane.

To illustrate the consequences of having a trigonal planar carbocation formed as a reactive intermediate, we examine the S_N1 reaction of a 3° alkyl halide **A** having the leaving group bonded to a stereogenic carbon.

Figure 7.15
An energy diagram for the S_N1 reaction:
 $(\text{CH}_3)_3\text{CBr} + \text{CH}_3\text{COO}^- \rightarrow (\text{CH}_3)_3\text{COCOCH}_3 + \text{Br}^-$



- Since the S_N1 mechanism has two steps, there are two energy barriers.
- $E_a[1] > E_a[2]$ since Step [1] involves bond breaking and Step [2] involves bond formation.
- In each step only one bond is broken or formed, so the transition state for each step has one partial bond.



Loss of the leaving group in Step [1] generates a planar carbocation that is now achiral. Attack of the nucleophile in Step [2] can occur from either side to afford two products, **B** and **C**. These two products are *different* compounds containing one stereogenic center. **B** and **C** are stereoisomers that are not superimposable—they are **enantiomers**. Because there is no preference for nucleophilic attack from either direction, an equal amount of the two enantiomers is formed—a **racemic mixture**. We say that **racemization** has occurred.


Nucleophilic attack from both sides of a planar carbocation occurs in $\text{S}_{\text{N}}1$ reactions, but we see the result of this phenomenon only when the leaving group is bonded to a stereogenic center.

- **Racemization** is the formation of equal amounts of two enantiomeric products from a single starting material.
- $\text{S}_{\text{N}}1$ reactions proceed with **racemization** at a single stereogenic center.

Two additional examples of racemization in $\text{S}_{\text{N}}1$ reactions are given in Figure 7.16.


Sample Problem 7.3

Draw the products (including stereochemistry) of the following $\text{S}_{\text{N}}1$ reaction.




Solution

Br^- is the leaving group and H_2O is the nucleophile. Loss of the leaving group generates a trigonal planar carbocation, which can react with the nucleophile from either direction to form two products.




In this example, the initial products of nucleophilic substitution bear a positive charge. They readily lose a proton to form neutral products. The overall process with a neutral nucleophile thus has **three steps**: the first two constitute the **two-step $\text{S}_{\text{N}}1$ mechanism** (loss of the leaving group and attack of the nucleophile), and the third is a **Brønsted–Lowry acid–base reaction** leading to a neutral organic product.

Figure 7.16
Two examples of racemization
in the S_N1 reaction




- Nucleophilic substitution of each starting material by an S_N1 mechanism forms a **racemic mixture** of two products.
- With H₂O, a neutral nucleophile, the initial product of nucleophilic substitution (ROH₂⁺) loses a proton to form the final neutral product, ROH (Section 7.6).



The two products in this reaction are nonsuperimposable mirror images—**enantiomers**. Because nucleophilic attack on the trigonal planar carbocation occurs with equal frequency from both directions, a **racemic mixture is formed**.


Problem 7.29 Draw the products of each S_N1 reaction and indicate the stereochemistry of any stereogenic centers.



7.13D The Identity of the R Group

How does the rate of an S_N1 reaction change as the alkyl group in the substrate alkyl halide changes from CH₃ → 1° → 2° → 3°?

- As the number of R groups on the carbon with the leaving group *increases*, the rate of an S_N1 reaction *increases*.



- 3° Alkyl halides undergo S_N1 reactions rapidly.
- 2° Alkyl halides react more slowly.
- Methyl and 1° alkyl halides do *not* undergo S_N1 reactions.

Table 7.6 summarizes the characteristics of the S_N1 mechanism.


This trend is exactly opposite to that observed for the S_N2 mechanism. To explain this result, we must examine the rate-determining step, the formation of the carbocation, and learn about the effect of alkyl groups on **carbocation stability**.

Table 7.6 Characteristics of the S_N1 Mechanism

Characteristic	Result
Kinetics	• First-order kinetics; rate = $k[\text{RX}]$
Mechanism	• Two steps
Stereochemistry	• Trigonal planar carbocation intermediate • Racemization at a single stereogenic center
Identity of R	• More substituted halides react fastest. • Rate: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X} > \text{CH}_3\text{X}$





7.14 Carbocation Stability

Carbocations are classified as **primary (1°)**, **secondary (2°)**, or **tertiary (3°)** by the number of R groups bonded to the charged carbon atom. As the number of R groups on the positively charged carbon atom increases, the stability of the carbocation **increases**.



When we speak of carbocation stability, we really mean *relative* stability. Tertiary carbocations are too unstable to isolate, but they are more stable than secondary carbocations. We will examine the reason for this order of stability by invoking two different principles: **inductive effects** and **hyperconjugation**.

Problem 7.30 Classify each carbocation as 1° , 2° , or 3° .




Problem 7.31 Draw the structure of a 1° , 2° , and 3° carbocation, each having molecular formula C_4H_9^+ . Rank the three carbocations in order of increasing stability.

7.14A Inductive Effects

Inductive effects are electronic effects that occur through σ bonds. In Section 2.5B, for example, we learned that more electronegative atoms stabilize a negative charge by an **electron-withdrawing inductive effect**.



To stabilize a positive charge, **electron-donating groups** are needed. **Alkyl groups are electron donor groups that stabilize a positive charge.** An alkyl group with several σ bonds is more polarizable than a hydrogen atom, and more able to donate electron density. Thus, as R groups successively replace the H atoms in CH_3^+ , **the positive charge is more dispersed on the electron donor R groups, and the carbocation is more stabilized.**

Electron donor groups (Z) stabilize a (+) charge; $\text{Z} \rightarrow \text{Y}^+$.
Electron-withdrawing groups (W) stabilize a (-) charge; $\text{W} \leftarrow \text{Y}^-$.



Electrostatic potential maps for four carbocations in Figure 7.17 illustrate the effect of increasing alkyl substitution on the positive charge of the carbocation.

Problem 7.32 Rank the following carbocations in order of increasing stability.



7.14B Hyperconjugation

A second explanation for the observed trend in carbocation stability is based on orbital overlap. A 3° carbocation is more stable than a 2° , 1° , or methyl carbocation because the positive charge is delocalized over more than one atom.

- Spreading out charge by the overlap of an empty p orbital with an adjacent σ bond is called **hyperconjugation**.

For example, CH_3^+ cannot be stabilized by hyperconjugation, but $(\text{CH}_3)_2\text{CH}^+$ can:




This carbocation has no opportunity for orbital overlap with the vacant p orbital.

Overlap of the C–H σ bond with the adjacent vacant p orbital stabilizes the carbocation.

Both carbocations contain an sp^2 hybridized carbon, so both are trigonal planar with a vacant p orbital extending above and below the plane. There are no adjacent C–H σ bonds with which the p orbital can overlap in CH_3^+ , but there *are* adjacent C–H σ bonds in $(\text{CH}_3)_2\text{CH}^+$. This overlap (the **hyperconjugation**) delocalizes the positive charge on the carbocation, spreading it over a larger volume, and this stabilizes the carbocation.

Figure 7.17

Electrostatic potential maps for different carbocations




- Dark blue areas in electrostatic potential plots indicate regions low in electron density. As alkyl substitution increases, the region of positive charge is less concentrated on carbon.

The larger the number of alkyl groups on the adjacent carbons, the greater the possibility for hyperconjugation, and the larger the stabilization. Hyperconjugation thus provides an alternate way of explaining why **carbocations with a larger number of R groups are more stabilized**.


7.15 The Hammond Postulate

The rate of an S_N1 reaction depends on the rate of formation of the carbocation (the product of the rate-determining step) via heterolysis of the C–X bond.

- The rate of an S_N1 reaction *increases* as the number of R groups on the carbon with the leaving group *increases*.
- The stability of a carbocation *increases* as the number of R groups on the positively charged carbon *increases*.



- Thus, the rate of an S_N1 reaction *increases* as the stability of the carbocation *increases*.




The rate of a reaction depends on the magnitude of E_a , and the stability of a product depends on ΔG° . The **Hammond postulate**, first proposed in 1955, **relates rate to stability**.

7.15A The General Features of the Hammond Postulate

The Hammond postulate provides a qualitative estimate of the energy of a transition state. Because the energy of the transition state determines the energy of activation and therefore the reaction rate, predicting the relative energy of two transition states allows us to determine the relative rates of two reactions.

According to the Hammond postulate, the transition state of a reaction resembles the structure of the species (reactant or product) to which it is closer in energy. In **endothermic reactions**, the transition state is closer in energy to the **products**. In **exothermic reactions**, the transition state is closer in energy to the **reactants**.



- Transition states in endothermic reactions resemble the products.
- Transition states in exothermic reactions resemble the reactants.

What happens to the reaction rate if the energy of the product is lowered? In an **endothermic reaction**, the transition state resembles the products, so anything that stabilizes the product stabilizes the transition state, too. **Lowering the energy of the transition state decreases the energy of activation (E_a), which increases the reaction rate.**

Suppose there are two possible products of an endothermic reaction, but one is more stable (lower in energy) than the other (Figure 7.18). According to the Hammond postulate, **the transition state to form the more stable product is lower in energy, so this reaction should occur faster.**

- Conclusion: In an endothermic reaction, the more stable product forms faster.

What happens to the reaction rate of an **exothermic reaction** if the energy of the product is lowered? The transition state resembles the reactants, so **lowering the energy of the products has little or no effect on the energy of the transition state.** If E_a is unaffected, then the reaction rate is unaffected, too, as shown in Figure 7.19.

- Conclusion: In an exothermic reaction, the more stable product may or may not form faster because E_a is similar for both products.

Figure 7.18
An endothermic reaction—
How the energies of the
transition state and products
are related





Figure 7.19


An exothermic reaction—How the energies of the transition state and products are related




7.15B The Hammond Postulate and the S_N1 Reaction

In the S_N1 reaction, the rate-determining step is the formation of the carbocation, an *endothermic* reaction. According to the Hammond postulate, the **stability of the carbocation determines the rate of its formation**.

For example, heterolysis of the C–Cl bond in $(\text{CH}_3)_2\text{CHCl}$ affords a less stable 2° carbocation, $(\text{CH}_3)_2\text{CH}^+$ (Equation [1]), whereas heterolysis of the C–Cl bond in $(\text{CH}_3)_3\text{CCl}$ affords a more stable 3° carbocation, $(\text{CH}_3)_3\text{C}^+$ (Equation [2]). The Hammond postulate states that Reaction [2] is faster than Reaction [1], because the transition state to form the more stable 3° carbocation is lower in energy. Figure 7.20 depicts an energy diagram comparing these two endothermic reactions.

**Figure 7.20**


Energy diagram for carbocation formation in two different S_N1 reactions



- Since $(\text{CH}_3)_2\text{CH}^+$ is less stable than $(\text{CH}_3)_3\text{C}^+$, $E_a[1] > E_a[2]$, and Reaction [1] is slower.

In conclusion, the Hammond postulate estimates the relative energy of transition states, and thus it can be used to predict the relative rates of two reactions.

Problem 7.33 Which alkyl halide in each pair reacts faster in an S_N1 reaction?





7.16 Application: S_N1 Reactions, Nitrosamines, and Cancer




Spam, a widely consumed canned meat in Alaska, Hawaii, and other parts of the United States, contains sodium nitrite.

Two common nitrosamines:




S_N1 reactions are thought to play a role in how **nitrosamines**, compounds having the general structure **R₂NN=O**, act as toxins and carcinogens. Nitrosamines are present in many foods, especially cured meats and smoked fish, and they are also found in tobacco smoke, alcoholic beverages, and cosmetics. Nitrosamines cause many forms of cancer.

Nitrosamines can be formed when amines that occur naturally in food react with sodium nitrite, NaNO₂, a preservative added to meats such as ham, bacon, and hot dogs to inhibit the growth of *Clostridium botulinum*, a bacterium responsible for a lethal form of food poisoning. Nitrosamines are also formed in vivo in the gastrointestinal tract when bacteria in the body convert nitrates (NO₃⁻) into nitrites (NO₂⁻), which then react with amines.



In the presence of acid or heat, nitrosamines are converted to **diazonium ions**, which contain a very good leaving group, N₂. With certain R groups, these diazonium compounds form carbocations, which then react with biological nucleophiles (such as DNA or an enzyme) in the cell. If this nucleophilic substitution reaction occurs at a crucial site in a biomolecule, it can disrupt normal cell function leading to cancer or cell death. This two-step process—loss of N₂ as a leaving group and reaction with a nucleophile—is an **S_N1 reaction**.



The use of sodium nitrite as a preservative is a classic example of the often delicate balance between risk and benefit. On the one hand, there is an enormous benefit in reducing the prevalence of fatal toxins in meats by the addition of sodium nitrite. On the other, there is the potential risk that sodium nitrite may increase the level of nitrosamines in certain foods. Nitrites are still used as food additives, but the allowable level of nitrites in cured meats has been reduced. Debate continues on whether nitrite preservatives used at their current low levels actually pose a risk to the consumer.

7.17 When Is the Mechanism S_N1 or S_N2 ?


Given a particular starting material and nucleophile, how do we know whether a reaction occurs by the S_N1 or S_N2 mechanism? Four factors are examined:

- The alkyl halide— CH_3X , RCH_2X , R_2CHX , or R_3CX
- The nucleophile—strong or weak
- The leaving group—good or poor
- The solvent—protic or aprotic

7.17A The Alkyl Halide—The Most Important Factor

The most important factor in determining whether a reaction follows the S_N1 or S_N2 mechanism is the *identity of the alkyl halide*.


- Increasing alkyl substitution favors S_N1 .
- Decreasing alkyl substitution favors S_N2 .



- Methyl and 1° halides (CH_3X and RCH_2X) undergo S_N2 reactions only.
- 3° Alkyl halides (R_3CX) undergo S_N1 reactions only.
- 2° Alkyl halides (R_2CHX) undergo both S_N1 and S_N2 reactions. Other factors determine the mechanism.

Examples are given in Figure 7.21.

Problem 7.34 What is the likely mechanism of nucleophilic substitution for each alkyl halide?




7.17B The Nucleophile

How does the strength of the nucleophile affect an S_N1 or S_N2 mechanism? The rate of the S_N1 reaction is unaffected by the identity of the nucleophile because the nucleophile does not appear in the rate equation ($rate = k[RX]$). The identity of the nucleophile *is* important for

Figure 7.21

Examples: The identity of RX and the mechanism of nucleophilic substitution




the S_N2 reaction, however, because the nucleophile does appear in the rate equation for this mechanism (rate = $k[\text{RX}][:\text{Nu}^-]$).


- Strong nucleophiles present in high concentration favor S_N2 reactions.
- Weak nucleophiles favor S_N1 reactions by decreasing the rate of any competing S_N2 reaction.

The most common nucleophiles in S_N2 reactions bear a net negative charge. The most common nucleophiles in S_N1 reactions are weak nucleophiles such as H₂O and ROH. The identity of the nucleophile is especially important in determining the mechanism and therefore the stereochemistry of nucleophilic substitution when 2° alkyl halides are starting materials.


Let's compare the substitution products formed when the 2° alkyl halide **A** (*cis*-1-bromo-4-methylcyclohexane) is treated with either the strong nucleophile ⁻OH or the weak nucleophile H₂O. Because a 2° alkyl halide can react by either mechanism, the strength of the nucleophile determines which mechanism takes place.



The **strong nucleophile** ⁻OH favors an S_N2 reaction, which occurs with **backside** attack of the nucleophile, resulting in **inversion of configuration**. Because the leaving group Br⁻ is above the plane of the ring, the nucleophile attacks from below, and a single product **B** is formed.







The **weak nucleophile H₂O** favors an S_N1 reaction, which occurs by way of an intermediate carbocation. Loss of the leaving group in **A** forms the carbocation, which undergoes nucleophilic attack from both above and below the plane of the ring to afford two products, **C** and **D**. Loss of a proton by proton transfer forms the final products, **B** and **E**. **B** and **E** are diastereomers of each other (**B** is a *trans* isomer and **E** is a *cis* isomer).





Thus, the mechanism of nucleophilic substitution determines the stereochemistry of the products formed.

Problem 7.35 For each alkyl halide and nucleophile: [1] Draw the product of nucleophilic substitution; [2] determine the likely mechanism (S_N1 or S_N2) for each reaction.



Problem 7.36 Draw the products (including stereochemistry) for each reaction.




7.17C The Leaving Group

How does the identity of the leaving group affect an S_N1 or S_N2 reaction?


- A better leaving group increases the rate of *both* S_N1 and S_N2 reactions.

Because the bond to the leaving group is partially broken in the transition state of the only step of the S_N2 mechanism and the slow step of the S_N1 mechanism, **a better leaving group increases the rate of both reactions.** The better the leaving group, the more willing it is to accept the electron pair in the C–X bond, and the faster the reaction.

Transition state of the S_N2 mechanism




Transition state of the rate-determining step of the S_N1 mechanism



A better leaving group is more able to accept the negative charge.

For alkyl halides, the following order of reactivity is observed for the S_N1 and the S_N2 mechanisms:



Problem 7.37 Which compound in each pair reacts faster in nucleophilic substitution?

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{I}$ c. $(\text{CH}_3)_3\text{COH}$ or $(\text{CH}_3)_3\text{COH}_2^+$
b. $(\text{CH}_3)_3\text{CBr}$ or $(\text{CH}_3)_3\text{CI}$ d. $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCOCH}_3$

7.17D The Solvent

See Section 7.8C to review the differences between polar protic solvents and polar aprotic solvents.

Polar protic solvents and polar aprotic solvents affect the rates of S_N1 and S_N2 reactions differently.

- Polar *protic* solvents are especially good for S_N1 reactions.
- Polar *aprotic* solvents are especially good for S_N2 reactions.

Polar protic solvents like H_2O and ROH solvate both cations and anions well, and this characteristic is important for the S_N1 mechanism, in which two ions (a carbocation and a leaving group) are formed by heterolysis of the C–X bond. The carbocation is solvated by ion–dipole interac-

tions with the polar solvent, and the leaving group is solvated by hydrogen bonding, in much the same way that Na⁺ and Br⁻ are solvated in Section 7.8C. These interactions stabilize the reactive intermediate. In fact, a polar protic solvent is generally needed for an S_N1 reaction.

Polar aprotic solvents exhibit dipole–dipole interactions but not hydrogen bonding, and as a result, they do not solvate anions well. This has a pronounced effect on the nucleophilicity of anionic nucleophiles. Because these nucleophiles are not “hidden” by strong interactions with the solvent, they are **more nucleophilic**. Because stronger nucleophiles favor S_N2 reactions, **polar aprotic solvents are especially good for S_N2 reactions**.

Problem 7.38

Which solvents favor S_N1 reactions and which favor S_N2 reactions?





- a. CH₃CH₂OH b. CH₃CN c. CH₃COOH d. CH₃CH₂OCH₂CH₃

Problem 7.39

For each reaction, use the identity of the alkyl halide and nucleophile to determine which substitution mechanism occurs. Then determine which solvent affords the faster reaction.

Summary of solvent effects:

- Polar protic solvents favor S_N1 reactions because the ionic intermediates are stabilized by solvation.
- Polar aprotic solvents favor S_N2 reactions because nucleophiles are not well solvated, and therefore are more nucleophilic.



7.17E Summary of Factors That Determine Whether the S_N1 or S_N2 Mechanism Occurs

Table 7.7 summarizes the factors that determine whether a reaction occurs by the S_N1 or S_N2 mechanism. Sample Problems 7.4 and 7.5 illustrate how these factors are used to determine the mechanism of a given reaction.

Table 7.7 Summary of Factors That Determine the S_N1 or S_N2 Mechanism


Alkyl halide	Mechanism	Other factors
CH ₃ X RCH ₂ X (1°)	S _N 2	Favored by <ul style="list-style-type: none"> • strong nucleophiles (usually a net negative charge) • polar aprotic solvents
R ₃ CX (3°)	S _N 1	Favored by <ul style="list-style-type: none"> • weak nucleophiles (usually neutral) • polar protic solvents
R ₂ CHX (2°)	S _N 1 or S _N 2	The mechanism depends on the conditions. <ul style="list-style-type: none"> • Strong nucleophiles favor the S_N2 mechanism over the S_N1 mechanism. For example, RO⁻ is a stronger nucleophile than ROH, so RO⁻ favors the S_N2 reaction and ROH favors the S_N1 reaction. • Protic solvents favor the S_N1 mechanism and aprotic solvents favor the S_N2 mechanism. For example, H₂O and CH₃OH are polar protic solvents that favor the S_N1 mechanism, whereas acetone [(CH₃)₂C=O] and DMSO [(CH₃)₂S=O] are polar aprotic solvents that favor the S_N2 mechanism.

Sample Problem 7.4 Determine the mechanism of nucleophilic substitution for each reaction and draw the products.




Solution


a. The alkyl halide is 1° , so it must react by an $\text{S}_{\text{N}}2$ mechanism with the nucleophile $\text{:C}\equiv\text{CH}^-$.



b. The alkyl halide is 2° , so it can react by either the $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$ mechanism. The strong nucleophile (:CN^-) favors the $\text{S}_{\text{N}}2$ mechanism.




Sample Problem 7.5 Determine the mechanism of nucleophilic substitution for each reaction and draw the products, including stereochemistry.




Solution

a. The 2° alkyl halide can react by either the $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$ mechanism. The strong nucleophile (:OCH_3^-) favors the $\text{S}_{\text{N}}2$ mechanism, as does the polar aprotic solvent (DMSO). $\text{S}_{\text{N}}2$ reactions proceed with inversion of configuration.



b. The alkyl halide is 3° , so it reacts by an $\text{S}_{\text{N}}1$ mechanism with the weak nucleophile CH_3OH . $\text{S}_{\text{N}}1$ reactions proceed with racemization at a single stereogenic center, so two products are formed.



Problem 7.40 Determine the mechanism and draw the products of each reaction. Include the stereochemistry at all stereogenic centers.







Table 7.8 Molecules Synthesized from R-X by the S_N2 Reaction

	Nucleophile (:Nu ⁻)	Product	Name
Oxygen compounds	⁻ OH	R-OH	alcohol
	⁻ OR'	R-OR'	ether
			ester
Carbon compounds	⁻ CN	R-CN	nitrile
	⁻ C≡C-H	R-C≡C-H	alkyne
Nitrogen compounds	N ₃ ⁻	R-N ₃	azide
	:NH ₃	R-NH ₂	amine
Sulfur compounds	⁻ SH	R-SH	thiol
	⁻ SR'	R-SR'	sulfide

↑
products of nucleophilic substitution

7.19A Background on Organic Synthesis


Chemists synthesize molecules for many reasons. Sometimes a **natural product**, a compound isolated from natural sources, has useful medicinal properties, but is produced by an organism in only minute quantities. Synthetic chemists then prepare this molecule from simpler starting materials so that it can be made available to a large number of people. **Taxol** (Section 5.5), the complex anticancer compound isolated in small amounts from the bark of the Pacific yew tree, is one such natural product. It can be synthesized from a compound isolated from the needles of the European yew.



Phenol, the starting material for the aspirin synthesis, is a petroleum product, like most of the starting materials used in large quantities in industrial syntheses. A shortage of petroleum reserves thus affects the availability not only of fuels for transportation, but also of raw materials needed for most chemical synthesis.

Sometimes, chemists prepare molecules that do not occur in nature (although they may be similar to those in nature), because these molecules have superior properties to their naturally occurring relatives. **Aspirin, or acetylsalicylic acid** (Section 2.7), is a well known example. Acetylsalicylic acid is prepared from phenol, a product of the petroleum industry, by a two-step procedure (Figure 7.22). Aspirin has become one of the most popular and widely used drugs in the world because it has excellent analgesic and anti-inflammatory properties, *and* it is cheap and readily available.

Figure 7.22
Synthesis of aspirin




- Aspirin is synthesized by a two-step procedure from simple, cheap starting materials.


7.19B Nucleophilic Substitution and Organic Synthesis

To carry out synthesis we must think backwards. We examine a compound and ask: **What starting material and reagent are needed to make it?** If we are using nucleophilic substitution, we must determine what alkyl halide and what nucleophile can be used to form a specific product. This is the simplest type of synthesis because it involves only one step. In Chapter 11 we will learn about multistep syntheses.

Suppose, for example, that we are asked to prepare $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$ (2-methyl-1-propanol) from an alkyl halide and any required reagents. To accomplish this synthesis, we must “fill in the boxes” for the starting material and reagent in the accompanying equation.




To determine the two components needed for the synthesis, remember that the carbon atoms come from the organic starting material, in this case a 1° alkyl halide $[(\text{CH}_3)_2\text{CHCH}_2\text{Br}]$. The functional group comes from the nucleophile, ^-OH in this case. With these two components, we can “fill in the boxes” to complete the synthesis.



After any synthesis is proposed, check to see if it is reasonable, given what we know about reactions. Will the reaction written give a high yield of product? The synthesis of $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$ is reasonable, because the starting material is a 1° alkyl halide and the nucleophile (^-OH) is strong, and both facts contribute to a successful $\text{S}_{\text{N}}2$ reaction.

Problem 7.42 What alkyl halide and nucleophile are needed to prepare each compound?



Problem 7.43 The ether, $\text{CH}_3\text{OCH}_2\text{CH}_3$, can be prepared by two different nucleophilic substitution reactions, one using CH_3O^- as nucleophile and the other using $\text{CH}_3\text{CH}_2\text{O}^-$ as nucleophile. Draw both routes.

KEY CONCEPTS


Alkyl Halides and Nucleophilic Substitution

General Facts about Alkyl Halides

- Alkyl halides contain a halogen atom X bonded to an sp^3 hybridized carbon (7.1).
- Alkyl halides are named as halo alkanes, with the halogen as a substituent (7.2).
- Alkyl halides have a polar C–X bond, so they exhibit dipole–dipole interactions but are incapable of intermolecular hydrogen bonding (7.3).
- The polar C–X bond containing an electrophilic carbon makes alkyl halides reactive towards nucleophiles and bases (7.5).

The Central Theme (7.6)

- Nucleophilic substitution is one of the two main reactions of alkyl halides. A nucleophile replaces a leaving group on an sp^3 hybridized carbon.



- One σ bond is broken and one σ bond is formed.
- There are two possible mechanisms: S_N1 and S_N2 .

S_N1 and S_N2 Mechanisms Compared

	S_N2 mechanism	S_N1 mechanism
[1] Mechanism	One step (7.11B)	Two steps (7.13B)
[2] Alkyl halide	Order of reactivity: $\text{CH}_3\text{X} > \text{RCH}_2\text{X} > \text{R}_2\text{CHX} > \text{R}_3\text{CX}$ (7.11D)	Order of reactivity: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X} > \text{CH}_3\text{X}$ (7.13D)
[3] Rate equation	Rate = $k[\text{RX}][\text{Nu}^-]$ Second-order kinetics (7.11A)	Rate = $k[\text{RX}]$ First-order kinetics (7.13A)
[4] Stereochemistry	Backside attack of the nucleophile (7.11C) Inversion of configuration at a stereogenic center	Trigonal planar carbocation intermediate (7.13C) Racemization at a single stereogenic center
[5] Nucleophile	Favored by stronger nucleophiles (7.17B)	Favored by weaker nucleophiles (7.17B)
[6] Leaving group	Better leaving group \rightarrow faster reaction (7.17C)	Better leaving group \rightarrow faster reaction (7.17C)
[7] Solvent	Favored by polar aprotic solvents (7.17D)	Favored by polar protic solvents (7.17D)

Important Trends

- The best leaving group is the weakest base. Leaving group ability increases left-to-right across a row and down a column of the periodic table (7.7).
- Nucleophilicity decreases left-to-right across a row of the periodic table (7.8A).
- Nucleophilicity decreases down a column of the periodic table in polar aprotic solvents (7.8C).
- Nucleophilicity increases down a column of the periodic table in polar protic solvents (7.8C).
- The stability of a carbocation increases as the number of R groups bonded to the positively charged carbon increases (7.14).









Important Principles

Principle	Example
Electron-donating groups (such as R groups) stabilize a positive charge (7.14A).	3° Carbocations (R_3C^+) are more stable than 2° carbocations (R_2CH^+), which are more stable than 1° carbocations (RCH_2^+).
Steric hindrance decreases nucleophilicity but not basicity (7.8B).	$(\text{CH}_3)_3\text{CO}^-$ is a stronger base but a weaker nucleophile than $\text{CH}_3\text{CH}_2\text{O}^-$.
Hammond postulate: In an endothermic reaction, the more stable product is formed faster. In an exothermic reaction, this is not necessarily true (7.15).	S_N1 reactions are faster when more stable (more substituted) carbocations are formed, because the rate-determining step is endothermic.
Planar, sp^2 hybridized atoms react with reagents from both sides of the plane (7.13C).	A trigonal planar carbocation reacts with nucleophiles from both sides of the plane.







PROBLEMS

Nomenclature

7.44 Give the IUPAC name for each compound.



7.45 Give the structure corresponding to each name.



7.46 Classify each alkyl halide in Problem 7.44 as 1°, 2°, or 3°. When a compound has more than one halogen, assign each separately.

7.47 Draw the eight constitutional isomers having the molecular formula $\text{C}_5\text{H}_{11}\text{Cl}$.




a. Give the IUPAC name for each compound (ignoring *R* and *S* designations).

b. Label any stereogenic centers.

c. For each constitutional isomer that contains a stereogenic center, draw all possible stereoisomers, and label each stereogenic center as *R* or *S*.


Physical Properties

7.48 Which compound in each pair has the higher boiling point?









General Nucleophilic Substitution, Leaving Groups, and Nucleophiles


7.49 Draw the substitution product that results when $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$ reacts with each nucleophile.



7.50 Draw the products of each nucleophilic substitution reaction.




7.51 Which of the following molecules contain a good leaving group?




7.52 Rank the species in each group in order of increasing leaving group ability.




7.53 Which of the following nucleophilic substitution reactions will take place?




7.54 What nucleophile is needed to convert **A** to each substitution product?



7.55 Rank the species in each group in order of increasing nucleophilicity.




7.56 Classify each solvent as protic or aprotic.



7.57 Why is the amine N atom more nucleophilic than the amide N atom in $\text{CH}_3\text{CONHCH}_2\text{CH}_2\text{CH}_2\text{NHCH}_3$?


The $\text{S}_{\text{N}}2$ Reaction

7.58 Consider the following $\text{S}_{\text{N}}2$ reaction:







- Draw a mechanism using curved arrows.
- Draw an energy diagram. Label the axes, the reactants, products, E_{a} , and ΔH° . Assume that the reaction is exothermic.
- Draw the structure of the transition state.
- What is the rate equation?
- What happens to the reaction rate in each of the following instances? [1] The leaving group is changed from Br^- to I^- ; [2] The solvent is changed from acetone to $\text{CH}_3\text{CH}_2\text{OH}$; [3] The alkyl halide is changed from $\text{CH}_3(\text{CH}_2)_4\text{Br}$ to $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{Br})\text{CH}_3$; [4] The concentration of CN^- is increased by a factor of five; and [5] The concentrations of both the alkyl halide and CN^- are increased by a factor of five.





7.59 Rank the alkyl halides in each group in order of increasing $\text{S}_{\text{N}}2$ reactivity.



7.60 Which S_N2 reaction in each pair is faster?




7.61 Draw the products of each S_N2 reaction and indicate the stereochemistry where appropriate.




Carbocations

7.62 Classify each carbocation as 1°, 2°, or 3°.




7.63 Rank the carbocations in each group in order of increasing stability.



7.64 The following order of stability is observed for three carbocations: $\text{CCl}_3\text{CH}_2^+ < \text{CH}_3\text{CH}_2^+ < \text{CH}_3\text{OCH}_2^+$; that is, $\text{CCl}_3\text{CH}_2^+$ is the least stable and $\text{CH}_3\text{OCH}_2^+$ is the most stable. Offer an explanation.


The S_N1 Reaction

7.65 Consider the following S_N1 reaction.




- Draw a mechanism for this reaction using curved arrows.
- Draw an energy diagram. Label the axes, starting material, product, E_a , and ΔH° . Assume that the starting material and product are equal in energy.
- Draw the structure of any transition states.
- What is the rate equation for this reaction?
- What happens to the reaction rate in each of the following instances? [1] The leaving group is changed from I^- to Cl^- ; [2] The solvent is changed from H_2O to DMF; [3] The alkyl halide is changed from $(\text{CH}_3)_2\text{C}(\text{I})\text{CH}_2\text{CH}_3$ to $(\text{CH}_3)_2\text{CHCH}(\text{I})\text{CH}_3$; [4] The concentration of H_2O is increased by a factor of five; and [5] The concentrations of both the alkyl halide and H_2O are increased by a factor of five.


7.66 Rank the alkyl halides in each group in order of increasing S_N1 reactivity.




7.67 Which S_N1 reaction in each pair is faster?



7.68 Draw the products of each S_N1 reaction and indicate the stereochemistry when necessary.




7.69 Draw a stepwise mechanism for the following reaction that illustrates why two substitution products are formed. Explain why 1-bromo-2-hexene reacts rapidly with a weak nucleophile (CH_3OH) under S_N1 reaction conditions, even though it is a 1° alkyl halide.




S_N1 and S_N2 Reactions


7.70 Determine the mechanism of nucleophilic substitution of each reaction and draw the products, including stereochemistry.




7.71 Draw the products of each nucleophilic substitution reaction.




7.72 Diphenhydramine, the antihistamine in Benadryl, can be prepared by the following two-step sequence. What is the structure of diphenhydramine?




7.73 Draw a stepwise, detailed mechanism for the following reaction. Use curved arrows to show the movement of electrons.



7.74 When a single compound contains both a nucleophile and a leaving group, an **intramolecular** reaction may occur. With this in mind, draw the product of the following reaction.



7.75 Nicotine can be made when the following ammonium salt is treated with Na_2CO_3 . Draw a stepwise mechanism for this reaction.



7.76 Explain each of the following statements.

- Hexane is not a common solvent for either $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$ reactions.
- $(\text{CH}_3)_3\text{CO}^-$ is a stronger base than $\text{CH}_3\text{CH}_2\text{O}^-$.
- $(\text{CH}_3)_3\text{CBr}$ is more reactive than $(\text{CH}_3)_2\text{C}(\text{CF}_3)\text{Br}$ in $\text{S}_{\text{N}}1$ reactions.
- $(\text{CH}_3)_3\text{CBr}$ reacts at the same rate with F^- and H_2O in substitution reactions even though F^- has a net negative charge.
- When optically active (2*R*)-2-bromobutane is added to a solution of NaBr in acetone, the solution gradually loses optical activity until it becomes optically inactive.





7.77 Draw a stepwise, detailed mechanism for the following reaction.




7.78 When (6*R*)-6-bromo-2,6-dimethylnonane is dissolved in CH_3OH , nucleophilic substitution yields an optically inactive solution. When the isomeric halide (5*R*)-2-bromo-2,5-dimethylnonane is dissolved in CH_3OH under the same conditions, nucleophilic substitution forms an optically active solution. Draw the products formed in each reaction, and explain why the difference in optical activity is observed.

Synthesis


7.79 Fill in the appropriate reagent or starting material in each of the following reactions.




7.80 Devise a synthesis of each compound from an alkyl halide using any other organic or inorganic reagents.




7.81 Benzalkonium chloride (**A**) is a weak germicide used in topical antiseptics and mouthwash. **A** can be prepared from amines **B** or **C** by $\text{S}_{\text{N}}2$ reaction with an alkyl chloride. (a) What alkyl chloride is needed to prepare **A** from **B**? (b) What alkyl chloride is needed to prepare **A** from **C**?



7.82 Suppose you have compounds **A–D** at your disposal. Using these compounds, devise two different ways to make **E**. Which one of these methods is preferred, and why?




7.83 Muscalure, the sex pheromone of the common housefly, can be prepared by a reaction sequence that uses two nucleophilic substitutions. Identify compounds **A–D** in the following synthesis of muscalure.




Challenge Problems


7.84 We will return often to nucleophilic substitution, in particular the $\text{S}_{\text{N}}2$ reaction, in subsequent chapters. In each instance we will concentrate on the *nucleophile*, rather than the alkyl halide, as we have done in this chapter. By using different nucleophiles, nucleophilic substitution allows the synthesis of a wide variety of organic compounds with many different functional groups. With this in mind, draw the products of each two-step sequence. (Hint: Step [1] in each part involves an acid–base reaction that removes the most acidic hydrogen from the starting material.)




- 7.85 Explain why quinuclidine is a much more reactive nucleophile than triethylamine, even though both compounds have N atoms surrounded by three R groups.




- 7.86 Draw a stepwise mechanism for the following reaction sequence.




- 7.87 As we will learn in Chapter 9, an epoxide is an ether with an oxygen atom in a three-membered ring. Epoxides can be made by intramolecular S_N2 reactions of intermediates that contain a nucleophile and a leaving group on adjacent carbons, as shown.



Assume that each of the following starting materials can be converted to an epoxide by this reaction. Draw the product formed (including stereochemistry) from each starting material. Why might some of these reactions be more difficult than others in yielding nucleophilic substitution products?




- 7.88 When trichloride **J** is treated with CH_3OH , nucleophilic substitution forms the dihalide **K**. Draw a mechanism for this reaction and explain why one Cl is much more reactive than the other two Cl's so that a single substitution product is formed.



8

Alkyl Halides and Elimination Reactions

- 8.1 General features of elimination
- 8.2 Alkenes—The products of elimination reactions
- 8.3 The mechanisms of elimination
- 8.4 The E2 mechanism
- 8.5 The Zaitsev rule
- 8.6 The E1 mechanism
- 8.7 S_N1 and E1 reactions
- 8.8 Stereochemistry of the E2 reaction
- 8.9 When is the mechanism E1 or E2?
- 8.10 E2 reactions and alkyne synthesis
- 8.11 When is the reaction S_N1 , S_N2 , E1, or E2?



DDE, **d**ichlorodiphenyl**d**ichloro**e**thylene, is formed by the elimination of HCl from the pesticide DDT. DDE and DDT accumulate in the fatty tissues of predator birds such as osprey that feed on fish contaminated with DDT. When DDE and DDT concentration is high, mother osprey produce eggs with very thin shells that are easily crushed, so fewer osprey chicks hatch. In Chapter 8, we learn about **elimination reactions**, the second general reaction of alkyl halides, which form alkenes like DDE.


Elimination reactions introduce π bonds into organic compounds, so they can be used to synthesize **alkenes** and **alkynes**—hydrocarbons that contain one and two π bonds, respectively. Like nucleophilic substitution, elimination reactions can occur by two different pathways, depending on the conditions. By the end of Chapter 8, therefore, you will have learned four different reaction mechanisms, two for nucleophilic substitution (S_N1 and S_N2) and two for elimination (E1 and E2).

The biggest challenge with this material is learning how to sort out two different reactions that follow four different mechanisms. **Will a particular alkyl halide undergo substitution or elimination with a given reagent, and by which of the four possible mechanisms?** To answer this question, we conclude Chapter 8 with a summary that allows you to predict which reaction and mechanism are likely for a given substrate.


8.1 General Features of Elimination

All **elimination reactions** involve loss of elements from the starting material to form a new π bond in the product.


- Alkyl halides undergo elimination reactions with Brønsted–Lowry bases. The elements of HX are lost and an alkene is formed.



Equations [1] and [2] illustrate examples of elimination reactions. In both reactions a base removes the elements of an acid, HBr or HCl, from the organic starting material.



Removal of the elements of HX, called **dehydrohalogenation**, is one of the most common methods to introduce a π bond and prepare an alkene. Dehydrohalogenation is an example of **β elimination**, because it involves loss of elements from two adjacent atoms: the **α carbon** bonded to the leaving group X, and the **β carbon** adjacent to it. Three curved arrows illustrate how four bonds are broken or formed in the process.



- The base (B:) removes a proton on the β carbon, thus forming H-B^+ .
- The electron pair in the β C–H bond forms the new π bond between the α and β carbons.
- The electron pair in the C–X bond ends up on halogen, forming the leaving group :X^- .

Table 8.1 Common Bases Used in Dehydrohalogenation



Structure	Name
$\text{Na}^+ \text{}^- \text{OH}$	sodium hydroxide
$\text{K}^+ \text{}^- \text{OH}$	potassium hydroxide
$\text{Na}^+ \text{}^- \text{OCH}_3$	sodium methoxide
$\text{Na}^+ \text{}^- \text{OCH}_2\text{CH}_3$	sodium ethoxide
$\text{K}^+ \text{}^- \text{OC}(\text{CH}_3)_3$	potassium <i>tert</i> -butoxide

The most common bases used in elimination reactions are negatively charged oxygen compounds such as $\text{}^- \text{OH}$ and its alkyl derivatives, $\text{}^- \text{OR}$, called **alkoxides**, listed in Table 8.1. Potassium *tert*-butoxide, $\text{K}^+ \text{}^- \text{OC}(\text{CH}_3)_3$, a bulky nonnucleophilic base, is especially useful (Section 7.8B).


To draw any product of dehydrohalogenation:

- Find the α carbon—the sp^3 hybridized carbon bonded to the leaving group.
- Identify all β carbons with H atoms.
- Remove the elements of H and X from the α and β carbons and form a π bond.




For example, 2-bromo-2-methylpropane has three β carbons (three CH_3 groups), but because all three are *identical*, only *one* alkene is formed upon elimination of HBr. In contrast, 2-bromobutane has two *different* β carbons (labeled β_1 and β_2), so elimination affords *two* constitutional isomers by loss of HBr across either the α and β_1 carbons, or the α and β_2 carbons. We learn about which product predominates and why in Section 8.5.



An elimination reaction is the first step in the slow degradation of the **pesticide DDT** (Chapter 8 opening paragraph and Section 7.4). Elimination of HCl from DDT forms the degradation product **DDE** (dichlorodiphenyldichloroethylene). This stable alkene is found in minute concentration in the fatty tissues of most adults in the United States.

**Problem 8.1**


Label the α and β carbons in each alkyl halide. Draw all possible elimination products formed when each alkyl halide is treated with $\text{K}^+ \text{}^- \text{OC}(\text{CH}_3)_3$.



8.2 Alkenes—The Products of Elimination Reactions


Because elimination reactions of alkyl halides form alkenes, let's review earlier material on alkene structure and learn some additional facts as well.

8.2A Bonding in a Carbon–Carbon Double Bond




Ethylene, the simplest alkene, is a hormone that regulates plant growth and fruit ripening. A ripe banana speeds up the ripening of green tomatoes because the banana gives off ethylene.

Recall from Section 1.9B that alkenes are hydrocarbons containing a carbon–carbon double bond. Each carbon of the double bond is sp^2 hybridized and trigonal planar, and all bond angles are 120° .



The double bond of an alkene consists of a σ bond and a π bond.



- The σ bond, formed by end-on overlap of the two sp^2 hybrid orbitals, lies in the plane of the molecule.
- The π bond, formed by side-by-side overlap of two $2p$ orbitals, lies perpendicular to the plane of the molecule. The π bond is formed during elimination.

Alkenes are classified according to the number of carbon atoms bonded to the carbons of the double bond. A **monosubstituted alkene** has one carbon atom bonded to the carbons of the double bond. A **disubstituted alkene** has two carbon atoms bonded to the carbons of the double bond, and so forth.





Figure 8.1 shows several alkenes and how they are classified. You must be able to classify alkenes in this way to determine the major and minor products of elimination reactions, when a mixture of alkenes is formed.


Figure 8.1

Classifying alkenes by the number of R groups bonded to the double bond



- Carbon atoms bonded to the double bond are screened in red.

Problem 8.2 Classify each alkene in the following vitamins by the number of carbon substituents bonded to the double bond.



8.2B Restricted Rotation

Figure 8.2 shows that there is free rotation about the carbon–carbon single bonds of butane, but *not* around the carbon–carbon double bond of 2-butene. Because of restricted rotation, two stereoisomers of 2-butene are possible.

- The *cis* isomer has two groups on the *same side* of the double bond.
- The *trans* isomer has two groups on *opposite sides* of the double bond.

The concept of *cis* and *trans* isomers was first introduced for disubstituted cycloalkanes in Chapter 4. In both cases, a ring or a double bond restricts motion, preventing the rotation of a group from one side of the ring or double bond to the other.








Figure 8.2

Rotation around C–C and C=C compared



These conformations **interconvert** by rotation.
They represent the **same** molecule.




These molecules **do not interconvert** by rotation.
They are **different** molecules.




cis-2-Butene and *trans*-2-butene are stereoisomers, but not mirror images of each other, so they are **diastereomers**.

The *cis* and *trans* isomers of 2-butene are a specific example of a general type of stereoisomer occurring at carbon–carbon double bonds. **Whenever the two groups on each end of a carbon–carbon double bond are different from each other, two diastereomers are possible.**


Stereoisomers on a C=C are possible when:





Problem 8.3 For which alkenes are stereoisomers possible?





Problem 8.4 Label each pair of alkenes as constitutional isomers, stereoisomers, or identical.





and




and



and




and



8.2C Stability of Alkenes


Some alkenes are more stable than others. For example, **trans alkenes are generally more stable than cis alkenes** because the groups bonded to the double bond carbons are farther apart, reducing steric interactions.

The *trans* isomer has the CH_3 groups farther away from each other.



more stable

Steric interactions of the CH_3 groups destabilize the *cis* isomer.



less stable

The stability of an alkene increases, moreover, as the **number of R groups bonded to the double bond carbons increases**.

least stable




most stable

Increasing number of R groups
Increasing stability


R groups increase the stability of an alkene because R groups are sp^3 hybridized, whereas the carbon atoms of the double bond are sp^2 hybridized. Recall from Sections 1.10B and 2.5D that

the percent s -character of a hybrid orbital increases from 25% to 33% in going from sp^3 to sp^2 . The higher the percent s -character, the more readily an atom accepts electron density. Thus, sp^2 hybridized carbon atoms are more able to *accept* electron density and sp^3 hybridized carbon atoms are more able to *donate* electron density.



- As a result, increasing the number of electron-donating R groups on a carbon atom able to accept electron density makes the alkene more stable.

Thus, *trans*-2-butene (a disubstituted alkene) is more stable than *cis*-2-butene (another disubstituted alkene), but both are more stable than 1-butene (a monosubstituted alkene).




Increasing stability

In summary:

- Trans alkenes are more stable than *cis* alkenes because they have fewer steric interactions.
- Increasing alkyl substitution stabilizes an alkene by an electron-donating inductive effect.

Problem 8.5 Which alkene in each pair is more stable?



Problem 8.6 While *cis*- and *trans*-2-butene differ in stability by only 3 kJ/mol, *cis*- and *trans*-2,2,5,5-tetramethyl-3-hexene differ in stability by 44 kJ/mol. Explain why the difference in the stability of the second pair of alkene stereoisomers is so much larger.

Problem 8.7 Several factors can affect alkene stability. Explain why alkene **A** is more stable than alkene **B** even though both contain disubstituted carbon-carbon double bonds.



8.3 The Mechanisms of Elimination

What is the mechanism for elimination? What is the order of bond breaking and bond making? Is the reaction a one-step process or does it occur in many steps?


There are two mechanisms for elimination—E2 and E1—just as there are two mechanisms for nucleophilic substitution—S_N2 and S_N1.

- The **E2 mechanism (bimolecular elimination)**
- The **E1 mechanism (unimolecular elimination)**

The E2 and E1 mechanisms differ in the timing of bond cleavage and bond formation, analogous to the S_N2 and S_N1 mechanisms. In fact, E2 and S_N2 reactions have some features in common, as do E1 and S_N1 reactions.

8.4 The E2 Mechanism

The most common mechanism for dehydrohalogenation is the E2 mechanism. For example, (CH₃)₃CBr reacts with ⁻OH to form (CH₃)₂C=CH₂ via an E2 mechanism.



8.4A Kinetics

An E2 reaction exhibits **second-order kinetics**; that is, the reaction is **bimolecular** and both the alkyl halide and the base appear in the rate equation.


- $\text{rate} = k[(\text{CH}_3)_3\text{CBr}][^-\text{OH}]$

8.4B A One-Step Mechanism

The most straightforward explanation for the second-order kinetics is a **concerted reaction: all bonds are broken and formed in a single step**, as shown in Mechanism 8.1.




Mechanism 8.1 The E2 Mechanism



- The base ⁻OH removes a proton from the β carbon, forming H₂O (a by-product).
- The electron pair in the β C-H bond forms the new π bond.
- The leaving group Br⁻ comes off with the electron pair in the C-Br bond.

An energy diagram for the reaction of (CH₃)₃CBr with ⁻OH is shown in Figure 8.3. The reaction has one step, so there is one energy barrier between reactants and products. Two bonds are broken (C-H and C-Br) and two bonds are formed (H-OH and the π bond) in a single step, so the transition state contains four partial bonds, with the negative charge distributed over the base and the leaving group. Entropy favors the products of an E2 reaction because two molecules of starting material form three molecules of product.


Problem 8.8 Use curved arrows to show the movement of electrons in the following E2 mechanism. Draw the structure of the transition state.



Problem 8.9 Why is (CH₃)₃CCH₂Br inert to an E2 elimination?

Figure 8.3

An energy diagram for an E2 reaction:
 $(\text{CH}_3)_3\text{CBr} + ^-\text{OH} \rightarrow$
 $(\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{H}_2\text{O} + \text{Br}^-$



- In the transition state, the C–H and C–Br bonds are partially broken, the O–H and π bonds are partially formed, and both the base and the departing leaving group bear a partial negative charge.

There are close parallels between the E2 and $\text{S}_{\text{N}}2$ mechanisms in how the identity of the base, the leaving group, and the solvent affect the rate.


The Base

- The base appears in the rate equation, so the rate of the E2 reaction increases as the strength of the base increases.


E2 reactions are generally run with strong, negatively charged bases like ^-OH and ^-OR . Two strong, sterically hindered nitrogen bases, called **DBN** and **DBU**, are also sometimes used. An example of an E2 reaction with DBN is shown in Figure 8.4.

The IUPAC names for **DBN** and **DBU** are rarely used because the names are complex. **DBN** stands for 1,5-diazabicyclo[4.3.0]non-5-ene, and **DBU** stands for 1,8-diazabicyclo[5.4.0]undec-7-ene.

Two useful bases for E2 reactions


**Figure 8.4**

An E2 elimination with DBN used as the base



The Leaving Group

- Because the bond to the leaving group is partially broken in the transition state, the better the leaving group the faster the E2 reaction.



The Solvent

- Polar aprotic solvents increase the rate of E2 reactions.


Because **polar aprotic solvents** like $(\text{CH}_3)_2\text{C}=\text{O}$ do not solvate anions well, a negatively charged base is not “hidden” by strong interactions with the solvent (Section 7.17D), and the base is stronger. **A stronger base increases the reaction rate.**

Problem 8.10 Consider an E2 reaction between $\text{CH}_3\text{CH}_2\text{Br}$ and $\text{KOC}(\text{CH}_3)_3$. What effect does each of the following changes have on the rate of elimination? (a) The base is changed to KOH . (b) The alkyl halide is changed to $\text{CH}_3\text{CH}_2\text{Cl}$.

8.4C The Identity of the Alkyl Halide

The $\text{S}_{\text{N}}2$ and E2 mechanisms differ in how the R group affects the reaction rate.


- As the number of R groups on the carbon with the leaving group increases, the rate of the E2 reaction increases.



This trend is exactly *opposite* to the reactivity of alkyl halides in $\text{S}_{\text{N}}2$ reactions, where increasing alkyl substitution decreases the rate of reaction (Section 7.11D).

Why does increasing alkyl substitution increase the rate of an E2 reaction? In the transition state, the double bond is partially formed, so increasing the stability of the double bond with alkyl substituents stabilizes the transition state (i.e., it lowers E_{a}), which increases the rate of the reaction.


Transition state for an E2 reaction with an alkoxide (^-OR) as base



The double bond is partially formed.

- Increasing the number of R groups on the carbon with the leaving group forms more highly substituted, more stable alkenes in E2 reactions.

For example, the E2 reaction of a 1° alkyl halide (1-bromobutane) forms a monosubstituted alkene, whereas the E2 reaction of a 3° alkyl halide (2-bromo-2-methylpropane) forms a disubstituted alkene. The disubstituted alkene is more stable, so the 3° alkyl halide reacts faster than the 1° alkyl halide.



Problem 8.11 Rank the alkyl halides in each group in order of increasing reactivity in an E2 reaction.




Table 8.2 summarizes the characteristics of the E2 mechanism, and Figure 8.5 illustrates two examples of E2 reactions used in the synthesis of **quinine** and **estradiol**, two naturally occurring compounds.

Problem 8.12 How does each of the following changes affect the rate of an E2 reaction?

- tripling [RX]
- halving [B:]
- changing the solvent from CH₃OH to DMSO
- changing the leaving group from I⁻ to Br⁻
- changing the base from ⁻OH to H₂O
- changing the alkyl halide from CH₃CH₂Br to (CH₃)₂CHBr

Table 8.2 Characteristics of the E2 Mechanism

Characteristic	Result
Kinetics	• Second order
Mechanism	• One step
Identity of R	• More substituted halides react fastest. • Rate: R ₃ CX > R ₂ CHX > RCH ₂ X
Base	• Favored by strong bases
Leaving group	• Better leaving group \longrightarrow faster reaction
Solvent	• Favored by polar aprotic solvents

8.5 The Zaitsev Rule


Recall from Section 8.1 that a mixture of alkenes can form from the dehydrohalogenation of alkyl halides having two or more different β carbon atoms. When this occurs, one of the products usually predominates. The **major product is the more stable product—the one with the more substituted double bond**. For example, elimination of the elements of H and I from

Figure 8.5

Two examples of the E2 reaction used in organic synthesis




Quinine, a natural product isolated from the bark of the cinchona tree native to the Andes Mountains, is a powerful antipyretic—that is, it reduces fever—and for centuries, it was the only effective treatment for malaria.



- Bonds and atoms in quinine and estradiol that originate in the alkene intermediate are shown in red.

1-iodo-1-methylcyclohexane yields two constitutional isomers: the trisubstituted alkene **A** (the major product) and the disubstituted alkene **B** (the minor product).




This phenomenon is called the **Zaitsev rule** (also called the **Saytzeff rule**, depending on the translation) for the Russian chemist who first noted this trend.


- The Zaitsev rule: The major product in β elimination has the more substituted double bond.

A reaction is **regioselective** when it yields predominantly or exclusively one constitutional isomer when more than one is possible. The E2 reaction is regioselective because the more substituted alkene predominates.

The Zaitsev rule results because the double bond is partially formed in the transition state for the E2 reaction. Thus, increasing the stability of the double bond by adding R groups lowers the energy of the transition state, which increases the reaction rate. For example, E2 elimination of HBr from 2-bromo-2-methylbutane yields alkenes **C** and **D**. **D**, having the more substituted double bond, is the major product, because the transition state leading to its formation is lower in energy.




When a mixture of stereoisomers is possible from dehydrohalogenation, the **major product is the more stable stereoisomer**. For example, dehydrohalogenation of alkyl halide **X** forms a mixture of trans and cis alkenes, **Y** and **Z**. The trans alkene **Y** is the major product because it is most stable.




A reaction is *stereoselective* when it forms predominantly or exclusively one stereoisomer when two or more are possible. The E2 reaction is stereoselective because one stereoisomer is formed preferentially.

Sample Problem 8.1 Predict the major product in the following E2 reaction.







Solution

The alkyl halide has two different β C atoms (labeled β_1 and β_2), so two different alkenes are possible: one formed by removal of HCl across the α and β_1 carbons, and one formed by removal of HCl across the α and β_2 carbons. Using the Zaitsev rule, the major product should be **A**, because it has the more substituted double bond.




Problem 8.13 What alkenes are formed from each alkyl halide by an E2 reaction? Use the Zaitsev rule to predict the major product.



8.6 The E1 Mechanism

The dehydrohalogenation of $(\text{CH}_3)_3\text{CI}$ with H_2O to form $(\text{CH}_3)_2\text{C}=\text{CH}_2$ can be used to illustrate the second general mechanism of elimination, the **E1 mechanism**.



8.6A Kinetics

An E1 reaction exhibits first-order kinetics.

- $\text{rate} = k[(\text{CH}_3)_3\text{CI}]$

Like the $\text{S}_{\text{N}}1$ mechanism, the kinetics suggest that the reaction mechanism involves more than one step, and that the slow step is **unimolecular**, involving *only* the alkyl halide.


8.6B A Two-Step Mechanism

The most straightforward explanation for the observed first-order kinetics is a **two-step reaction**: **the bond to the leaving group breaks first before the π bond is formed**, as shown in Mechanism 8.2.



Mechanism 8.2 The E1 Mechanism

Step [1] The C–I bond is broken.



- **Heterolysis of the C–I bond** forms an intermediate **carbocation**. This is the same first step as the $\text{S}_{\text{N}}1$ mechanism. It is responsible for the first-order kinetics because it is rate-determining.

Step [2] A C–H bond is cleaved and the π bond is formed.




- **A base** (such as H_2O or I^-) **removes a proton from a carbon adjacent to the carbocation** (a β carbon). The electron pair in the C–H bond is used to form the new π bond.

The E1 and E2 mechanisms both involve the same number of bonds broken and formed. **The only difference is the timing.**

- In an E1 reaction, the leaving group comes off *before* the β proton is removed, and the reaction occurs in two steps.
- In an E2 reaction, the leaving group comes off as the β proton is removed, and the reaction occurs in one step.

Figure 8.6

Energy diagram
for an E1 reaction:
 $(\text{CH}_3)_3\text{CI} + \text{H}_2\text{O} \rightarrow$
 $(\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{H}_3\text{O}^+ + \text{I}^-$




- Since the E1 mechanism has two steps, there are two energy barriers.
- Step [1] is rate-determining.

An energy diagram for the reaction of $(\text{CH}_3)_3\text{CI} + \text{H}_2\text{O}$ is shown in Figure 8.6. Each step has its own energy barrier, with a transition state at each energy maximum. Because its transition state is higher in energy, Step [1] is rate-determining. ΔH° for Step [1] is positive because only bond breaking occurs, whereas ΔH° of Step [2] is negative because two bonds are formed and only one is broken.


Problem 8.14

Draw an E1 mechanism for the following reaction. Draw the structure of the transition state for each step.

**8.6C Other Characteristics of E1 Reactions**

Three other features of E1 reactions are worthy of note.

[1] **The rate of an E1 reaction increases as the number of R groups on the carbon with the leaving group increases.**



Increasing alkyl substitution has the same effect on the rate of *both* an E1 and E2 reaction; increasing rate of the E1 and E2 reactions: RCH_2X (1°) < R_2CHX (2°) < R_3CX (3°).

Like an $\text{S}_\text{N}1$ reaction, more substituted alkyl halides yield more substituted (and more stable) carbocations in the rate-determining step. Increasing the stability of a carbocation, in turn,

decreases E_a for the slow step, which increases the rate of the E1 reaction according to the Hammond postulate.

[2] Because the base does not appear in the rate equation, weak bases favor E1 reactions.

The strength of the base usually determines whether a reaction follows the E1 or E2 mechanism.

- Strong bases like OH^- and OR^- favor E2 reactions, whereas weaker bases like H_2O and ROH favor E1 reactions.

[3] E1 reactions are regioselective, favoring formation of the more substituted, more stable alkene.

The Zaitsev rule applies to E1 reactions, too. For example, E1 elimination of HBr from 1-bromo-1-methylcyclopentane yields alkenes **A** and **B**. **A**, having the more substituted double bond, is the major product.





Table 8.3 summarizes the characteristics of E1 reactions.

Problem 8.15 What alkenes are formed from each alkyl halide by an E1 reaction? Use the Zaitsev rule to predict the major product.



Problem 8.16 How does each of the following changes affect the rate of an E1 reaction?


- doubling $[\text{RX}]$
- doubling $[\text{B}^-]$
- changing the halide from $(\text{CH}_3)_3\text{CBr}$ to $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$
- changing the leaving group from Cl^- to Br^-
- changing the solvent from DMSO to CH_3OH

Table 8.3 Characteristics of the E1 Mechanism

Characteristic	Result
Kinetics	<ul style="list-style-type: none"> • First order
Mechanism	<ul style="list-style-type: none"> • Two steps
Identity of R	<ul style="list-style-type: none"> • More substituted halides react fastest. • Rate: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X}$
Base	<ul style="list-style-type: none"> • Favored by weaker bases such as H_2O and ROH
Leaving group	<ul style="list-style-type: none"> • A better leaving group makes the reaction faster because the bond to the leaving group is partially broken in the rate-determining step.
Solvent	<ul style="list-style-type: none"> • Polar protic solvents that solvate the ionic intermediates are needed.

8.7 S_N1 and E1 Reactions

S_N1 and E1 reactions have exactly the same first step—formation of a carbocation. They differ in what happens to the carbocation.



- In an S_N1 reaction, a nucleophile attacks the carbocation, forming a substitution product.
- In an E1 reaction, a base removes a proton, forming a new π bond.

The same conditions that favor substitution by an S_N1 mechanism also favor elimination by an E1 mechanism: a 3° alkyl halide as substrate, a weak nucleophile or base as reagent, and a polar protic solvent. As a result, both reactions usually occur in the same reaction mixture to afford a mixture of products, as illustrated in Sample Problem 8.2.

Sample Problem 8.2


Draw the S_N1 and E1 products formed in the reaction of (CH₃)₃CBr with H₂O.

Solution

The first step in both reactions is heterolysis of the C–Br bond to form a **carbocation**.




Reaction of the carbocation with H₂O as a nucleophile affords the substitution product (Reaction [1]). Alternatively, H₂O acts as a base to remove a proton, affording the elimination product (Reaction [2]). **Two products are formed.**



Because E1 reactions often occur with a competing S_N1 reaction, **E1 reactions of alkyl halides are much less useful than E2 reactions.**

Problem 8.17

Draw both the S_N1 and E1 products of each reaction.




8.8 Stereochemistry of the E2 Reaction

Although the E2 reaction does not produce products with tetrahedral stereogenic centers, its transition state consists of four atoms that react at the same time, and they react only if they possess a particular stereochemical arrangement.

8.8A General Stereochemical Features

The transition state of an E2 reaction consists of **four atoms** from the alkyl halide—one hydrogen atom, two carbon atoms, and the leaving group (X)—**all aligned in a plane**. There are two ways for the C–H and C–X bonds to be coplanar.



The dihedral angle for the C–H and C–X bonds equals 0° for the syn periplanar arrangement and 180° for the anti periplanar arrangement.

- The H and X atoms can be oriented on the same side of the molecule. This geometry is called *syn periplanar*.
- The H and X atoms can be oriented on opposite sides of the molecule. This geometry is called *anti periplanar*.

All evidence suggests that **E2 elimination occurs most often in the anti periplanar geometry**. This arrangement allows the molecule to react in the lower energy *staggered* conformation. It also allows two electron-rich species, the incoming base and the departing leaving group, to be farther away from each other, as illustrated in Figure 8.7.


Anti periplanar geometry is the preferred arrangement for any alkyl halide undergoing E2 elimination, regardless of whether it is cyclic or acyclic. This stereochemical requirement has important consequences for compounds containing six-membered rings.

Problem 8.18 Draw the anti periplanar geometry for the E2 reaction of $(\text{CH}_3)_2\text{CHCH}_2\text{Br}$ with base. Then draw the product that results after elimination of HBr.

Problem 8.19 Given that an E2 reaction proceeds with anti periplanar stereochemistry, draw the products of each elimination. The alkyl halides in (a) and (b) are diastereomers of each other. How are the products of these two reactions related? Recall from Section 3.2A that C_6H_5^- is a phenyl group, a benzene ring bonded to another group.




Figure 8.7
Two possible geometries for the E2 reaction




8.8B Anti Periplanar Geometry and Halocyclohexanes

Recall from Section 4.13 that cyclohexane exists as two chair conformations that rapidly interconvert, and that substituted cyclohexanes are more stable with substituents in the roomier equatorial position. Thus, chlorocyclohexane exists as two chair conformations, but **X** is preferred because the Cl group is equatorial.



For E2 elimination, **the C–Cl bond must be anti periplanar to a C–H bond on a β carbon**, and this occurs only when the H and Cl atoms are both in the **axial** position. This requirement for **trans diaxial geometry** means that E2 elimination must occur from the less stable conformation **Y**, as shown in Figure 8.8.

Sometimes this rigid stereochemical requirement affects the regioselectivity of the E2 reaction of substituted cyclohexanes. Dehydrohalogenation of *cis*- and *trans*-1-chloro-2-methylcyclohexane via an E2 mechanism illustrates this phenomenon.



The **cis isomer** exists as two conformations (**A** and **B**), each of which has one group axial and one group equatorial. E2 reaction must occur from conformation **B**, which contains an axial Cl atom.





Figure 8.8


The trans diaxial geometry for the E2 elimination in chlorocyclohexane

Conformation X (equatorial Cl):




no reaction with this conformation

Conformation Y (axial Cl):




The H and Cl are **trans diaxial**.




- In conformation **X** (**equatorial** Cl group), a β C–H bond and a C–Cl bond are never anti periplanar; therefore, no E2 elimination can occur.
- In conformation **Y** (**axial** Cl group), a β C–H bond and a C–Cl bond are **trans diaxial**; therefore, E2 elimination occurs.


Because conformation **B** has two different axial β H atoms, labeled H_a and H_b , E2 reaction occurs in two different directions to afford two alkenes. **The major product contains the more stable trisubstituted double bond, as predicted by the Zaitsev rule.**



The **trans isomer** exists as two conformations, **C**, having two equatorial substituents, and **D**, having two axial substituents. E2 reaction must occur from conformation **D**, which contains an axial Cl atom.




Because conformation **D** has **only one axial β H**, E2 reaction occurs in only one direction to afford a **single product**, having the disubstituted double bond. This is *not* predicted by the Zaitsev rule. E2 reaction requires H and Cl to be trans and diaxial, and with the trans isomer, this is possible only when the less stable alkene is formed as product.



- In conclusion, with substituted cyclohexanes, E2 elimination must occur with a trans diaxial arrangement of H and X, and as a result of this requirement, the more substituted alkene is not necessarily the major product.


Sample Problem 8.3

Draw the major E2 elimination product formed from the following alkyl halide.




Solution

To draw the elimination products, locate the β carbons and look for H atoms that are trans to the leaving group. The given alkyl chloride has two different β carbons, labeled β_1 and β_2 . Elimination can occur only when the leaving group (Cl) and a H atom on the β carbon are trans.



Since the β_1 C has a H atom *trans* to Cl, E2 elimination occurs to form a disubstituted alkene. Since there is no *trans* H on the β_2 C, E2 elimination cannot occur in this direction, and the more stable trisubstituted alkene is *not* formed. Although this result is not predicted by the Zaitsev rule, it is consistent with the requirement that the H and X atoms in an E2 elimination must be located *trans* to each other.

Problem 8.20 Draw the major E2 elimination products from each of the following alkyl halides.



Problem 8.21 Explain why *cis*-1-chloro-2-methylcyclohexane undergoes E2 elimination much faster than its *trans* isomer.

8.9 When Is the Mechanism E1 or E2?

Given a particular starting material and base, how do we know whether a reaction occurs by the E1 or E2 mechanism?

Because the rate of *both* the E1 and E2 reactions increases as the number of R groups on the carbon with the leaving group increases, you cannot use the identity of the alkyl halide to decide which elimination mechanism occurs. This makes determining the mechanisms for substitution and elimination very different processes.


- The strength of the base is the most important factor in determining the mechanism for elimination. Strong bases favor the E2 mechanism. Weak bases favor the E1 mechanism.

Table 8.4 compares the E1 and E2 mechanisms.

Table 8.4 A Comparison of the E1 and E2 Mechanisms

Mechanism	Comment
E2 mechanism	<ul style="list-style-type: none"> • Much more common and useful • Favored by strong, negatively charged bases, especially OH^- and OR^- • The reaction occurs with 1°, 2°, and 3° alkyl halides. Order of reactivity: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X}$.
E1 mechanism	<ul style="list-style-type: none"> • Much less useful because a mixture of $\text{S}_{\text{N}}1$ and E1 products usually results • Favored by weaker, neutral bases, such as H_2O and ROH • This mechanism does not occur with 1° RX because they form highly unstable 1° carbocations.

Problem 8.22 Which mechanism, E1 or E2, will occur in each reaction?



8.10 E2 Reactions and Alkyne Synthesis


Recall from Section 1.9C that the carbon–carbon triple bond of alkynes consists of one σ and two π bonds.

A single elimination reaction produces the π bond of an alkene. Two consecutive elimination reactions produce the two π bonds of an alkyne.




- Alkynes are prepared by two successive dehydrohalogenation reactions.

Two elimination reactions are needed to remove two moles of HX from a **dihalide** as substrate. Two different starting materials can be used.



- A **vicinal dihalide** has two X atoms on *adjacent* carbon atoms.
- A **geminal dihalide** has two X atoms on the *same* carbon atom.

Equations [1] and [2] illustrate how two moles of HX can be removed from these dihalides with base. Two equivalents of strong base are used and each step follows an E2 mechanism.




The word *geminal* comes from the Latin *geminus*, meaning *twin*.

The relative strength of C–H bonds depends on the hybridization of the carbon atom: $sp > sp^2 > sp^3$. For more information, review Section 1.10B.


Stronger bases are needed to synthesize alkynes by dehydrohalogenation than are needed to synthesize alkenes. The typical base is **amide** ($^-\text{NH}_2$), used as the sodium salt NaNH_2 (sodium amide). $\text{KOC}(\text{CH}_3)_3$ can also be used with DMSO as solvent. Because DMSO is a polar aprotic solvent, the anionic base is not well solvated, thus **increasing its basicity** and making it strong enough to remove two equivalents of HX. Examples are given in Figure 8.9.

Figure 8.9
Example of
dehydrohalogenation
of dihalides to afford alkynes



The strongly basic conditions needed for alkyne synthesis result from the difficulty of removing the second equivalent of HX from the intermediate vinyl halide, $\text{RCH}=\text{C}(\text{R})\text{X}$. Since H and X are both bonded to sp^2 hybridized carbons, these bonds are shorter and stronger than the sp^3 hybridized C–H and C–X bonds of an alkyl halide, necessitating the use of a stronger base.

Problem 8.23 Draw the alkynes formed in each reaction. Two equivalents of each base are used.




8.11 When Is the Reaction $\text{S}_{\text{N}}1$, $\text{S}_{\text{N}}2$, E1, or E2?

We have now considered two different kinds of reactions (substitution and elimination) and four different mechanisms ($\text{S}_{\text{N}}1$, $\text{S}_{\text{N}}2$, E1, and E2) that begin with one class of compounds (alkyl halides). How do we know if a given alkyl halide will undergo substitution or elimination with a given base or nucleophile, and by what mechanism?

Unfortunately, there is no easy answer, and often mixtures of products result. Two generalizations help to determine whether substitution or elimination occurs.


[1] Good nucleophiles that are weak bases favor substitution over elimination.

Certain anions generally give products of substitution because they are good nucleophiles but weak bases. These include: I^- , Br^- , HS^- , CN^- , and CH_3COO^- .



[2] Bulky, nonnucleophilic bases favor elimination over substitution.

$\text{KOC}(\text{CH}_3)_3$, DBU, and DBN are too sterically hindered to attack a tetravalent carbon, but are able to remove a small proton, favoring elimination over substitution.




Most often, however, we will have to rely on other criteria to predict the outcome of these reactions. To determine the product of a reaction with an alkyl halide:

- [1] Classify the alkyl halide as 1° , 2° , or 3° .
- [2] Classify the base or nucleophile as strong, weak, or bulky.

Predicting the substitution and elimination products of a reaction can then be organized by the type of alkyl halide, as shown in Figure 8.10.


Sample Problems 8.4–8.6 illustrate how to apply the information in Figure 8.10 to specific alkyl halides.

Sample Problem 8.4 Draw the products of the following reaction.




Solution

- [1] Classify the halide as 1° , 2° , or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 3° and the reagent (H_2O) is a weak base and nucleophile, so products of both S_N1 and E1 mechanisms are formed.
- [2] To draw the products of substitution and elimination:



Sample Problem 8.5 Draw the products of the following reaction.



Solution

- [1] Classify the halide as 1° , 2° , or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 2° and the reagent (CH_3O^-) is a strong base and nucleophile, so products of both S_N2 and E2 mechanisms are formed.
- [2] To draw the products of substitution and elimination:





Figure 8.10 Determining whether an alkyl halide reacts by an S_N1, S_N2, E1, or E2 mechanism

[1] 3° Alkyl halides (R₃CX react by all mechanisms except S_N2.)

- With strong bases

- Elimination occurs by an E2 mechanism.**
- Rationale: A strong base or nucleophile favors an S_N2 or E2 mechanism, but 3° halides are too sterically hindered to undergo an S_N2 reaction, so only E2 elimination occurs.


- Example:



- With weak nucleophiles or bases

- A mixture of S_N1 and E1 products results.**
- Rationale: A weak base or nucleophile favors S_N1 and E1 mechanisms, and both occur.

- Example:




[2] 1° Alkyl halides (RCH₂X react by S_N2 and E2 mechanisms.)

- With strong nucleophiles

- Substitution occurs by an S_N2 mechanism.**
- Rationale: A strong base or nucleophile favors S_N2 or E2, but 1° halides are the least reactive halide type in elimination; therefore, only an S_N2 reaction occurs.


- Example:



- With strong, sterically hindered bases

- Elimination occurs by an E2 mechanism.**
- Rationale: A strong, sterically hindered base cannot act as a nucleophile, so elimination occurs and the mechanism is E2.

- Example:




[3] 2° Alkyl halides (R₂CHX react by all mechanisms.)

- With strong bases and nucleophiles


- A mixture of S_N2 and E2 products results.**
- Rationale: A strong base that is also a strong nucleophile gives a mixture of S_N2 and E2 products.

- Example:




- With strong, sterically hindered bases

- **Elimination occurs by an E2 mechanism.**
- Rationale: A strong, sterically hindered base cannot act as a nucleophile, so elimination occurs and the mechanism is E2.
- Example:




- With weak nucleophiles or bases

- **A mixture of S_N1 and E1 products results.**
- Rationale: A weak base or nucleophile favors S_N1 and E1 mechanisms, and both occur.
- Example:




Sample Problem 8.6

Draw the products of the following reaction, and include the mechanism showing how each product is formed.




Solution


- [1] Classify the halide as 1°, 2°, or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 3° and the reagent (CH₃OH) is a weak base and nucleophile, so products of both **S_N1** and **E1** mechanisms are formed.
- [2] Draw the steps of the mechanisms to give the products. Both mechanisms begin with the same first step: loss of the leaving group to form a carbocation.



- **For S_N1:** The carbocation reacts with a nucleophile. Nucleophilic attack of CH₃OH on the carbocation generates a positively charged intermediate that loses a proton to afford the neutral S_N1 product.




- **For E1:** The carbocation reacts with a base (CH₃OH or Br⁻). Two different products of elimination can form because the carbocation has two different β carbons.




In this problem, three products are formed: one from an S_N1 reaction and two from E1 reactions.

Problem 8.24 Draw the products in each reaction.



Problem 8.25 Draw a stepwise mechanism for the following reaction.




KEY CONCEPTS


Alkyl Halides and Elimination Reactions

A Comparison Between Nucleophilic Substitution and β Elimination

Nucleophilic substitution—A nucleophile attacks a carbon atom (7.6).



β Elimination—A base attacks a proton (8.1).



Similarities

- In both reactions RX acts as an electrophile, reacting with an electron-rich reagent.
- Both reactions require a **good leaving group** X^- that can accept the electron density in the C–X bond.


Differences

- In substitution, a nucleophile attacks a single carbon atom.
- In elimination, a Brønsted–Lowry base removes a proton to form a π bond, and two carbons are involved in the reaction.

The Importance of the Base in E2 and E1 Reactions (8.9)

The strength of the base determines the mechanism of elimination.

- Strong bases favor E2 reactions.
- Weak bases favor E1 reactions.



E1 and E2 Mechanisms Compared

	E2 mechanism	E1 mechanism
Mechanism	• One step (8.4B)	• Two steps (8.6B)
Alkyl halide	• Rate: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X}$ (8.4C)	• Rate: $\text{R}_3\text{CX} > \text{R}_2\text{CHX} > \text{RCH}_2\text{X}$ (8.6C)
Rate equation	• Rate = $k[\text{RX}][\text{B}^-]$ • Second-order kinetics (8.4A)	• Rate = $k[\text{RX}]$ • First-order kinetics (8.6A)
Stereochemistry	• Anti periplanar arrangement of H and X (8.8)	• Trigonal planar carbocation intermediate (8.6B)
Base	• Favored by strong bases (8.4B)	• Favored by weak bases (8.6C)
Leaving group	• Better leaving group \rightarrow faster reaction (8.4B)	• Better leaving group \rightarrow faster reaction (Table 8.3)
Solvent	• Favored by polar aprotic solvents (8.4B)	• Favored by polar protic solvents (Table 8.3)
Product	• More substituted alkene favored (Zaitsev rule, 8.5)	• More substituted alkene favored (Zaitsev rule, 8.6C)


Summary Chart on the Four Mechanisms: $\text{S}_{\text{N}}1$, $\text{S}_{\text{N}}2$, E1, or E2

Alkyl halide type	Conditions	Mechanism
$1^\circ \text{RCH}_2\text{X}$	strong nucleophile \rightarrow	$\text{S}_{\text{N}}2$
	strong bulky base \rightarrow	E2
$2^\circ \text{R}_2\text{CHX}$	strong base and nucleophile \rightarrow	$\text{S}_{\text{N}}2 + \text{E2}$
	strong bulky base \rightarrow	E2
	weak base and nucleophile \rightarrow	$\text{S}_{\text{N}}1 + \text{E1}$
$3^\circ \text{R}_3\text{CX}$	weak base and nucleophile \rightarrow	$\text{S}_{\text{N}}1 + \text{E1}$
	strong base \rightarrow	E2


PROBLEMS

General Elimination

8.26 Draw all possible constitutional isomers formed by dehydrohalogenation of each alkyl halide.





8.27 What alkyl halide forms each of the following alkenes as the *only* product in an elimination reaction?








Alkenes

8.28 Which double bonds in the following natural products can exhibit stereoisomerism? Farnesene is found in the waxy coating of apple skins and geranial is isolated from lemon grass.




8.29 Label each pair of alkenes as constitutional isomers, stereoisomers, or identical.







8.30 Draw all isomers of molecular formula C_2H_2BrCl . Label pairs of diastereomers and constitutional isomers.



8.31 $PGF_{2\alpha}$ is a prostaglandin, a group of compounds that are responsible for inflammation (Section 19.6). (a) How many tetrahedral stereogenic centers does $PGF_{2\alpha}$ contain? (b) How many double bonds can exist as *cis* and *trans* isomers? (c) Considering both double bonds and tetrahedral stereogenic centers, what is the maximum number of stereoisomers that can exist for $PGF_{2\alpha}$?



8.32 Rank the alkenes in each group in order of increasing stability.









8.33 ΔH° values obtained for a series of similar reactions are one set of experimental data used to determine the relative stability of alkenes. Explain how the following data suggest that *cis*-2-butene is more stable than 1-butene (Section 12.3A).




E2 Reaction

8.34 Draw all constitutional isomers formed in each E2 reaction and predict the major product using the Zaitsev rule.




8.35 For each of the following alkenes, draw the structure of two different alkyl halides that yield the given alkene as the only product of dehydrohalogenation.




8.36 Explain why $(\text{CH}_3)_2\text{CHCH}(\text{Br})\text{CH}_2\text{CH}_3$ reacts faster than $(\text{CH}_3)_2\text{CHCH}_2\text{CH}(\text{Br})\text{CH}_3$ in an E2 reaction, even though both alkyl halides are 2° .

8.37 Consider the following E2 reaction.




- Draw the by-products of the reaction and use curved arrows to show the movement of electrons.
- What happens to the reaction rate with each of the following changes? [1] The solvent is changed to DMF. [2] The concentration of $^-\text{OC}(\text{CH}_3)_3$ is decreased. [3] The base is changed to ^-OH . [4] The halide is changed to $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{Br})\text{CH}_3$. [5] The leaving group is changed to I^- .

8.38 Dehydrohalogenation of 1-chloro-1-methylcyclopropane affords two alkenes (**A** and **B**) as products. Explain why **A** is the major product despite the fact that it contains the less substituted double bond.




8.39 What is the major stereoisomer formed in each reaction?




E1 Reaction

8.40 What alkene is the major product formed from each alkyl halide in an E1 reaction?




E1 and E2


8.41 Draw all constitutional isomers formed in each elimination reaction. Label the mechanism as E2 or E1.



8.42 Rank the alkyl halides in each group in order of increasing E2 reactivity. Then do the same for E1 reactivity.




8.43 Which elimination reaction in each pair is faster?




8.44 In the dehydrohalogenation of bromocyclodecane, the major product is *cis*-cyclodecene rather than *trans*-cyclodecene. Offer an explanation.

8.45 Explain the following observation. Treatment of alkyl chloride **A** with $\text{NaOCH}_2\text{CH}_3$ yields only one product **B**, whereas treatment of **A** with very dilute base in $\text{CH}_3\text{CH}_2\text{OH}$ yields a mixture of alkenes **B** and **C**, with **C** predominating.




Stereochemistry and the E2 Reaction

8.46 What is the major E2 elimination product formed from each halide?



8.47 Taking into account anti periplanar geometry, predict the major E2 product formed from each starting material.




8.48 a. Draw three-dimensional representations for all stereoisomers of 2-chloro-3-methylpentane, and label pairs of enantiomers.
 b. Considering dehydrohalogenation across C2 and C3 only, draw the E2 product that results from each of these alkyl halides. How many different products have you drawn?
 c. How are these products related to each other?




8.49 Which stereoisomer—*cis*- or *trans*-1-bromo-3-*tert*-butylcyclohexane—will react faster in an E2 elimination reaction? Offer an explanation.

Alkynes

8.50 Draw the products of each reaction.



8.51 Draw the structure of a dihalide that could be used to prepare each alkyne. There may be more than one possible dihalide.













8.52 Under certain reaction conditions, 2,3-dibromobutane reacts with two equivalents of base to give three products, each of which contains two new π bonds. Product **A** has two sp hybridized carbon atoms, product **B** has one sp hybridized carbon atom, and product **C** has none. What are the structures of **A**, **B**, and **C**?

S_N1 , S_N2 , E1, and E2 Mechanisms





8.53 For which reaction mechanisms— S_N1 , S_N2 , E1, or E2—are each of the following statements true? A statement may be true for one or more mechanisms.

- The mechanism involves carbocation intermediates.
- The mechanism has two steps.
- The reaction rate increases with better leaving groups.
- The reaction rate increases when the solvent is changed from CH_3OH to $(\text{CH}_3)_2\text{SO}$.
- The reaction rate depends on the concentration of the alkyl halide only.
- The mechanism is concerted.
- The reaction of $\text{CH}_3\text{CH}_2\text{Br}$ with NaOH occurs by this mechanism.
- Racemization at a stereogenic center occurs.
- Tertiary (3°) alkyl halides react faster than 2° or 1° alkyl halides.
- The reaction follows a second-order rate equation.



8.54 Draw the organic products formed in each reaction.



8.55 What is the major product formed when each alkyl halide is treated with each of the following reagents: [1] NaOCOCH_3 ; [2] NaOCH_3 ; [3] $\text{KOC}(\text{CH}_3)_3$? If it is not possible to predict the major product, identify the products in the mixture and the mechanism by which each is formed.



8.56




cis-1-bromo-4-*tert*-butylcyclohexane


trans-1-bromo-4-*tert*-butylcyclohexane

- The *cis* and *trans* isomers of 1-bromo-4-*tert*-butylcyclohexane react at different rates with $\text{KOC}(\text{CH}_3)_3$ to afford the same mixture of enantiomers **A** and **B**. Draw the structures of **A** and **B**.
- Which isomer reacts faster with $\text{KOC}(\text{CH}_3)_3$? Offer an explanation for this difference in reactivity.
- Reaction of *cis*-1-bromo-4-*tert*-butylcyclohexane with NaOCH_3 affords **C** as the major product, whereas reaction of *trans*-1-bromo-4-*tert*-butylcyclohexane affords **D** as the major product. Draw the structures for **C** and **D**.
- The *cis* and *trans* isomers react at different rates with NaOCH_3 . Which isomer reacts faster? Offer an explanation for the difference in reactivity.
- Why are different products formed from these alkyl halides when two different alkoxides are used as reagents?


8.57 Draw all products, including stereoisomers, in each reaction.




8.58 Draw all of the substitution and elimination products formed from the following alkyl halide with each reagent. Indicate the stereochemistry around the stereogenic centers present in the products, as well as the mechanism by which each product is formed.




8.59 The following reactions do not afford the major product that is given. Explain why this is so, and draw the structure of the major product actually formed.




8.60 Draw a stepwise, detailed mechanism for each reaction.

8.61 Draw the major product formed when (3R)-1-chloro-3-methylpentane is treated with each reagent: (a) $\text{NaOCH}_2\text{CH}_3$; (b) KCN ; (c) DBU.

8.62 Draw a stepwise, detailed mechanism for the following reaction.


8.63 Explain why the reaction of 2-bromopropane with NaOCOCH_3 gives $(\text{CH}_3)_2\text{CHOCOCH}_3$ exclusively as product, but the reaction of 2-bromopropane with $\text{NaOCH}_2\text{CH}_3$ gives a mixture of $(\text{CH}_3)_2\text{CHOCH}_2\text{CH}_3$ (20%) and $\text{CH}_3\text{CH}=\text{CH}_2$ (80%).

8.64 Draw a stepwise detailed mechanism that illustrates how four organic products are formed in the following reaction.




Challenge Problems


- 8.65** Although there are nine stereoisomers of 1,2,3,4,5,6-hexachlorocyclohexane, one stereoisomer reacts 7000 times more slowly than any of the others in an E2 elimination. Draw the structure of this isomer and explain why this is so.
- 8.66** Explain the selectivity observed in the following reactions.



- 8.67** Draw a stepwise mechanism for the following reaction. The four-membered ring in the starting material and product is called a β -lactam. This functional group confers biological activity on penicillin and many related antibiotics, as is discussed in Chapter 22. (Hint: The mechanism begins with β elimination and involves only two steps.)




- 8.68** Although dehydrohalogenation occurs with anti periplanar geometry, some eliminations have syn periplanar geometry. Examine the starting material and product of each elimination, and state whether the elimination occurs with syn or anti periplanar geometry.



9

Alcohols, Ethers, and Epoxides

- 9.1 Introduction
- 9.2 Structure and bonding
- 9.3 Nomenclature
- 9.4 Physical properties
- 9.5 Interesting alcohols, ethers, and epoxides
- 9.6 Preparation of alcohols, ethers, and epoxides
- 9.7 General features—
Reactions of alcohols, ethers, and epoxides
- 9.8 Dehydration of alcohols to alkenes
- 9.9 Carbocation rearrangements
- 9.10 Dehydration using POCl_3 and pyridine
- 9.11 Conversion of alcohols to alkyl halides with HX
- 9.12 Conversion of alcohols to alkyl halides with SOCl_2 and PBr_3
- 9.13 Tosylate—Another good leaving group
- 9.14 Reaction of ethers with strong acid
- 9.15 Reactions of epoxides
- 9.16 Application: Epoxides, leukotrienes, and asthma
- 9.17 Application: Benzo[a]pyrene, epoxides, and cancer




Palytoxin ($\text{C}_{129}\text{H}_{223}\text{N}_3\text{O}_{54}$), first isolated from marine soft corals of the genus *Palythoa*, is a potent poison that contains several hydroxy (OH) groups. Historically used by ancient Hawaiians to poison their spears, palytoxin was isolated in 1971 at the University of Hawai'i at Mānoa, and its structure determined simultaneously by two different research groups in 1981. Its many functional groups and stereogenic centers made it a formidable synthetic target, but in 1994, Harvard chemists synthesized palytoxin in the laboratory. In Chapter 9 we learn about the properties of alcohols like palytoxin, as well as related oxygen-containing functional groups.

In Chapter 9, we take the principles learned in Chapters 7 and 8 about leaving groups, nucleophiles, and bases, and apply them to **alcohols, ethers, and epoxides**, three new functional groups that contain polar C–O bonds. In the process, you will discover that all of the reactions in Chapter 9 follow one of the four mechanisms introduced in Chapters 7 and 8— S_N1 , S_N2 , E1, or E2—so there are **no new general mechanisms to learn**.


Although alcohols, ethers, and epoxides share many characteristics, each functional group has its own distinct reactivity, making each unique and different from the alkyl halides studied in Chapters 7 and 8. Appreciate the similarities but pay attention to the differences.

9.1 Introduction


Alcohols, ethers, and epoxides are three functional groups that contain carbon–oxygen σ bonds.




Alcohols contain a hydroxy group (OH group) bonded to an sp^3 hybridized carbon atom. Alcohols are classified as **primary (1°)**, **secondary (2°)**, or **tertiary (3°)** based on the number of carbon atoms bonded to the carbon with the OH group.



Compounds having a hydroxy group on an sp^2 hybridized carbon atom—**enols** and **phenols**—undergo different reactions than alcohols and are discussed in Chapters 11 and 19, respectively. **Enols** have an OH group on a carbon of a C–C double bond. **Phenols** have an OH group on a benzene ring.



Ethers have two alkyl groups bonded to an oxygen atom. An ether is **symmetrical** if the two alkyl groups are the same, and **unsymmetrical** if they are different. Both alcohols and ethers are organic derivatives of H_2O , formed by replacing one or both of the hydrogens on the oxygen atom by R groups, respectively.



Epoxides are ethers having the oxygen atom in a three-membered ring. Epoxides are also called **oxiranes**.




Problem 9.1

Draw all constitutional isomers having molecular formula $C_4H_{10}O$. Classify each compound as a 1° , 2° , or 3° alcohol, or a symmetrical or unsymmetrical ether.

Figure 9.1


Electrostatic potential maps for a simple alcohol, ether, and epoxide



- Electron-rich regions are shown by the red around the O atoms.


Problem 9.2

Classify each OH group in cortisol as 1°, 2°, or 3°. Cortisol is a hormone produced by the adrenal gland that increases blood pressure and blood glucose levels, and acts as an anti-inflammatory agent.




9.2 Structure and Bonding

Alcohols, ethers, and epoxides each contain an oxygen atom surrounded by two atoms and two nonbonded electron pairs, making the O atom **tetrahedral** and sp^3 hybridized. Because only two of the four groups around O are atoms, alcohols and ethers have a **bent** shape like H_2O .



The bond angle around the O atom in an alcohol or ether is similar to the tetrahedral bond angle of 109.5° . In contrast, the C–O–C bond angle of an epoxide must be 60° , a considerable deviation from the tetrahedral bond angle. For this reason, **epoxides have angle strain**, making them much more reactive than other ethers.



Because oxygen is much more electronegative than carbon or hydrogen, the C–O and O–H bonds are all polar, with the O atom electron rich and the C and H atoms electron poor. The electrostatic potential maps in Figure 9.1 show these polar bonds for all three functional groups.

9.3 Nomenclature


To name an alcohol, ether, or epoxide using the IUPAC system, we must learn how to name the functional group either as a substituent or by using a suffix added to the parent name.

9.3A Naming Alcohols


In the IUPAC system, alcohols are identified by the suffix *-ol*.

HOW TO Name an Alcohol Using the IUPAC System

Example Give the IUPAC name of the following alcohol:




Step [1] Find the longest carbon chain containing the carbon bonded to the OH group.




- Change the **-e** ending of the parent alkane to the suffix **-ol**.

Step [2] Number the carbon chain to give the OH group the lower number, and apply all other rules of nomenclature.

a. **Number** the chain.



b. **Name and number** the substituents.




$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ is named as 1-butanol using the 1979 IUPAC recommendations and butan-1-ol using the 1993 IUPAC recommendations. The first convention is more widely used, so we follow it in this text.

When an OH group is bonded to a ring, the **ring is numbered beginning with the OH group**. Because the functional group is always at C1, the "1" is usually omitted from the name. The ring is then numbered in a clockwise or counterclockwise fashion to give the next substituent the lower number. Representative examples are given in Figure 9.2.

Common names are often used for simple alcohols. To assign a common name:


- Name all the carbon atoms of the molecule as a single **alkyl group**.
- Add the word **alcohol**, separating the words with a space.




Compounds with two hydroxy groups are called **diols** (using the IUPAC system) or **glycols**. Compounds with three hydroxy groups are called **triols**, and so forth. To name a diol, for example, the suffix **-diol** is added to the name of the parent alkane, and numbers are used in the prefix to indicate the location of the two OH groups.

Figure 9.2

Examples: Naming cyclic alcohols




3-methylcyclohexanol




2,5,5-trimethylcyclohexanol

[The OH group is at C1; the second substituent (CH_3) gets the lower number.] [The OH group is at C1; the second substituent (CH_3) gets the lower number.]



Problem 9.3 Give the IUPAC name for each compound.




Problem 9.4 Give the structure corresponding to each name.

- a. 7,7-dimethyl-4-octanol c. 2-*tert*-butyl-3-methylcyclohexanol
 b. 5-methyl-4-propyl-3-heptanol d. *trans*-1,2-cyclohexanediol

9.3B Naming Ethers


Simple ethers are usually assigned common names. To do so, **name both alkyl groups** bonded to the oxygen, arrange these names alphabetically, and add the word **ether**. For symmetrical ethers, name the alkyl group and add the prefix **di-**.



[Alphabetize the **b** of **butyl** before the **m** of **methyl**.]


More complex ethers are named using the IUPAC system. One alkyl group is named as a hydrocarbon chain, and the other is named as part of a substituent bonded to that chain.

- Name the simpler alkyl group + O atom as an **alkoxy** substituent by changing the **-yl** ending of the alkyl group to **-oxy**.




- Name the remaining alkyl group as an alkane, with the alkoxy group as a substituent bonded to this chain.

Sample Problem 9.1 Give the IUPAC name for the following ether.




Solution

[1] Name the longer chain as an alkane and the shorter chain as an alkoxy group.






[2] Apply the other nomenclature rules to complete the name.



Answer: 4-ethoxyoctane

Problem 9.5 Name each of the following ethers.



Problem 9.6 Name each simple ether as an alkoxy alkane: (a) $(\text{CH}_3\text{CH}_2)_2\text{O}$, an anesthetic commonly called diethyl ether; (b) $(\text{CH}_3)_3\text{COCH}_3$, a gasoline additive commonly referred to as MTBE.




Cyclic ethers have an O atom in a ring. A common cyclic ether is **tetrahydrofuran (THF)**, a polar aprotic solvent used in nucleophilic substitution (Section 7.8C) and many other organic reactions.

9.3C Naming Epoxides


Any cyclic compound containing a heteroatom is called a **heterocycle**.

Epoxides are named in three different ways—**epoxyalkanes**, **oxiranes**, or **alkene oxides**.


To name an epoxide as an **epoxyalkane**, first name the alkane chain or ring to which the oxygen is attached, and use the prefix **epoxy** to name the epoxide as a substituent. Use two numbers to designate the location of the atoms to which the O's are bonded.



1,2-epoxycyclohexane




1,2-epoxy-2-methylpropane



cis-2,3-epoxypentane


Epoxides bonded to a chain of carbon atoms can also be named as derivatives of **oxirane**, the simplest epoxide having two carbons and one oxygen atom in a ring. The oxirane ring is numbered to put the O atom at position "1," and the first substituent at position "2." No number is used for a substituent in a monosubstituted oxirane.

Number the ring beginning at the O atom.




oxirane


position 2




2,2-dimethyloxirane

Epoxides are also named as **alkene oxides**, since they are often prepared by adding an O atom to an alkene (Chapter 12). To name an epoxide this way, mentally replace the epoxide oxygen by a double bond, name the alkene (Section 10.3), and then add the word **oxide**. For example, the common name for oxirane is ethylene oxide, since it is an epoxide derived from the alkene ethylene. We will use this method of naming epoxides after the details of alkene nomenclature are presented in Chapter 10.




Problem 9.7 Name each epoxide.**9.4 Physical Properties**

Alcohols, ethers, and epoxides exhibit dipole–dipole interactions because they have a bent structure with two polar bonds. **Alcohols are also capable of intermolecular hydrogen bonding**, because they possess a hydrogen atom on an oxygen, making alcohols much more polar than ethers and epoxides.



Steric factors affect the extent of hydrogen bonding. Although all alcohols can hydrogen bond, increasing the number of R groups around the carbon atom bearing the OH group decreases the extent of hydrogen bonding. Thus, 3° alcohols are least able to hydrogen bond, whereas 1° alcohols are most able to.




How these factors affect the physical properties of alcohols, ethers, and epoxides is summarized in Table 9.1.

Table 9.1 Physical Properties of Alcohols, Ethers, and Epoxides

Property	Observation
Boiling point (bp) and melting point (mp)	<ul style="list-style-type: none"> For compounds of comparable molecular weight, the stronger the intermolecular forces, the higher the bp or mp. <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ $\text{CH}_3\text{OCH}_2\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ VDW VDW, DD VDW, DD, HB bp 0 °C bp 11 °C bp 97 °C </div> <p style="text-align: center;">→ Increasing boiling point</p> <hr/> <ul style="list-style-type: none"> Bp's increase as the extent of hydrogen bonding increases. <div style="text-align: center;"> $(\text{CH}_3)_3\text{C-OH}$ $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{-OH}$ 3° 2° 1° bp 83 °C bp 98 °C bp 118 °C </div> <p style="text-align: center;">→ Increasing ability to hydrogen bond Increasing boiling point</p>
Solubility	<ul style="list-style-type: none"> Alcohols, ethers, and epoxides having ≤ 5 C's are H₂O soluble because they each have an oxygen atom capable of hydrogen bonding to H₂O (Section 3.4C). Alcohols, ethers, and epoxides having > 5 C's are H₂O insoluble because the nonpolar alkyl portion is too large to dissolve in H₂O. Alcohols, ethers, and epoxides of any size are soluble in organic solvents.

Key: VDW = van der Waals forces; DD = dipole–dipole; HB = hydrogen bonding

Problem 9.8 Rank the following compounds in order of increasing boiling point.



Problem 9.9 Explain why dimethyl ether ($\text{CH}_3)_2\text{O}$ and ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) are both water soluble, but the boiling point of ethanol (78°C) is much higher than the boiling point of dimethyl ether (-24°C).

9.5 Interesting Alcohols, Ethers, and Epoxides

A large number of alcohols, ethers, and epoxides have interesting and useful properties.

9.5A Interesting Alcohols

The structure and properties of three simple alcohols—methanol, 2-propanol, and ethylene glycol—are given in Figure 9.3. **Ethanol** ($\text{CH}_3\text{CH}_2\text{OH}$), formed by the fermentation of the carbohydrates in grains, grapes, and potatoes, is the alcohol present in alcoholic beverages. It is perhaps the first organic compound synthesized by humans, because alcohol production has been known for at least 4000 years. Ethanol depresses the central nervous system, increases the production of stomach acid, and dilates blood vessels, producing a flushed appearance. Ethanol is also a common laboratory solvent, which is sometimes made unfit to ingest by adding small amounts of benzene or methanol (both of which are toxic).


Ethanol is a common gasoline additive, widely touted as an environmentally friendly fuel source. Two common gasoline–ethanol fuels are gasohol, which contains 10% ethanol, and E-85, which contains 85% ethanol. Ethanol is now routinely prepared from the carbohydrates in corn (Figure 9.4). Starch, a complex carbohydrate polymer, can be hydrolyzed to the simple sugar glucose, which forms ethanol by the process of fermentation. Combining ethanol with gasoline forms a usable fuel, which combusts to form CO_2 , H_2O , and a great deal of energy.

Since green plants use sunlight to convert CO_2 and H_2O to carbohydrates during photosynthesis, next year's corn crop removes CO_2 from the atmosphere to make new molecules of starch as the corn grows. While in this way ethanol is a renewable fuel source, the need for large-scale farm equipment and the heavy reliance on fertilizers and herbicides make ethanol expensive to produce. Moreover, many criticize the use of valuable farmland for an energy-producing crop rather than for food production. As a result, discussion continues on ethanol as an alternative to fossil fuels.


Figure 9.3
Some simple alcohols



CH_3OH




$(\text{CH}_3)_2\text{CHOH}$



$\text{HOCH}_2\text{CH}_2\text{OH}$

- **Methanol (CH_3OH)** is also called wood alcohol, because it can be obtained by heating wood at high temperatures in the absence of air. Methanol is extremely toxic because of the oxidation products formed when it is metabolized in the liver (Section 12.14). Ingestion of as little as 15 mL causes blindness, and 100 mL causes death.
- **2-Propanol [$(\text{CH}_3)_2\text{CHOH}$]** is the major component of rubbing alcohol. When rubbed on the skin it evaporates readily, producing a pleasant cooling sensation. Because it has weak antibacterial properties, 2-propanol is used to clean skin before minor surgery and to sterilize medical instruments.
- **Ethylene glycol ($\text{HOCH}_2\text{CH}_2\text{OH}$)** is the major component of antifreeze. It is readily prepared from ethylene oxide by reactions discussed in Section 9.15. It is sweet tasting but toxic.

Figure 9.4 Ethanol from corn, a renewable fuel source

- Hydrolysis of starch and fermentation of the resulting simple sugars (Step [1]) yield ethanol, which is mixed with hydrocarbons from petroleum refining (Step [2]) to form usable fuels.
- Combustion of this ethanol–hydrocarbon fuel forms CO_2 and releases a great deal of energy (Step [3]).
- Photosynthesis converts atmospheric CO_2 back to plant carbohydrates in Step [4], and the cycle continues.

9.5B Interesting Ethers




This painting by Robert Hinckley depicts a public demonstration of the use of diethyl ether as an anesthetic at the Massachusetts General Hospital in Boston in the 1840s.

The discovery that **diethyl ether** ($\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$) is a general anesthetic revolutionized surgery in the nineteenth century. For years, a heated controversy existed over who first discovered diethyl ether's anesthetic properties and recognized the enormous benefit in its use. Early experiments were performed by a dentist, Dr. William Morton, resulting in a public demonstration of diethyl ether as an anesthetic in Boston in 1846. In fact, Dr. Crawford Long, a Georgia physician, had been using diethyl ether in surgery and obstetrics for several years, but had not presented his findings to a broader audience.

Diethyl ether is an imperfect anesthetic, but considering the alternatives in the nineteenth century, it was a miracle drug. It is safe, easy to administer, and causes little patient mortality, but it is highly flammable and causes nausea in many patients. For these reasons, it has largely been replaced by halothane (Figure 7.4), which is non-flammable and causes little patient discomfort.


Recall from Section 3.7B that some cyclic **polyethers**—compounds with two or more ether linkages—contain cavities that can complex specific-sized cations. For example, 18-crown-6 binds K^+ , whereas 12-crown-4 binds Li^+ .



- A crown ether–cation complex is called a *host–guest* complex. The crown ether is the *host* and the cation is the *guest*.
- The ability of a host molecule to bind specific guests is called *molecular recognition*.

Figure 9.5 KCN is insoluble in nonpolar solvents alone, but with 18-crown-6:

The use of crown ethers in nucleophilic substitution reactions



Recall from Section 3.7B that crown ethers are named as **x-crown-y**, where **x** is the total number of atoms in the ring and **y** is the number of O atoms.



The ability of crown ethers to complex cations can be exploited in nucleophilic substitution reactions, as shown in Figure 9.5. Nucleophilic substitution reactions are usually run in polar solvents to dissolve both the polar organic substrate and the ionic nucleophile. With a crown ether, though, the reaction can be run in a nonpolar solvent under conditions that enhance nucleophilicity.

When 18-crown-6 is added to the reaction of $\text{CH}_3\text{CH}_2\text{Br}$ with KCN, for example, the crown ether forms a tight complex with K^+ that has nonpolar C–H bonds on the outside, making the complex soluble in nonpolar solvents like benzene (C_6H_6) or hexane. When the crown ether/ K^+ complex dissolves in the nonpolar solvent, it carries the CN^- along with it to maintain electrical neutrality. The result is a solution of tightly complexed cation and relatively unsolvated anion (nucleophile). The anion, therefore, is extremely nucleophilic because it is not hidden from the substrate by solvent molecules.

Problem 9.10 Which mechanism is favored by the use of crown ethers in nonpolar solvents, $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$?

9.5C Interesting Epoxides


Although epoxides occur less widely in natural products than alcohols or ethers, interesting and useful epoxides are also known. As an example, two recently introduced drugs that contain an epoxide are eplerenone and tiotropium bromide. Eplerenone (trade name Inspra) is prescribed to reduce cardiovascular risk in patients who have already had a heart attack. Tiotropium bromide (trade name Spiriva) is a long-acting bronchodilator used to treat the chronic obstructive pulmonary disease of smokers and those routinely exposed to secondhand smoke.



Problem 9.11 Predict the solubility of eplerenone and tiotropium bromide in water and organic solvents.


9.6 Preparation of Alcohols, Ethers, and Epoxides

Alcohols and ethers are both common products of nucleophilic substitution. They are synthesized from alkyl halides by $\text{S}_{\text{N}}2$ reactions using strong nucleophiles. As in all $\text{S}_{\text{N}}2$ reactions, highest yields of products are obtained with unhindered methyl and 1° alkyl halides.




The preparation of ethers by this method is called the **Williamson ether synthesis**, and, although it was first reported in the 1800s, it is still the most general method to prepare an ether. Unsymmetrical ethers can be synthesized in two different ways, but often one path is preferred.


For example, isopropyl methyl ether can be prepared from CH_3O^- and 2-bromopropane (Path [a]), or from $(\text{CH}_3)_2\text{CHO}^-$ and bromomethane (Path [b]). Because the mechanism is S_N2 , **the preferred path uses the less sterically hindered halide, CH_3Br —Path [b].**



Problem 9.12 Draw the organic product of each reaction and classify the product as an alcohol, symmetrical ether, or unsymmetrical ether.




Problem 9.13 Draw two different routes to each ether and state which route, if any, is preferred.




NaH is an especially good base for forming an alkoxide, because the by-product of the reaction, H_2 , is a gas that just bubbles out of the reaction mixture.

A **hydroxide** nucleophile is needed to synthesize an alcohol, and salts such as **NaOH** and **KOH** are inexpensive and commercially available. An **alkoxide** salt is needed to make an ether. Simple alkoxides such as sodium methoxide (NaOCH_3) can be purchased, but others are prepared from alcohols by a Brønsted–Lowry acid–base reaction. For example, **sodium ethoxide ($\text{NaOCH}_2\text{CH}_3$)** is prepared by treating ethanol with NaH .

An acid–base reaction




Sample Problem 9.2 Draw the product of the following two-step reaction sequence.




Solution

[1] The base removes a proton from the OH group, forming an alkoxide.




[2] The alkoxide acts as a nucleophile in an $\text{S}_{\text{N}}2$ reaction, forming an ether.




- This two-step sequence converts an alcohol to an ether.

When an organic compound contains both a hydroxy group and a halogen atom on adjacent carbon atoms, an *intramolecular* version of this reaction forms an epoxide. The starting material for this two-step sequence, a **halohydrin**, is prepared from an alkene, as we will learn in Chapter 10.

Epoxide synthesis—A two-step procedure



Problem 9.14 Draw the products of each reaction.




9.7 General Features—Reactions of Alcohols, Ethers, and Epoxides

We begin our discussion of the chemical reactions of alcohols, ethers, and epoxides with a look at the general reactive features of each functional group.


9.7A Alcohols

Unlike many families of molecules, the reactions of alcohols do *not* fit neatly into a single reaction class. In Chapter 9, we discuss only the substitution and β elimination reactions of alcohols. Alcohols are also key starting materials in oxidation reactions (Chapter 12), and their polar O–H bond makes them more acidic than many other organic compounds, a feature we will explore in Chapter 19.

Alcohols are similar to alkyl halides in that both contain an electronegative element bonded to an sp^3 hybridized carbon atom. Alkyl halides contain a good leaving group (X^-), however, whereas alcohols do not. Nucleophilic substitution with ROH as starting material would displace ^-OH , a **strong base and therefore a poor leaving group**.




For an alcohol to undergo a nucleophilic substitution or elimination reaction, the **OH group must be converted into a better leaving group**. This can be done by reaction with acid. Treatment of an alcohol with a strong acid like HCl or H_2SO_4 protonates the O atom via an acid–base reaction. This transforms the ^-OH leaving group into H_2O , a **weak base and therefore a good leaving group**.




If the OH group of an alcohol is made into a good leaving group, alcohols *can* undergo β elimination and nucleophilic substitution, as described in Sections 9.8–9.12.

Because the pK_a of $(\text{ROH}_2)^+$ is ~ -2 , protonation of an alcohol occurs only with very strong acids—namely, those having a $pK_a \leq -2$.




9.7B Ethers and Epoxides

Like alcohols, **ethers do not contain a good leaving group**, which means that nucleophilic substitution and β elimination do not occur directly. Ethers undergo fewer useful reactions than alcohols.




Epoxides don't have a good leaving group either, but they have one characteristic that neither alcohols nor ethers have: **the "leaving group" is contained in a strained three-membered ring**. Nucleophilic attack opens the three-membered ring and relieves angle strain, making nucleophilic attack a favorable process that occurs even with the poor leaving group. Specific examples are presented in Section 9.15.



9.8 Dehydration of Alcohols to Alkenes

The dehydrohalogenation of alkyl halides, discussed in Chapter 8, is one way to introduce a π bond into a molecule. Another way is to eliminate water from an alcohol in a **dehydration**

reaction. **Dehydration, like dehydrohalogenation, is a β elimination reaction in which the elements of OH and H are removed from the α and β carbon atoms, respectively.**





Dehydration is typically carried out using H_2SO_4 and other strong acids, or phosphorus oxychloride (POCl_3) in the presence of an amine base. We consider dehydration in acid first, followed by dehydration with POCl_3 in Section 9.10.


9.8A General Features of Dehydration in Acid

Alcohols undergo dehydration in the presence of strong acid to afford alkenes, as illustrated in Equations [1] and [2]. Typical acids used for this conversion are H_2SO_4 or *p*-toluenesulfonic acid (abbreviated as TsOH).


Recall from Section 2.6 that *p*-toluenesulfonic acid is a strong organic acid ($\text{p}K_a = -7$).



More substituted alcohols dehydrate more readily, giving rise to the following order of reactivity:




When an alcohol has two or three different β carbons, dehydration is regioselective and follows the Zaitsev rule. **The more substituted alkene is the major product when a mixture of constitutional isomers is possible.** For example, elimination of H and OH from 2-methyl-2-butanol yields two constitutional isomers: the trisubstituted alkene **A** as major product and the disubstituted alkene **B** as minor product.




Problem 9.15

Draw the products formed when each alcohol undergoes dehydration with TsOH, and label the major product when a mixture results.



Problem 9.16 Rank the alcohols in each group in order of increasing reactivity when dehydrated with H_2SO_4 .



9.8B The E1 Mechanism for the Dehydration of 2° and 3° Alcohols


The mechanism of dehydration depends on the structure of the alcohol: **2° and 3° alcohols react by an E1 mechanism, whereas 1° alcohols react by an E2 mechanism.** Regardless of the type of alcohol, however, strong acid is *always* needed to protonate the O atom to form a good leaving group.

The E1 dehydration of 2° and 3° alcohols is illustrated with $(\text{CH}_3)_3\text{COH}$ (a 3° alcohol) as starting material to form $(\text{CH}_3)_2\text{C}=\text{CH}_2$ as product (Mechanism 9.1). The mechanism consists of **three steps**.




Mechanism 9.1 Dehydration of 2° and 3° ROH—An E1 Mechanism

Step [1] The O atom is protonated.




- **Protonation of the oxygen atom** of the alcohol converts a poor leaving group (^-OH) into a good leaving group (H_2O).

Step [2] The C–O bond is broken.



- **Heterolysis of the C–O bond forms a carbocation.** This step is rate-determining because it involves only bond cleavage.

Step [3] A C–H bond is cleaved and the π bond is formed.



- **A base (such as HSO_4^- or H_2O) removes a proton** from a carbon adjacent to the carbocation (a β carbon). The electron pair in the C–H bond is used to form the new π bond.

Thus, **dehydration of 2° and 3° alcohols occurs via an E1 mechanism with an added first step.**

- Step [1] protonates the OH group to make a good leaving group.
- Steps [2] and [3] are the two steps of an E1 mechanism: loss of a leaving group (H_2O in this case) to form a carbocation, followed by removal of a β proton to form a π bond.
- The acid used to protonate the alcohol in Step [1] is regenerated upon removal of the proton in Step [3], so dehydration is **acid-catalyzed**.

The E1 dehydration of 2° and 3° alcohols with acid gives clean elimination products without by-products formed from an $\text{S}_{\text{N}}1$ reaction. This makes the E1 dehydration of alcohols much more synthetically useful than the E1 dehydrohalogenation of alkyl halides (Section 8.7). Clean elimination takes place because the reaction mixture contains no good nucleophile to react with the intermediate carbocation, so no competing $\text{S}_{\text{N}}1$ reaction occurs.

Problem 9.17

Draw the structure of each transition state in the three-step mechanism for the reaction,
 $(\text{CH}_3)_3\text{COH} + \text{H}_2\text{SO}_4 \rightarrow (\text{CH}_3)_2\text{C}=\text{CH}_2 + \text{H}_2\text{O}$.

9.8C The E2 Mechanism for the Dehydration of 1° Alcohols

Because 1° carbocations are highly unstable, the dehydration of 1° alcohols cannot occur by an E1 mechanism involving a carbocation intermediate. With 1° alcohols, therefore, **dehydration follows an E2 mechanism**. This two-step process for the conversion of CH₃CH₂CH₂OH (a 1° alcohol) to CH₃CH=CH₂ with H₂SO₄ as acid catalyst is shown in Mechanism 9.2.




Mechanism 9.2 Dehydration of a 1° ROH—An E2 Mechanism

Step [1] The O atom is protonated.



- **Protonation of the oxygen atom** of the alcohol converts a poor leaving group (⁻OH) into a good leaving group (H₂O).

Step [2] The C–H and C–O bonds are broken and the π bond is formed.



- **Two bonds are broken and two bonds are formed in a single step:** the base (HSO₄⁻ or H₂O) removes a proton from the β carbon; the electron pair in the β C–H bond forms the new π bond; the leaving group (H₂O) comes off with the electron pair in the C–O bond.

The dehydration of a 1° alcohol begins with the protonation of the OH group to form a good leaving group, just as in the dehydration of a 2° or 3° alcohol. With 1° alcohols, however, loss of the leaving group and removal of a β proton occur at the same time, so that no highly unstable 1° carbocation is generated.

Problem 9.18 Draw the structure of each transition state in the two-step mechanism for the reaction, CH₃CH₂CH₂OH + H₂SO₄ → CH₃CH=CH₂ + H₂O.

9.8D Le Châtelier's Principle

Although entropy favors product formation in dehydration (one molecule of reactant forms two molecules of products), enthalpy does not, because the two σ bonds broken in the reactant are stronger than the σ and π bonds formed in the products. For example, ΔH° for the dehydration of CH₃CH₂OH to CH₂=CH₂ is +38 kJ/mol (Figure 9.6).

Figure 9.6

The dehydration of CH₃CH₂OH to CH₂=CH₂—An endothermic reaction



ΔH° calculation:

[1] Bonds broken

	ΔH° (kJ/mol)
CH ₃ CH ₂ –OH	+393
HOCH ₂ CH ₂ –H	+410
Total	+803 kJ/mol

Energy needed to break bonds.

[2] Bonds formed

	ΔH° (kJ/mol)
CH ₂ =CH ₂ π bond	–267
H–OH	–498
Total	–765 kJ/mol

Energy released in forming bonds.

[3] Overall ΔH° =

sum in Step [1]
+
sum in Step [2]

+803 kJ/mol

–765 kJ/mol

ΔH° = +38 kJ/mol

The reaction is endothermic.

[Values taken from Appendix C.]


According to **Le Châtelier's principle**, a system at equilibrium will react to counteract any disturbance to the equilibrium. Thus, removing a product from a reaction mixture as it is formed drives the equilibrium to the right, forming more product.

Le Châtelier's principle can be used to favor products in dehydration reactions because the alkene product has a lower boiling point than the alcohol reactant. Thus, the alkene can be distilled from the reaction mixture as it is formed, leaving the alcohol and acid to react further, forming more product.

9.9 Carbocation Rearrangements

Sometimes “unexpected” products are formed in dehydration; that is, the carbon skeletons of the starting material and product might be different, or the double bond might be in an unexpected location. For example, the dehydration of 3,3-dimethyl-2-butanol yields two alkenes, whose carbon skeletons do not match the carbon framework of the starting material.


The carbon skeletons of the reactant and products are different.



This phenomenon sometimes occurs when carbocations are reactive intermediates. A **less stable carbocation can rearrange to a more stable carbocation by shift of a hydrogen atom or an alkyl group**. These rearrangements are called **1,2-shifts**, because they involve migration of an alkyl group or hydrogen atom from one carbon to an adjacent carbon atom. The migrating group moves with the two electrons that bonded it to the carbon skeleton.

Because the migrating group in a 1,2-shift moves with two bonding electrons, the carbon it leaves behind now has only three bonds (six electrons), giving it a net positive (+) charge.

Carbocation rearrangement




- Movement of a hydrogen atom is called a **1,2-hydride shift**.
- Movement of an alkyl group is called a **1,2-alkyl shift**.

The dehydration of 3,3-dimethyl-2-butanol illustrates the rearrangement of a 2° to a 3° carbocation by a **1,2-methyl shift**, as shown in Mechanism 9.3. The carbocation rearrangement occurs in Step [3] of the four-step mechanism.

Steps [1], [2], and [4] in the mechanism for the dehydration of 3,3-dimethyl-2-butanol are exactly the same steps previously seen in dehydration: protonation, loss of H₂O, and loss of a proton. Only Step [3], rearrangement of the less stable 2° carbocation to the more stable 3° carbocation, is new.

- **1,2-Shifts convert a less stable carbocation to a more stable carbocation.**

For example, 2° carbocation **A** rearranges to the more stable 3° carbocation by a 1,2-hydride shift, whereas carbocation **B** does not rearrange because it is 3° to begin with.




Sample Problem 9.3 illustrates a dehydration reaction that occurs with a **1,2-hydride shift**.




Mechanism 9.3 A 1,2-Methyl Shift—Carbocation Rearrangement During Dehydration

Steps [1] and [2] Formation of a 2° carbocation




- **Protonation of the oxygen atom** on the alcohol in Step [1] forms a good leaving group (H₂O), and loss of H₂O in Step [2] forms a 2° carbocation.

Step [3] Rearrangement of the carbocation by a 1,2-CH₃ shift




- **1,2-Shift** of a CH₃ group from one carbon to the adjacent carbon converts the 2° carbocation to a more stable 3° carbocation.

Step [4] Loss of a proton to form the π bond




- **Loss of a proton** from a β carbon (labeled β₁ and β₂) forms two different alkenes.


Sample Problem 9.3 Show how the dehydration of alcohol **X** forms alkene **Y** using a 1,2-hydride shift.



Solution

Steps [1] and [2] Protonation of **X** and loss of H₂O form a 2° carbocation.






Problem 9.19 What other alkene is also formed along with **Y** in Sample Problem 9.3? What alkenes would form from **X** if no carbocation rearrangement occurred?

Rearrangements are not unique to dehydration reactions. **Rearrangements can occur whenever a carbocation is formed as reactive intermediate**, meaning any $\text{S}_{\text{N}}1$ or $\text{E}1$ reaction. In fact, the formation of rearranged products often indicates the presence of a carbocation intermediate.



Problem 9.20 Show how a 1,2-shift forms a more stable carbocation from each intermediate.



Problem 9.21 Explain how the reaction of $(\text{CH}_3)_2\text{CHCH}(\text{Cl})\text{CH}_3$ with H_2O yields two substitution products, $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_3$ and $(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CH}_3$.

9.10 Dehydration Using POCl_3 and Pyridine

Because some organic compounds decompose in the presence of strong acid, other methods that avoid strong acid have been developed to convert alcohols to alkenes. A common method uses phosphorus oxychloride (POCl_3) and pyridine (an amine base) in place of H_2SO_4 or TsOH . For example, the treatment of cyclohexanol with POCl_3 and pyridine forms cyclohexene in good yield.



POCl_3 serves much the same role as strong acid does in acid-catalyzed dehydration. **It converts a poor leaving group (OH) into a good leaving group.** Dehydration then proceeds by an **$\text{E}2$ mechanism**, as shown in Mechanism 9.4. Pyridine is the base that removes a β proton during elimination.


- Steps [1] and [2] of the mechanism convert the OH group to a good leaving group.
- In Step [3], the $\text{C}-\text{H}$ and $\text{C}-\text{O}$ bonds are broken and the π bond is formed.
- No rearrangements occur during dehydration with POCl_3 , suggesting that carbocations are not formed as intermediates in this reaction.

We have now learned about two different reagents for alcohol dehydration—strong acid (H_2SO_4 or TsOH) and POCl_3 + pyridine. The best dehydration method for a given alcohol is often hard to know ahead of time, and this is why organic chemists develop more than one method for a given type of transformation. Two examples of dehydration reactions used in the synthesis of natural products are given in Figure 9.7.




Mechanism 9.4 Dehydration Using POCl_3 + Pyridine—An E2 Mechanism

Steps [1] and [2] Conversion of OH to a good leaving group



- A two-step process converts an OH group into OPOCl_2 , a **good leaving group**: reaction of the OH group with POCl_3 followed by removal of a proton.

Step [3] The C–H and C–O bonds are broken and the π bond is formed.




- **Two bonds are broken and two bonds are formed in a single step**: the base (pyridine) removes a proton from the β carbon; the electron pair in the β C–H bond forms the new π bond; the leaving group (OPOCl_2) comes off with the electron pair from the C–O bond.

Figure 9.7


Dehydration reactions in the synthesis of two natural products

Patchouli alcohol, obtained from the patchouli plant native to Malaysia, has been used in perfumery because of its exotic fragrance. In the 1800s shawls imported from India were often packed with patchouli leaves to ward off insects, thus permeating the clothing with the distinctive odor.





9.11 Conversion of Alcohols to Alkyl Halides with HX


Alcohols undergo nucleophilic substitution reactions only if the OH group is converted into a better leaving group before nucleophilic attack. Thus, substitution does *not* occur when an alcohol is treated with X^- because OH^- is a poor leaving group (Reaction [1]), but substitution *does* occur on treatment of an alcohol with HX because H_2O is now the leaving group (Reaction [2]).




The reaction of alcohols with HX (X = Cl, Br, I) is a general method to prepare 1°, 2°, and 3° alkyl halides.



More substituted alcohols usually react more rapidly with HX:




Problem 9.22 Draw the products of each reaction.



9.11A Two Mechanisms for the Reaction of ROH with HX

How does the reaction of ROH with HX occur? Acid–base reactions are very fast, so the strong acid HX protonates the OH group of the alcohol, forming a **good leaving group** (H_2O) and a **good nucleophile** (the conjugate base, X^-). Both components are needed for nucleophilic substitution. The mechanism of substitution of X^- for H_2O then depends on the structure of the R group.

Whenever there is an oxygen-containing reactant and a strong acid, the first step in the mechanism is protonation of the oxygen atom.




- Methyl and 1° ROH form RX by an $\text{S}_{\text{N}}2$ mechanism.
- Secondary (2°) and 3° ROH form RX by an $\text{S}_{\text{N}}1$ mechanism.

The reaction of $\text{CH}_3\text{CH}_2\text{OH}$ with HBr illustrates the $\text{S}_{\text{N}}2$ mechanism of a 1° alcohol (Mechanism 9.5). Nucleophilic attack on the protonated alcohol occurs in one step: **the bond to the nucleophile X^- is formed as the bond to the leaving group (H_2O) is broken.**




Mechanism 9.5 Reaction of a 1° ROH with HX—An S_N2 Mechanism

Step [1] The O atom is protonated.



Step [2] The C–O bond is broken as the C–Br bond is formed.




The reaction of (CH₃)₃COH with HBr illustrates the S_N1 mechanism of a 3° alcohol (Mechanism 9.6). Nucleophilic attack on the protonated alcohol occurs in two steps: **the bond to the leaving group (H₂O) is broken before the bond to the nucleophile X[−] is formed.**




Mechanism 9.6 Reaction of 2° and 3° ROH with HX—An S_N1 Mechanism

Step [1] The O atom is protonated.




Steps [2] and [3] The C–O bond is broken, and then the C–Br bond is formed.




Both mechanisms begin with the same first step—protonation of the O atom to form a good leaving group—and both mechanisms give an alkyl halide (RX) as product. The mechanisms differ only in the timing of bond breaking and bond making.



The reactivity of hydrogen halides increases with increasing acidity:



Because Cl[−] is a poorer nucleophile than Br[−] or I[−], the reaction of 1° alcohols with HCl occurs only when an additional Lewis acid catalyst, usually ZnCl₂, is added. ZnCl₂ complexes with the O atom of the alcohol in a Lewis acid–base reaction, making an especially good leaving group and facilitating the S_N2 reaction.




Knowing the mechanism allows us to predict the stereochemistry of the products when reaction occurs at a stereogenic center.



- The 1° alcohol **A** reacts with HBr via an S_N2 mechanism to yield the alkyl bromide **B** with **inversion** of stereochemistry at the stereogenic center.
- The 3° alcohol **C** reacts with HCl via an S_N1 mechanism to yield a **racemic mixture** of alkyl chlorides **D** and **E**, because a trigonal planar carbocation intermediate is formed.


Problem 9.23 Draw the products of each reaction, indicating the stereochemistry around any stereogenic centers.



9.11B Carbocation Rearrangement in the S_N1 Reaction

Because carbocations are formed in the S_N1 reaction of 2° and 3° alcohols with HX, **carbocation rearrangements are possible**, as illustrated in Sample Problem 9.4.


Sample Problem 9.4 Draw a stepwise mechanism for the following reaction.




Solution

A 2° alcohol reacts with HBr by an S_N1 mechanism. Because substitution converts a 2° alcohol to a 3° alkyl halide in this example, a carbocation rearrangement must occur.




Steps [1] and [2] Protonation of the O atom and then loss of H₂O form a 2° carbocation.



Steps [3] and [4] **Rearrangement of the 2° carbocation by a 1,2-hydride shift forms a more stable 3° carbocation.** Nucleophilic attack forms the substitution product.



Problem 9.24 What is the major product formed when each alcohol is treated with HCl ?



9.12 Conversion of Alcohols to Alkyl Halides with SOCl_2 and PBr_3


Primary (1°) and 2° alcohols can be converted to alkyl halides using SOCl_2 and PBr_3 .

- SOCl_2 (thionyl chloride) converts alcohols into alkyl chlorides.
- PBr_3 (phosphorus tribromide) converts alcohols into alkyl bromides.

Both reagents convert ^-OH into a good leaving group *in situ*—that is, directly in the reaction mixture—as well as provide the **nucleophile**, either Cl^- or Br^- , to displace the leaving group.

9.12A Reaction of ROH with SOCl_2

The treatment of a 1° or 2° alcohol with thionyl chloride, SOCl_2 , and pyridine forms an alkyl chloride, with SO_2 and HCl as by-products.




The mechanism for this reaction consists of two parts: **conversion of the OH group into a better leaving group, and nucleophilic attack by Cl^- via an $\text{S}_{\text{N}}2$ reaction**, as shown in Mechanism 9.7.

Problem 9.25 If the reaction of an alcohol with SOCl_2 and pyridine follows an $\text{S}_{\text{N}}2$ mechanism, what is the stereochemistry of the alkyl chloride formed from (2*R*)-2-butanol?




Mechanism 9.7 Reaction of ROH with SOCl₂ + Pyridine—An S_N2 Mechanism

Steps [1] and [2] The OH group is converted into a good leaving group.



- Reaction of the alcohol with SOCl₂ forms an intermediate that loses a proton by reaction with pyridine in Step [2]. This two-step process converts the OH group into OSOCl, a **good leaving group**, and also generates the **nucleophile (Cl⁻)** needed for Step [3].

Step [3] The C–O bond is broken as the C–Cl bond is formed.



- **Nucleophilic attack of Cl⁻ and loss of the leaving group (SO₂ + Cl⁻)** occur in a single step.



9.12B Reaction of ROH with PBr₃

In a similar fashion, the treatment of a 1° or 2° alcohol with phosphorus tribromide, PBr₃, forms an alkyl bromide.

General reaction



Examples




1° and 2° RBr are formed.

The mechanism for this reaction also consists of two parts: **conversion of the OH group into a better leaving group, and nucleophilic attack by Br⁻ via an S_N2 reaction**, as shown in Mechanism 9.8.



Mechanism 9.8 Reaction of ROH with PBr₃—An S_N2 Mechanism

Step [1] The OH group is converted into a good leaving group.



- Reaction of the alcohol with PBr₃ converts the OH group into a **better leaving group**, and also generates the **nucleophile (Br⁻)** needed for Step [2].

Step [2] The C–O bond is broken as the C–Br bond is formed.



- **Nucleophilic attack of Br⁻ and loss of the leaving group (HOPBr₂)** occur in a single step.


Table 9.2 summarizes the methods for converting an alcohol to an alkyl halide presented in Sections 9.11 and 9.12.

Table 9.2 Summary of Methods for $\text{ROH} \rightarrow \text{RX}$

Overall reaction	Reagent	Comment
$\text{ROH} \rightarrow \text{RCl}$	HCl	<ul style="list-style-type: none"> Useful for all ROH An $\text{S}_{\text{N}}1$ mechanism for 2° and 3° ROH; an $\text{S}_{\text{N}}2$ mechanism for CH_3OH and 1° ROH
	SOCl_2	<ul style="list-style-type: none"> Best for CH_3OH, and 1° and 2° ROH An $\text{S}_{\text{N}}2$ mechanism
$\text{ROH} \rightarrow \text{RBr}$	HBr	<ul style="list-style-type: none"> Useful for all ROH An $\text{S}_{\text{N}}1$ mechanism for 2° and 3° ROH; an $\text{S}_{\text{N}}2$ mechanism for CH_3OH and 1° ROH
	PBr_3	<ul style="list-style-type: none"> Best for CH_3OH, and 1° and 2° ROH An $\text{S}_{\text{N}}2$ mechanism
$\text{ROH} \rightarrow \text{RI}$	HI	<ul style="list-style-type: none"> Useful for all ROH An $\text{S}_{\text{N}}1$ mechanism for 2° and 3° ROH; an $\text{S}_{\text{N}}2$ mechanism for CH_3OH and 1° ROH

Problem 9.26 If the reaction of an alcohol with PBr_3 follows an $\text{S}_{\text{N}}2$ mechanism, what is the stereochemistry of the alkyl bromide formed from (2*R*)-2-butanol?

Problem 9.27 Draw the organic products formed in each reaction, and indicate the stereochemistry of products that contain stereogenic centers.




9.12C The Importance of Making RX from ROH

We have now learned two methods to prepare an alkyl chloride and two methods to prepare an alkyl bromide from an alcohol. If there is one good way to carry out a reaction, why search for more? A particular reagent might work well for one starting material, but not so well for another. Thus, organic chemists try to devise several different ways to perform the same overall reaction. For now, though, concentrate on one or two of the most general methods, so you can better understand the underlying concepts.


Why are there so many ways to convert an alcohol to an alkyl halide? Alkyl halides are versatile starting materials in organic synthesis, as shown in Sample Problem 9.5.

Sample Problem 9.5 Convert 1-propanol to butanenitrile (**A**).




Solution

Direct conversion of 1-propanol to **A** using CN^- as a nucleophile is not possible because OH^- is a poor leaving group. However, conversion of the OH group to a Br atom forms a good leaving group, which can then readily undergo an $\text{S}_{\text{N}}2$ reaction with CN^- to yield **A**. **This two-step sequence is our first example of a multistep synthesis.**



Problem 9.29 Draw the products of each reaction, and indicate the stereochemistry at any stereogenic centers.




9.13B Reactions of Alkyl Tosylates


Because alkyl tosylates have good leaving groups, **they undergo both nucleophilic substitution and β elimination**, exactly as alkyl halides do. Generally, alkyl tosylates are treated with strong nucleophiles and bases, so that the mechanism of substitution is $\text{S}_{\text{N}}2$ and the mechanism of elimination is **E2**.

For example, ethyl tosylate, which has the leaving group on a 1° carbon, reacts with NaOCH_3 to yield ethyl methyl ether, the product of nucleophilic substitution by an $\text{S}_{\text{N}}2$ mechanism. It reacts with $\text{KOC}(\text{CH}_3)_3$, a strong bulky base, to yield ethylene by an E2 mechanism.


Two reactions of ethyl tosylate



Because substitution occurs via an $\text{S}_{\text{N}}2$ mechanism, **inversion of configuration** results when the leaving group is bonded to a stereogenic center.




Sample Problem 9.6 Draw the product of the following reaction, including stereochemistry.




Solution

The 1° alkyl tosylate and the strong nucleophile both favor substitution by an S_N2 mechanism, which proceeds by backside attack, resulting in **inversion** of configuration at the stereogenic center.




Problem 9.30 Draw the products of each reaction, and include the stereochemistry at any stereogenic centers in the products.



9.13C The Two-Step Conversion of an Alcohol to a Substitution Product


We now have another **two-step method to convert an alcohol to a substitution product**: reaction of an alcohol with TsCl and pyridine to form an alkyl tosylate (Step [1]), followed by nucleophilic attack on the tosylate (Step [2]).



Let's look at the stereochemistry of this two-step process.

- Step [1], formation of the tosylate, proceeds with **retention** of configuration at a stereogenic center because the C–O bond remains intact.
- Step [2] is an S_N2 reaction, so it proceeds with **inversion of configuration** because the nucleophile attacks from the back side.
- Overall there is a **net inversion of configuration** at a stereogenic center.


For example, the treatment of *cis*-3-methylcyclohexanol with *p*-toluenesulfonyl chloride and pyridine forms a *cis* tosylate **A**, which undergoes backside attack by the nucleophile OCH_3^- to yield the *trans* ether **B**.



Problem 9.31 Draw the products formed when (2*S*)-2-butanol is treated with TsCl and pyridine, followed by NaOH. Label the stereogenic center in each compound as *R* or *S*. What is the stereochemical relationship between the starting alcohol and the final product?

Figure 9.8

Summary: Nucleophilic substitution and β elimination reactions of alcohols



9.13D A Summary of Substitution and Elimination Reactions of Alcohols

The reactions of alcohols in Sections 9.8–9.13C share two similarities:

- The OH group is converted into a better leaving group by treatment with acid or another reagent.
- The resulting product undergoes either elimination or substitution, depending on the reaction conditions.

Figure 9.8 summarizes these reactions with cyclohexanol as starting material.

Problem 9.32



Draw the product formed when $(\text{CH}_3)_2\text{CHOH}$ is treated with each reagent.

- SOCl_2 , pyridine
- TsCl , pyridine
- H_2SO_4
- HBr
- PBr_3 , then NaCN
- POCl_3 , pyridine

9.14 Reaction of Ethers with Strong Acid

Because ethers are so unreactive, diethyl ether and tetrahydrofuran (THF) are often used as solvents for organic reactions.

Recall from Section 9.7B that ethers have a poor leaving group, so they cannot undergo nucleophilic substitution or β elimination reactions directly. Instead, they must first be converted to a good leaving group by reaction with strong acids. Only **HBr** and **HI** can be used, though, because they are strong acids that are also sources of good nucleophiles (Br^- and I^- , respectively). **When ethers react with HBr or HI, both C–O bonds are cleaved and two alkyl halides are formed as products.**



HBr or HI serves as a strong acid that both protonates the O atom of the ether and is the source of a good nucleophile (Br^- or I^-). Because both C–O bonds in the ether are broken, **two successive nucleophilic substitution reactions occur.**


- The mechanism of ether cleavage is $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$, depending on the identity of R.
- With 2° or 3° alkyl groups bonded to the ether oxygen, the C–O bond is cleaved by an $\text{S}_{\text{N}}1$ mechanism involving a carbocation; with methyl or 1° R groups, the C–O bond is cleaved by an $\text{S}_{\text{N}}2$ mechanism.

For example, cleavage of $(\text{CH}_3)_3\text{COCH}_3$ with HI occurs at two bonds, as shown in Mechanism 9.9. The 3° alkyl group undergoes nucleophilic substitution by an $\text{S}_\text{N}1$ mechanism, resulting in the cleavage of one C–O bond. The methyl group undergoes nucleophilic substitution by an $\text{S}_\text{N}2$ mechanism, resulting in the cleavage of the second C–O bond.




Mechanism 9.9 Mechanism of Ether Cleavage in Strong Acid— $(\text{CH}_3)_3\text{COCH}_3 + \text{HI} \rightarrow (\text{CH}_3)_3\text{CI} + \text{CH}_3\text{I} + \text{H}_2\text{O}$

Part [1] Cleavage of the 3° C–O bond by an $\text{S}_\text{N}1$ mechanism




- **Protonation** of the O atom forms a good leaving group in Step [1]. Cleavage of the C–O bond then occurs in two steps: the bond to the leaving group is broken to form a **carbocation**, and then the bond to the nucleophile (I^-) is formed. This generates one of the alkyl iodides, **$(\text{CH}_3)_3\text{CI}$** .

Part [2] Cleavage of the CH_3 –O bond by an $\text{S}_\text{N}2$ mechanism






- **Protonation of the OH group** forms a good leaving group (H_2O), and then nucleophilic attack by I^- forms the second alkyl iodide, **CH_3I** , and **H_2O** .




The mechanism illustrates the central role of HX in the reaction:

- HX protonates the ether oxygen, thus making a good leaving group.
- HX provides a source of X^- for nucleophilic attack.

Problem 9.33 What alkyl halides are formed when each ether is treated with HBr?




Problem 9.34 Explain why the treatment of anisole with HBr yields phenol and CH₃Br, but not bromobenzene.




9.15 Reactions of Epoxides

Although epoxides do not contain a good leaving group, they contain a strained three-membered ring with two polar bonds. **Nucleophilic attack opens the strained three-membered ring**, making it a favorable process even with the poor leaving group.




This reaction occurs readily with strong nucleophiles, and with acids like HZ, where Z is a nucleophilic atom.



Problem 9.35 Explain why cyclopropane, which has a strained three-membered ring like an epoxide, does not react readily with nucleophiles.

9.15A Opening of Epoxide Rings with Strong Nucleophiles


Virtually all strong nucleophiles open an epoxide ring by a two-step reaction sequence:



- **Step [1]:** The nucleophile attacks an electron-deficient carbon of the epoxide, thus cleaving a C–O bond and relieving the strain of the three-membered ring.
- **Step [2]:** Protonation of the alkoxide with water generates a neutral product with two functional groups on adjacent atoms.


Common nucleophiles that open epoxide rings include [−]OH, [−]OR, [−]CN, [−]SR, and NH₃. With these strong nucleophiles, the reaction occurs via an **S_N2 mechanism**, resulting in two consequences:

- The nucleophile opens the epoxide ring from the back side.




Other examples of the nucleophilic opening of epoxide rings are presented in Sections 12.6 and 20.14.

- In an unsymmetrical epoxide, the nucleophile attacks at the less substituted carbon atom.




Problem 9.36


Draw the product of each reaction, indicating the stereochemistry at any stereogenic centers.



1,2-Epoxycyclohexane is a symmetrical epoxide that is achiral because it possesses a plane of symmetry. It reacts with ⁻OCH₃, however, to yield two *trans*-1,2-disubstituted cyclohexanes, **A** and **B**, which are **enantiomers**; each has two stereogenic centers.



In this case, **nucleophilic attack of ⁻OCH₃ occurs from the back side at either C–O bond, because both ends are equally substituted**. Because attack at either side occurs with equal probability, an equal amount of the two enantiomers is formed—a **racemic mixture**. This is a specific example of a general rule concerning the stereochemistry of products obtained from an achiral reactant.




- Whenever an achiral reactant yields a product with stereogenic centers, the product must be achiral (meso) or racemic.

This general rule can be restated in terms of optical activity. Recall from Section 5.12 that achiral compounds and racemic mixtures are optically inactive.

- Optically inactive starting materials give optically inactive products.


Problem 9.37

The *cis* and *trans* isomers of 2,3-dimethyloxirane both react with OH^- to give 2,3-butanediol. One stereoisomer gives a single achiral product, and one gives two chiral enantiomers. Which epoxide gives one product and which gives two?



9.15B Reaction with Acids HZ


Acids **HZ** that contain a nucleophile **Z** also open epoxide rings by a two-step reaction sequence:



- **Step [1]:** Protonation of the epoxide oxygen with HZ makes the epoxide oxygen into a good leaving group (OH). It also provides a source of a good nucleophile (Z^-) to open the epoxide ring.
- **Step [2]:** The nucleophile Z^- then opens the protonated epoxide ring by **backside** attack.

These two steps—**protonation followed by nucleophilic attack**—are the exact reverse of the opening of epoxide rings with strong nucleophiles, where nucleophilic attack precedes protonation.

HCl, **HBr**, and **HI** all open an epoxide ring in this manner. **H₂O** and **ROH** can, too, but acid must also be added. Regardless of the reaction, the product has an OH group from the epoxide on one carbon and a new functional group **Z** from the nucleophile on the adjacent carbon. With epoxides fused to rings, ***trans*-1,2-disubstituted cycloalkanes** are formed.




Although backside attack of the nucleophile suggests that this reaction follows an $\text{S}_{\text{N}}2$ mechanism, the regioselectivity of the reaction with unsymmetrical epoxides does not.

- With unsymmetrical epoxides, nucleophilic attack occurs at the *more* substituted carbon atom.


Figure 9.9

Opening of an unsymmetrical epoxide ring with HCl



- Transition state **A** is lower in energy because the partial positive charge (δ^+) is located on the more substituted carbon. In this case, therefore, nucleophilic attack occurs from the back side (an S_N2 characteristic) at the more substituted carbon (an S_N1 characteristic).

For example, the treatment of 2,2-dimethyloxirane with HCl results in nucleophilic attack at the carbon with two methyl groups.



Backside attack of the nucleophile suggests an S_N2 mechanism, but attack at the more substituted carbon suggests an S_N1 mechanism. To explain these results, the **mechanism of nucleophilic attack is thought to be somewhere in between S_N1 and S_N2 .**

Figure 9.9 illustrates two possible pathways for the reaction of 2,2-dimethyloxirane with HCl. Backside attack of Cl^- at the more substituted carbon proceeds via transition state **A**, whereas backside attack of Cl^- at the less substituted carbon proceeds via transition state **B**. **Transition state A has a partial positive charge on a more substituted carbon, making it more stable.** Thus, the preferred reaction path takes place by way of the lower energy transition state **A**.

Opening of an epoxide ring with either a strong nucleophile $:\text{Nu}^-$ or an acid HZ is **regioselective**, because one constitutional isomer is the major or exclusive product. The **site selectivity of these two reactions, however, is exactly the opposite.**





Figure 9.10

The synthesis of two bronchodilators uses the opening of an epoxide ring.







- A key step in each synthesis is the opening of an epoxide ring with a nitrogen nucleophile to form a new C–N bond.

- With a strong nucleophile, :Nu^- attacks at the less substituted carbon.
- With an acid HZ , the nucleophile attacks at the more substituted carbon.

The reaction of epoxide rings with nucleophiles is important for the synthesis of many biologically active compounds, including **salmeterol** and **albuterol**, two bronchodilators used in the treatment of asthma (Figure 9.10).

Problem 9.38

Draw the product of each reaction.

**9.16 Application: Epoxides, Leukotrienes, and Asthma**


The opening of epoxide rings with nucleophiles is a key step in some important biological processes.

9.16A Asthma and Leukotrienes

Asthma is an obstructive lung disease that affects millions of Americans. Because it involves episodic constriction of small airways, bronchodilators such as albuterol (Figure 9.10) are used to treat symptoms by widening airways. Because asthma is also characterized by chronic inflammation, inhaled steroids that reduce inflammation are also commonly used.


The **leukotrienes** are molecules that contribute to the asthmatic response. A typical example, **leukotriene C₄**, is shown. Although its biological activity was first observed in the 1930s, the chemical structure of leukotriene C₄ was not determined until 1979. Structure determination and chemical synthesis were difficult because leukotrienes are highly unstable and extremely potent, and are therefore present in tissues in exceedingly small amounts.

Leukotrienes were first synthesized in 1980 in the laboratory of Professor E. J. Corey, the 1990 recipient of the Nobel Prize in Chemistry.




9.16B Leukotriene Synthesis and Asthma Drugs

Leukotrienes are synthesized in cells by the oxidation of **arachidonic acid** to 5-HPETE, which is then converted to an epoxide, **leukotriene A₄**. Opening of the epoxide ring with a sulfur nucleophile **RSH** yields leukotriene C₄.



New asthma drugs act by blocking the synthesis of leukotriene C₄ from arachidonic acid. For example, **zileuton** (trade name: Zylflo) inhibits the enzyme (called a lipoxygenase) needed for the first step of this process. By blocking the synthesis of leukotriene C₄, a compound responsible for the disease, zileuton treats the **cause of asthma**, not just its symptoms.




Generic name: zileuton
Trade name: Zylflo
anti-asthma drug

9.17 Application: Benzo[a]pyrene, Epoxides, and Cancer




The sooty exhaust from trucks and buses contains PAHs such as benzo[a]pyrene.

Benzo[a]pyrene is a widespread environmental pollutant, produced during the combustion of all types of organic material—gasoline, fuel oil, wood, garbage, and cigarettes. It is a **polycyclic aromatic hydrocarbon (PAH)**, a class of compounds that are discussed further in Chapter 17.



After this nonpolar and water-insoluble hydrocarbon is inhaled or ingested, it is oxidized in the liver to a diol epoxide. Oxidation is a common fate of foreign substances that are not useful nutrients for the body. The oxidation product has three oxygen-containing functional groups, making it much more water soluble, and more readily excreted in urine. It is also a potent carcinogen. The strained three-membered ring of the epoxide reacts readily with biological nucleophiles (such as DNA or an enzyme), leading to ring-opened products that often disrupt normal cell function, causing cancer or cell death.



These examples illustrate the central role of the nucleophilic opening of epoxide rings in two well-defined cellular processes.

KEY CONCEPTS


Alcohols, Ethers, and Epoxides

General Facts about ROH, ROR, and Epoxides

- All three compounds contain an O atom that is sp^3 hybridized and tetrahedral (9.2).
- All three compounds have polar C–O bonds, but only alcohols have an O–H bond for intermolecular hydrogen bonding (9.4).
- Alcohols and ethers do not contain a good leaving group. Nucleophilic substitution can occur only after the OH (or OR) group is converted to a better leaving group (9.7A).
- Epoxides have a leaving group located in a strained three-membered ring, making them reactive to strong nucleophiles and acids HZ that contain a nucleophilic atom Z (9.15).

A New Reaction of Carbocations (9.9)


- Less stable carbocations rearrange to more stable carbocations by the shift of a hydrogen atom or an alkyl group.



- Besides rearranging, a carbocation can also react with a nucleophile (7.13) and a base (8.6).

Preparation of Alcohols, Ethers, and Epoxides (9.6)

[1] Preparation of alcohols



- The mechanism is $\text{S}_{\text{N}}2$.
- The reaction works best for CH_3X and 1°RX .


[2] Preparation of alkoxides—A Brønsted–Lowry acid–base reaction



[3] Preparation of ethers (Williamson ether synthesis)



- The mechanism is $\text{S}_{\text{N}}2$.
- The reaction works best for CH_3X and 1°RX .


[4] Preparation of epoxides—Intramolecular $\text{S}_{\text{N}}2$ reaction

- A two-step reaction sequence:
 - [1] The removal of a proton with base forms an alkoxide.
 - [2] An intramolecular $\text{S}_{\text{N}}2$ reaction forms the epoxide.


Reactions of Alcohols

[1] Dehydration to form alkenes


a. Using strong acid (9.8, 9.9)




- Order of reactivity: $\text{R}_3\text{COH} > \text{R}_2\text{CHOH} > \text{RCH}_2\text{OH}$.
- The mechanism for 2° and 3°ROH is $\text{E}1$ —carbocations are intermediates and rearrangements occur.
- The mechanism for 1°ROH is $\text{E}2$.
- The Zaitsev rule is followed.

b. Using POCl_3 and pyridine (9.10)

- The mechanism is $\text{E}2$.
- No carbocation rearrangements occur.


[2] Reaction with HX to form RX (9.11)

- Order of reactivity: $\text{R}_3\text{COH} > \text{R}_2\text{CHOH} > \text{RCH}_2\text{OH}$.
- The mechanism for 2° and 3°ROH is $\text{S}_{\text{N}}1$ —carbocations are intermediates and rearrangements occur.
- The mechanism for CH_3OH and 1°ROH is $\text{S}_{\text{N}}2$.

[3] Reaction with other reagents to form RX (9.12)

- Reactions occur with CH_3OH and 1° and 2°ROH .
- The reactions follow an $\text{S}_{\text{N}}2$ mechanism.


[4] Reaction with tosyl chloride to form alkyl tosylates (9.13A)



- The C–O bond is not broken, so the configuration at a stereogenic center is retained.


Reactions of Alkyl Tosylates

Alkyl tosylates undergo either substitution or elimination, depending on the reagent (9.13B).




Reactions of Ethers

Only one reaction is useful: cleavage with strong acids (9.14).



Reactions of Epoxides

Epoxide rings are opened with nucleophiles :Nu^- and acids HZ (9.15).









PROBLEMS

Structure and Nomenclature

9.39 a. Draw the structure of a 1° , 2° , and 3° alcohol with molecular formula $\text{C}_4\text{H}_8\text{O}$.







b. Draw the structure of an enol with molecular formula $\text{C}_4\text{H}_8\text{O}$.

9.40 Give the IUPAC name for each alcohol.



[Also label the stereogenic centers as *R* or *S*.]

9.41 Name each ether and epoxide.



9.42 Give the structure corresponding to each name.

a. 4-ethyl-3-heptanol

b. *trans*-2-methylcyclohexanol

c. 2,3,3-trimethyl-2-butanol

d. 6-*sec*-butyl-7,7-diethyl-4-decanol

e. 3-chloro-1,2-propanediol

f. diisobutyl ether

g. 1,2-epoxy-1,3,3-trimethylcyclohexane

h. 1-ethoxy-3-ethylheptane

i. (2*R*,3*S*)-3-isopropyl-2-hexanol

j. (2*S*)-2-ethoxy-1,1-dimethylcyclopentane

9.43 Draw the eight constitutional isomers with molecular formula $C_5H_{12}O$ that contain an OH group. Give the IUPAC name for each compound. Classify each alcohol as 1°, 2°, or 3°.

Physical Properties

9.44 Rank each group of compounds in order of:

a. increasing boiling point: $CH_3CH_2CH_2OH$, $(CH_3)_2CHOH$, $CH_3CH_2OCH_3$

b. increasing water solubility: $CH_3(CH_2)_5OH$, $HO(CH_2)_6OH$, $CH_3(CH_2)_4CH_3$

9.45 Explain the observed trend in the melting points of the following three isomeric alcohols: $(CH_3)_2CHCH_2OH$ ($-108^\circ C$), $CH_3CH_2CH_2CH_2OH$ ($-90^\circ C$), $(CH_3)_3COH$ ($26^\circ C$).

9.46 Why is the boiling point of 1,3-propanediol ($HOCH_2CH_2CH_2OH$) higher than the boiling point of 1,2-propanediol [$HOCH_2CH(OH)CH_3$] ($215^\circ C$ vs. $187^\circ C$)? Why do both diols have a higher boiling point than 1-butanol ($CH_3CH_2CH_2CH_2OH$, $118^\circ C$)?

Alcohols

9.47 Draw the organic product(s) formed when $CH_3CH_2CH_2OH$ is treated with each reagent.

a. H_2SO_4

d. HBr

g. $TsCl$, pyridine

b. NaH

e. $SOCl_2$, pyridine

h. [1] NaH ; [2] CH_3CH_2Br

c. $HCl + ZnCl_2$

f. PBr_3

i. [1] $TsCl$, pyridine; [2] $NaSH$

9.48 Draw the organic product(s) formed when 1-methylcyclohexanol is treated with each reagent. In some cases no reaction occurs.

a. NaH

c. HBr

e. H_2SO_4

g. [1] NaH ; [2] CH_3CH_2Br




b. $NaCl$

d. HCl


f. $NaHCO_3$

h. $POCl_3$, pyridine

9.49 What alkenes are formed when each alcohol is dehydrated with $TsOH$? Label the major product when a mixture results.



d. $CH_3CH_2CH_2CH_2OH$



9.50 What three alkenes are formed when $CH_3CH_2CH_2CH(OH)CH_3$ is treated with H_2SO_4 ? Label the major product.

9.51 Draw the products formed when $CH_3CH_2CH_2CH_2OTs$ is treated with each reagent.





a. CH_3SH

b. $NaOCH_2CH_3$

c. $NaOH$



d. $KOC(CH_3)_3$

9.52 Draw the products of each reaction and indicate stereochemistry around stereogenic centers.




9.53 Draw the substitution product formed (including stereochemistry) when (2*R*)-2-hexanol is treated with each series of reagents: (a) NaH , followed by CH_3I ; (b) $TsCl$ and pyridine, followed by $NaOCH_3$; (c) PBr_3 , followed by $NaOCH_3$. Which two routes produce identical products?


9.54 Draw a stepwise mechanism for each reaction.




9.55 Consider the four compounds **A**, **B**, **C**, and **D**. (a) Draw all possible alkenes formed when each compound undergoes β elimination, and label the major product when one product predominates. Assume alcohols are dehydrated with H_2SO_4 and alkyl halides are treated with $\text{KOC}(\text{CH}_3)_3$. (b) Which compound would be the best starting material for the synthesis of 3,3-dimethylcyclopentene?



9.56 Although alcohol **V** gives a single alkene **W** when treated with POCl_3 and pyridine, three isomeric alkenes (**X–Z**) are formed on dehydration with H_2SO_4 . Draw a stepwise mechanism for each reaction and explain why the difference occurs.



9.57 Sometimes carbocation rearrangements can change the size of a ring. Draw a stepwise, detailed mechanism for the following reaction.




9.58 Indicate the stereochemistry of the alkyl halide formed when (3*S*)-3-hexanol is treated with (a) HBr ; (b) PBr_3 ; (c) HCl ; (d) SOCl_2 and pyridine.

9.59 Explain the following observation. When 3-methyl-2-butanol is treated with HBr , a single alkyl bromide is isolated, resulting from a 1,2-shift. When 2-methyl-1-propanol is treated with HBr , no rearrangement occurs to form an alkyl bromide.


9.60 To convert a 1° alcohol into a 1° alkyl chloride with HCl , a Lewis acid such as ZnCl_2 must be added to the reaction mixture. Explain why it is possible to omit the Lewis acid if a polar aprotic solvent such as HMPA, $[(\text{CH}_3)_2\text{N}]_3\text{P}=\text{O}$, is used.

9.61 An allylic alcohol contains an OH group on a carbon atom adjacent to a $\text{C}-\text{C}$ double bond. Treatment of allylic alcohol **A** with HCl forms a mixture of two allylic chlorides, **B** and **C**. Draw a stepwise mechanism that illustrates how both products are formed.




9.62 When $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ is treated with $\text{H}_2\text{SO}_4 + \text{NaBr}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$ is the major product, and $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ are isolated as minor products. Draw a mechanism that accounts for the formation of each of these products.

9.63 Draw a stepwise, detailed mechanism for the following reaction.





9.64 Draw a stepwise, detailed mechanism for the following intramolecular reaction that forms a cyclic ether.




Ethers



9.65 Draw two different routes to each of the following ethers using a Williamson ether synthesis. Indicate the preferred route (if there is one).





9.66 Explain why it is not possible to prepare the following ether using a Williamson ether synthesis.




9.67 Draw the products formed when each ether is treated with two equivalents of HBr.



9.68 Draw a stepwise mechanism for each reaction.



9.69 Draw a stepwise, detailed mechanism for the following reaction.



Epoxides

9.70 Draw the products formed when ethylene oxide is treated with each reagent.

a. HBr

b. H_2O (H_2SO_4)





c. [1] $\text{CH}_3\text{CH}_2\text{O}^-$; [2] H_2O

d. [1] $\text{HC}\equiv\text{C}^-$; [2] H_2O




e. [1] ^-OH ; [2] H_2O

f. [1] CH_3S^- ; [2] H_2O


9.71 Draw the products of each reaction.



9.72 When each halohydrin is treated with NaH, a product of molecular formula $\text{C}_4\text{H}_8\text{O}$ is formed. Draw the structure of the product and indicate its stereochemistry.




9.73 Devise a stepwise mechanism for the following reaction.












General Problems

9.74 Answer the following questions about alcohol **A**.



- Give the IUPAC name for **A**, including *R,S* designations for stereogenic centers.
- Classify **A** as a 1°, 2°, or 3° alcohol.
- Draw a stereoisomer for **A** and give its IUPAC name.
- Draw a constitutional isomer that contains an OH group and give its IUPAC name.
- Draw a constitutional isomer that contains an ether and give its IUPAC name.
- Draw the products formed (including stereochemistry) when **A** is treated with each reagent: [1] NaH; [2] H₂SO₄; [3] POCl₃, pyridine; [4] HCl; [5] SOCl₂, pyridine; [6] TsCl, pyridine.

9.75 Draw the products of each reaction, and indicate the stereochemistry where appropriate.




9.76 Prepare each compound from CH₃CH₂CH₂CH₂OH. More than one step may be needed.

- a. CH₃CH₂CH₂CH₂Br b. CH₃CH₂CH₂CH₂Cl c. CH₃CH₂CH₂CH₂OCH₂CH₃ d. CH₃CH₂CH₂CH₂N₃


9.77 Prepare each compound from cyclopentanol. More than one step may be needed.

- a.
- b.
- c.
- d.

9.78 Identify the reagents (a–h) needed to carry out each reaction.




9.79 Propranolol, an antihypertensive agent used in the treatment of high blood pressure, can be prepared from 1-naphthol, epichlorohydrin, and isopropylamine using two successive nucleophilic substitution reactions. Devise a stepwise synthesis of propranolol from these starting materials.




9.80 Palytoxin, the chapter-opening molecule, is a potent poison first isolated from marine soft corals obtained from a tidal pool on the Hawaiian island of Maui.

- Ignoring the OH group in blue, label all 2° OH groups that are located on stereogenic centers in palytoxin.
- The OH group in blue is part of a hemiacetal, a functional group that has the general structure $R_2C(OH)OR'$; that is, a hemiacetal contains a hydroxyl (OH) and alkoxy group (OR') bonded to the same carbon. Draw the carbocation that results from protonation and loss of H_2O from a hemiacetal. Explain why hemiacetals are more reactive than other 2° alcohols towards loss of H_2O in the presence of acid. We discuss hemiacetals in greater detail in Chapter 21.




Challenge Problems


9.81 Epoxides are converted to allylic alcohols with nonnucleophilic bases such as lithium diethylamide $[LiN(CH_2CH_3)_2]$. Draw a stepwise mechanism for the conversion of 1,2-epoxycyclohexane to 2-cyclohexen-1-ol with this base. Explain why a strong bulky base must be used in this reaction.




- 9.82** Rearrangements can occur during the dehydration of 1° alcohols even though no 1° carbocation is formed—that is, a 1,2-shift occurs as the C–OH₂⁺ bond is broken, forming a more stable 2° or 3° carbocation, as shown. Using this information, draw a stepwise mechanism that shows how CH₃CH₂CH₂CH₂OH is dehydrated with H₂SO₄ to form a mixture of CH₃CH₂CH=CH₂ and the cis and trans isomers of CH₃CH=CHCH₃. We will see another example of this type of rearrangement in Section 18.5C.



- 9.83** 1,2-Diols are converted to carbonyl compounds when treated with strong acids, in a reaction called the *pinacol rearrangement*. Draw a stepwise mechanism for this reaction.




- 9.84** Draw a stepwise mechanism for the following reaction.



10

Alkenes

- 10.1 Introduction
- 10.2 Calculating degrees of unsaturation
- 10.3 Nomenclature
- 10.4 Physical properties
- 10.5 Interesting alkenes
- 10.6 Lipids—Part 2
- 10.7 Preparation of alkenes
- 10.8 Introduction to addition reactions
- 10.9 Hydrohalogenation—Electrophilic addition of HX
- 10.10 Markovnikov's rule
- 10.11 Stereochemistry of electrophilic addition of HX
- 10.12 Hydration—Electrophilic addition of water
- 10.13 Halogenation—Addition of halogen
- 10.14 Stereochemistry of halogenation
- 10.15 Halohydrin formation
- 10.16 Hydroboration–oxidation
- 10.17 Keeping track of reactions
- 10.18 Alkenes in organic synthesis



Stearic acid and **oleic acid** are fatty acids, compounds that contain a carboxy group (COOH) attached to the end of a long carbon chain. Stearic acid is a **saturated fatty acid** because each carbon atom in its long chain has the maximum number of bonds to hydrogen. Oleic acid is an **unsaturated fatty acid** because its carbon chain contains one (cis) double bond. The presence of a double bond greatly affects the chemical and physical properties of these fatty acids. In Chapter 10 we learn about alkenes, organic compounds that contain carbon-carbon double bonds.


In Chapters 10 and 11 we turn our attention to **alkenes** and **alkynes**, compounds that contain one and two π bonds, respectively. Because π bonds are easily broken, alkenes and alkynes undergo **addition**, the third general type of organic reaction. These multiple bonds also make carbon atoms electron rich, so alkenes and alkynes react with a wide variety of electrophilic reagents in addition reactions.

In Chapter 10 we review the properties and synthesis of alkenes first, and then concentrate on reactions. **Every new reaction in Chapter 10 is an addition reaction.** The most challenging part is learning the reagents, mechanism, and stereochemistry that characterize each individual reaction.


10.1 Introduction

Alkenes are also called **olefins**.


Alkenes are compounds that contain a carbon–carbon double bond. **Terminal alkenes** have the double bond at the end of the carbon chain, whereas **internal alkenes** have at least one carbon atom bonded to each end of the double bond. **Cycloalkenes** contain a double bond in a ring.



The double bond of an alkene consists of one σ bond and one π bond. Each carbon is sp^2 hybridized and trigonal planar, and all bond angles are approximately 120° (Section 8.2A).



Bond dissociation energies of the C–C bonds in ethane (a σ bond only) and ethylene (one σ and one π bond) can be used to estimate the strength of the π component of the double bond. If we assume that the σ bond in ethylene is similar in strength to the σ bond in ethane (368 kJ/mol), then the π bond is worth 267 kJ/mol.



- The π bond is much weaker than the σ bond of a C–C double bond, making it much more easily broken. As a result, alkenes undergo many reactions that alkanes do not.

Other features of the carbon–carbon double bond, which were presented in Chapter 8, are summarized in Table 10.1.

Cycloalkenes having fewer than eight carbon atoms have a *cis* geometry. A *trans* cycloalkene must have a carbon chain long enough to connect the ends of the double bond without introducing too much strain. *trans*-Cyclooctene is the smallest, isolable *trans* cycloalkene, but it is considerably less stable than *cis*-cyclooctene, making it one of the few alkenes having a higher energy *trans* isomer.





Table 10.1 Properties of the Carbon–Carbon Double Bond

Property	Result
Restricted rotation	<ul style="list-style-type: none"> The rotation around the C–C double bond is restricted. Rotation can occur only if the π bond breaks and then re-forms, a process that is unfavorable (Section 8.2B).
Stereoisomerism	<ul style="list-style-type: none"> Whenever the two groups on each end of a C=C are different from each other, two diastereomers are possible. <i>Cis</i>- and <i>trans</i>-2-butene (drawn at the bottom of Table 10.1) are diastereomers (Section 8.2B).
Stability	<ul style="list-style-type: none"> Trans alkenes are generally more stable than cis alkenes. The stability of an alkene increases as the number of R groups on the C=C increases (Section 8.2C).



Problem 10.1 Draw the six alkenes of molecular formula C_5H_{10} . Label one pair of diastereomers.

10.2 Calculating Degrees of Unsaturation

An acyclic alkene has the general molecular formula C_nH_{2n} , giving it two fewer hydrogens than an acyclic alkane with the same number of carbons.

- Alkenes are *unsaturated hydrocarbons* because they have fewer than the maximum number of hydrogen atoms per carbon.

In Chapter 12 we will learn how to use the hydrogenation of π bonds to determine how many degrees of unsaturation result from π bonds and how many result from rings.

Cycloalkanes also have the general molecular formula C_nH_{2n} . Thus, **each π bond or ring removes two hydrogen atoms from a molecule, and this introduces one degree of unsaturation**. The number of degrees of unsaturation for a given molecular formula can be calculated by comparing the actual number of H atoms in a compound and the maximum number of H atoms possible. Remember that for n carbons, the maximum number of H atoms is $2n + 2$ (Section 4.1). This procedure gives the total number of rings and π bonds in a molecule.

Sample Problem 10.1

Calculate the number of degrees of unsaturation in a compound of molecular formula C_4H_6 , and propose possible structures.

Solution

[1] Calculate the maximum number of H's possible.

- For n carbons, the maximum number of H's is $2n + 2$; in this example, $2n + 2 = 2(4) + 2 = 10$.

[2] Subtract the actual number of H's from the maximum number and divide by two.

- 10 H's (maximum) – 6 H's (actual) = 4 H's fewer than the maximum number.

$$\frac{4 \text{ H's fewer than the maximum}}{2 \text{ H's removed for each degree of unsaturation}} =$$

two degrees of unsaturation

A compound with two degrees of unsaturation has:

two rings



or

two π bonds

or

one ring and one π bond

Possible structures
for C_4H_6 :



This procedure can be extended to compounds that contain heteroatoms such as oxygen, nitrogen, and halogen, as illustrated in Sample Problem 10.2.

Sample Problem 10.2

Calculate the number of degrees of unsaturation for each molecular formula: (a) C_5H_8O ; (b) $C_6H_{11}Cl$; (c) C_8H_9N . Propose one possible structure for each compound.

Solution

a. You can ignore the presence of an O atom when calculating degrees of unsaturation; that is, use only the given number of C's and H's for the calculation (C_5H_8).

[1] For 5 C's, the maximum number of H's = $2n + 2 = 2(5) + 2 = 12$.

[2] Because the compound contains only 8 H's, it has $12 - 8 = 4$ H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.

Answer: two degrees of unsaturation

b. A compound with a halogen atom is equivalent to a hydrocarbon having one more H; that is, $C_6H_{11}Cl$ is equivalent to C_6H_{12} when calculating degrees of unsaturation.

[1] For 6 C's, the maximum number of H's = $2n + 2 = 2(6) + 2 = 14$.

[2] Because the compound contains only 12 H's, it has $14 - 12 = 2$ H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.

Answer: one degree of unsaturation

c. A compound with a nitrogen atom is equivalent to a hydrocarbon having one fewer H; that is, C_8H_9N is equivalent to C_8H_8 when calculating degrees of unsaturation.




[1] For 8 C's, the maximum number of H's = $2n + 2 = 2(8) + 2 = 18$.

[2] Since the compound contains only 8 H's, it has $18 - 8 = 10$ H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.

Answer: five degrees of unsaturation

Possible structures:



Problem 10.2

How many degrees of unsaturation are present in each compound?

- a. C_2H_2 b. C_6H_6 c. C_8H_{18} d. C_7H_8O e. $C_7H_{11}Br$ f. C_5H_9N

Problem 10.3

Give an example of a compound with molecular formula C_6H_{10} that satisfies each criterion.

- a. a compound with two π bonds c. a compound with two rings
b. a compound with one ring and one π bond d. a compound with one triple bond


10.3 Nomenclature

- In the IUPAC system, an alkene is identified by the suffix *-ene*.


10.3A General IUPAC Rules

HOW TO Name an Alkene

Example Give the IUPAC name of the following alkene:



Step [1] Find the longest chain that contains *both* carbon atoms of the double bond.




6 C's in the longest chain
hexane ----> hexene

- Change the *-ane* ending of the parent alkane to *-ene*.

Step [2] Number the carbon chain to give the double bond the lower number, and apply all other rules of nomenclature.


a. **Number** the chain, and name using the **first number** assigned to the C=C.



- Number the chain to put the C=C at C2, not C4.

2-hexene

b. **Name and number** the substituents.




three methyl groups at C2, C3, and C5

Answer: 2,3,5-trimethyl-2-hexene

Compounds with two double bonds are named as **dienes** by changing the *-ane* ending of the parent alkane to the suffix *-adiene*. Compounds with three double bonds are named as **trienes**, and so forth. Always choose the longest chain that contains *both* atoms of the double bond. In Figure 10.1, the alkene is named as a derivative of heptene because the seven-carbon chain contains both atoms of the double bond, but the eight-carbon chain does not.

Figure 10.1


Naming an alkene in which the longest carbon chain does not contain both atoms of the double bond



7 C's ----> heptene

Both C's of the C=C are contained in this long chain.

Correct: 2-ethyl-1-heptene




8 C's

Both C's of the C=C are NOT contained in this long chain.

Incorrect


Figure 10.2
Examples of cycloalkene nomenclature



$\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ is named as 1-butene using the 1979 IUPAC recommendations and but-1-ene using the 1993 IUPAC recommendations. The first convention is more widely used, so we follow it in this text.




In naming cycloalkenes, the double bond is located between C1 and C2, and the “1” is usually omitted in the name. The ring is numbered clockwise or counterclockwise to give the first substituent the lower number. Representative examples are given in Figure 10.2.

Compounds that contain both a double bond and a hydroxy group are named as **alkenols**, and the chain (or ring) is numbered to give the OH group the lower number.






Problem 10.4 Give the IUPAC name for each alkene.

- a. $\text{CH}_2=\text{CHCH}(\text{CH}_3)\text{CH}_2\text{CH}_3$
b. $(\text{CH}_3\text{CH}_2)_2\text{C}=\text{CHCH}_2\text{CH}_2\text{CH}_3$



Problem 10.5 Give the IUPAC name for each polyfunctional compound.




10.3B Naming Stereoisomers

A prefix is needed to distinguish two alkenes when diastereomers are possible.

Using Cis and Trans as Prefixes


An alkene having one alkyl group bonded to each carbon atom can be named using the prefixes **cis** and **trans** to designate the relative location of the two alkyl groups. For example, *cis*-3-hexene has two ethyl groups on the **same side** of the double bond, whereas *trans*-3-hexene has two ethyl groups on **opposite sides** of the double bond.



Using the Prefixes E and Z

Although the prefixes *cis* and *trans* can be used to distinguish diastereomers when two alkyl groups are bonded to the C=C, they cannot be used when there are three or four alkyl groups bonded to the C=C.

E stands for the German word *entgegen* meaning “opposite.” *Z* stands for the German word *zusammen*, meaning “together.” Using *E,Z* nomenclature, a *cis* isomer has the *Z* configuration and a *trans* isomer has the *E* configuration.




For example, alkenes **A** and **B** are two *different* compounds that are both called 3-methyl-2-pentene. In **A** the two CH_3 groups are *cis*, whereas in **B** the CH_3 and CH_2CH_3 groups are *cis*. The ***E,Z* system of nomenclature** has been devised to unambiguously name these kinds of alkenes.


HOW TO Assign the Prefixes *E* and *Z* to an Alkene

Step [1] Assign priorities to the two substituents on each end of the $\text{C}=\text{C}$ by using the priority rules for *R,S* nomenclature (Section 5.6).

- Divide the double bond in half, and assign the numbers **1** and **2** to indicate the relative priority of the two groups on each end—the higher priority group is labeled **1**, and the lower priority group is labeled **2**.






Step [2] Assign *E* or *Z* based on the location of the two higher priority groups (**1**).



- The ***E*** isomer has the two higher priority groups on the **opposite** sides.
- The ***Z*** isomer has the two higher priority groups on the **same** side.

Problem 10.6 Label each $\text{C}-\text{C}$ double bond as *E* or *Z*. Kavain is a naturally occurring relaxant isolated from kava root.



kavain

Problem 10.7 Give the IUPAC name for each alkene.






Figure 10.3
Naming alkenes with common substituent names




Problem 10.8 Draw the structure corresponding to each IUPAC name.
a. (3*Z*)-4-ethyl-3-heptene b. (2*E*)-3,5,6-trimethyl-2-octene c. (1*Z*)-2-bromo-1-iodo-1-hexene

Problem 10.9 Draw and name all the stereoisomers of $\text{CH}_3\text{CH}=\text{CH}-\text{CH}=\text{CHCH}_3$.

10.3C Common Names

The simplest alkene, $\text{CH}_2=\text{CH}_2$, named in the IUPAC system as **ethene**, is often called **ethylene**, its common name. The common names for three **alkyl groups** derived from alkenes are also used. Two examples of naming organic molecules using these common names are shown in Figure 10.3.




10.4 Physical Properties


Most alkenes exhibit only weak van der Waals interactions, so their physical properties are similar to alkanes of comparable molecular weight.

- Alkenes have low melting points and boiling points.
- Melting points and boiling points increase as the number of carbons increases because of increased surface area.
- Alkenes are soluble in organic solvents and insoluble in water.

cis- and *trans*-alkenes often have somewhat different physical properties. For example, *cis*-2-butene has a higher boiling point (4 °C) than *trans*-2-butene (1 °C). This difference arises because the C–C single bond between an alkyl group and one of the double-bond carbons of an alkene is slightly polar. **The sp^3 hybridized alkyl carbon donates electron density to the sp^2 hybridized alkenyl carbon.**




The bond dipole places a partial negative charge on the alkenyl carbon (sp^2) relative to the alkyl carbon (sp^3) because an sp^2 hybridized orbital has greater percent *s*-character (33%) than an sp^3 hybridized orbital (25%). **In a *cis* isomer, the two $\text{C}_{sp^3}-\text{C}_{sp^2}$ bond dipoles reinforce each other, yielding a small net molecular dipole. In a *trans* isomer, the two bond dipoles cancel.**



Related arguments involving $\text{C}_{sp^3}-\text{C}_{sp^2}$ bonds were used in Section 8.2C to explain why the stability of an alkene increases with increasing alkyl substitution.

- A *cis* alkene is more polar than a *trans* alkene, giving it a slightly higher boiling point and making it more soluble in polar solvents.

Problem 10.10 Rank the following isomers in order of increasing boiling point.



10.5 Interesting Alkenes

The alkenes in Figure 10.5 all originate from the same five-carbon starting material in nature, as we will learn in Chapter 29.

Ethylene is prepared from petroleum by a process called **cracking**. Ethylene is the most widely produced organic chemical, serving as the starting material not only for the polymer **polyethylene**, a widely used plastic, but also for many other useful organic compounds, as shown in Figure 10.4.

Numerous organic compounds containing carbon–carbon double bonds have been isolated from natural sources (Figure 10.5).

Problem 10.11 Using Figure 10.5, draw the structure of (*S*)-limonene, which is isolated from pine needles and has a turpentine-like odor. Explain how it can smell differently from the *R* isomer, which has a citrus fragrance.

10.6 Lipids—Part 2

Understanding the geometry of C–C double bonds provides an insight into the properties of **triacylglycerols**, the most abundant lipids. Triacylglycerols contain three ester groups, each having a long carbon chain (abbreviated as R, R', and R'') bonded to a carbonyl group (C=O).

Figure 10.4
Ethylene, an industrial starting material for many useful products








Table 10.2 The Effect of Double Bonds on the Melting Point of Fatty Acids

Name	Structure	Mp (°C)
Stearic acid (0 C=C)		69
Oleic acid (1 C=C)		4
Linoleic acid (2 C=C)		-5
Linolenic acid (3 C=C)		-11

Increasing number of double bonds

Linoleic and linolenic acids are essential fatty acids, meaning they cannot be synthesized in the body and must therefore be obtained in the diet. A common source of these essential fatty acids is whole milk. Babies fed a diet of nonfat milk in their early months do not thrive because they do not obtain enough of these essential fatty acids.

- Saturated fatty acids have no double bonds in their long hydrocarbon chains, and unsaturated fatty acids have one or more double bonds in their hydrocarbon chains.
- Double bonds in naturally occurring fatty acids have the Z configuration.

Table 10.2 lists the structure and melting point of four fatty acids containing 18 carbon atoms. Stearic acid is one of the two most common saturated fatty acids, and oleic and linoleic acids are the most common unsaturated ones. The data show the effect of Z double bonds on the melting point of fatty acids.

- As the number of double bonds in the fatty acid *increases*, the melting point *decreases*.


The three-dimensional structures of the fatty acids in Figure 10.6 illustrate how Z double bonds introduce kinks in the long hydrocarbon chain, decreasing the ability of the fatty acid to pack well in a crystalline lattice. **The larger the number of Z double bonds, the more kinks in the hydrocarbon chain, and the lower the melting point.**

Problem 10.12


Linolenic acid (Table 10.2) and stearidonic acid are omega-3 fatty acids, unsaturated fatty acids that contain the first double bond located at C3, when numbering begins at the methyl end of the chain. Predict how the melting point of stearidonic acid compares with the melting points of linolenic and

Figure 10.6

Three-dimensional structure of four C₁₈ fatty acids



stearic acids. A current avenue of research is examining the use of soybean oil enriched in stearidonic acid as a healthier alternative to vegetable oils that contain fewer degrees of unsaturation.



10.6B Fats and Oils



Canola, soybeans, and flaxseed are excellent dietary sources of linolenic acid, an essential fatty acid. Oils derived from omega-3 fatty acids (Problem 10.12) are currently thought to be especially beneficial for individuals at risk of developing coronary artery disease.

Fats and oils are triacylglycerols with different physical properties.

- Fats have higher melting points—they are *solids* at room temperature.
- Oils have lower melting points—they are *liquids* at room temperature.

The identity of the three fatty acids in the triacylglycerol determines whether it is a fat or an oil. **Increasing the number of double bonds in the fatty acid side chains decreases the melting point of the triacylglycerol.**

- Fats are derived from fatty acids having few double bonds.
- Oils are derived from fatty acids having a larger number of double bonds.


Saturated fats are typically obtained from animal sources, whereas unsaturated oils are common in vegetable sources. Thus, butter and lard are high in saturated triacylglycerols, and olive oil and safflower oil are high in unsaturated triacylglycerols. An exception to this generalization is coconut oil, which is composed largely of saturated alkyl side chains.

Considerable evidence suggests that an elevated cholesterol level is linked to an increased risk of heart disease. Saturated fats stimulate cholesterol synthesis in the liver, thus increasing the cholesterol concentration in the blood.


10.7 Preparation of Alkenes

Recall from Chapters 8 and 9 that alkenes can be prepared from alkyl halides and alcohols via elimination reactions. For example, **dehydrohalogenation of alkyl halides with strong base yields alkenes via an E2 mechanism** (Sections 8.4 and 8.5).


- Typical bases include OH^- and OR^- [especially $\text{OC}(\text{CH}_3)_3^-$], and nonnucleophilic bases such as DBU and DBN.
- Alkyl tosylates can also be used as starting materials under similar reaction conditions (Section 9.13).




The acid-catalyzed dehydration of alcohols with H_2SO_4 or TsOH yields alkenes, too (Sections 9.8 and 9.9). The reaction occurs via an E1 mechanism for 2° and 3° alcohols, and an E2 mechanism for 1° alcohols. E1 reactions involve carbocation intermediates, so rearrangements are possible. **Dehydration can also be carried out with POCl_3 and pyridine** by an E2 mechanism (Section 9.10).



These elimination reactions are stereoselective and regioselective, so the most stable alkene is usually formed as the major product.




Problem 10.13 Draw the products of each elimination reaction.



10.8 Introduction to Addition Reactions

Because the C–C π bond of an alkene is much weaker than a C–C σ bond, the characteristic reaction of alkenes is **addition: the π bond is broken and two new σ bonds are formed.**



Alkenes are electron rich, as seen in the electrostatic potential plot in Figure 10.7. The electron density of the π bond is concentrated above and below the plane of the molecule, making the π bond more exposed than the σ bond.


What kinds of reagents add to the weak, electron-rich π bond of alkenes? There are many of them, and that can make alkene chemistry challenging. To help you organize this information, keep in mind the following:

- Every reaction of alkenes involves addition: the π bond is always broken.
- Because alkenes are electron rich, simple alkenes do *not* react with nucleophiles or bases, reagents that are themselves electron rich. Alkenes react with *electrophiles*.

The addition reactions of alkenes are discussed in Sections 10.9–10.16 and in Chapter 12 (Oxidation and Reduction).


Figure 10.7

Electrostatic potential plot of ethylene



- The red electron-rich region of the π bond is located above and below the plane of the molecule. Because the plane of the alkene depicted in this electrostatic potential plot is tipped, only the red region above the molecule is visible.

The stereochemistry of addition is often important in delineating a reaction's mechanism. Because the carbon atoms of a double bond are both trigonal planar, the elements of X and Y can be added to them from the **same side** or from **opposite sides**.




- *Syn addition* takes place when both X and Y are added from the *same* side.
- *Anti addition* takes place when X and Y are added from *opposite* sides.

Five reactions of alkenes are discussed in Chapter 10 and each is illustrated in Figure 10.8, using cyclohexene as the starting material.

10.9 Hydrohalogenation—Electrophilic Addition of HX

Hydrohalogenation is the addition of hydrogen halides HX (X = Cl, Br, and I) to alkenes to form alkyl halides.



Hydrohalogenation of an alkene to form an alkyl halide is the reverse of the dehydrohalogenation of an alkyl halide to form an alkene, a reaction discussed in detail in Sections 8.4 and 8.5.

Two bonds are broken in this reaction—the weak π bond of the alkene and the HX bond—and two new σ bonds are formed—one to H and one to X. Because X is more electronegative than H, the H–X bond is polarized, with a partial positive charge on H. Because the electrophilic (H) end of HX is attracted to the electron-rich double bond, these reactions are called **electrophilic additions**.

Figure 10.8
Five addition reactions of cyclohexene






Figure 10.9

The addition of HBr to $\text{CH}_2=\text{CH}_2$, an exothermic reaction



To draw the products of an addition reaction:

- **Locate the C–C double bond.**
- **Identify the σ bond of the reagent that breaks—namely, the H–X bond in hydrohalogenation.**
- **Break the π bond of the alkene and the σ bond of the reagent, and form two new σ bonds to the C atoms of the double bond.**



Addition reactions are exothermic because the two σ bonds formed in the product are stronger than the σ and π bonds broken in the reactants. For example, ΔH° for the addition of HBr to ethylene is -60 kJ/mol , as illustrated in Figure 10.9.

The mechanism of electrophilic addition of HX consists of **two steps**: addition of H^+ to form a carbocation, followed by nucleophilic attack of X^- . The mechanism is illustrated for the reaction of *cis*-2-butene with HBr in Mechanism 10.1.


The mechanism of electrophilic addition consists of two successive Lewis acid–base reactions. In Step [1], the **alkene is the Lewis base** that donates an electron pair to **H–Br, the Lewis acid**, while in Step [2], **Br[−] is the Lewis base** that donates an electron pair to the **carbocation, the Lewis acid**.

An energy diagram for the reaction of $\text{CH}_3\text{CH}=\text{CHCH}_3$ with HBr is given in Figure 10.10. Each step has its own energy barrier with a transition state at each energy maximum. Because Step [1] has a higher energy transition state, it is rate-determining. ΔH° for Step [1] is positive because more bonds are broken than formed, whereas ΔH° for Step [2] is negative because only bond making occurs.




Mechanism 10.1 Electrophilic Addition of HX to an Alkene

Step [1] Addition of the electrophile (H^+) to the π bond






- The π bond attacks the H atom of HBr, thus forming a new C–H bond while breaking the H–Br bond. Because the remaining carbon atom of the original double bond is left with only six electrons, a **carbocation** intermediate is formed. This step is **rate-determining** because two bonds are broken but only one bond is formed.

Step [2] Nucleophilic attack of Br^-



- Nucleophilic attack of Br^-** on the carbocation forms the new C–Br bond.

Problem 10.14 What product is formed when each alkene is treated with HCl?



Problem 10.15 Draw a stepwise mechanism for the following reaction. Draw the transition state for each step.





Figure 10.10


Energy diagram for electrophilic addition: $\text{CH}_3\text{CH}=\text{CHCH}_3 + \text{HBr} \rightarrow \text{CH}_3\text{CH}_2\text{CH}(\text{Br})\text{CH}_3$



- The mechanism has two steps, so there are two energy barriers.
- Step [1] is rate-determining.

10.10 Markovnikov's Rule


With an unsymmetrical alkene, HX can add to the double bond to give two constitutional isomers.



For example, HCl addition to propene could in theory form 1-chloropropane by addition of H and Cl to C2 and C1, respectively, and 2-chloropropane by addition of H and Cl to C1 and C2, respectively. In fact, **electrophilic addition forms only 2-chloropropane**. This is a specific example of a general trend called **Markovnikov's rule**, named for the Russian chemist who first determined the regioselectivity of electrophilic addition of HX.

- In the addition of HX to an unsymmetrical alkene, the H atom bonds to the less substituted carbon atom—that is, the carbon that has more H atoms to begin with.

The basis of Markovnikov's rule is the formation of a carbocation in the rate-determining step of the mechanism. With propene, there are two possible paths for this first step, depending on which carbon atom of the double bond forms the new bond to hydrogen.




The Hammond postulate was first introduced in Section 7.15 to explain the relative rate of $\text{S}_{\text{N}}1$ reactions with 1° , 2° , and 3° RX.

Path [1] forms a highly unstable 1° carbocation, whereas Path [2] forms a **more stable 2° carbocation**. According to the Hammond postulate, Path [2] is faster because formation of the carbocation is an endothermic process, so **the transition state to form the more stable 2° carbocation is lower in energy** (Figure 10.11).

Figure 10.11

Electrophilic addition and the Hammond postulate




- The E_a for formation of the more stable 2° carbocation is lower than the E_a for formation of the 1° carbocation. The 2° carbocation is formed faster.

- In the addition of HX to an unsymmetrical alkene, the H atom is added to the less substituted carbon to form the more stable, more substituted carbocation.

Similar results are seen in any electrophilic addition involving an intermediate carbocation: **the more stable, more substituted carbocation is formed by addition of the electrophile to the less substituted carbon.**


Problem 10.16 Draw the products formed when each alkene is treated with HCl.



Problem 10.17 Use the Hammond postulate to explain why $(\text{CH}_3)_2\text{C}=\text{CH}_2$ reacts faster than $\text{CH}_3\text{CH}=\text{CH}_2$ in electrophilic addition of HX.

Because carbocations are formed as intermediates in hydrohalogenation, carbocation rearrangements can occur, as illustrated in Sample Problem 10.3.


Sample Problem 10.3 Draw a stepwise mechanism for the following reaction.




Solution

Because the carbon skeletons of the starting material and product are different—the alkene reactant has a 4° carbon and the product alkyl halide does not—a carbocation rearrangement must have occurred.

Step [1] Markovnikov addition of HBr adds H^+ to the less substituted end of the double bond, forming a 2° carbocation.



Steps [2] Rearrangement of the 2° carbocation by a 1,2-methyl shift forms a more stable 3° and [3] carbocation. Nucleophilic attack of Br^- forms the product, a 3° alkyl halide.



Problem 10.18 Treatment of 3-methylcyclohexene with HCl yields two products, 1-chloro-3-methylcyclohexane and 1-chloro-1-methylcyclohexane. Draw a mechanism to explain this result.

Problem 10.19 Addition of HBr to which of the following alkenes will lead to a rearrangement?


- a. $\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}_2\text{CH}_3$ b. $\text{CH}_3\text{CH}=\text{CHCH}_2\text{CH}_3$ c. $\text{CH}_3\text{CH}=\text{CHCH}(\text{CH}_3)_2$

10.11 Stereochemistry of Electrophilic Addition of HX



To understand the stereochemistry of electrophilic addition, recall two stereochemical principles learned in Chapters 7 and 9.

- Trigonal planar atoms react with reagents from two directions with equal probability (Section 7.13C).
- Achiral starting materials yield achiral or racemic products (Section 9.15).


Many hydrohalogenation reactions begin with an **achiral reactant** and form an **achiral product**. For example, the addition of HBr to cyclohexene, an achiral alkene, forms bromocyclohexane, an achiral alkyl halide.




Because addition converts sp^2 hybridized carbons to sp^3 hybridized carbons, however, sometimes new stereogenic centers are formed from hydrohalogenation. For example, Markovnikov addition of HCl to 2-ethyl-1-pentene, an achiral alkene, forms one constitutional isomer, 3-chloro-3-methylhexane. Because this product now has a stereogenic center at one of the newly formed sp^3 hybridized carbons, **an equal amount of two enantiomers—a racemic mixture**—must form.



The mechanism of hydrohalogenation illustrates why two enantiomers are formed. Initial addition of the electrophile H^+ (from HCl) occurs from **either side of the planar double bond** to form a carbocation. Both modes of addition (from above and below) generate the same **achiral carbocation**. Either representation of this carbocation can then be used to draw the second step of the mechanism.



Nucleophilic attack of Cl^- on the trigonal planar carbocation also occurs from two different directions, forming two products, **A** and **B**, having a new stereogenic center. **A** and **B** are not superimposable, so they are **enantiomers**. Because attack from either direction occurs with equal probability, a **racemic mixture** of **A** and **B** is formed.



Because hydrohalogenation begins with a **planar** double bond and forms a **planar** carbocation, addition of H and Cl occurs in two different ways. The elements of H and Cl can both be added from the same side of the double bond—that is, **syn addition**—or they can be added from opposite sides—that is, **anti addition**. *Both* modes of addition occur in this two-step reaction mechanism.

- Hydrohalogenation occurs with syn and anti addition of HX.

The terms **cis** and **trans** refer to the arrangement of groups in a particular compound, usually an alkene or a disubstituted cycloalkane. The terms **syn** and **anti** describe the stereochemistry of a process—for example, how two groups are added to a double bond.

Finally, addition of HCl to 1,2-dimethylcyclohexene forms **two** new stereogenic centers. Initial addition of H⁺ (from HCl) forms two enantiomeric carbocations that react with the Cl⁻ nucleophile from two different directions, forming **four stereoisomers, A–D—two pairs of enantiomers** (Figure 10.12).





Figure 10.12

Reaction of 1,2-dimethylcyclohexene with HCl



Four stereoisomers are formed:


- Compounds **A** and **D** are enantiomers formed in equal amounts.
- Compounds **B** and **C** are enantiomers formed in equal amounts.

Table 10.3 summarizes the characteristics of electrophilic addition of HX to alkenes.

Table 10.3 Summary: Electrophilic Addition of HX to Alkenes

Observation	
Mechanism	<ul style="list-style-type: none"> The mechanism involves two steps. The rate-determining step forms a carbocation. Rearrangements can occur.
Regioselectivity	<ul style="list-style-type: none"> Markovnikov's rule is followed. In unsymmetrical alkenes, H bonds to the less substituted C to form the more stable carbocation.
Stereochemistry	<ul style="list-style-type: none"> Syn and anti addition occur.




Problem 10.20 Draw the products, including stereochemistry, of each reaction.



Problem 10.21 Which compounds (A–D) in Figure 10.12 are formed by syn addition of HCl and which are formed by anti addition?

10.12 Hydration—Electrophilic Addition of Water

Hydration is the addition of water to an alkene to form an alcohol. H_2O itself is too weak an acid to protonate an alkene, but with added H_2SO_4 , H_3O^+ is formed and addition readily occurs.



Hydration is simply another example of **electrophilic addition**. The first two steps of the mechanism are similar to those of electrophilic addition of HX—that is, addition of H^+ (from H_3O^+) to generate a carbocation, followed by nucleophilic attack of H_2O . Mechanism 10.2 illustrates the addition of H_2O to cyclohexene to form cyclohexanol.

There are three consequences to the formation of carbocation intermediates:

- In unsymmetrical alkenes, H adds to the less substituted carbon to form the more stable carbocation; that is, Markovnikov's rule holds.
- Addition of H and OH occurs in both a syn and anti fashion.
- Carbocation rearrangements can occur.


Alcohols add to alkenes, forming ethers, using the same mechanism. Addition of CH_3OH to 2-methylpropene, for example, forms *tert*-butyl methyl ether (MTBE), a high octane fuel additive described in Section 3.4C.

Hydration of an alkene to form an alcohol is the reverse of the dehydration of an alcohol to form an alkene, a reaction discussed in detail in Section 9.8.




Mechanism 10.2 Electrophilic Addition of H₂O to an Alkene—Hydration

Step [1] Addition of the electrophile (H⁺) to the π bond




- The π bond attacks H₃O⁺, thus forming a new C–H bond while breaking the H–O bond. Because the remaining carbon atom of the original double bond is left with only six electrons, a **carbocation** intermediate is formed. This step is **rate-determining** because two bonds are broken but only one bond is formed.

Step [2] Nucleophilic attack of H₂O




- Nucleophilic attack of H₂O** on the carbocation forms the new C–O bond.


Step [3] Loss of a proton



- Removal of a proton with a base (H₂O) forms a neutral alcohol. Because the acid used in Step [1] is regenerated in Step [3], hydration is **acid-catalyzed**.




Problem 10.22 What two alkenes give rise to each alcohol as the major product of acid-catalyzed hydration?



Problem 10.23 What stereoisomers are formed when 1-pentene is treated with H₂O and H₂SO₄?

10.13 Halogenation—Addition of Halogen


Halogenation is the addition of halogen X₂ (X = Cl or Br) to an alkene, forming a vicinal dihalide.



Halogenation is synthetically useful only with Cl₂ and Br₂. The dichlorides and dibromides formed in this reaction serve as starting materials for the synthesis of alkynes, as we learned in Section 8.10.




Bromination is a simple chemical test for the presence of π bonds in unknown compounds. When bromine, a fuming red liquid, is added to an alkene dissolved in the solvent CCl_4 , the bromine adds to the double bond and the red color disappears. The disappearance of the red color is therefore a positive test for π bonds.



Halogens add to π bonds because halogens are **polarizable**. The electron-rich double bond induces a dipole in an approaching halogen molecule, making one halogen atom electron deficient and the other electron rich ($\text{X}^{\delta+}-\text{X}^{\delta-}$). **The electrophilic halogen atom is then attracted to the nucleophilic double bond**, making addition possible.

Two facts demonstrate that halogenation follows a different mechanism from that of hydrohalogenation or hydration. First, no rearrangements occur, and second, only anti addition of X_2 is observed. For example, treatment of cyclohexene with Br_2 yields two **trans** enantiomers formed by **anti addition**.




These facts suggest that **carbocations are not intermediates in halogenation**. Unstable carbocations rearrange, and both syn and anti addition is possible with carbocation intermediates. The accepted mechanism for halogenation comprises **two steps**, but it does *not* proceed with formation of a carbocation, as shown in Mechanism 10.3.




Mechanism 10.3 Addition of X_2 to an Alkene—Halogenation

Step [1] Addition of the electrophile (X^+) to the π bond




- Four bonds are broken or formed in this step: the electron pair in the π bond and a lone pair on a halogen atom are used to form two new $\text{C}-\text{X}$ bonds. The $\text{X}-\text{X}$ bond is also cleaved heterolytically, forming X^- . This step is rate-determining.
- The three-membered ring containing a positively charged halogen atom is called a **bridged halonium ion**. This strained three-membered ring is highly unstable, making it amenable to opening of the ring in the second step.

Step [2] Nucleophilic attack of X^-




- **Nucleophilic attack of X^-** opens the ring of the halonium ion, forming a new $\text{C}-\text{X}$ bond and relieving the strain in the three-membered ring.

Bridged halonium ions resemble carbocations in that they are short-lived intermediates that react readily with nucleophiles. Carbocations are inherently unstable because only six electrons surround carbon, whereas **halonium ions are unstable because they contain a strained three-membered ring** with a positively charged halogen atom.




Problem 10.24 Draw the transition state for each step in the general mechanism for the halogenation of an alkene.

Problem 10.25 Draw the products of each reaction, including stereochemistry.




10.14 Stereochemistry of Halogenation

How does the proposed mechanism invoking a bridged halonium ion intermediate explain the observed **trans products of halogenation**? For example, chlorination of cyclopentene affords both enantiomers of *trans*-1,2-dichlorocyclopentane, with *no cis* products.




Initial addition of the electrophile Cl⁺ (from Cl₂) occurs from either side of the planar double bond to form the bridged chloronium ion. In this example, both modes of addition (from above and below) generate the same **achiral** intermediate, so either representation can be used to draw the second step.



In the second step, **nucleophilic attack of Cl⁻ must occur from the back side**—that is, from the side of the five-membered ring opposite to the side having the bridged chloronium ion. Because the nucleophile attacks from below in this example and the leaving group departs from above, the two Cl atoms in the product are oriented **trans** to each other. Backside attack occurs with equal probability at either carbon of the three-membered ring to yield an equal amount of two enantiomers—a **racemic mixture**.

The opening of bridged halonium ion intermediates resembles the opening of epoxide rings with nucleophiles discussed in Section 9.15.



In summary, the mechanism for halogenation of alkenes occurs in two steps:

- Addition of X^+ forms an unstable bridged halonium ion in the rate-determining step.
- Nucleophilic attack of X^- occurs from the back side to form trans products. The overall result is anti addition of X_2 across the double bond.

Because halogenation occurs exclusively in an anti fashion, cis and trans alkenes yield different stereoisomers. Halogenation of alkenes is a **stereospecific reaction**.

- A reaction is stereospecific when each of two specific stereoisomers of a starting material yields a particular stereoisomer of a product.

cis-2-Butene yields two enantiomers, whereas *trans*-2-butene yields a single achiral meso compound, as shown in Figure 10.13.

Problem 10.26

Draw all stereoisomers formed in each reaction.





Figure 10.13

Halogenation of *cis*- and *trans*-2-butene





To draw the products of halogenation:

- Add Br_2 in an **anti** fashion across the double bond, leaving all other groups in their original orientations. Draw the products such that a given Br atom is above the plane in one product and below the plane in the other product.
- Sometimes this reaction produces two stereoisomers, as in the case of *cis*-2-butene, which forms an equal amount of two enantiomers. Sometimes it produces a single compound, as in the case of *trans*-2-butene, where a meso compound is formed.

Problem 10.27 Draw a stepwise mechanism for the conversion of *trans*-2-butene to the meso dibromide in Figure 10.13.

10.15 Halohydrin Formation

Treatment of an alkene with a halogen X_2 and H_2O forms a **halohydrin** by addition of the elements of X and OH to the double bond.




The mechanism for halohydrin formation is similar to the mechanism for halogenation: addition of the electrophile X^+ (from X_2) to form a **bridged halonium ion**, followed by nucleophilic attack by H_2O from the back side on the three-membered ring (Mechanism 10.4). Even though X^- is formed in Step [1] of the mechanism, its concentration is small compared to H_2O (often the solvent), so H_2O and *not* X^- is the nucleophile.




Mechanism 10.4 Addition of X and OH—Halohydrin Formation

Step [1] Addition of the electrophile (X^+) to the π bond



- Four bonds are broken or formed in this step: the electron pair in the π bond and a lone pair on a halogen atom are used to form two new C–X bonds in the bridged halonium ion. The X–X bond is also cleaved heterolytically, forming X^- . This step is rate-determining.


Steps [2] and [3] Nucleophilic attack of H_2O and loss of a proton



- **Nucleophilic attack of H_2O** opens the halonium ion ring, forming a new C–O bond. Subsequent loss of a proton forms the neutral halohydrin.


Although the combination of Br_2 and H_2O effectively forms **bromohydrins** from alkenes, other reagents can also be used. Bromohydrins are also formed with *N*-**bromosuccinimide** (abbreviated as **NBS**) in **aqueous DMSO** [$(CH_3)_2S=O$]. NBS serves as a source of Br_2 , which then goes on to form a bromohydrin by the same reaction mechanism.

Recall from Section 7.8C that DMSO (dimethyl sulfoxide) is a polar aprotic solvent.




10.15A Stereochemistry and Regioselectivity of Halohydrin Formation

Because the bridged halonium ion ring is opened by backside attack of H₂O, addition of X and OH occurs in an **anti** fashion and **trans** products are formed.




Sample Problem 10.4 Draw the products of the following reaction, including stereochemistry.




Solution

The reagent (Br₂ + H₂O) adds the elements of Br and OH to a double bond in an **anti** fashion—that is, from **opposite** sides. To draw two products of anti addition: add Br from above and OH from below in one product; then add Br from below and OH from above in the other product. In this example, the two products are nonsuperimposable mirror images—**enantiomers**.



With unsymmetrical alkenes, two constitutional isomers are possible from addition of X and OH, but only one is formed. **The preferred product has the electrophile X⁺ bonded to the less substituted carbon atom**—that is, the carbon that has more H atoms to begin with in the reacting alkene. Thus, the **nucleophile (H₂O) bonds to the more substituted carbon**.



The electrophile (Br⁺) ends up on the **less substituted C**.

This result is reminiscent of the opening of epoxide rings with acids HZ (Z = a nucleophile), which we encountered in Section 9.15B. As in the opening of an epoxide ring, **nucleophilic attack occurs at the more substituted carbon end of the bridged halonium ion** because that carbon is better able to accommodate a partial positive charge in the transition state.

Halohydrin formation in an unsymmetrical alkene




Table 10.4 summarizes the characteristics of halohydrin formation.

Table 10.4 Summary: Conversion of Alkenes to Halohydrins

	Observation
Mechanism	<ul style="list-style-type: none"> The mechanism involves three steps. The rate-determining step forms a bridged halonium ion. No rearrangements can occur.
Regioselectivity	<ul style="list-style-type: none"> The electrophile X^+ bonds to the less substituted carbon.
Stereochemistry	<ul style="list-style-type: none"> Anti addition occurs.

Problem 10.28 Draw the products of each reaction and indicate their stereochemistry.



10.15B Halohydrins: Useful Compounds in Organic Synthesis

Because halohydrins are easily converted to epoxides by intramolecular S_N2 reaction (Section 9.6), they have been used in the synthesis of many naturally occurring compounds. Key steps in the synthesis of estrone, a female sex hormone, are illustrated in Figure 10.14.

10.16 Hydroboration–Oxidation

Hydroboration–oxidation is a two-step reaction sequence that converts an alkene to an alcohol.






Figure 10.14
The synthesis of estrone
from a chlorohydrin




- Chlorohydrin **B**, prepared from alkene **A** by addition of Cl and OH, is converted to epoxide **C** with base. **C** is converted to estrone in one step.

- **Hydroboration** is the addition of borane (BH_3) to an alkene, forming an alkylborane.
- **Oxidation** converts the C–B bond of the alkylborane to a C–O bond.

Hydroboration–oxidation results in **addition of H_2O** to an alkene.



Borane (BH_3) is a reactive gas that exists mostly as the dimer, diborane (B_2H_6). Borane is a strong **Lewis acid** that reacts readily with Lewis bases. For ease in handling in the laboratory, it is commonly used as a complex with tetrahydrofuran (THF).




Problem 10.29 Borane is sold for laboratory use as a complex with many other Lewis bases. Draw the Lewis acid–base complex that forms between BH_3 and each compound.

a. $(\text{CH}_3)_2\text{S}$ b. $(\text{CH}_3\text{CH}_2)_3\text{N}$ c. $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_3\text{P}$

10.16A Hydroboration

The first step in hydroboration–oxidation is **addition of the elements of H and BH_2** to the π bond of the alkene, forming an intermediate alkylborane.



Because syn addition to the double bond occurs and no carbocation rearrangements are observed, carbocations are *not* formed during hydroboration, as shown in Mechanism 10.5. The proposed mechanism involves a **concerted addition of H and BH_2 from the same side of the planar double bond**: the π bond and H– BH_2 bond are broken as two new σ bonds are formed. Because four atoms are involved, the transition state is said to be **four-centered**.



Mechanism 10.5 Addition of H and BH_2 —Hydroboration

One step The π bond and H– BH_2 bonds break as the C–H and C–B bonds form.






Figure 10.15
Conversion of BH_3 to a trialkylborane with three equivalents of $\text{CH}_2=\text{CH}_2$




- We often draw hydroboration as if addition stopped after one equivalent of alkene reacts with BH_3 . Instead, all three B–H bonds actually react with three equivalents of an alkene to form a trialkylborane. The term **organoborane** is used for any compound with a carbon–boron bond.

Because the alkylborane formed by reaction with one equivalent of alkene still has two B–H bonds, it can react with two more equivalents of alkene to form a trialkylborane. This is illustrated in Figure 10.15 for the reaction of $\text{CH}_2=\text{CH}_2$ with BH_3 .

Because only one B–H bond is needed for hydroboration, commercially available dialkylboranes having the general structure R_2BH are sometimes used instead of BH_3 . A common example is 9-borabicyclo[3.3.1]nonane (**9-BBN**). 9-BBN undergoes hydroboration in the same manner as BH_3 .



Hydroboration is regioselective. **With unsymmetrical alkenes, the boron atom bonds to the less substituted carbon atom.** For example, addition of BH_3 to propene forms an alkylborane with the B bonded to the terminal carbon atom.



Because H is more electronegative than B, the B–H bond is polarized to give boron a partial positive charge ($\text{H}^{\delta-} - \text{B}^{\delta+}$), making BH_2 the electrophile in hydroboration.

Steric factors explain this regioselectivity. The larger boron atom bonds to the less sterically hindered, more accessible carbon atom.

Electronic factors are also used to explain this regioselectivity. If bond breaking and bond making are not completely symmetrical, boron bears a partial negative charge in the transition state and carbon bears a partial positive charge. Because alkyl groups stabilize a positive charge, the more stable transition state has the partial positive charge on the more substituted carbon, as illustrated in Figure 10.16.

- In hydroboration, the boron atom bonds to the less substituted carbon.

Problem 10.30

What alkylborane is formed from hydroboration of each alkene?






Figure 10.16
Hydroboration of an
unsymmetrical alkene



10.16B Oxidation of the Alkylborane

Because alkylboranes react rapidly with water and spontaneously burn when exposed to the air, they are oxidized, without isolation, with basic hydrogen peroxide (H_2O_2 , HO^-). **Oxidation replaces the C–B bond with a C–O bond, forming a new OH group with retention of configuration;** that is, the OH group replaces the BH_2 group in the same position relative to the other three groups on carbon.




Thus, to draw the product of a hydroboration–oxidation reaction, keep in mind two stereochemical facts:

- Hydroboration occurs with syn addition.
- Oxidation occurs with retention of configuration.

The overall result of this two-step sequence is **syn addition of the elements of H and OH** to a double bond, as illustrated in Sample Problem 10.5. The OH group bonds to the less substituted carbon.


Sample Problem 10.5

Draw the product of the following reaction sequence, including stereochemistry.






Solution

In Step [1], **syn addition of BH_3 to the unsymmetrical alkene adds the BH_2 group to the less substituted carbon from above and below the planar double bond.** Two enantiomeric alkylboranes are formed. In Step [2], oxidation replaces the BH_2 group with OH in each enantiomer with **retention of configuration** to yield two alcohols that are also enantiomers.



Hydroboration–oxidation results in the **addition of H and OH in a syn fashion** across the double bond. The achiral alkene is converted to an equal mixture of two enantiomers—that is, a **racemic mixture of alcohols**.

Problem 10.31 Draw the products formed when each alkene is treated with BH_3 followed by $\text{H}_2\text{O}_2, \text{HO}^-$. Include the stereochemistry at all stereogenic centers.



Problem 10.32 What alkene can be used to prepare each alcohol as the exclusive product of a two-step hydroboration–oxidation sequence?






Table 10.5 summarizes the features of hydroboration–oxidation.

Hydroboration–oxidation is a very common method for adding H_2O across a double bond. One example is shown in the synthesis of **artemisinin** (or **qinghaosu**), the active component of **qinghao**, a Chinese herbal remedy used for the treatment of malaria (Figure 10.17).

Table 10.5 Summary: Hydroboration–Oxidation of Alkenes


	Observation
Mechanism	<ul style="list-style-type: none"> The addition of H and BH_2 occurs in one step. No rearrangements can occur.
Regioselectivity	<ul style="list-style-type: none"> The OH group bonds to the less substituted carbon atom.
Stereochemistry	<ul style="list-style-type: none"> Syn addition occurs. OH replaces BH_2 with retention of configuration.

Figure 10.17

An example of hydroboration–oxidation in synthesis



Artemisia annua, source of the antimalarial agent artemisinin



- The carbon atoms of artemisinin that come from alcohol **A** are indicated in red.


10.16C A Comparison of Hydration Methods

Hydration (H_2O , H^+) and hydroboration–oxidation (BH_3 followed by H_2O_2 , HO^-) both add the elements of H_2O across a double bond. Despite their similarities, these reactions often form different constitutional isomers, as shown in Sample Problem 10.6.


Sample Problem 10.6 Draw the product formed when $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$ is treated with (a) H_2O , H_2SO_4 ; and (b) BH_3 followed by H_2O_2 , HO^- .

Solution

With $\text{H}_2\text{O} + \text{H}_2\text{SO}_4$, electrophilic addition of H and OH places the **H atom on the less substituted carbon** of the alkene to yield a **2° alcohol**. In contrast, addition of BH_3 gives an alkylborane with the **BH_2 group on the less substituted terminal carbon** of the alkene. Oxidation replaces BH_2 by OH to yield a **1° alcohol**.



Problem 10.33 Draw the constitutional isomer formed when the following alkenes are treated with each set of reagents: [1] H_2O , H_2SO_4 ; or [2] BH_3 followed by H_2O_2 , OH^- .



10.17 Keeping Track of Reactions

Chapters 7–10 have introduced three basic kinds of organic reactions: **nucleophilic substitution**, **β elimination**, and **addition**. In the process, many specific reagents have been discussed and the stereochemistry that results from many different mechanisms has been examined. **How can we keep track of all the reactions?**


To make the process easier, **remember that most organic molecules undergo only one or two different kinds of reactions**. For example:

- Alkyl halides undergo substitution and elimination because they have good leaving groups.
- Alcohols also undergo substitution and elimination, but can do so only when OH is made into a good leaving group.
- Alkenes undergo addition because they have easily broken π bonds.

You must still learn many reaction details, and in truth, there is no one method to learn them. **You must practice these reactions over and over again, not by merely looking at them, but by writing them.** Some students do this by making a list of specific reactions for each functional group, and then rewriting them with different starting materials. Others make flash cards: index cards that have the starting material and reagent on one side and the product on the other. Whatever method you choose, **the details must become second nature**, much like the answers to simple addition problems, such as, what is the sum of $2 + 2$?

To solve this problem we must:

- Work backwards from the product by asking: What type of reactions introduce the functional groups in the product?
- Work forwards from the starting material by asking: What type of reactions does the starting material undergo?




In Chapter 11 we will learn about retrosynthetic analysis in more detail.

Working backwards from the product to determine the starting material from which it is made is called *retrosynthetic analysis*.

We know reactions that answer each of these questions.


Working backwards:

- [1] 1,2-Dibromocyclohexane, a vicinal dibromide, can be prepared by the addition of Br_2 to **cyclohexene**.




Working forwards:

- [2] Cyclohexanol can undergo acid-catalyzed dehydration to form **cyclohexene**.




A **reactive intermediate** is an unstable intermediate like a carbocation, which is formed during the conversion of a stable starting material to a stable product. A **synthetic intermediate** is a stable compound that is the product of one step and the starting material of another in a multistep synthesis.

Cyclohexene is called a **synthetic intermediate**, or simply an **intermediate**, because it is the **product of one step and the starting material of another**. We now have a two-step sequence to convert cyclohexanol to 1,2-dibromocyclohexane, and the synthesis is complete. Take note of the central role of the alkene in this synthesis.



Problem 10.35

Devise a synthesis of each compound from the indicated starting material.



KEY CONCEPTS

Alkenes


General Facts About Alkenes

- Alkenes contain a carbon-carbon double bond consisting of a stronger σ bond and a weaker π bond. Each carbon is sp^2 hybridized and trigonal planar (10.1).
- Alkenes are named using the suffix **-ene** (10.3).
- Alkenes with different groups on each end of the double bond exist as a pair of diastereomers, identified by the prefixes *E* and *Z* (10.3B).
- Alkenes have weak intermolecular forces, giving them low mp's and bp's, and making them water insoluble. A *cis* alkene is more polar than a *trans* alkene, giving it a slightly higher boiling point (10.4).
- Because a π bond is electron rich and much weaker than a σ bond, alkenes undergo addition reactions with electrophiles (10.8).

Stereochemistry of Alkene Addition Reactions (10.8)


A reagent XY adds to a double bond in one of three different ways:

- Syn addition**—X and Y add from the same side.




- Syn addition occurs in **hydroboration**.

- Anti addition**—X and Y add from opposite sides.



- Anti addition occurs in **halogenation** and **halohydrin formation**.


- Both syn and anti addition** occur when carbocations are intermediates.



- Syn and anti addition occur in **hydrohalogenation** and **hydration**.



Addition Reactions of Alkenes

[1] Hydrohalogenation—Addition of HX (X = Cl, Br, I) (10.9–10.11)




- The mechanism has two steps.
- Carbocations are formed as intermediates.
- Carbocation rearrangements are possible.
- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- Syn and anti addition occur.

[2] Hydration and related reactions (Addition of H₂O or ROH) (10.12)




For both reactions:


- The mechanism has three steps.
- Carbocations are formed as intermediates.
- Carbocation rearrangements are possible.
- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- Syn and anti addition occur.

[3] Halogenation (Addition of X_2 ; $X = \text{Cl}$ or Br) (10.13–10.14)

- The mechanism has two steps.
- Bridged halonium ions are formed as intermediates.
- No rearrangements can occur.
- Anti addition occurs.

[4] Halohydrin formation (Addition of OH and X ; $\text{X} = \text{Cl}, \text{Br}$) (10.15)

- The mechanism has three steps.
- Bridged halonium ions are formed as intermediates.
- No rearrangements can occur.
- X bonds to the less substituted C.
- Anti addition occurs.
- NBS in DMSO and H_2O adds Br and OH in the same fashion.

[5] Hydroboration–oxidation (Addition of H_2O) (10.16)



- Hydroboration has a one-step mechanism.
- No rearrangements can occur.
- OH bonds to the less substituted C.
- Syn addition of H_2O results.

PROBLEMS

Degrees of Unsaturation









10.36 Calculate the number of degrees of unsaturation for each molecular formula.

- a. C_3H_4 c. $\text{C}_{40}\text{H}_{56}$ e. $\text{C}_{10}\text{H}_{16}\text{O}_2$ g. $\text{C}_8\text{H}_9\text{ClO}$ i. $\text{C}_7\text{H}_{11}\text{N}$
 b. C_6H_8 d. $\text{C}_8\text{H}_8\text{O}$ f. $\text{C}_8\text{H}_9\text{Br}$ h. $\text{C}_7\text{H}_9\text{Br}$ j. $\text{C}_4\text{H}_8\text{BrN}$

10.37 How many rings and π bonds does a compound with molecular formula $\text{C}_{10}\text{H}_{14}$ possess? List all possibilities.10.38 The fertility drug clomiphene (trade name Clomid) is sold as a mixture of diastereomers, enclomiphene and zuclomiphene. Designate each alkene as an *E* or *Z* isomer. What is wrong with the statement, “Clomid is a mixture of two isomers, 30% *cis* and the remainder *trans*”?

Nomenclature and Stereochemistry

10.39 Give the IUPAC name for each compound.




10.40 Give the structure corresponding to each name.

- | | |
|------------------------------------|---|
| a. (3 <i>E</i>)-4-ethyl-3-heptene | e. (2 <i>Z</i>)-3-isopropyl-2-heptene |
| b. 3,3-dimethylcyclopentene | f. <i>cis</i> -3,4-dimethylcyclopentene |
| c. <i>cis</i> -4-octene | g. <i>trans</i> -2-heptene |
| d. 4-vinylcyclopentene | h. 1-isopropyl-4-propylcyclohexene |

10.41 (a) Draw all possible stereoisomers of 4-methyl-2-nonene, and name each isomer, including its *E,Z* and *R,S* prefixes. (b) Label two pairs of enantiomers. (c) Label four pairs of diastereomers.

10.42 (a) Draw the structure of (1*E*,4*R*)-1,4-dimethylcyclodecene. (b) Draw the enantiomer and name it, including its *E,Z* and *R,S* prefixes. (c) Draw two diastereomers and name them, including the *E,Z* and *R,S* prefixes.


10.43 Now that you have learned how to name alkenes in Section 10.3, name each of the following epoxides as an alkene oxide, as described in Section 9.3.



10.44 Each of the following names is incorrect. Explain why it is incorrect and give the correct IUPAC name.

- | | | |
|--------------------------------------|--|-------------------------|
| a. 2-butyl-3-methyl-1-pentene | d. 5-methylcyclohexene | g. 1-cyclohexen-4-ol |
| b. (<i>Z</i>)-2-methyl-2-hexene | e. 4-isobutyl-2-methylcyclohexene | h. 3-ethyl-3-octen-5-ol |
| c. (<i>E</i>)-1-isopropyl-1-butene | f. 1- <i>sec</i> -butyl-2-cyclopentene | |

10.45




Bongkreic acid is a toxic compound produced by *Pseudomonas cocovenenans*, and isolated from a mold that grows on bongkrek, a fermented Indonesian coconut dish. (a) Label each double bond in bongkreic acid as *E* or *Z*. (b) Label each tetrahedral stereogenic center as *R* or *S*. (c) How many stereoisomers are possible for bongkreic acid?


10.46 Draw all stereoisomers having molecular formula C_6H_{12} that contain one double bond and a five-carbon chain with a one-carbon branch. Name each compound, including its *E,Z* or *R,S* designation when necessary.

Lipids

10.47 Although naturally occurring unsaturated fatty acids have the *Z* configuration, elaidic acid, a C_{18} fatty acid having an *E* double bond, is present in processed foods such as margarine and cooking oils. Predict how the melting point of elaidic acid compares with the melting points of stearic and oleic acids (Table 10.2).



10.48 Eleostearic acid is an unsaturated fatty acid obtained from the seeds of the tung oil tree (*Aleurites fordii*), a deciduous tree native to China. (a) Draw the structure of a stereoisomer that has a higher melting point than eleostearic acid. (b) Draw the structure of a stereoisomer that has a lower melting point.



10.49 (a) Draw two possible triacylglycerols formed from one molecule of stearic acid and two molecules of oleic acid. (b) One of these molecules contains a tetrahedral stereogenic center. Draw both enantiomers, and label the stereogenic center as *R* or *S*.

Energy Diagram and ΔH° Calculations

10.50 Draw an energy diagram for the two-step mechanism for the addition of Br_2 to $CH_2=CH_2$ to form 1,2-dibromoethane. Draw the structure of the transition state for each step.

10.51 By using the bond dissociation energies in Appendix C, calculate ΔH° for the addition of HCl and HI to ethylene to form chloroethane and iodoethane, respectively. Assuming entropy changes for both reactions are similar, which reaction has the larger K_{eq} ?

Reactions of Alkenes


10.52 Draw the products formed when cyclohexene is treated with each reagent.

- a. HBr
 b. HI
 c. H₂O, H₂SO₄
 d. CH₃CH₂OH, H₂SO₄
 e. Cl₂
 f. Br₂, H₂O
 g. NBS (aqueous DMSO)
 h. [1] BH₃; [2] H₂O₂, HO⁻
 i. [1] 9-BBN; [2] H₂O₂, HO⁻


10.53 Repeat Problem 10.52 with (CH₃)₂C=CH₂ as the starting material.

10.54 Draw the product formed when 1-butene is treated with each reagent: (a) Br₂; (b) Br₂ in H₂O; (c) Br₂ in CH₃OH.


10.55 What alkene can be used to prepare each alkyl halide or dihalide as the exclusive or major product of an addition reaction?



10.56 Which alcohols can be prepared as a single product by hydroboration-oxidation of an alkene? Which alcohols can be prepared as a single product by the acid-catalyzed addition of H₂O to an alkene?




10.57 Draw the constitutional isomer formed in each reaction.




10.58 What three alkenes (excluding stereoisomers) can be used to prepare 3-chloro-3-methylhexane by addition of HCl?

10.59 Draw all stereoisomers formed in each reaction.



10.60 Draw the products of each reaction, including stereoisomers.



10.61 The elements of Br and Cl are added to a double bond with Br₂ + NaCl. Draw the product formed when an unsymmetrical alkene such as 2-methyl-1-propene is used as the starting material.

10.62 (a) Which diastereomer of 4-octene yields a mixture of two enantiomers, (4*R*,5*R*)- and (4*S*,5*S*)-4,5-dibromooctane on reaction with Br₂? (b) Which diastereomer of 4-octene yields a single meso compound, (4*R*,5*S*)-4,5-dibromooctane?


10.63 Using *cis*- and *trans*-3-hexene, demonstrate that the addition of HCl is not a stereospecific reaction. Draw the structure of the stereoisomers formed from each alkene.

10.64 Explain each of the following differences in alkene reactivity in electrophilic addition reactions.


- $\text{C}_6\text{H}_5\text{CH}=\text{CHC}_6\text{H}_5$ reacts with HBr faster than $\text{CH}_3\text{CH}=\text{CHCH}_3$, even though both compounds are 1,2-disubstituted alkenes.
- When treated with H_2O in the presence of acid, $\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}_2\text{OCH}_3$ reacts more slowly than $\text{CH}_2=\text{C}(\text{CH}_3)_2$.

Mechanisms


10.65 Draw a stepwise mechanism for each reaction.



10.66 Draw a stepwise mechanism for each reaction.



10.67 Draw a stepwise mechanism that shows how all three alcohols are formed from the bicyclic alkene.




10.68 Less stable alkenes can be isomerized to more stable alkenes by treatment with strong acid. For example, 2,3-dimethyl-1-butene is converted to 2,3-dimethyl-2-butene when treated with H_2SO_4 . Draw a stepwise mechanism for this isomerization process.

10.69 When 1,3-butadiene ($\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$) is treated with HBr , two constitutional isomers are formed, $\text{CH}_3\text{CHBrCH}=\text{CH}_2$ and $\text{BrCH}_2\text{CH}=\text{CHCH}_3$. Draw a stepwise mechanism that accounts for the formation of both products.

10.70 Explain why the addition of HBr to alkenes **A** and **C** is regioselective, forming addition products **B** and **D**, respectively.




10.71 Bromoetherification, the addition of the elements of Br and OR to a double bond, is a common method for constructing rings containing oxygen atoms. This reaction has been used in the synthesis of the polyether antibiotic monensin (Problem 21.40). Draw a stepwise mechanism for the following intramolecular bromoetherification reaction.




Synthesis


10.72 Devise a synthesis of each product from the given starting material. More than one step is required.




10.73 Devise a synthesis of each compound from cyclohexene as the starting material. More than one step is needed.



10.74 Devise a synthesis of each compound from $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$. You may use any needed organic or inorganic reagents. More than one step may be required.




10.75 Devise a synthesis of each epoxide (**B** and **C**) from alcohol **A**. (Hint: Determine what alkene is needed to make each epoxide, and begin each synthesis with a reaction that yields the needed alkene from an elimination reaction.)




Challenge Problems


10.76 Alkene **A** can be isomerized to isocomene, a natural product isolated from goldenrod, by treatment with TsOH. Draw a stepwise mechanism for this conversion. (Hint: Look for a carbocation rearrangement.)




10.77 Lactones, cyclic esters such as compound **A**, are prepared by halolactonization, an addition reaction to an alkene. For example, iodolactonization of **B** forms lactone **C**, a key intermediate in the synthesis of prostaglandin $\text{PGF}_{2\alpha}$ (Section 4.15). Draw a stepwise mechanism for this addition reaction.



10.78 Like other electrophiles, carbocations add to alkenes to form new carbocations, which can then undergo substitution or elimination reactions depending on the reaction conditions. With this in mind, consider the following reactions of nerol, a natural product isolated from lemon grass and other plant sources. Treatment of nerol with TsOH forms α -terpineol as the major product, whereas treatment of nerol with chlorosulfonic acid, HSO_3Cl , forms a constitutional isomer, α -cyclogeraniol. Write stepwise mechanisms for both processes. Each mechanism involves the addition of an electrophile—a carbocation—to a double bond.



10.79 Draw a stepwise mechanism for the following reaction. This reaction combines two processes together: the opening of an epoxide ring with a nucleophile and the addition of an electrophile to a carbon-carbon double bond. (Hint: Begin the mechanism by protonating the epoxide ring.)



Alkynes

11

- 11.1 Introduction
- 11.2 Nomenclature
- 11.3 Physical properties
- 11.4 Interesting alkynes
- 11.5 Preparation of alkynes
- 11.6 Introduction to alkyne reactions
- 11.7 Addition of hydrogen halides
- 11.8 Addition of halogen
- 11.9 Addition of water
- 11.10 Hydroboration–oxidation
- 11.11 Reaction of acetylide anions
- 11.12 Synthesis




Ethynylestradiol is a synthetic compound whose structure closely resembles the carbon skeleton of female estrogen hormones. Because it is more potent than its naturally occurring analogues, it is a component of several widely used oral contraceptives. Ethynylestradiol and related compounds with similar biological activity contain a carbon–carbon triple bond. In Chapter 11 we learn about alkynes, hydrocarbons that contain triple bonds.

In Chapter 11 we continue our focus on organic molecules with electron-rich functional groups by examining *alkynes*, compounds that contain a carbon–carbon triple bond. Like alkenes, **alkynes are nucleophiles with easily broken π bonds**, and as such, they undergo **addition** reactions with electrophilic reagents.

Alkynes also undergo a reaction that has no analogy in alkene chemistry. Because a C–H bond of an alkyne is more acidic than a C–H bond in an alkene or an alkane, alkynes are readily deprotonated with strong base. The resulting nucleophiles react with electrophiles to form new carbon–carbon σ bonds, so that complex molecules can be prepared from simple starting materials. The study of alkynes thus affords an opportunity to learn more about organic synthesis.

11.1 Introduction


Alkynes contain a carbon–carbon triple bond. A **terminal alkyne** has the triple bond at the end of the carbon chain, so that a hydrogen atom is directly bonded to a carbon atom of the triple bond. An **internal alkyne** has a carbon atom bonded to each carbon atom of the triple bond.




An alkyne has the general molecular formula $\text{C}_n\text{H}_{2n-2}$, giving it four fewer hydrogens than the maximum number possible. Because every degree of unsaturation removes two hydrogens, a **triple bond introduces two degrees of unsaturation**.

Problem 11.1 Draw structures for the three alkynes having molecular formula C_5H_8 and classify each as an internal or terminal alkyne.

Each carbon of a triple bond is *sp* hybridized and **linear**, and all bond angles are **180°** (Section 1.9C).




The triple bond of an alkyne consists of one σ bond and two π bonds.



- The σ bond is formed by end-on overlap of the two *sp* hybrid orbitals.
- Each π bond is formed by side-by-side overlap of two *2p* orbitals.


Bond dissociation energies of the C–C bonds in ethylene (one σ and one π bond) and acetylene (one σ and two π bonds) can be used to estimate the strength of the second π bond of the triple bond. If

we assume that the σ bond and first π bond in acetylene are similar in strength to the σ and π bonds in ethylene (368 and 267 kJ/mol, respectively), then the second π bond is worth 202 kJ/mol.




- Both π bonds of a C–C triple bond are weaker than a C–C σ bond, making them much more easily broken. As a result, alkynes undergo many addition reactions.
- Alkynes are more polarizable than alkenes because the electrons in their π bonds are more loosely held.

Skeletal structures for alkynes may look somewhat unusual, but they follow the customary convention: a carbon atom is located at the intersection of any two lines and at the end of any line; thus,




Like trans cycloalkenes, cycloalkynes with small rings are unstable. The carbon chain must be long enough to connect the two ends of the triple bond without introducing too much strain. Cyclooctyne is the smallest isolated cycloalkyne, though it decomposes upon standing at room temperature after a short time.



To accommodate the triple bond in a ring, bending occurs around the sp hybridized C's, destabilizing the molecule.

Problem 11.2

Santalbic acid, a fatty acid isolated from the seeds of the sandalwood tree, is an unusual fatty acid that contains a carbon–carbon triple bond. What orbitals are used to form each of the three indicated single bonds in santalbic acid? Rank these σ bonds in order of increasing bond strength.



Problem 11.3


Would you predict an internal or terminal alkyne to be more stable? Why?

11.2 Nomenclature

Alkynes are named in the same way that alkenes were named in Section 10.3.


- In the IUPAC system, change the *-ane* ending of the parent alkane to the suffix *-yne*.
- Choose the longest carbon chain that contains both atoms of the triple bond and number the chain to give the triple bond the lower number.
- Compounds with two triple bonds are named as *diynes*, those with three are named as *triyne*s, and so forth.
- Compounds with both a double and a triple bond are named as *enynes*. The chain is numbered to give the first site of unsaturation (either C=C or C≡C) the lower number.

Sample Problem 11.1 Give the IUPAC name for the following alkyne.



Solution


[1] Find the longest chain that contains both carbons of the triple bond.



8 C's in the longest chain

octane ----> octyne

[2] Number the long chain; then name and number the substituents.




two methyl groups at C6

Answer: 6,6-dimethyl-3-octyne

The simplest alkyne, $\text{HC}\equiv\text{CH}$, named in the IUPAC system as **ethyne**, is more often called **acetylene**, its common name. The two-carbon alkyl group derived from acetylene is called an **ethynyl group** ($\text{HC}\equiv\text{C}-$). Examples of alkyne nomenclature are shown in Figure 11.1.

Problem 11.4 Give the IUPAC name for each compound.

- $\text{H}-\text{C}\equiv\text{C}-\text{CH}_2\text{C}(\text{CH}_2\text{CH}_2\text{CH}_3)_3$
- $\text{CH}_3\text{C}\equiv\text{CC}(\text{CH}_3)\text{CICH}_2\text{CH}_3$
- $\text{CH}_2=\text{CHCH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{C}\equiv\text{CC}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{CH}_3$



Problem 11.5 Give the structure corresponding to each of the following names.

- trans*-2-ethynylcyclopentanol
- 4-*tert*-butyl-5-decyne
- 3-methylcyclononyne

11.3 Physical Properties

The physical properties of alkynes resemble those of hydrocarbons having a similar shape and molecular weight.

- Alkynes have low melting points and boiling points.
- Melting points and boiling points increase as the number of carbons increases.
- Alkynes are soluble in organic solvents and insoluble in water.

Problem 11.6 Explain why an alkyne often has a slightly higher boiling point than an alkene of similar molecular weight. For example, the bp of 1-pentyne is 39 °C, and the bp of 1-pentene is 30 °C.

11.4 Interesting Alkynes

Acetylene, $\text{HC}\equiv\text{CH}$, is a colorless gas with an ethereal odor that burns in oxygen to form CO_2 and H_2O . Because the combustion of acetylene releases more energy per mole of product formed than other hydrocarbons, it burns with a very hot flame, making it an excellent fuel for welding torches.

Ethinylestradiol, the molecule that opened Chapter 11, and **norethindrone** are two components of oral contraceptives that contain a carbon–carbon triple bond (Figure 11.2). Both molecules are synthetic analogues of the naturally occurring female hormones estradiol and progesterone, but

Figure 11.1
Examples of alkyne nomenclature








Figure 11.2 How oral contraceptives work




Monthly cycles of hormones from the pituitary gland cause ovulation, the release of an egg from an ovary. To prevent pregnancy, the two synthetic hormones in many oral contraceptives have different effects on the female reproductive system.

A: The elevated level of **ethynylestradiol**, a synthetic estrogen, “fools” the pituitary gland into thinking a woman is pregnant, so ovulation does not occur.

B: The elevated level of **norethindrone**, a synthetic progesterone, stimulates the formation of a thick layer of mucus in the cervix, making it difficult for sperm to reach the uterus.

are more potent so they can be administered in lower doses. Most oral contraceptives contain two of these synthetic hormones. They act by artificially elevating hormone levels in a woman, thereby preventing pregnancy.

Naturally occurring female sex hormones



Two other synthetic hormones with alkynyl appendages are **RU 486** and **levonorgestrel**. RU 486 blocks the effects of progesterone, and because of this, prevents implantation of a fertilized egg. RU 486 is used to induce abortions within the first few weeks of pregnancy. Levonorgestrel interferes with ovulation, and so it prevents pregnancy if taken within a few days of unprotected sex.





Figure 11.3
Histrionicotoxin





- Histrionicotoxin is a defensive toxin that protects *Dendrobates histrionicus* from potential predators. These small “poison dart” frogs inhabit the moist humid floor of tropical rainforests, and are commonly found in western Ecuador and Colombia. Histrionicotoxin acts by interfering with nerve transmission in mammals, resulting in prolonged muscle contraction.

Histrionicotoxin is a diyne isolated in small quantities from the skin of *Dendrobates histrionicus*, a colorful South American frog (Figure 11.3). This toxin, secreted by the frog as a natural defense mechanism, was used as a poison on arrow tips by the Choco tribe of South America.


11.5 Preparation of Alkynes

Alkynes are prepared by elimination reactions, as discussed in Section 8.10. **A strong base removes two equivalents of HX from a vicinal or geminal dihalide to yield an alkyne** by two successive E2 eliminations.



Because vicinal dihalides are synthesized by adding halogens to alkenes, an alkene can be converted to an alkyne by the two-step process illustrated in Sample Problem 11.2.


Sample Problem 11.2 Convert alkene **A** into alkyne **B** by a stepwise method.



Solution

A two-step method is needed:

- **Addition of X₂** forms a vicinal dihalide.
- **Elimination** of two equivalents of HX forms two π bonds.



- **This two-step process introduces one degree of unsaturation:** an alkene with one π bond is converted to an alkyne with two π bonds.

Problem 11.7 Convert each compound to 1-hexyne, $\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$.


- a. $\text{Br}_2\text{CH}(\text{CH}_2)_4\text{CH}_3$ b. $\text{CH}_2=\text{CCl}(\text{CH}_2)_3\text{CH}_3$ c. $\text{CH}_2=\text{CH}(\text{CH}_2)_3\text{CH}_3$

11.6 Introduction to Alkyne Reactions

All reactions of alkynes occur because they contain **easily broken π bonds** or, in the case of terminal alkynes, an **acidic, sp hybridized C–H bond**.

11.6A Addition Reactions

Like alkenes, **alkynes undergo addition reactions because they contain weak π bonds**. Two sequential reactions take place: addition of one equivalent of reagent forms an alkene, which then adds a second equivalent of reagent to yield a product having **four new bonds**.




The oxidation and reduction of alkynes, reactions that also involve addition, are discussed in Chapter 12.

Alkynes are electron rich, as shown in the electrostatic potential map of acetylene in Figure 11.4. The two π bonds form a cylinder of electron density between the two sp hybridized carbon atoms, and this exposed electron density makes a triple bond nucleophilic. As a result, **alkynes react with electrophiles**. Four addition reactions are discussed in Chapter 11 and illustrated in Figure 11.5 with 1-butyne as the starting material.

11.6B Terminal Alkynes—Reaction as an Acid


Because **sp hybridized C–H bonds are more acidic than sp^2 and sp^3 hybridized C–H bonds**, terminal alkynes are readily deprotonated with strong base in a Brønsted–Lowry acid–base reaction. The resulting anion is called an **acetylide anion**.



What bases can be used for this reaction? Because an acid–base equilibrium favors the weaker acid and base, only **bases having conjugate acids with pK_a values higher than the terminal**

Figure 11.4


Electrostatic potential map of acetylene



- The red electron-rich region is located between the two carbon atoms, forming a cylinder of electron density.

Figure 11.5


Four addition reactions of 1-butyne



Recall from Section 2.5D that the acidity of a C-H bond increases as the percent s-character of C increases. Thus, the following order of relative acidity results:
 $\text{C}_{\text{sp}^3}\text{-H} < \text{C}_{\text{sp}^2}\text{-H} < \text{C}_{\text{sp}}\text{-H}$.

alkyne—that is, $\text{p}K_{\text{a}}$ values > 25—are strong enough to form a significant concentration of acetylide anion. As shown in Table 11.1, NH_2^- and H^- are strong enough to deprotonate a terminal alkyne, but OH^- and OR^- are not.

Why is this reaction useful? The acetylide anions formed by deprotonating terminal alkynes are **strong nucleophiles** that can react with a variety of electrophiles, as shown in Section 11.11.



Problem 11.8

Which bases can deprotonate acetylene? The $\text{p}K_{\text{a}}$ values of the conjugate acids are given in parentheses.


- a. CH_3NH^- ($\text{p}K_{\text{a}} = 40$) b. CO_3^{2-} ($\text{p}K_{\text{a}} = 10.2$) c. $\text{CH}_2=\text{CH}^-$ ($\text{p}K_{\text{a}} = 44$) d. $(\text{CH}_3)_3\text{CO}^-$ ($\text{p}K_{\text{a}} = 18$)

Table 11.1 A Comparison of Bases for Alkyne Deprotonation

	Base	$\text{p}K_{\text{a}}$ of the conjugate acid
These bases are strong enough to deprotonate an alkyne.	NH_2^- H^-	38 35
These bases are not strong enough to deprotonate an alkyne.	OH^- OR^-	15.7 15.5–18


11.7 Addition of Hydrogen Halides

Alkynes undergo **hydrohalogenation, the addition of hydrogen halides, HX** ($\text{X} = \text{Cl, Br, I}$). Two equivalents of HX are usually used: addition of one mole forms a **vinyl halide**, which then reacts with a second mole of HX to form a **geminal dihalide**.




Addition of HX to an alkyne is another example of **electrophilic addition**, because the electrophilic (H) end of the reagent is attracted to the electron-rich triple bond.

- With two equivalents of HX, both H atoms bond to the *same* carbon.
- With a terminal alkyne, both H atoms bond to the *terminal* carbon; that is, the hydrohalogenation of alkynes follows Markovnikov's rule.



- With only one equivalent of HX, the reaction stops with formation of the vinyl halide.




One currently accepted mechanism for the addition of two equivalents of HX to an alkyne involves **two steps for each addition of HX**: addition of H^+ (from HX) to form a carbocation, followed by nucleophilic attack of X^- . Mechanism 11.1 illustrates the addition of HBr to 1-butyne to yield 2,2-dibromobutane. Each two-step mechanism is similar to the two-step addition of HBr to *cis*-2-butene discussed in Section 10.9.




Mechanism 11.1 Electrophilic Addition of HX to an Alkyne

Part [1] Addition of HBr to form a vinyl halide



- The π bond attacks the H atom of HBr to form a new C-H bond, generating a **vinyl carbocation**. Addition follows Markovnikov's rule: H^+ adds to the less substituted carbon atom to form the **more substituted, more stable carbocation**. Nucleophilic attack of Br^- then forms a vinyl bromide; one mole of HBr has now been added.

Part [2] Addition of HBr to form a geminal dihalide




- **The second addition of HBr occurs in the same two-step manner.** Addition of H^+ to the π bond of the vinyl bromide generates a carbocation. Nucleophilic attack of Br^- then forms a geminal dibromide (2,2-dibromobutane), and two moles of HBr have now been added.


The formation of both carbocations (in Steps [1] and [3]) deserves additional scrutiny. **The vinyl carbocation formed in Step [1] is sp hybridized and therefore less stable than a 2° sp^3**

Because of the instability of a vinyl carbocation, other mechanisms for HX addition that avoid formation of a discrete carbocation have been proposed. It is likely that more than one mechanism occurs, depending in part on the identity of the alkyne substrate.


hybridized carbocation (Section 7.18). This makes electrophilic addition of HX to an alkyne *slower* than electrophilic addition of HX to an alkene, even though alkynes are more polarizable and have more loosely held π electrons than alkenes.



In Step [3] two carbocations are possible but only one is formed. Markovnikov addition in Step [3] places the H on the terminal carbon (C1) to form the more substituted carbocation **A**, rather than the less substituted carbocation **B**. Because the more stable carbocation is formed faster—another example of the Hammond postulate—carbocation **A** must be more stable than carbocation **B**.




Why is carbocation **A**, having a positive charge on a carbon that also has a Br atom, more stable? Shouldn't the electronegative Br atom withdraw electron density from the positive charge, and thus destabilize it? It turns out that **A is stabilized by resonance** but **B** is not. Two resonance structures can be drawn for carbocation **A**, but only one Lewis structure can be drawn for carbocation **B**.




- Resonance stabilizes a molecule by delocalizing charge and electron density.
- Thus, halogens stabilize an adjacent positive charge by resonance.


Markovnikov's rule applies to the addition of HX to vinyl halides because **addition of H^+ forms a resonance-stabilized carbocation**. As a result, addition of each equivalent of HX to a triple bond forms the more stable carbocation, so that both H atoms bond to the less substituted C.

Problem 11.9 Draw the organic products formed when each alkyne is treated with two equivalents of HBr.




Problem 11.10 Draw an additional resonance structure for each cation.





Internal alkynes undergo hydration with concentrated acid, whereas terminal alkynes require the presence of an additional Hg^{2+} catalyst—usually $HgSO_4$ —to yield methyl ketones by **Markovnikov addition of H_2O** .

Examples



Markovnikov addition of H_2O


H adds to the terminal C.

Because an enol contains both a $C=C$ and a hydroxy group, the name **enol** comes from **alkene** + **alcohol**.

$HgSO_4$ is often used in the hydration of internal alkynes as well, because hydration can be carried out under milder reaction conditions.

Let's first examine the conversion of a general enol **A** to the carbonyl compound **B**. **A** and **B** are called **tautomers**: **A** is the **enol form** and **B** is the **keto form** of the tautomer.

- **Tautomers** are constitutional isomers that differ in the location of a double bond and a hydrogen atom. Two tautomers are in equilibrium with each other.



Tautomers differ in the position of a double bond and a hydrogen atom. In Chapter 23 an in-depth discussion of keto-enol tautomers is presented.


- An enol tautomer has an $O-H$ group bonded to a $C=C$.
- A keto tautomer has a $C=O$ and an additional $C-H$ bond.

Equilibrium favors the keto form largely because a $C=O$ is much stronger than a $C=C$. Tautomerization, the process of converting one tautomer into another, is catalyzed by both acid




Mechanism 11.3 Tautomerization in Acid

Step [1] Protonation of the enol double bond



Step [2] Deprotonation of the OH group




and base. Under the strongly acidic conditions of hydration, tautomerization of the enol to the keto form occurs rapidly by a two-step process: **protonation**, followed by **deprotonation** as shown in Mechanism 11.3.

Hydration of an internal alkyne with strong acid forms an enol by a mechanism similar to that of the acid-catalyzed hydration of an alkene (Section 10.12). Mechanism 11.4 illustrates the hydration of 2-butyne with H_2O and H_2SO_4 . Once formed, the enol then tautomerizes to the more stable keto form by protonation followed by deprotonation.




Mechanism 11.4 Hydration of an Alkyne

Step [1] Addition of the electrophile (H^+) to a π bond




- Addition of H^+ (from H_3O^+) forms an sp hybridized **vinyl carbocation**.

Steps [2] and [3] Nucleophilic attack of H_2O and loss of a proton



- **Nucleophilic attack of H_2O** on the carbocation followed by loss of a proton forms the enol.


Steps [4] and [5] Tautomerization



- **Tautomerization of the enol to the keto form** occurs by protonation of the double bond to form a carbocation. Loss of a proton from this **resonance-stabilized carbocation** generates the more stable keto form.


Sample Problem 11.3

Draw the enol intermediate and the ketone product formed in the following reaction.




Solution

First, form the enol by adding H_2O to the triple bond with the **H bonded to the less substituted terminal carbon**.




To convert the enol to the keto tautomer, add a proton to the $\text{C}=\text{C}$ and remove a proton from the OH group. In tautomerization, the $\text{C}-\text{OH}$ bond is converted to a $\text{C}=\text{O}$, and a new $\text{C}-\text{H}$ bond is formed on the other enol carbon.




- The overall result is the addition of H_2O to a triple bond to form a ketone.

Problem 11.13 Draw the keto tautomer of each enol.



Problem 11.14 What two enols are formed when 2-pentyne is treated with H_2O , H_2SO_4 , and HgSO_4 ? Draw the ketones formed from these enols after tautomerization.

Problem 11.15 (a) Draw two different enol tautomers of 2-methylcyclohexanone. (b) Draw two constitutional isomers that are not tautomers, but contain a $\text{C}=\text{C}$ and an OH group.




2-methylcyclohexanone

11.10 Hydroboration–Oxidation


Hydroboration–oxidation is a two-step reaction sequence that converts an alkyne to a carbonyl compound.

Hydroboration–oxidation



- Addition of borane forms an organoborane.
- Oxidation with basic H_2O_2 forms an enol.
- Tautomerization of the enol forms a carbonyl compound.
- The overall result is addition of H_2O to a triple bond.

Hydroboration–oxidation of an *internal* alkyne forms a ketone. **Hydroboration of a terminal alkyne adds BH_2 to the less substituted, terminal carbon.** After oxidation to the enol, tautomerization yields an **aldehyde**, a carbonyl compound having a hydrogen atom bonded to the carbonyl carbon.




Hydration (H_2O , H_2SO_4 , and HgSO_4) and **hydroboration–oxidation** (BH_3 followed by H_2O_2 , HO^-) both **add the elements of H_2O across a triple bond**. Sample Problem 11.4 shows that different constitutional isomers are formed from terminal alkynes in these two reactions despite their similarities.

Sample Problem 11.4

Draw the product formed when $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CH}$ is treated with each of the following sets of reagents: (a) H_2O , H_2SO_4 , HgSO_4 ; and (b) BH_3 , followed by H_2O_2 , HO^- .

Solution



(a) With $\text{H}_2\text{O} + \text{H}_2\text{SO}_4 + \text{HgSO}_4$, electrophilic addition of H and OH places the **H atom on the less substituted carbon** of the alkyne to form a **ketone** after tautomerization. (b) In contrast, addition of BH_3 places the **BH_2 group on the less substituted terminal carbon** of the alkyne. Oxidation and tautomerization yield an **aldehyde**.



- Addition of H_2O using H_2O , H_2SO_4 , and HgSO_4 forms methyl ketones from terminal alkynes.
- Addition of H_2O using BH_3 , then H_2O_2 , HO^- forms aldehydes from terminal alkynes.


Problem 11.16

Draw the products formed when the following alkynes are treated with each set of reagents: [1] H_2O , H_2SO_4 , HgSO_4 ; or [2] BH_3 followed by H_2O_2 , HO^- .




11.11 Reaction of Acetylide Anions

Terminal alkynes are readily converted to acetylide anions with strong bases such as NaNH_2 and NaH . These anions are strong nucleophiles, capable of reacting with electrophiles such as alkyl halides and epoxides.




11.11A Reaction of Acetylide Anions with Alkyl Halides

Acetylide anions react with unhindered alkyl halides to yield products of nucleophilic substitution.



Because acetylide anions are strong nucleophiles, the mechanism of nucleophilic substitution is $\text{S}_\text{N}2$, and thus the reaction is fastest with CH_3X and 1° alkyl halides. Terminal alkynes (Reaction [1]) or internal alkynes (Reaction [2]) can be prepared depending on the identity of the acetylide anion.




- Nucleophilic substitution with acetylide anions forms new carbon-carbon bonds.

Because organic compounds consist of a carbon framework, reactions that form carbon-carbon bonds are especially useful. In Reaction [2], for example, nucleophilic attack of a three-carbon acetylide anion on a two-carbon alkyl halide yields a five-carbon alkyne as product.


Although nucleophilic substitution with acetylide anions is a very valuable carbon-carbon bond-forming reaction, it has the same limitations as any $\text{S}_\text{N}2$ reaction. **Steric hindrance around the leaving group causes 2° and 3° alkyl halides to undergo elimination by an $\text{E}2$ mechanism**, as shown with 2-bromo-2-methylpropane. Thus, nucleophilic substitution with acetylide anions forms new carbon-carbon bonds in high yield only with unhindered CH_3X and 1° alkyl halides.

Steric hindrance prevents an $\text{S}_\text{N}2$ reaction.

The acetylide anion acts as a base instead.




Sample Problem 11.5 Draw the organic products formed in each reaction.




Solution


a. Because the alkyl halide is 1° and the acetylide anion is a strong nucleophile, substitution occurs by an $\text{S}_{\text{N}}2$ mechanism, resulting in a new C–C bond.



b. Because the alkyl halide is 2° , the major product is formed from elimination by an $\text{E}2$ mechanism.



Problem 11.17 Draw the organic products formed in each reaction.



Problem 11.18 What acetylide anion and alkyl halide can be used to prepare each alkyne? Indicate all possibilities when more than one route will work.




Because acetylene has two sp hybridized C–H bonds, two sequential reactions can occur to form **two new carbon–carbon bonds**, as shown in Sample Problem 11.6.

Sample Problem 11.6 Identify the terminal alkyne **A** and the internal alkyne **B** in the following reaction sequence.




Solution


In each step, the base NH_2^- removes a proton on an sp hybridized carbon, and the resulting acetylide anion reacts as a nucleophile with an alkyl halide to yield an $\text{S}_{\text{N}}2$ product. The first two-step reaction sequence forms the **terminal alkyne A** by nucleophilic attack of the acetylide anion on CH_3Br .



The second two-step reaction sequence forms the **internal alkyne B** by nucleophilic attack of the acetylide anion on $\text{CH}_3\text{CH}_2\text{Cl}$.



Sample Problem 11.6 illustrates how a five-carbon product can be prepared from three smaller molecules by forming two new carbon–carbon bonds.

The soft coral *Capnella imbricata*

Carbon-carbon bond formation with acetylide anions is a valuable reaction used in the synthesis of numerous natural products. Two examples include **capnellene**, isolated from the soft coral *Capnella imbricata*, and **niphatoxin B**, isolated from a red sea sponge, as shown in Figure 11.6.

Problem 11.19


Show how $\text{HC}\equiv\text{CH}$, $\text{CH}_3\text{CH}_2\text{Br}$, and $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Br}$ can be used to prepare $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$. Show all reagents, and use curved arrows to show movement of electron pairs.

Problem 11.20

Explain why 2,2,5,5-tetramethyl-3-hexyne can't be made using acetylide anions.

11.11B Reaction of Acetylide Anions with Epoxides


Acetylide anions are strong nucleophiles that open epoxide rings by an $\text{S}_{\text{N}}2$ mechanism. This reaction also results in the formation of a **new carbon-carbon bond**. Backside attack occurs at the less substituted end of the epoxide.




Opening of epoxide rings with strong nucleophiles was first discussed in Section 9.15A.

Figure 11.6

Use of acetylide anion reactions in the synthesis of two marine natural products



Problem 11.21 Draw the products of each reaction.



Problem 11.22 Draw the products formed when $\text{CH}_3\text{CH}_2\text{C}\equiv\text{C}^- \text{Na}^+$ reacts with each compound.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ d. $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$
 b. $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{Cl}$ e. ethylene oxide followed by H_2O
 c. $(\text{CH}_3\text{CH}_2)_3\text{CCl}$ f. propene oxide followed by H_2O

11.12 Synthesis


The reactions of acetylide anions give us an opportunity to examine organic synthesis more systematically. Performing a multistep synthesis can be difficult. Not only must you know the reactions for a particular functional group, but you must also put these reactions in a logical order, a process that takes much practice to master.

11.12A General Terminology and Conventions

To plan a synthesis of more than one step, we use the process of **retrosynthetic analysis**—that is, working backwards from the desired product to determine the starting materials from which it is made (Section 10.18). To write a synthesis working backwards from the product to the starting material, an **open arrow** (\Rightarrow) is used to indicate that the product is drawn on the left and the starting material on the right.

The product of a synthesis is often called the **target compound**. Using retrosynthetic analysis, we must determine what compound can be converted to the target compound by a single reaction. That is, **what is the immediate precursor of the target compound?** After an appropriate precursor is identified, this process is continued until we reach a specified starting material. Sometimes multiple retrosynthetic pathways are examined before a particular route is decided upon.

Carefully read the directions for each synthesis problem. Sometimes a starting material is specified, whereas at other times you must begin with a compound that meets a particular criterion; for example, you may be asked to synthesize a compound from alcohols having five or fewer carbon atoms. These limitations are meant to give you some direction in planning a multistep synthesis.



In designing a synthesis, reactions are often divided into two categories:

- Those that form new carbon–carbon bonds.
- Those that convert one functional group into another—that is, functional group interconversions.

Carbon–carbon bond-forming reactions are central to organic synthesis because simpler and less valuable starting materials can be converted to more complex products. Keep in mind that whenever the product of a synthesis has more carbon–carbon bonds than the starting material, the synthesis must contain at least one of these reactions.

Appendix D lists the carbon–carbon bond-forming reactions encountered in this text.

HOW TO Develop a Retrosynthetic Analysis

Step [1] Compare the carbon skeletons of the starting material and product.

- If the product has more carbon-carbon σ bonds than the starting material, the synthesis must form one or more C-C bonds. If not, only functional group interconversion occurs.
- Match the carbons in the starting material with those in the product, to see where new C-C bonds must be added or where functional groups must be changed.

Step [2] Concentrate on the functional groups in the starting material and product and ask:

- What methods introduce the functional groups in the product?
- What kind of reactions does the starting material undergo?

Step [3] Work backwards from the product and forwards from the starting material.

- Ask: What is the immediate precursor of the product?
- Compare each precursor to the starting material to determine if there is a one-step reaction that converts one to the other. Continue this process until the starting material is reached.
- Always generate simpler precursors when working backwards.
- Use fewer steps when multiple routes are possible.
- Keep in mind that you may need to evaluate several different precursors for a given compound.

Step [4] Check the synthesis by writing it in the synthetic direction.

- To check a retrosynthetic analysis, write out the steps beginning with the starting material, indicating all necessary reagents.

11.12B Examples of Multistep Synthesis


Retrosynthetic analysis with acetylide anions is illustrated in Sample Problems 11.7 and 11.8.

Sample Problem 11.7

Devise a synthesis of $\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$ from $\text{HC}\equiv\text{CH}$ and any other organic or inorganic reagents.

Retrosynthetic Analysis

The two C's in the starting material match up with the two sp hybridized C's in the product, so a three-carbon unit must be added.




Thinking backwards . . .

- [1] Form a new C-C bond using an acetylide anion and a 1° alkyl halide.
- [2] Prepare the acetylide anion from acetylene by treatment with base.


Synthesis

Deprotonation of $\text{HC}\equiv\text{CH}$ with NaH forms the acetylide anion, which undergoes $\text{S}_{\text{N}}2$ reaction with an alkyl halide to form the target compound, a five-carbon alkyne.

A two-step process:




Sample Problem 11.8 Devise a synthesis of the following compound from starting materials having two carbons or fewer.



Retrosynthetic Analysis

A carbon-carbon bond-forming reaction must be used to convert the two-carbon starting materials to the four-carbon product.




Thinking backwards . . .

- [1] Form the carbonyl group by hydration of a triple bond.
- [2] Form a new C-C bond using an acetylide anion and a 1° alkyl halide.
- [3] Prepare the acetylide anion from acetylene by treatment with base.

Synthesis

Three steps are needed to complete the synthesis. Treatment of $\text{HC}\equiv\text{CH}$ with NaH forms the acetylide anion, which undergoes an $\text{S}_{\text{N}}2$ reaction with an alkyl halide to form a four-carbon terminal alkyne. Hydration of the alkyne with H_2O , H_2SO_4 , and HgSO_4 yields the target compound.



These examples illustrate the synthesis of organic compounds by multistep routes. In Chapter 12 we will learn other useful reactions that expand our capability to do synthesis.

Problem 11.23 Use retrosynthetic analysis to show how 3-hexyne can be prepared from acetylene and any other organic and inorganic compounds. Then draw the synthesis in the synthetic direction, showing all needed reagents.

Problem 11.24 Devise a synthesis of $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ from two-carbon starting materials.

KEY CONCEPTS


Alkynes

General Facts About Alkynes


- Alkynes contain a carbon-carbon triple bond consisting of a strong σ bond and two weak π bonds. Each carbon is sp hybridized and linear (11.1).
- Alkynes are named using the suffix **-yne** (11.2).
- Alkynes have weak intermolecular forces, giving them low mp's and low bp's, and making them water insoluble (11.3).
- Because its weaker π bonds make an alkyne electron rich, alkynes undergo addition reactions with electrophiles (11.6).

Addition Reactions of Alkynes


[1] Hydrohalogenation—Addition of HX (X = Cl, Br, I) (11.7)




- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.

[2] Halogenation—Addition of X₂ (X = Cl or Br) (11.8)

- Bridged halonium ions are formed as intermediates.
- Anti addition of X₂ occurs.

[3] Hydration—Addition of H₂O (11.9)


- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- An unstable enol is first formed, which rearranges to a carbonyl group.

[4] Hydroboration–oxidation—Addition of H₂O (11.10)

- The unstable enol, first formed after oxidation, rearranges to a carbonyl group.


Reactions Involving Acetylide Anions

[1] Formation of acetylide anions from terminal alkynes (11.6B)




- Typical bases used for the reaction are NaNH₂ and NaH.

[2] Reaction of acetylide anions with alkyl halides (11.11A)



- The reaction follows an S_N2 mechanism.
- The reaction works best with CH₃X and RCH₂X.

[3] Reaction of acetylide anions with epoxides (11.11B)





- The reaction follows an S_N2 mechanism.
- Opening of the ring occurs from the back side at the less substituted end of the epoxide.









PROBLEMS

Nomenclature




11.25 Answer the following questions about erlotinib and phomallenic acid C. Erlotinib, sold under the trade name Tarceva, was introduced in 2004 for the treatment of lung cancer. Phomallenic acid C is an inhibitor of bacterial fatty acid synthesis.



- Which C–H bond in erlotinib is most acidic?
 - What orbitals are used to form the shortest C–C single bond in erlotinib?
 - Which H atom in phomallenic acid C is most acidic?
 - How many sp hybridized carbons are contained in phomallenic acid C?
 - Rank the labeled bonds in phomallenic acid C in order of increasing bond strength.
- 11.26** Draw the seven isomeric alkynes having molecular formula C_6H_{10} , and give the IUPAC name for each compound. Consider constitutional isomers only.
- 11.27** Give the IUPAC name for each alkyne.







11.28 Give the structure corresponding to each name.






Tautomers




11.29 Which of the following pairs of compounds represent keto–enol tautomers?




11.30 Draw the enol form of each keto tautomer.




11.31 Draw the keto form of each enol tautomer.




11.32 How is each compound related to **A**? Choose from tautomers, constitutional isomers but not tautomers, or neither.




11.33 Draw a stepwise mechanism for the conversion of cyclopentanone (**A**) to its enol tautomer (**B**) in the presence of acid.



11.34 Conversion of an enol to a ketone also occurs in the presence of base. Draw a stepwise mechanism for the following tautomerization.




11.35 Enamines and imines are tautomers that contain N atoms. Draw a stepwise mechanism for the acid-catalyzed conversion of enamine **X** to imine **Y**.



Reactions

11.36 Draw the products of each acid–base reaction. Indicate whether equilibrium favors the starting materials or the products.



11.37 Draw the products formed when 1-hexyne is treated with each reagent.



11.38 Draw the products formed when 3-hexyne is treated with each reagent.




11.39 What reagents are needed to convert $(\text{CH}_3\text{CH}_2)_3\text{CC}\equiv\text{CH}$ to each compound?




11.40 Explain the apparent paradox. Although the addition of one equivalent of HX to an alkyne is more exothermic than the addition of HX to an alkene, an alkene reacts faster with HX.



11.41 What alkyne gives each of the following ketones as the only product after hydration with H_2O , H_2SO_4 , and HgSO_4 ?



11.42 What two different alkynes yield 2-butanone from hydration with H_2O , H_2SO_4 , and HgSO_4 ?














11.43 What alkyne gives each compound as the only product after hydroboration–oxidation?




11.44 Explain why butyllithium, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2^-\text{Li}^+$ is an effective base for converting alkynes to acetylide anions.


11.45 Draw the organic products formed in each reaction.







11.46 Draw the structure of compounds **A–E** in the following reaction scheme.






11.47 When alkyne **A** is treated with NaNH_2 followed by CH_3I , a product having molecular formula $\text{C}_6\text{H}_{10}\text{O}$ is formed, but it is *not* compound **B**. What is the structure of the product and why is it formed?





11.48 Draw the products formed in each reaction and indicate stereochemistry.



11.49 What reactions are needed to convert alcohol **A** into either alkyne **B** or alkyne **C**?




Synthesis


- 11.58** What reagents are needed to prepare $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$ from each starting material?
 a. $\text{CH}_3(\text{CH}_2)_4\text{CHCl}_2$ b. $\text{CH}_3(\text{CH}_2)_3\text{CH}=\text{CH}_2$ c. $\text{CH}_3(\text{CH}_2)_5\text{OH}$
- 11.59** What steps are needed to prepare phenylacetylene, $\text{C}_6\text{H}_5\text{C}\equiv\text{CH}$, from each compound: (a) $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{Br}$; (b) $\text{C}_6\text{H}_5\text{CHBrCH}_3$; (c) $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{OH}$?
- 11.60** What acetylide anion and alkyl halide are needed to synthesize each alkyne?
 a. $\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$ b. $\text{CH}_3-\text{C}\equiv\text{C}-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_2\text{CH}_3$ c. 
- 11.61** Synthesize each compound from acetylene. You may use any other organic or inorganic reagents.
 a. $(\text{CH}_3)_2\text{CHCH}_2\text{C}\equiv\text{CH}$ c. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$ e. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CCl}_2\text{CH}_3$
 b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$ d. $\text{CH}_3\text{CH}_2\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$ f. $\text{CH}_3\text{CH}_2\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$
- 11.62** Devise a synthesis of each compound using $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2$ as the starting material. You may use any other organic compounds or inorganic reagents.
 a. $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CH}$ d. $\text{CH}_3\text{CH}_2\text{CHBrCH}_2\text{Br}$
 b. $\text{CH}_3\text{CH}_2\text{CBr}_2\text{CH}_3$ e. $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$
 c. $\text{CH}_3\text{CH}_2\text{CCl}_2\text{CHCl}_2$ f. $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OH}$
 g.  (+ enantiomer)
- 11.63** Devise a synthesis of each compound. You may use $\text{HC}\equiv\text{CH}$, ethylene oxide, and alkyl halides as organic starting materials and any inorganic reagents.
 a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$ c. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OH}$
 b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_3$ d. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$
- 11.64** Devise a synthesis of the ketone 3-hexanone, $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_2\text{CH}_3$, from $\text{CH}_3\text{CH}_2\text{Br}$ as the only organic starting material; that is, all the carbon atoms in 3-hexanone must come from $\text{CH}_3\text{CH}_2\text{Br}$. You may use any other needed reagents.
- 11.65** Devise a synthesis of each compound using $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ as the only organic starting material: (a) $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$; (b) $\text{CH}_3\text{C}\equiv\text{CCH}_2\text{CH}(\text{OH})\text{CH}_3$. You may use any other needed inorganic reagents.
- 11.66** Devise a synthesis of each compound from $\text{CH}_3\text{CH}_2\text{OH}$ as the only organic starting material: (a) $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OH}$; (b) $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$. You may use any other needed reagents.

Challenge Problems


- 11.67** Draw a stepwise mechanism for the following reaction.



- 11.68** Why is compound **X** formed in the following reaction, instead of its constitutional isomer **Y**?



- 11.69** Draw a stepwise mechanism for the following intramolecular reaction.



12

Oxidation and Reduction

- 12.1 Introduction
- 12.2 Reducing agents
- 12.3 Reduction of alkenes
- 12.4 Application:
Hydrogenation of oils
- 12.5 Reduction of alkynes
- 12.6 The reduction of polar
C–X σ bonds
- 12.7 Oxidizing agents
- 12.8 Epoxidation
- 12.9 Dihydroxylation
- 12.10 Oxidative cleavage of
alkenes
- 12.11 Oxidative cleavage of
alkynes
- 12.12 Oxidation of alcohols
- 12.13 Green chemistry
- 12.14 Application: The
oxidation of ethanol
- 12.15 Sharpless epoxidation



Throughout history, humans have ingested alcoholic beverages for their pleasant taste and the feeling of euphoria they impart. Wine, beer, and similar products contain **ethanol** ($\text{CH}_3\text{CH}_2\text{OH}$), a 1° alcohol that is quickly absorbed in the stomach and small intestines and rapidly transported in the bloodstream to other organs. Like other 1° alcohols, ethanol is easily oxidized, and as a result, ethanol is metabolized in the body by a series of enzyme-catalyzed oxidation reactions that take place in the liver. In Chapter 12, we learn about oxidation and reduction reactions of organic molecules like ethanol.

In Chapter 12, we discuss the oxidation and reduction of **alkenes** and **alkynes**, as well as compounds with **polar C–X σ bonds**—alcohols, alkyl halides, and epoxides. Although there will be many different reagents and mechanisms, discussing these reactions as a group allows us to more easily compare and contrast them.

The word *mechanism* will often be used loosely here. In contrast to the S_N1 reaction of alkyl halides or the electrophilic addition reactions of alkenes, the details of some of the mechanisms presented in Chapter 12 are known with less certainty. For example, although the identity of a particular intermediate might be confirmed by experiment, other details of the mechanism are suggested by the structure or stereochemistry of the final product.

Oxidation and reduction reactions are very versatile, and knowing them allows us to design many more complex organic syntheses.

12.1 Introduction

Recall from Section 4.14 that the way to determine whether an organic compound has been oxidized or reduced is to compare the **relative number of C–H and C–Z bonds** ($Z =$ an element *more electronegative* than carbon) in the starting material and product.

- **Oxidation** results in an *increase* in the number of C–Z bonds (usually C–O bonds) or a *decrease* in the number of C–H bonds.
- **Reduction** results in a *decrease* in the number of C–Z bonds (usually C–O bonds) or an *increase* in the number of C–H bonds.

Two components are always present in an oxidation or reduction reaction—**one component is oxidized and one is reduced**. When an organic compound is *oxidized* by a reagent, the reagent itself must be *reduced*. Similarly, when an organic compound is *reduced* by a reagent, the reagent becomes *oxidized*.

Thus, an organic compound such as CH_4 can be oxidized by replacing C–H bonds with C–O bonds, as shown in Figure 12.1. Reduction is the opposite of oxidation, so Figure 12.1 also shows how a compound can be reduced by replacing C–O bonds with C–H bonds. The symbols **[O]** and **[H]** indicate oxidation and reduction, respectively.

Sometimes two carbon atoms are involved in a single oxidation or reduction reaction, and the net change in the number of C–H or C–Z bonds at *both* atoms must be taken into account. The conversion of an **alkyne to an alkene** and an **alkene to an alkane** are examples of reduction, because each process adds two new C–H bonds to the starting material, as shown in Figure 12.2.

Problem 12.1 Classify each reaction as oxidation, reduction, or neither.




Figure 12.1

A general scheme for the oxidation and reduction of a carbon compound





Figure 12.2


Oxidation and reduction of hydrocarbons




12.2 Reducing Agents

All reducing agents provide the equivalent of two hydrogen atoms, but **there are three types of reductions**, differing in how H_2 is added. The simplest reducing agent is molecular H_2 . Reductions of this sort are carried out in the presence of a metal catalyst that acts as a surface on which reaction occurs.


The second way to deliver H_2 in a reduction is to add two protons and two electrons to a substrate—that is, $\text{H}_2 = 2 \text{H}^+ + 2 \text{e}^-$. Reducing agents of this sort use alkali metals as a source of electrons and liquid ammonia (NH_3) as a source of protons. Reductions with **Na in NH_3** are called **dissolving metal reductions**.



The third way to deliver the equivalent of two hydrogen atoms is to add **hydride (H^-)** and a **proton (H^+)**. The most common hydride reducing agents contain a hydrogen atom bonded to boron or aluminum. Simple examples include **sodium borohydride (NaBH_4)** and **lithium aluminum hydride (LiAlH_4)**. These reagents deliver H^- to a substrate, and then a proton is added from H_2O or an alcohol.




- Metal hydride reagents act as a source of H^- because they contain polar metal-hydrogen bonds that place a partial negative charge on hydrogen.





12.3 Reduction of Alkenes

Reduction of an alkene forms an alkane by addition of H_2 . Two bonds are broken—the **weak π bond** of the alkene and the H_2 σ bond—and two new $\text{C}-\text{H}$ σ bonds are formed.




Hydrogenation catalysts are insoluble in common solvents, thus creating a **heterogeneous** reaction mixture. This insolubility has a practical advantage. These catalysts contain expensive metals, but they can be filtered away from the other reactants after the reaction is complete, and then reused.

The addition of H_2 occurs only in the presence of a **metal catalyst**, and thus, the reaction is called **catalytic hydrogenation**. The catalyst consists of a metal—usually Pd, Pt, or Ni—adsorbed onto a finely divided inert solid, such as charcoal. For example, the catalyst 10% Pd on carbon is composed of 10% Pd and 90% carbon, by weight. H_2 adds in a **syn** fashion, as shown in Equation [2].



Problem 12.2 What alkane is formed when each alkene is treated with H_2 and a Pd catalyst?



Problem 12.3 Draw all alkenes that react with one equivalent of H_2 in the presence of a palladium catalyst to form each alkane. Consider constitutional isomers only.




12.3A Hydrogenation and Alkene Stability

Hydrogenation reactions are **exothermic** because the bonds in the product are stronger than the bonds in the starting materials, making them similar to other alkene addition reactions. The ΔH° for hydrogenation, called the **heat of hydrogenation**, can be used as a measure of the relative stability of two different alkenes that are hydrogenated to the same alkane.


For example, both *cis*- and *trans*-2-butene are hydrogenated to butane, and the heat of hydrogenation for the *trans* isomer is less than that for the *cis* isomer. **Because less energy is released in converting the *trans* alkene to butane, it must be lower in energy (more stable) to begin with.** The relative energies of the butene isomers are illustrated in Figure 12.3.

Recall from Chapter 8 that **trans alkenes are generally more stable than cis alkenes.**




- When hydrogenation of two alkenes gives the same alkane, the more stable alkene has the *smaller* heat of hydrogenation.

Figure 12.3

Relative energies of *cis*- and *trans*-2-butene

Problem 12.4 Which alkene in each pair has the larger heat of hydrogenation?



Problem 12.5 Explain why heats of hydrogenation cannot be used to determine the relative stability of 2-methyl-2-pentene and 3-methyl-1-pentene.


12.3B The Mechanism of Catalytic Hydrogenation

In the generally accepted mechanism for catalytic hydrogenation, the surface of the metal catalyst binds both H_2 and the alkene, and H_2 is transferred to the π bond in a rapid but stepwise process (Mechanism 12.1).




Mechanism 12.1 Addition of H_2 to an Alkene—Hydrogenation

Steps [1] and [2] Complexation of H_2 and the alkene to the catalyst



- H_2 adsorbs to the catalyst surface with partial or complete cleavage of the H–H bond.
- The π bond of the alkene complexes with the metal.


Steps [3] and [4] Sequential addition of the elements of H_2




- **Two H atoms are transferred sequentially** to the π bond in Steps [3] and [4], forming the alkane.
- Because the product alkane no longer has a π bond with which to complex to the metal, it is released from the catalyst surface.

The mechanism explains two facts about hydrogenation:




- Rapid, sequential addition of H_2 occurs from the side of the alkene complexed to the metal surface, resulting in *syn* addition.
- Less crowded double bonds complex more readily to the catalyst surface, resulting in faster reaction.



Problem 12.6 What product is formed when limonene is treated with one equivalent of H_2 and a Pd catalyst?



Problem 12.7 Given that syn addition of H_2 occurs from both sides of a trigonal planar double bond, draw all stereoisomers formed when each alkene is treated with H_2 .



12.3C Hydrogenation Data and Degrees of Unsaturation

Recall from Section 10.2 that the **number of degrees of unsaturation gives the total number of rings and π bonds in a molecule**. Because H_2 adds to π bonds but does *not* add to the $\text{C}-\text{C}$ σ bonds of rings, hydrogenation allows us to determine how many degrees of unsaturation are due to π bonds and how many are due to rings. This is done by comparing the number of degrees of unsaturation before and after a molecule is treated with H_2 , as illustrated in Sample Problem 12.1.

Sample Problem 12.1 How many rings and π bonds are contained in a compound of molecular formula C_8H_{12} that is hydrogenated to a compound of molecular formula C_8H_{14} ?

Solution

[1] Determine the number of degrees of unsaturation in the compounds before and after hydrogenation.

Before H_2 addition— C_8H_{12}

- The maximum number of H's possible for n C's is $2n + 2$; in this example, $2n + 2 = 2(8) + 2 = 18$.
- 18 H's (maximum) – 12 H's (actual) = 6 H's fewer than the maximum number.

$$\frac{6 \text{ H's fewer than the maximum}}{2 \text{ H's removed for each degree of unsaturation}} =$$

three degrees of unsaturation

After H_2 addition— C_8H_{14}

- The maximum number of H's possible for n C's is $2n + 2$; in this example, $2n + 2 = 2(8) + 2 = 18$.
- 18 H's (maximum) – 14 H's (actual) = 4 H's fewer than the maximum number.

$$\frac{4 \text{ H's fewer than the maximum}}{2 \text{ H's removed for each degree of unsaturation}} =$$

two degrees of unsaturation

[2] Assign the number of degrees of unsaturation to rings or π bonds as follows:

- The number of degrees of unsaturation that remain in the product after H_2 addition = the **number of rings** in the starting material.
- The number of degrees of unsaturation that react with H_2 = the **number of π bonds**.

In this example, **two** degrees of unsaturation remain after hydrogenation, so the starting material has **two** rings. Thus:

Before H₂ addition:After H₂ addition:

three degrees of unsaturation — two degrees of unsaturation = one degree of unsaturation that reacted with H₂


three rings or π bonds in C₈H₁₂ = two rings + one π bond ANSWER

Problem 12.8 Complete the missing information for compounds **A**, **B**, and **C**, each subjected to hydrogenation. The number of rings and π bonds refers to the reactant (**A**, **B**, or **C**) prior to hydrogenation.

Compound	Molecular formula before hydrogenation	Molecular formula after hydrogenation	Number of rings	Number of π bonds
A	C ₁₀ H ₁₂	C ₁₀ H ₁₆	?	?
B	?	C ₄ H ₁₀	0	1
C	C ₆ H ₈	?	1	?

12.3D Hydrogenation of Other Double Bonds

Compounds that contain a carbonyl group also react with H₂ and a metal catalyst. For example, **aldehydes and ketones are reduced to 1° and 2° alcohols**, respectively. We return to this reaction in Chapter 20.



12.4 Application: Hydrogenation of Oils




S
5 INGREDIENTS: ROASTED PEANUTS, SUGAR, PARTIALLY HYDROGENATED VEGETABLE OILS (RAPESEED, COTTONSEED AND SOYBEAN) TO PREVENT SEPARATION, SALT.

Peanut butter is a common consumer product that contains partially hydrogenated vegetable oil.

Several ingredients are added to make margarine more closely resemble butter: orange β -carotene (Section 10.5) is often added for color, salt for flavor, and 3-hydroxy-2-butanone [CH₃COCH(OH)CH₃] or 2,3-butanedione (CH₃COCOCH₃) to mimic the flavor of butter.

Many processed foods, such as peanut butter, margarine, and some brands of crackers, contain *partially hydrogenated* vegetable oils. These oils are produced by hydrogenating the long hydrocarbon chains of triacylglycerols.

In Section 10.6 we learned that **fats and oils are triacylglycerols that differ in the number of degrees of unsaturation** in their long alkyl side chains.




triacylglycerol

The number of double bonds in the R groups makes a triacylglycerol either a fat or an oil.

- **Fats**—usually animal in origin—are solids with triacylglycerols having few degrees of unsaturation.
- **Oils**—usually vegetable in origin—are liquids with triacylglycerols having a larger number of degrees of unsaturation.

When an unsaturated vegetable oil is treated with hydrogen, some (or all) of the π bonds add H₂, decreasing the number of degrees of unsaturation (Figure 12.4). This increases the melting point of the oil. For example, margarine is prepared by partially hydrogenating vegetable oil to give a product having a semi-solid consistency that more closely resembles butter. This process is sometimes called **hardening**.

Figure 12.4 Partial hydrogenation of the double bonds in a vegetable oil




- **Decreasing** the number of degrees of unsaturation **increases** the melting point. Only one long chain of the triacylglycerol is drawn.
- When an oil is *partially* hydrogenated, some double bonds react with H₂, whereas some double bonds remain in the product.
- Partial hydrogenation **decreases** the number of allylic sites (shown in blue), making a triacylglycerol **less** susceptible to oxidation, thereby increasing its shelf life.

If unsaturated oils are healthier than saturated fats, why does the food industry hydrogenate oils? There are two reasons—aesthetics and shelf life. Consumers prefer the semi-solid consistency of margarine to a liquid oil. Imagine pouring vegetable oil on a piece of toast or pancakes.


Furthermore, unsaturated oils are more susceptible than saturated fats to oxidation at the **allylic carbon atoms**—the carbons adjacent to the double bond carbons—a process discussed in Chapter 15. Oxidation makes the oil rancid and inedible. Hydrogenating the double bonds reduces the number of allylic carbons (also illustrated in Figure 12.4), thus reducing the likelihood of oxidation and increasing the shelf life of the food product. This process reflects a delicate balance between providing consumers with healthier food products, while maximizing shelf life to prevent spoilage.

One other fact is worthy of note. Because the steps in hydrogenation are reversible and H atoms are added in a sequential rather than concerted fashion, a *cis* double bond can be isomerized to a *trans* double bond. After addition of one H atom (Step [3] in Mechanism 12.1), an intermediate can lose a hydrogen atom to re-form a double bond with either the *cis* or *trans* configuration.

As a result, some of the *cis* double bonds in vegetable oils are converted to *trans* double bonds during hydrogenation, forming so-called “*trans* fats.” The shape of the resulting fatty acid chain is very different, closely resembling the shape of a *saturated* fatty acid chain. Consequently, *trans* fats are thought to have the same negative effects on blood cholesterol levels as saturated fats; that is, *trans* fats stimulate cholesterol synthesis in the liver, thus increasing blood cholesterol levels, a factor linked to increased risk of heart disease.



Problem 12.9 Draw the products formed when triacylglycerol **A** is treated with each reagent, forming compounds **B** and **C**. Rank **A**, **B**, and **C** in order of increasing melting point.




- a. H_2 (excess), Pd-C (Compound **B**)
 b. H_2 (1 equiv), Pd-C (Compound **C**)


12.5 Reduction of Alkynes

Reduction of an alkyne adds H_2 to one or both of the π bonds. There are three different ways by which the elements of H_2 can be added to a triple bond.


- Adding two equivalents of H_2 forms an alkane.



- Adding one equivalent of H_2 in a syn fashion forms a cis alkene.




- Adding one equivalent of H_2 in an anti fashion forms a trans alkene.



12.5A Reduction of an Alkyne to an Alkane



When an alkyne is treated with two or more equivalents of H_2 and a Pd catalyst, reduction of *both* π bonds occurs. **Syn addition** of one equivalent of H_2 forms a cis alkene, which adds a second equivalent of H_2 to form an **alkane**. **Four new C–H bonds are formed.** By using a Pd-C catalyst, it is not possible to stop the reaction after addition of only one equivalent of H_2 .

12.5C Reduction of an Alkyne to a Trans Alkene



NH_3 has a boiling point of $-33\text{ }^\circ\text{C}$, making it a gas at room temperature. To carry out a Na, NH_3 reduction, NH_3 gas is condensed into a flask kept at $-78\text{ }^\circ\text{C}$ by a cooling bath of solid CO_2 in acetone. When Na is added to the liquid NH_3 , a brilliant blue solution is formed.

Although catalytic hydrogenation is a convenient method for preparing cis alkenes from alkynes, it cannot be used to prepare trans alkenes. With a dissolving metal reduction (such as Na in NH_3), however, the elements of H_2 are added in an **anti** fashion to the triple bond, thus forming a **trans alkene**. For example, 2-butyne reacts with Na in NH_3 to form *trans*-2-butene.




The **mechanism** for the dissolving metal reduction using Na in NH_3 features sequential addition of electrons and protons to the triple bond. Half-headed arrows denoting the movement of a single electron must be used in two steps when Na donates *one* electron. The mechanism can be divided conceptually into two parts, each of which consists of two steps: **addition of an electron followed by protonation of the resulting negative charge**, as shown in Mechanism 12.2.




Mechanism 12.2 Dissolving Metal Reduction of an Alkyne to a Trans Alkene

Steps [1] and [2] Addition of one electron and one proton to form a radical




- Addition of an electron to the triple bond in Step [1] forms a **radical anion**, a species containing *both* a negative charge and an unpaired electron.
- Protonation of the anion with the solvent NH_3 in Step [2] yields a **radical**. The net effect of Steps [1] and [2] is to add one hydrogen atom (H^\bullet) to the triple bond.

Steps [3] and [4] Addition of one electron and one proton to form the trans alkene



- Addition of a second electron to the radical in Step [3] forms a **carbanion**.
- Protonation of the carbanion in Step [4] forms the trans alkene. These last two steps add the second hydrogen atom (H^\bullet) to the triple bond.

Although the vinyl carbanion formed in Step [3] could have two different arrangements of its R groups, only the trans alkene is formed from the more stable vinyl carbanion; this carbanion has the larger R groups farther away from each other to avoid steric interactions. Protonation of this anion leads to the more stable trans product.




Dissolving metal reduction of a triple bond with Na in NH_3 is a **stereoselective reaction** because it forms a trans product exclusively.

- Dissolving metal reductions always form the more stable trans product preferentially.

Figure 12.5

Summary: Three methods to reduce a triple bond




The three methods to reduce a triple bond are summarized in Figure 12.5 using 3-hexyne as starting material.

- Problem 12.13** What product is formed when $\text{CH}_3\text{OCH}_2\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}(\text{CH}_3)_2$ is treated with each reagent: (a) H_2 (excess), Pd-C; (b) H_2 (1 equiv), Lindlar catalyst; (c) H_2 (excess), Lindlar catalyst; (d) Na, NH_3 ?
- Problem 12.14** Deuterium is introduced into a molecule by using reducing agents that contain D atoms instead of H atoms. Draw the products formed when 2-hexyne is treated with each reagent: (a) D_2 , Pd-C; (b) D_2 , Lindlar catalyst; (c) Na, ND_3 .
- Problem 12.15** A chiral alkyne **A** with molecular formula C_6H_{10} is reduced with H_2 and Lindlar catalyst to **B** having the *R* configuration at its stereogenic center. What are the structures of **A** and **B**?

12.6 The Reduction of Polar C–X σ Bonds

Compounds containing polar C–X σ bonds that react with strong nucleophiles are reduced with metal hydride reagents, most commonly lithium aluminum hydride. Two functional groups possessing both of these characteristics are **alkyl halides** and **epoxides**. Alkyl halides are reduced to alkanes with loss of X^- as the leaving group. Epoxide rings are opened to form alcohols.



Reduction of these C–X σ bonds is another example of nucleophilic substitution, in which LiAlH_4 serves as a source of a hydride nucleophile (H^-). Because H^- is a strong nucleophile, the reaction follows an **$\text{S}_{\text{N}}2$ mechanism**, illustrated for the one-step reduction of an alkyl halide in Mechanism 12.3.



Mechanism 12.3 Reduction of RX with LiAlH_4

One step The nucleophile H^- substitutes for X^- in a single step.





Figure 12.6

Examples of reduction of C–X
σ bonds with LiAlH₄




Because the reaction follows an S_N2 mechanism:

- Unhindered CH₃X and 1° alkyl halides are more easily reduced than more substituted 2° and 3° halides.
- In unsymmetrical epoxides, nucleophilic attack of H[−] (from LiAlH₄) occurs at the less substituted carbon atom.

Examples are shown in Figure 12.6.

Problem 12.16 Draw the products of each reaction.



12.7 Oxidizing Agents

Oxidizing agents fall into two main categories:



- Reagents that contain an oxygen–oxygen bond
- Reagents that contain metal–oxygen bonds

Oxidizing agents containing an O–O bond include O₂, O₃ (ozone), H₂O₂ (hydrogen peroxide), (CH₃)₃COOH (*tert*-butyl hydroperoxide), and peroxyacids. **Peroxyacids**, a group of reagents with the general structure RCO₃H, have one more O atom than carboxylic acids (RCO₂H). Some peroxyacids are commercially available whereas others are prepared and used without isolation. Examples are shown in Figure 12.7. All of these reagents contain a weak O–O bond that is cleaved during oxidation.

The most common oxidizing agents with metal–oxygen bonds contain either chromium in the +6 oxidation state (six Cr–O bonds) or manganese in the +7 oxidation state (seven Mn–O bonds). Common Cr⁶⁺ reagents include chromium(VI) oxide (CrO₃) and sodium or potassium dichromate (Na₂Cr₂O₇ and K₂Cr₂O₇). **These reagents are strong oxidants** used in the presence of a strong aqueous acid such as H₂SO₄. **Pyridinium chlorochromate (PCC)**, a Cr⁶⁺ reagent that is soluble in halogenated organic solvents, can be used without strong acid present. This makes it a **more selective Cr⁶⁺ oxidant**, as described in Section 12.12.

Figure 12.7

Common peroxyacids



peroxyacetic acid





*meta*-chloroperoxybenzoic acid
mCPBAmagnesium monoperoxyphthalate
MMPP

Figure 12.8
Oxidation reactions of alkenes,
alkynes, and alcohols




The most common Mn^{7+} reagent is **KMnO₄** (potassium permanganate), a strong, water-soluble oxidant. Other oxidizing agents that contain metals include **OsO₄** (osmium tetroxide) and **Ag₂O** [silver(I) oxide].


In the remainder of Chapter 12, the oxidation of alkenes, alkynes, and alcohols—three functional groups already introduced in this text—is presented (Figure 12.8). Addition reactions to alkenes and alkynes that increase the number of C–O bonds are described in Sections 12.8–12.11. Oxidation of alcohols to carbonyl compounds appears in Sections 12.12–12.14.

12.8 Epoxidation

Epoxidation is the addition of a single oxygen atom to an alkene to form an epoxide.



The weak π bond of the alkene is broken and two new C–O σ bonds are formed. Epoxidation is typically carried out with a peroxyacid, resulting in cleavage of the weak O–O bond of the reagent.




Epoxidation occurs via the concerted addition of one oxygen atom of the peroxyacid to the π bond as shown in Mechanism 12.4. Epoxidation resembles the formation of the bridged halonium ion in Section 10.13, in that two bonds in a three-membered ring are formed in one step.






Mechanism 12.4 Epoxidation of an Alkene with a Peroxyacid

One step All bonds are broken or formed in a single step.



- Two C–O bonds are formed to one O atom with one electron pair from the peroxyacid and one from the π bond.
- The weak O–O bond is broken.

Problem 12.17 What epoxide is formed when each alkene is treated with mCPBA?



12.8A The Stereochemistry of Epoxidation

Epoxidation occurs via **syn addition** of an O atom from either side of the planar double bond, so that both C–O bonds are formed on the same side. The relative position of substituents in the alkene reactant is **retained** in the epoxide product.


- A **cis** alkene gives an epoxide with **cis** substituents. A **trans** alkene gives an epoxide with **trans** substituents.

Epoxidation is a **stereospecific** reaction because **cis** and **trans** alkenes yield different stereoisomers as products, as illustrated in Sample Problem 12.2.

Sample Problem 12.2 Draw the stereoisomers formed when *cis*- and *trans*-2-butene are epoxidized with mCPBA.


Solution

To draw each product of epoxidation, add an O atom from either side of the alkene, and keep all substituents in their *original* orientations. The **cis** methyl groups in *cis*-2-butene become **cis** substituents in the epoxide. Addition of an O atom from either side of the trigonal planar alkene leads to the same compound—an **achiral meso compound that contains two stereogenic centers**.




Epoxidation of *cis*- and *trans*-2-butene illustrates the general rule about the stereochemistry of reactions: **an achiral starting material gives achiral or racemic products**.

The **trans** methyl groups in *trans*-2-butene, become **trans** substituents in the epoxide. Addition of an O atom from either side of the trigonal planar alkene yields an equal mixture of two enantiomers—a **racemic mixture**.



Problem 12.18

Draw all stereoisomers formed when each alkene is treated with mCPBA.




12.8B The Synthesis of Disparlure



In 1869, the gypsy moth was introduced into New England in an attempt to develop a silk industry. Some moths escaped into the wild and the population flourished. Mature gypsy moth caterpillars eat an average of one square foot of leaf surface per day, defoliating shade trees and entire forests. Many trees die after a single defoliation.

Disparlure, the sex pheromone of the female gypsy moth, is synthesized by a stepwise reaction sequence that uses an epoxidation reaction as the final step. Retrosynthetic analysis of disparlure illustrates three key operations:



- **Step [1]** The cis epoxide in disparlure is prepared from a cis alkene **A** by epoxidation.
- **Step [2]** **A** is prepared from an internal alkyne **B** by reduction.
- **Step [3]** **B** is prepared from acetylene and two 1° alkyl halides (**C** and **D**) by using S_N2 reactions with acetylide anions.

Figure 12.9 illustrates the synthesis of disparlure beginning with acetylene. The synthesis is conceptually divided into three parts:




- **Part [1]** Acetylene is converted to an internal alkyne **B** by forming two C–C bonds. Each bond is formed by treating an alkyne with base (NaNH_2) to form an acetylide anion, which reacts with an alkyl halide (**C** or **D**) in an S_N2 reaction (Section 11.11).
- **Part [2]** The internal alkyne **B** is reduced to a cis alkene **A** by syn addition of H_2 using the Lindlar catalyst (Section 12.5B).
- **Part [3]** The cis alkene **A** is epoxidized to disparlure using a peroxyacid such as mCPBA.

How to separate a racemic mixture into its component enantiomers is discussed in Section 28.3.

Epoxidation of the cis alkene **A** from two different sides of the double bond affords two cis epoxides in the last step—a racemic mixture of two enantiomers. Thus, half of the product is the desired pheromone disparlure, but the other half is its biologically inactive enantiomer. Separating the desired from the undesired enantiomer is difficult and expensive, because both compounds have identical physical properties. A reaction that affords a chiral epoxide from an achiral precursor without forming a racemic mixture is discussed in Section 12.15.

Figure 12.9

The synthesis of disparlure


Part [1] Formation of two C–C bonds using acetylide anions**Part [2]** Reduction of alkyne B to form cis alkene A**Part [3]** Epoxidation of A to form disparlure

- Disparlure has been used to control the spread of the gypsy moth caterpillar, a pest that has periodically devastated forests in the northeastern United States by defoliating many shade and fruit-bearing trees. The active pheromone is placed in a trap containing a poison or sticky substance, and the male moth is lured to the trap by the pheromone. Such a species-specific method presents a new way of controlling an insect population that avoids the widespread use of harmful, nonspecific pesticides.


12.9 Dihydroxylation

Dihydroxylation is the addition of two hydroxy groups to a double bond, forming a **1,2-diol** or **glycol**. Depending on the reagent, the two new OH groups can be added to the opposite sides (**anti** addition) or the same side (**syn** addition) of the double bond.

Dihydroxylation—General reaction




Stereochemistry




12.9A Anti Dihydroxylation

Anti dihydroxylation is achieved in two steps—epoxidation followed by opening of the ring with OH^- or H_2O . Cyclohexene, for example, is converted to a racemic mixture of two *trans*-1,2-cyclohexanediols by anti addition of two OH groups.



The stereochemistry of the products can be understood by examining the stereochemistry of each step.




Epoxidation of cyclohexene adds an O atom from either above or below the plane of the double bond to form a single **achiral epoxide**, so only one representation is shown. Opening of the epoxide ring then occurs with **backside attack at either C–O bond**. Because the epoxide is drawn above the plane of the six-membered ring, nucleophilic attack occurs from **below** the plane. This reaction is a specific example of the opening of epoxide rings with strong nucleophiles, first presented in Section 9.15.

Because one OH group of the 1,2-diol comes from the epoxide and one OH group comes from the nucleophile (OH^-), the overall result is **anti addition of two OH groups** to an alkene.


Problem 12.19 Draw the products formed when both *cis*- and *trans*-2-butene are treated with a peroxyacid followed by OH^- (in H_2O). Explain how these reactions illustrate that anti dihydroxylation is stereospecific.

12.9B Syn Dihydroxylation

Syn dihydroxylation results when an alkene is treated with either KMnO_4 or OsO_4 .




Each reagent adds two oxygen atoms to the same side of the double bond—that is, in a **syn** fashion—to yield a cyclic intermediate. Hydrolysis of the cyclic intermediate cleaves the metal–oxygen bonds, forming the *cis*-1,2-diol. With OsO_4 , sodium bisulfite (NaHSO_3) is also added in the hydrolysis step.




Although KMnO_4 is inexpensive and readily available, its use is limited by its insolubility in organic solvents. To prevent further oxidation of the product 1,2-diol, the reaction mixture must be kept basic with added OH^- .

Although OsO_4 is a more selective oxidant than KMnO_4 and is soluble in organic solvents, it is toxic and expensive. To overcome these limitations, dihydroxylation can be carried out by using a *catalytic* amount of OsO_4 , if the oxidant *N*-methylmorpholine *N*-oxide (NMO) is also added.

NMO is an **amine oxide**. It is not possible to draw a Lewis structure of an amine oxide having only neutral atoms.




In the catalytic process, dihydroxylation of the double bond converts the Os^{8+} oxidant into an Os^{6+} product, which is then re-oxidized by NMO to Os^{8+} . This Os^{8+} reagent can then be used for dihydroxylation once again, and the catalytic cycle continues.




Problem 12.20 Draw the products formed when both *cis*- and *trans*-2-butene are treated with OsO_4 , followed by hydrolysis with $\text{NaHSO}_3 + \text{H}_2\text{O}$. Explain how these reactions illustrate that syn dihydroxylation is stereospecific.

12.10 Oxidative Cleavage of Alkenes

Oxidative cleavage of an alkene breaks both the σ and π bonds of the double bond to form two carbonyl groups. Depending on the number of R groups bonded to the double bond, oxidative cleavage yields either **ketones** or **aldehydes**.




One method of oxidative cleavage relies on a two-step procedure using **ozone (O_3) as the oxidant** in the first step. Cleavage with ozone is called **ozonolysis**.



Addition of ozone to the π bond of the alkene forms an unstable intermediate called a **molozone**, which then rearranges to an **ozonide** by a stepwise process. The unstable ozonide is then reduced without isolation to afford carbonyl compounds. **Zn (in H_2O) or dimethyl sulfide (CH_3SCH_3)** are two common reagents used to convert the ozonide to carbonyl compounds.

The key intermediates in ozonolysis




The pungent odor around a heavily used photocopier machine is O_3 produced from O_2 during the process. O_3 at ground level is an unwanted atmospheric pollutant. In the stratosphere, however, it protects us from harmful ultraviolet radiation, as discussed in Chapter 15.

To draw the product of any oxidative cleavage:

- **Locate all π bonds in the molecule.**
- **Replace each $\text{C}=\text{C}$ by two $\text{C}=\text{O}$ bonds.**

Sample Problem 12.3


Draw the products when each alkene is treated with O_3 followed by CH_3SCH_3 .



Solution


a. **Cleave the double bond and replace it with two carbonyl groups.**

Break both the σ and π bonds.





b. **For a cycloalkene, oxidative cleavage results in a single molecule with two carbonyl groups—a dicarbonyl compound.**


Break both the σ and π bonds.



Problem 12.21 Draw the products formed when each alkene is treated with O_3 followed by Zn, H_2O .




Ozonolysis of dienes (and other polyenes) results in oxidative cleavage of all $C=C$ bonds. The number of carbonyl groups formed in the products is *twice* the number of double bonds in the starting material.




Oxidative cleavage is a valuable tool for structure determination of unknown compounds. The ability to determine what alkene gives rise to a particular set of oxidative cleavage products is thus a useful skill, illustrated in Sample Problem 12.4.

Sample Problem 12.4 What alkene forms the following products after reaction with O_3 followed by CH_3SCH_3 ?




Solution





To draw the starting material, ignore the O atoms in the carbonyl groups and join the carbonyl carbons together by a $C=C$.



Problem 12.22 What alkene yields each set of oxidative cleavage products?




Problem 12.23 Draw the products formed when each diene is treated with O_3 followed by CH_3SCH_3 .




12.11 Oxidative Cleavage of Alkynes


Alkynes also undergo oxidative cleavage of the σ bond and both π bonds of the triple bond. Internal alkynes are oxidized to **carboxylic acids (RCOOH)**, whereas terminal alkynes afford carboxylic acids and CO_2 from the sp hybridized $C-H$ bond.



Oxidative cleavage is commonly carried out with O_3 , followed by cleavage of the intermediate ozonide with H_2O .



Problem 12.24 Draw the products formed when each alkyne is treated with O_3 followed by H_2O .




Problem 12.25 What alkyne (or diyne) yields each set of oxidative cleavage products?

- a. $\text{CO}_2 + \text{CH}_3(\text{CH}_2)_8\text{CO}_2\text{H}$ c. $\text{CH}_3\text{CH}_2\text{CO}_2\text{H}, \text{HO}_2\text{CCH}_2\text{CO}_2\text{H}, \text{CH}_3\text{CO}_2\text{H}$
 b. $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{CO}_2\text{H}$ only d. $\text{HO}_2\text{C}(\text{CH}_2)_{14}\text{CO}_2\text{H}$


12.12 Oxidation of Alcohols

Alcohols are oxidized to a variety of carbonyl compounds, depending on the type of alcohol and reagent. Oxidation occurs by replacing the $\text{C}-\text{H}$ bonds *on the carbon bearing the OH group* by $\text{C}-\text{O}$ bonds.


- **1° Alcohols** are oxidized to either **aldehydes** or **carboxylic acids** by replacing either one or two $\text{C}-\text{H}$ bonds by $\text{C}-\text{O}$ bonds.



- **2° Alcohols** are oxidized to **ketones** by replacing the one $\text{C}-\text{H}$ bond by a $\text{C}-\text{O}$ bond.



- **3° Alcohols** have **no H atoms on the carbon with the OH group**, so they are not easily oxidized.




The oxidation of alcohols to carbonyl compounds is typically carried out with Cr^{6+} oxidants, which are reduced to Cr^{3+} products.

- CrO_3 , $\text{Na}_2\text{Cr}_2\text{O}_7$, and $\text{K}_2\text{Cr}_2\text{O}_7$ are **strong, nonselective oxidants** used in aqueous acid ($\text{H}_2\text{SO}_4 + \text{H}_2\text{O}$).
- **PCC** (Section 12.7) is soluble in CH_2Cl_2 (dichloromethane), and can be used without strong acid present, making it a **more selective, milder oxidant**.

12.12A Oxidation of 2° Alcohols

Any of the Cr^{6+} oxidants effectively oxidize 2° alcohols to ketones.




The mechanism for alcohol oxidation has two key parts: **formation of a chromate ester and loss of a proton**. Mechanism 12.5 is drawn for the oxidation of a general 2° alcohol with CrO_3 .




Mechanism 12.5 Oxidation of an Alcohol with CrO_3

Steps [1] and [2] Formation of the chromate ester



- Nucleophilic attack of the alcohol on the electrophilic metal followed by proton transfer forms a **chromate ester**. The C–H bond in the starting material (the 2° alcohol) is still present in the chromate ester, so there is no net oxidation in Steps [1] and [2].

Step [3] Removal of a proton to form the carbonyl group




- In Step [3], a base (H_2O or a molecule of the starting alcohol) removes a proton, with the electron pair in the C–H bond forming the new π bond of the $\text{C}=\text{O}$. **Oxidation at carbon occurs in this step** because the number of C–H bonds decreases and the number of C–O bonds increases.

These three steps convert the Cr^{6+} oxidant to a Cr^{4+} product, which is then further reduced to a Cr^{3+} product by a series of steps.

12.12B Oxidation of 1° Alcohols

1° Alcohols are oxidized to either aldehydes or carboxylic acids, depending on the reagent.

- 1° Alcohols are oxidized to aldehydes (RCHO) under mild reaction conditions—using **PCC** in CH_2Cl_2 .
- 1° Alcohols are oxidized to carboxylic acids (RCOOH) under harsher reaction conditions: $\text{Na}_2\text{Cr}_2\text{O}_7$, $\text{K}_2\text{Cr}_2\text{O}_7$, or CrO_3 in the presence of H_2O and H_2SO_4 .




The mechanism for the oxidation of 1° alcohols to aldehydes parallels the oxidation of 2° alcohols to ketones detailed in Section 12.12A. Oxidation of a 1° alcohol to a carboxylic acid requires three operations: **oxidation first to the aldehyde, reaction with water**, and then further **oxidation to the carboxylic acid**, as shown in Mechanism 12.6.




Mechanism 12.6 Oxidation of a 1° Alcohol to a Carboxylic Acid

Part [1] Oxidation of a 1° alcohol to an aldehyde




- Oxidation of a 1° alcohol to an aldehyde occurs by the three-step mechanism detailed in Mechanism 12.5.

Part [2] Addition of H₂O to form a hydrate




- The aldehyde reacts with H₂O to form a **hydrate**, a compound with two OH groups on the same carbon atom. Hydrates are discussed in greater detail in Section 21.13.

Part [3] Oxidation of the hydrate to a carboxylic acid




- The C–H bond of the hydrate is then oxidized with the Cr⁶⁺ reagent, following Mechanism 12.5. Because the hydrate contains two OH groups, the product of oxidation is a carboxylic acid.

Cr⁶⁺ oxidations are characterized by a color change, as the **red-orange Cr⁶⁺ reagent** is reduced to **green Cr³⁺**. The first devices used to measure blood alcohol content in individuals suspected of “driving under the influence” made use of this color change. Oxidation of CH₃CH₂OH, the 1° alcohol in alcoholic beverages, with orange K₂Cr₂O₇ forms CH₃COOH and green Cr³⁺.



Blood alcohol level can be determined by having an individual blow into a tube containing K₂Cr₂O₇, H₂SO₄, and an inert solid. The alcohol in the exhaled breath is oxidized by the Cr⁶⁺ reagent, which turns green in the tube (Figure 12.10). The higher the concentration of CH₃CH₂OH


Figure 12.10
Blood alcohol screening



- The oxidation of CH₃CH₂OH with K₂Cr₂O₇ to form CH₃COOH and Cr³⁺ was the first available method for the routine testing of alcohol concentration in exhaled air. Some consumer products for alcohol screening are still based on this technology.

in the breath, the more Cr^{6+} is reduced, and the farther the green Cr^{3+} color extends down the length of the sample tube. This value is then correlated with blood alcohol content to determine if an individual has surpassed the legal blood alcohol limit.

Problem 12.26 Draw the organic products in each of the following reactions.




12.13 Green Chemistry

Several new methods of oxidation are based on green chemistry. **Green chemistry is the use of environmentally benign methods to synthesize compounds.** Its purpose is to use safer reagents and less solvent, and develop reactions that form fewer by-products and generate less waste. Although the concept of designing green chemical processes was only formally established by the Environmental Protection Agency in the early 1990s, developing new chemical methods that minimize environmental impact has been in practice much longer.


Green polymer synthesis using starting materials derived from renewable resources (rather than petroleum) is discussed in Section 30.8.

Since many oxidation methods use toxic reagents (such as OsO_4 and O_3) and corrosive acids (such as H_2SO_4), or they generate carcinogenic by-products (such as Cr^{3+}), alternative reactions have been developed. One method uses a polymer-supported Cr^{6+} reagent— HCrO_4^- -Amberlyst A-26 resin—that avoids the use of strong acid, and forms a Cr^{3+} by-product that can be easily removed from the product by filtration.


The Amberlyst A-26 resin consists of a complex hydrocarbon network with cationic ammonium ion appendages that serve as counterions to the anionic chromium oxidant, HCrO_4^- . Heating the insoluble polymeric reagent with an alcohol results in oxidation to a carbonyl compound, with formation of an insoluble Cr^{3+} by-product. Not only can the metal by-product be removed by filtration without added solvent, it can also be regenerated and reused in a subsequent reaction.



With HCrO_4^- -Amberlyst A-26 resin, 1° alcohols are oxidized to aldehydes and 2° alcohols are oxidized to ketones.



Many other green approaches to oxidation that avoid the generation of metal by-products entirely are also under active investigation.




K. Barry Sharpless, currently at The Scripps Research Institute, reasoned that using a chiral reagent might make it possible to favor the formation of one enantiomer over the other.


K. Barry Sharpless shared the 2001 Nobel Prize in Chemistry for his work on chiral oxidation reactions.

- An *enantioselective* reaction affords predominantly or exclusively one enantiomer.
- A reaction that converts an achiral starting material into predominantly one enantiomer is also called an *asymmetric reaction*.

The Sharpless asymmetric epoxidation is an enantioselective reaction that oxidizes alkenes to epoxides. Only the double bonds of **allylic alcohols**—that is, alcohols having a hydroxy group on the carbon adjacent to a C=C—are oxidized in this reaction.




The **Sharpless reagent** consists of three components: *tert*-butyl hydroperoxide, (CH₃)₃COOH; a titanium catalyst—usually titanium(IV) isopropoxide, Ti[OCH(CH₃)₂]₄; and **diethyl tartrate (DET)**. There are two different chiral diethyl tartrate isomers, labeled as (+)-DET or (–)-DET to indicate the direction in which they rotate polarized light.



The identity of the DET isomer determines which enantiomer is the major product obtained in the epoxidation of an allylic alcohol with the Sharpless reagent.



(+)-DET is prepared from (+)-(*R,R*)-tartaric acid [HO₂CCH(OH)CH(OH)CO₂H], a naturally occurring carboxylic acid found in grapes and sold as a by-product of the wine industry.




Enantiomeric excess = **ee** =
% of one enantiomer – % of
the other enantiomer.

The degree of enantioselectivity of a reaction is measured by its enantiomeric excess (**ee**) (Section 5.12D). Reactions [1] and [2] are highly enantioselective because each has an enantiomeric excess of 95% (97.5% of the major enantiomer – 2.5% of the minor enantiomer).


To determine which enantiomer is formed for a given isomer of DET, draw the allylic alcohol in a plane, with the **C=C horizontal and the OH group in the upper right corner**; then:

- Epoxidation with (–)-DET adds an oxygen atom from above the plane.
- Epoxidation with (+)-DET adds an oxygen atom from below the plane.



Sample Problem 12.5

Predict the major product in each epoxidation.




Solution


To draw an epoxidation product:

- Draw the allylic alcohol with the **C=C horizontal and the OH group in the upper right corner of the alkene**. Re-draw the alkene if necessary.
- **(+)-DET** adds the O atom from **below**, and **(–)-DET** adds the O atom from **above**.

a. Since the C=C is drawn horizontal with the OH group in the upper right corner, it is not necessary to re-draw the alkene. With **(+)-DET**, the O atom is added from **below**.



b. The allylic alcohol must be re-drawn with the C=C horizontal and the OH group in the upper right corner. Because **(–)-DET** is used, the O atom is then added from **above**.



Problem 12.30

Draw the products of each Sharpless epoxidation.





Figure 12.11


The synthesis of chiral insect pheromones using asymmetric epoxidation



- The bonds in the products that originate from the epoxide intermediate are indicated in red.

The Sharpless epoxidation has been used to synthesize many chiral natural products, including two insect pheromones—(+)- α -multistriatin and (-)-frontalin, as shown in Figure 12.11.

Problem 12.31 Explain why only one C=C of geraniol is epoxidized with the Sharpless reagent.



KEY CONCEPTS

Oxidation and Reduction

Summary: Terms that Describe Reaction Selectivity

- A **regioselective reaction** forms predominantly or exclusively one constitutional isomer (Section 8.5).
- A **stereoselective reaction** forms predominantly or exclusively one stereoisomer (Section 8.5).
- An **enantioselective reaction** forms predominantly or exclusively one enantiomer (Section 12.15).

Definitions of Oxidation and Reduction (12.1)

Oxidation reactions result in:


- an increase in the number of C–Z bonds, or
- a decrease in the number of C–H bonds

Reduction reactions result in:

- a decrease in the number of C–Z bonds, or
- an increase in the number of C–H bonds


Reduction Reactions

[1] Reduction of alkenes—Catalytic hydrogenation (12.3)




- **Syn addition** of H₂ occurs.
- Increasing alkyl substitution on the C=C decreases the rate of reaction.


[2] Reduction of alkynes



- Two equivalents of H₂ are added and four new C-H bonds are formed (12.5A).




- **Syn addition** of H₂ occurs, forming a **cis** alkene (12.5B).
- The Lindlar catalyst is deactivated; reaction stops after one equivalent of H₂ has added.




- **Anti addition** of H₂ occurs, forming a **trans** alkene (12.5C).

[3] Reduction of alkyl halides (12.6)



- The reaction follows an **S_N2** mechanism.
- CH₃X and RCH₂X react faster than a more substituted RX.

[4] Reduction of epoxides (12.6)




- The reaction follows an **S_N2** mechanism.
- In unsymmetrical epoxides, H⁻ (from LiAlH₄) attacks at the less substituted carbon.

Oxidation Reactions


[1] Oxidation of alkenes

a. Epoxidation (12.8)




- The mechanism has **one step**.
- **Syn addition** of an O atom occurs.
- The reaction is stereospecific.

b. Anti dihydroxylation (12.9A)




- Opening of an epoxide ring intermediate with ⁻OH or H₂O forms a 1,2-diol with two OH groups added in an **anti** fashion.

c. Syn dihydroxylation (12.9B)




- Each reagent adds two new C-O bonds to the C=C in a **syn** fashion.

d. Oxidative cleavage (12.10)




- Both the σ and π bonds of the alkene are cleaved to form two carbonyl groups.


[2] Oxidative cleavage of alkynes (12.11)




- The σ bond and both π bonds of the alkyne are cleaved.




[3] Oxidation of alcohols (12.12, 12.13)



- Oxidation of a 1° alcohol with PCC or HCrO_4^- -Amberlyst A-26 resin stops at the aldehyde stage. Only one C-H bond is replaced by a C-O bond.




- Oxidation of a 1° alcohol under harsher reaction conditions— CrO_3 (or $\text{Na}_2\text{Cr}_2\text{O}_7$ or $\text{K}_2\text{Cr}_2\text{O}_7$) + H_2O + H_2SO_4 —leads to a RCOOH . Two C-H bonds are replaced by two C-O bonds.



- Because a 2° alcohol has only one C-H bond on the carbon bearing the OH group, all Cr^{6+} reagents—PCC, CrO_3 , $\text{Na}_2\text{Cr}_2\text{O}_7$, $\text{K}_2\text{Cr}_2\text{O}_7$, or HCrO_4^- (Amberlyst A-26 resin)—oxidize a 2° alcohol to a ketone.




[4] Asymmetric epoxidation of allylic alcohols (12.15)



PROBLEMS





Classifying Reactions as Oxidation or Reduction

12.32 Label each reaction as oxidation, reduction, or neither.



Hydrogenation

12.33 Draw the organic products formed when each alkene is treated with H_2 , Pd-C. Indicate the three-dimensional structure of all stereoisomers formed.




12.34 Match each alkene to its heat of hydrogenation.

Alkenes: 3-methyl-1-butene, 2-methyl-1-butene, 2-methyl-2-butene
 ΔH° (hydrogenation) kJ/mol: -119, -127, -112

12.35 How many rings and π bonds are contained in compounds **A–C**? Draw one possible structure for each compound.


- Compound **A** has molecular formula C_5H_8 and is hydrogenated to a compound having molecular formula C_5H_{10} .
- Compound **B** has molecular formula $\text{C}_{10}\text{H}_{16}$ and is hydrogenated to a compound having molecular formula $\text{C}_{10}\text{H}_{18}$.
- Compound **C** has molecular formula C_8H_8 and is hydrogenated to a compound having molecular formula C_8H_{16} .

12.36 For alkenes **A**, **B**, and **C**: (a) Rank **A**, **B**, and **C** in order of increasing heat of hydrogenation; (b) rank **A**, **B**, and **C** in order of increasing rate of reaction with H_2 , Pd-C; (c) draw the products formed when each alkene is treated with ozone, followed by Zn , H_2O .



12.37 A chiral compound **X** having the molecular formula C_6H_{12} is converted to 3-methylpentane with H_2 , Pd-C. Draw all possible structures for **X**.

12.38 Stearidonic acid ($\text{C}_{18}\text{H}_{28}\text{O}_2$) is an unsaturated fatty acid obtained from oils isolated from hemp and blackcurrant (see also Problem 10.12).



- What fatty acid is formed when stearidonic acid is hydrogenated with excess H_2 and a Pd catalyst?
- What fatty acids are formed when stearidonic acid is hydrogenated with one equivalent of H_2 and a Pd catalyst?
- Draw the structure of a possible product formed when stearidonic acid is hydrogenated with one equivalent of H_2 and a Pd catalyst, and one double bond is isomerized to a trans isomer.
- How do the melting points of the following fatty acids compare: stearidonic acid; one of the products formed in part (b); the product drawn in part (c)?

Reactions—General


12.39 Draw the organic products formed when cyclopentene is treated with each reagent. With some reagents, no reaction occurs.

- | | | |
|---|--|--|
| a. $\text{H}_2 + \text{Pd-C}$ | e. [1] $\text{CH}_3\text{CO}_3\text{H}$; [2] H_2O , HO^- | i. [1] O_3 ; [2] CH_3SCH_3 |
| b. $\text{H}_2 + \text{Lindlar catalyst}$ | f. [1] $\text{OsO}_4 + \text{NMO}$; [2] NaHSO_3 , H_2O | j. $(\text{CH}_3)_3\text{COOH}$, $\text{Ti}[\text{OCH}(\text{CH}_3)_2]_4$, (-)-DET |
| c. Na , NH_3 | g. KMnO_4 , H_2O , HO^- | k. mCPBA |
| d. $\text{CH}_3\text{CO}_3\text{H}$ | h. [1] LiAlH_4 ; [2] H_2O | l. Product in (k); then [1] LiAlH_4 ; [2] H_2O |

12.40 Draw the organic products formed when 4-octyne is treated with each reagent.


- | | | | |
|---------------------------------|---|--------------------------------|--|
| a. H_2 (excess) + Pd-C | b. $\text{H}_2 + \text{Lindlar catalyst}$ | c. Na , NH_3 | d. [1] O_3 ; [2] H_2O |
|---------------------------------|---|--------------------------------|--|

12.41 Draw the organic products formed when allylic alcohol **A** is treated with each reagent.






- | | |
|--|---|
| a. $\text{H}_2 + \text{Pd-C}$ | e. $(\text{CH}_3)_3\text{COOH}, \text{Ti}[\text{OCH}(\text{CH}_3)_2]_4, (+)\text{-DET}$ |
| b. mCPBA | f. $(\text{CH}_3)_3\text{COOH}, \text{Ti}[\text{OCH}(\text{CH}_3)_2]_4, (-)\text{-DET}$ |
| c. PCC | g. [1] PBr_3 ; [2] LiAlH_4 ; [3] H_2O |
| d. $\text{CrO}_3, \text{H}_2\text{SO}_4, \text{H}_2\text{O}$ | h. HCrO_4^- -Amberlyst A-26 resin |

12.42 Draw the products formed when allylic alcohol **B** is treated with each reagent. Indicate the stereochemistry of any stereoisomers formed.






- | | |
|--|---|
| a. $\text{H}_2 + \text{Pd-C}$ | e. [1] OsO_4 ; [2] $\text{NaHSO}_3, \text{H}_2\text{O}$ |
| b. $\text{Na}_2\text{Cr}_2\text{O}_7, \text{H}_2\text{SO}_4, \text{H}_2\text{O}$ | f. [1] HCO_3H ; [2] $\text{H}_2\text{O}, \text{HO}^-$ |
| c. PCC | g. $(\text{CH}_3)_3\text{COOH}, \text{Ti}[\text{OCH}(\text{CH}_3)_2]_4, (+)\text{-DET}$ |
| d. $\text{CF}_3\text{CO}_3\text{H}$ | h. $\text{KMnO}_4, \text{H}_2\text{O}, \text{HO}^-$ |


12.43 Draw the organic products formed in each reaction.






12.44 Draw the organic products formed in each reaction.




12.45 Identify the reagents needed to carry out each transformation.




12.46 What alkene is needed to synthesize each 1,2-diol using [1] OsO_4 followed by NaHSO_3 in H_2O ; or [2] $\text{CH}_3\text{CO}_3\text{H}$ followed by OH^- in H_2O ?



12.47 Draw a stepwise mechanism for the reduction of epoxide **A** to alcohol **B** using LiAlH_4 . What product would be formed if LiAlD_4 were used as reagent? Indicate the stereochemistry of all stereogenic centers in the product using wedges and dashes.







- 12.48 Draw the products formed after Steps [1] and [2] in the following three-step sequence. Then draw stepwise mechanisms for each step.





Oxidative Cleavage



- 12.49 Draw the products formed in each oxidative cleavage.




- 12.50 What alkene yields each set of products after treatment with O_3 followed by CH_3SCH_3 ?






- 12.51 Identify the starting material in each reaction.



- 12.52 What alkyne gives each set of products after treatment with O_3 followed by H_2O ?



- 12.53 Draw the products formed when each naturally occurring compound is treated with O_3 followed by $\text{Zn, H}_2\text{O}$.



Identifying Compounds from Reactions


- 12.54 Identify compounds **A**, **B**, and **C**.

a. Compound **A** has molecular formula C_8H_{12} and reacts with two equivalents of H_2 . **A** gives $\text{HCOCH}_2\text{CH}_2\text{CHO}$ as the only product of oxidative cleavage with O_3 followed by CH_3SCH_3 .


b. Compound **B** has molecular formula C_6H_{10} and gives $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{CH}_3$ when treated with excess H_2 in the presence of Pd. **B** reacts with NaNH_2 and CH_3I to form compound **C** (molecular formula C_7H_{12}).

- 12.55 Oximene and myrcene, two hydrocarbons isolated from alfalfa that have the molecular formula $\text{C}_{10}\text{H}_{16}$, both yield 2,6-dimethyloctane when treated with H_2 and a Pd catalyst. Ozonolysis of oximene forms $(\text{CH}_3)_2\text{C}=\text{O}$, $\text{CH}_2=\text{O}$, $\text{CH}_2(\text{CHO})_2$, and CH_3COCHO . Ozonolysis of myrcene yields $(\text{CH}_3)_2\text{C}=\text{O}$, $\text{CH}_2=\text{O}$ (two equiv), and $\text{HCOCH}_2\text{CH}_2\text{COCHO}$. Identify the structures of oximene and myrcene.

- 12.56** An achiral hydrocarbon **A** of molecular formula C_7H_{12} reacts with two equivalents of H_2 in the presence of Pd-C to form $CH_3CH_2CH_2CH_2CH(CH_3)_2$. One oxidative cleavage product formed by the treatment of **A** with O_3 is CH_3COOH . Reaction of **A** with H_2 and Lindlar catalyst forms **B**, and reaction of **A** with Na, NH_3 forms **C**. (a) Identify compounds **A**, **B**, and **C**. (b) Explain why **A** does not react with NaH.
- 12.57** An unknown compound **A** of molecular formula $C_{10}H_{18}O$ reacts with H_2SO_4 to form two compounds (**B** and **C**) of molecular formula $C_{10}H_{16}$. **B** and **C** both react with H_2 in the presence of Pd-C to form decalin. Ozonolysis of **B** forms **D**, and ozonolysis of **C** forms a diketone **E** of molecular formula $C_{10}H_{16}O_2$. Identify the structures of compounds **A**, **B**, **C**, and **E**.





decalin

**D**


- 12.58** DHA is a fatty acid derived from fish oil and an abundant fatty acid in vertebrate brains. Hydrogenation of DHA forms docosanoic acid [$CH_3(CH_2)_{20}CO_2H$] and ozonolysis forms CH_3CH_2CHO , $CH_2(CHO)_2$ (five equivalents), and $OHCCH_2CH_2CO_2H$. What is the structure of DHA if all double bonds have the *Z* configuration?
- 12.59** One compound that contributes to the “seashore smell” at beaches in Hawaii is dictyopterene D', a component of a brown edible seaweed called limu lipoa. Hydrogenation of dictyopterene D' with excess H_2 in the presence of a Pd catalyst forms butylcycloheptane. Ozonolysis with O_3 followed by $(CH_3)_2S$ forms $CH_2(CHO)_2$, $OHCCH_2CH(CHO)_2$, and CH_3CH_2CHO . What are possible structures of dictyopterene D'?

Sharpless Asymmetric Epoxidation

- 12.60** Draw the product of each asymmetric epoxidation reaction.





- 12.61** Epoxidation of the following allylic alcohol using the Sharpless reagent with (-)-DET gives two epoxy alcohols in a ratio of 87:13.







- a. Assign structures to the major and minor product.
b. What is the enantiomeric excess in this reaction?

- 12.62** What allylic alcohol and DET isomer are needed to make each chiral epoxide using a Sharpless asymmetric epoxidation reaction?




Synthesis

- 12.63** Devise a synthesis of each hydrocarbon from acetylene, and any other needed reagents.







- 12.64** Devise a synthesis of muscalure, the sex pheromone of the common housefly, from acetylene and any other required reagents.





- 12.65** It is sometimes necessary to isomerize a *cis* alkene to a *trans* alkene in a synthesis, a process that cannot be accomplished in a single step. Using the reactions you have learned in Chapters 8–12, devise a stepwise method to convert *cis*-2-butene to *trans*-2-butene.

12.66 Devise a synthesis of each compound from acetylene and any other required reagents.






12.67 Devise a synthesis of each compound from styrene ($C_6H_5CH=CH_2$). You may use any inorganic or organic reagents. More than one step is required.







12.68 Devise a synthesis of (2*E*)-2-hexene from 1-pentene and any needed organic compounds or inorganic reagents.





12.69 Devise a synthesis of each compound from the indicated starting material and any other required reagents.



12.70 Devise a synthesis of each compound from acetylene and organic compounds containing two carbons or fewer. You may use any other required reagents.





12.71 Devise a synthesis of each compound from ethynylcyclohexane. You may use any other required reagents.




12.72 Devise a synthesis of (3*R*,4*S*)-3,4-dichlorohexane from acetylene and any needed organic compounds or inorganic reagents.

12.73 Devise a synthesis of each compound from CH_3CH_2OH as the only organic starting material; that is, every carbon in the product must come from a molecule of ethanol. You may use any other needed inorganic reagents.




Challenge Problems


12.74 The Birch reduction is a dissolving metal reaction that converts substituted benzenes to 1,4-cyclohexadienes using Li and liquid ammonia in the presence of an alcohol. Draw a stepwise mechanism for the following Birch reduction.




12.75 In the Cr^{6+} oxidation of cyclohexanols, it is generally true that sterically hindered alcohols react faster than unhindered alcohols. Which of the following alcohols should be oxidized more rapidly?



12.76 Draw a stepwise mechanism for the following reaction.



12.77 Although reagents can always add to an alkene from either side, sometimes one side of the double bond is more sterically hindered, so an unequal mixture of addition products results. For example, when **X** is treated with H_2 in the presence of a Pd catalyst, 80% of the product mixture contains the cis isomer **Y** and only 20% is the trans isomer **Z**. Thus, addition of H_2 occurs predominantly on the side opposite to the bulky *tert*-butyl group, resulting in a new equatorial C–H bond. Keeping this in mind, what is the major epoxidation product formed from **X** under each of the following reaction conditions: (a) $mCPBA$; or (b) Br_2 , H_2O followed by NaH ?



Mass Spectrometry and Infrared Spectroscopy

13

- 13.1 Mass spectrometry
- 13.2 Alkyl halides and the M + 2 peak
- 13.3 Fragmentation
- 13.4 Other types of mass spectrometry
- 13.5 Electromagnetic radiation
- 13.6 Infrared spectroscopy
- 13.7 IR absorptions
- 13.8 IR and structure determination



The serendipitous discovery of **penicillin** by Scottish bacteriologist Sir Alexander Fleming in 1928 is considered one of the single most important events in the history of medicine. Penicillin G and related compounds are members of the β -lactam family of antibiotics, all of which contain a strained four-membered amide ring that is responsible for their biological activity. Penicillin was first used to cure a streptococcal infection in 1942, and by 1944 penicillin production was given high priority by the United States government, because it was needed to treat the many injured soldiers in World War II. The unusual structure of penicillin was elucidated by modern instrumental methods in the 1940s. In Chapter 13, we learn about mass spectrometry and infrared spectroscopy, two techniques for characterizing organic compounds like penicillin.

Whether a compound is prepared in the laboratory or isolated from a natural source, a chemist must determine its identity. Seventy years ago, determining the structure of an organic compound involved a series of time-consuming operations: measuring physical properties (melting point, boiling point, solubility, and density), identifying the functional groups using a series of chemical tests, and converting an unknown compound into another compound whose physical and chemical properties were then characterized as well.

Although still a challenging task, structure determination has been greatly simplified by modern instrumental methods. These techniques have both decreased the time needed for compound characterization, and increased the complexity of compounds whose structures can be completely determined.

In Chapter 13 we examine **mass spectrometry (MS)**, which is used to determine the molecular weight and molecular formula of a compound, and **infrared (IR) spectroscopy**, a tool used to identify a compound's functional groups. Chapter 14 is devoted to **nuclear magnetic resonance (NMR) spectroscopy**, which is used to identify the carbon–hydrogen framework in a compound, making it the most powerful spectroscopic tool for organic structure analysis. **Each of these methods relies on the interaction of an energy source with a molecule to produce a change that is recorded in a spectrum.**


13.1 Mass Spectrometry

Mass spectrometry is a technique used for measuring the molecular weight and determining the molecular formula of an organic molecule.

13.1A General Features


In the most common type of **mass spectrometer**, a molecule is vaporized and ionized, usually by bombardment with a beam of high-energy electrons, as shown in Figure 13.1. The energy of these electrons is typically about 6400 kJ, or 70 electron volts (eV). Because it takes ~400 kJ of energy to cleave a typical σ bond, 6400 kJ is an enormous amount of energy to come into contact with a molecule. This electron beam ionizes a molecule by causing it to eject an electron.

Figure 13.1
Schematic of a mass spectrometer



In a mass spectrometer, a sample is vaporized and bombarded by a beam of electrons to form an unstable radical cation, which then decomposes to smaller fragments. The positively charged ions are accelerated toward a negatively charged plate, and then passed through a curved analyzer tube in a magnetic field, where they are deflected by different amounts depending on their ratio of mass to charge (m/z). A mass spectrum plots the intensity of each ion versus its m/z ratio.


The term **spectroscopy** is usually used for techniques that use electromagnetic radiation as an energy source. Because the energy source in MS is a beam of electrons, the term **mass spectrometry** is used instead.



The species formed is a **radical cation**, symbolized $\text{M}^{+\cdot}$. It is a radical because it has an unpaired electron, and it is a cation because it has one fewer electron than it started with.

- The radical cation $\text{M}^{+\cdot}$ is called the *molecular ion* or the *parent ion*.


A single electron has a negligible mass, so the **mass of $\text{M}^{+\cdot}$ represents the molecular weight of M**. Because the molecular ion $\text{M}^{+\cdot}$ is inherently unstable, it decomposes. Single bonds break to form **fragments, radicals and cations having a lower molecular weight than the molecular ion**. A mass spectrometer analyzes the masses of cations only. The cations are accelerated in an electric field and deflected in a curved path in a magnetic field, thus sorting the molecular ion and its fragments by their **mass-to-charge (m/z) ratio**. Because z is almost always +1, m/z actually measures the mass (m) of the individual ions.



- A **mass spectrum** plots the amount of each cation (its relative abundance) versus its mass.

The whole-number mass of CH_4 is $(1 \text{ C} \times 12 \text{ amu}) + (4 \text{ H} \times 1 \text{ amu}) = 16 \text{ amu}$; amu = atomic mass unit.

A mass spectrometer analyzes the masses of *individual* molecules, not the weighted average mass of a group of molecules, so the whole-number masses of the most common individual isotopes must be used to calculate the mass of the molecular ion. Thus, the mass of the molecular ion for CH_4 should be 16. As a result, the mass spectrum of CH_4 shows a line for the molecular ion—the parent peak or **M peak**—at $m/z = 16$.



The tallest peak in a mass spectrum is called the **base peak**. For CH_4 , the base peak is also the **M peak**, although this may *not* always be the case for all organic compounds.

The mass spectrum of CH_4 consists of more peaks than just the **M peak**. What is responsible for the peaks at $m/z < 16$? Because the molecular ion is unstable, it fragments into other cations and radical cations containing one, two, three, or four fewer hydrogen atoms than methane itself. Thus, the peaks at $m/z = 15, 14, 13,$ and 12 , are due to these lower molecular weight fragments. The decomposition of a molecular ion into lower molecular weight fragments is called **fragmentation**.





Figure 13.2

Mass spectrum of hexane
($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$)



- The molecular ion for hexane (molecular formula C_6H_{14}) is at $m/z = 86$.
- The base peak (relative abundance = 100) occurs at $m/z = 57$.
- A small $M + 1$ peak occurs at $m/z = 87$.

What is responsible for the small peak at $m/z = 17$ in the mass spectrum of CH_4 ? Although most carbon atoms have an atomic mass of 12, 1.1% of them have an additional neutron in the nucleus, giving them an atomic mass of 13. When one of these carbon-13 isotopes forms methane, it gives a molecular ion peak at $m/z = 17$ in the mass spectrum. This peak is called the $M + 1$ peak.


These key features—the molecular ion, the base peak, and the $M + 1$ peak—are illustrated in the mass spectrum of hexane in Figure 13.2.



13.1B Analyzing Unknowns Using the Molecular Ion

Because the **mass of the molecular ion equals the molecular weight of a compound**, a mass spectrum can be used to distinguish between compounds that have similar physical properties but different molecular weights, as illustrated in Sample Problem 13.1.

Sample Problem 13.1

Pentane, 1-pentene, and 1-pentyne are low-boiling hydrocarbons that have different molecular ions in their mass spectra. Match each hydrocarbon to its mass spectrum.





Solution

To solve this problem, first determine the molecular formula and molecular weight of each compound. Then, because the molecular weight of the compound equals the mass of the molecular ion, match the molecular weight to m/z for the molecular ion:


Compound	Molecular formula	Molecular weight = m/z of molecular ion	Spectrum
pentane, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$	C_5H_{12}	72	[3]
1-pentene, $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_3$	C_5H_{10}	70	[2]
1-pentyne, $\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$	C_5H_8	68	[1]

Problem 13.1


What is the mass of the molecular ion formed from compounds having each molecular formula: (a) $\text{C}_3\text{H}_6\text{O}$; (b) $\text{C}_{10}\text{H}_{20}$; (c) $\text{C}_8\text{H}_8\text{O}_2$; (d) methamphetamine ($\text{C}_{10}\text{H}_{15}\text{N}$)?

How to use the mass of the molecular ion to propose molecular formulas for an unknown is shown in Sample Problem 13.2. In this process, keep in mind the following useful fact. Hydrocarbons like methane (CH_4) and hexane (C_6H_{14}), as well as compounds that contain only C, H, and O atoms, always have a molecular ion with an *even* mass. An odd molecular ion generally indicates that a compound contains nitrogen.

The effect of N atoms on the mass of the molecular ion in a mass spectrum is called the **nitrogen rule: A compound that contains an odd number of N atoms gives an odd molecular ion.** Conversely, a compound that contains an *even* number of N atoms (including *zero*) gives an *even* molecular ion. Two “street” drugs that mimic the effects of heroin illustrate this principle: 3-methylfentanyl (two N atoms, even molecular weight) and MPPP (one N atom, odd molecular weight).



3-methylfentanyl
 $C_{23}H_{30}N_2O$
 molecular weight = 350



MPPP
 (1-methyl-4-phenyl-4-propionyloxypiperidine)
 $C_{15}H_{21}NO_2$
 molecular weight = 247

Sample Problem 13.2 Propose possible molecular formulas for a compound with a molecular ion at $m/z = 86$.

Solution

Because the molecular ion has an **even** mass, the compound likely contains C, H, and possibly O atoms. Begin by determining the molecular formula for a hydrocarbon having a molecular ion at 86. Then, because the mass of an O atom is 16 (the mass of CH_4), replace CH_4 by O to give a molecular formula containing one O atom. Repeat this last step to give possible molecular formulas for compounds with two or more O atoms.

For a molecular ion at $m/z = 86$:


Possible hydrocarbons:

- Divide 86 by 12 (mass of 1 C atom). This gives the maximum number of C's possible.

$$\frac{86}{12} = 7 \text{ C's maximum} \rightarrow \boxed{C_7H_2}$$


(remainder = 2)

- Replace 1 C by 12 H's for another possible molecular formula.




Possible compounds with C, H, and O:

- Substitute 1 O for CH_4 . (This can't be done for C_7H_2 .)



- Repeat the process.



Problem 13.2 Propose two molecular formulas for each of the following molecular ions: (a) 72; (b) 100; (c) 73.

13.2 Alkyl Halides and the $M + 2$ Peak

Most of the elements found in organic compounds, such as carbon, hydrogen, oxygen, nitrogen, sulfur, phosphorus, fluorine, and iodine, have one major isotope. **Chlorine** and **bromine**, on the other hand, have two, giving characteristic patterns to the mass spectra of their compounds.

Chlorine has two common isotopes, ^{35}Cl and ^{37}Cl , which occur naturally in a 3:1 ratio. Thus, **there are two peaks in a 3:1 ratio for the molecular ion of an alkyl chloride**. The larger peak—the **M** peak—corresponds to the compound containing ^{35}Cl , and the smaller peak—the **M + 2** peak—corresponds to the compound containing ^{37}Cl .

- When the molecular ion consists of two peaks (M and $M + 2$) in a 3:1 ratio, a Cl atom is present.

Sample Problem 13.3 What molecular ions will be present in a mass spectrum of 2-chloropropane, $(\text{CH}_3)_2\text{CHCl}$?


Solution

Calculate the molecular weight using each of the common isotopes of Cl.

Molecular formula	Mass of molecular ion (m/z)
$C_3H_7^{35}\text{Cl}$	78 (M peak)
$C_3H_7^{37}\text{Cl}$	80 (M + 2 peak)

There should be two peaks in a ratio of 3:1, at $m/z = 78$ and 80, as illustrated in the mass spectrum of 2-chloropropane in Figure 13.3.

Figure 13.3
Mass spectrum of
2-chloropropane [(CH₃)₂CHCl]



Bromine has two common isotopes, ⁷⁹Br and ⁸¹Br, which occur naturally in a 1:1 ratio. Thus, **there are two peaks in a 1:1 ratio for the molecular ion of an alkyl bromide.** In the mass spectrum of 2-bromopropane (Figure 13.4), for example, there is an *M* peak at *m/z* = 122 and an *M* + 2 peak at *m/z* = 124.

- When the molecular ion consists of two peaks (*M* and *M* + 2) in a 1:1 ratio, a Br atom is present in the molecule.

Problem 13.3 What molecular ions would you expect for compounds having each of the following molecular formulas: (a) C₄H₉Cl; (b) C₃H₇F; (c) C₆H₁₁Br; (d) C₄H₁₁N; (e) C₄H₄N₂?

13.3 Fragmentation

While many chemists use a mass spectrum to determine only a compound's molecular weight and molecular formula, additional useful structural information can be obtained from fragmentation patterns. Although each organic compound fragments in a unique way, a particular functional group exhibits common fragmentation patterns.

13.3A General Features of Fragmentation

As an example, consider hexane, whose mass spectrum was shown in Figure 13.2. When hexane is bombarded by an electron beam, it forms a highly unstable radical cation (*m/z* = 86) that can decompose by cleavage of any of the C–C bonds. Thus, cleavage of the terminal C–C bond

Figure 13.4
Mass spectrum of
2-bromopropane [(CH₃)₂CHBr]






Figure 13.5
Identifying fragments in the
mass spectrum of hexane



- Cleavage of C–C bonds (labeled [1]–[4]) in hexane forms lower molecular weight fragments that correspond to lines in the mass spectrum. Although the mass spectrum is complex, possible structures can be assigned to some of the fragments, as shown.

forms $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2^+$ and CH_3^\bullet . Fragmentation generates a cation and a radical, and cleavage generally yields the more stable, more substituted carbocation.



- Loss of a CH_3 group always forms a fragment with a mass 15 units less than the molecular ion.


As a result, the mass spectrum of hexane shows a peak at $m/z = 71$ due to $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2^+$. Figure 13.5 illustrates how cleavage of other C–C bonds in hexane gives rise to other fragments that correspond to peaks in its mass spectrum.

Sample Problem 13.4

The mass spectrum of 2,3-dimethylpentane $[(\text{CH}_3)_2\text{CHCH}(\text{CH}_3)\text{CH}_2\text{CH}_3]$ shows fragments at $m/z = 85$ and 71. Propose possible structures for the ions that give rise to these peaks.

Solution

To solve a problem of this sort, first calculate the mass of the molecular ion. Draw out the structure of the compound, break a C–C bond, and calculate the mass of the resulting fragments. Repeat this process on different C–C bonds until fragments of the desired mass-to-charge ratio are formed.




In this example, 2,3-dimethylpentane has a molecular ion at $m/z = 100$. Cleavage of bond [1] forms a 2° carbocation with $m/z = 85$ and CH_3^\bullet . Cleavage of bond [2] forms another 2° carbocation with $m/z = 71$ and $\text{CH}_3\text{CH}_2^\bullet$. Thus, the fragments at $m/z = 85$ and 71 are possibly due to the two carbocations drawn.

Problem 13.4 The mass spectrum of 2,3-dimethylpentane also shows peaks at $m/z = 57$ and 43 . Propose possible structures for the ions that give rise to these peaks.


Problem 13.5 The base peak in the mass spectrum of 2,2,4-trimethylpentane $[(\text{CH}_3)_3\text{CCH}_2\text{CH}(\text{CH}_3)_2]$ occurs at $m/z = 57$. What ion is responsible for this peak and why is this ion the most abundant fragment?

13.3B Fragmentation Patterns of Some Common Functional Groups


Each functional group exhibits characteristic fragmentation patterns that help to analyze a mass spectrum. For example, aldehydes and ketones often undergo the process of α cleavage, breaking the bond between the carbonyl carbon and the carbon adjacent to it. Cleavage yields a neutral radical and a resonance-stabilized acylium ion.



Alcohols undergo fragmentation in two different ways— α cleavage and dehydration. Alpha (α) cleavage occurs by breaking a bond between an alkyl group and the carbon that bears the OH group, resulting in an alkyl radical and a resonance-stabilized carbocation.



Likewise, alcohols undergo dehydration, the elimination of H_2O , from two adjacent atoms. Unlike fragmentations discussed thus far, dehydration results in the cleavage of two bonds and forms H_2O and the radical cation derived from an alkene.




- Loss of H_2O from an alcohol always forms a fragment with a mass 18 units less than the molecular ion.

Sample Problem 13.5 What mass spectral fragments are formed from α cleavage of 2-pentanone, $\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_3$?

Solution




Alpha (α) cleavage breaks the bond between the carbonyl carbon and the carbon adjacent to it, yielding a neutral radical and a resonance-stabilized acylium ion. A ketone like 2-pentanone with two different alkyl groups bonded to the carbonyl carbon has two different pathways for α cleavage.



As a result, two fragments are formed by α cleavage of 2-pentanone, giving peaks at $m/z = 71$ and 43.

Problem 13.6 (a) What mass spectral fragments are formed by α cleavage of 2-butanol, $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$?
(b) What fragments are formed by dehydration of 2-butanol?

Problem 13.7 What cations are formed in the mass spectrometer by α cleavage of each of the following compounds?



13.4 Other Types of Mass Spectrometry

Recent advances have greatly expanded the information obtained from mass spectrometry.

13.4A High-Resolution Mass Spectrometry

The mass spectra described thus far have been low-resolution spectra; that is, they report m/z values to the nearest whole number. As a result, the mass of a given molecular ion can correspond to many different molecular formulas, as shown in Sample Problem 13.2.

High-resolution mass spectrometers measure m/z ratios to four (or more) decimal places. This is valuable because except for carbon-12, whose mass is defined as 12.0000, the masses of all other nuclei are very close to—but not exactly—whole numbers. Table 13.1 lists the exact mass values of a few common nuclei. Using these values it is possible to determine the single molecular formula that gives rise to a molecular ion.

Table 13.1 Exact Masses of Some Common Isotopes

Isotope	Mass
^{12}C	12.0000
^1H	1.00783
^{16}O	15.9949
^{14}N	14.0031

For example, a compound having a molecular ion at $m/z = 60$ using a low-resolution mass spectrometer could have the following molecular formulas:

Formula	Exact mass
$\text{C}_3\text{H}_8\text{O}$	60.0575
$\text{C}_2\text{H}_4\text{O}_2$	60.0211
$\text{C}_2\text{H}_8\text{N}_2$	60.0688

If the molecular ion had an exact mass of 60.0578, the compound's molecular formula is $\text{C}_3\text{H}_8\text{O}$, because its mass is closest to the observed value.


Problem 13.8 The low-resolution mass spectrum of an unknown analgesic **X** had a molecular ion of 151. Possible molecular formulas include $\text{C}_7\text{H}_5\text{NO}_3$, $\text{C}_8\text{H}_9\text{NO}_2$, and $\text{C}_{10}\text{H}_{17}\text{N}$. High-resolution mass spectrometry gave an exact mass of 151.0640. What is the molecular formula of **X**?

13.4B Gas Chromatography–Mass Spectrometry (GC–MS)

Two analytical tools—**gas chromatography (GC)** and **mass spectrometry (MS)**—can be combined into a single instrument (**GC–MS**) to analyze mixtures of compounds (Figure 13.6a). The gas chromatograph separates the mixture, and then the mass spectrometer records a spectrum of the individual components.

A gas chromatograph consists of a thin capillary column containing a viscous, high-boiling liquid, all housed in an oven. When a sample is injected into the GC, it is vaporized and swept by an inert gas through the column. The components of the mixture travel through the column at different rates, often separated by boiling point, with lower boiling compounds exiting the column before higher boiling compounds. Each compound then enters the mass spectrometer, where it is ionized to form its molecular ion and lower molecular weight fragments. The GC–MS records a gas chromatogram for the mixture, which plots the amount of each component versus its **retention time**—that is, the time required to travel through the column. Each component of a mixture is characterized by its retention time in the gas chromatogram and its molecular ion in the mass spectrum (Figure 13.6b).

GC–MS is widely used for characterizing mixtures containing environmental pollutants. It is also used to analyze urine and hair samples for the presence of illegal drugs or banned substances thought to improve athletic performance.



b. GC trace of a three-component mixture. The mass spectrometer gives a spectrum for each component.





Figure 13.7

Mass spectrum of tetrahydrocannabinol (THC)



Like other controlled substances, the tetrahydrocannabinol from marijuana leaves can be detected in minute amounts by GC-MS.


To analyze a urine sample for THC (tetrahydrocannabinol), the principal psychoactive component of marijuana, the organic compounds are extracted from urine, purified, concentrated, and injected into the GC-MS. THC appears as a GC peak with a characteristic retention time (for a given set of experimental parameters), and gives a molecular ion at 314, its molecular weight, as shown in Figure 13.7.

Problem 13.9


Benzene, toluene, and *p*-xylene (BTX) are often added to gasoline to boost octane ratings. What would be observed if a mixture of these three compounds were subjected to GC-MS analysis? How many peaks would be present in the gas chromatogram? What would be the relative order of the peaks? What molecular ions would be observed in the mass spectra?



benzene



toluene

*p*-xylene**13.4C Mass Spectra of High Molecular Weight Biomolecules**

Dr. John Fenn, professor emeritus of chemical engineering of Yale University, shared the 2002 Nobel Prize in Chemistry for his development of ESI mass spectrometry.

Until the 1980s mass spectra were limited to molecules that could be readily vaporized with heat under vacuum, and thus had molecular weights of < 800 . In the last 25 years, new methods have been developed to generate gas phase ions of large molecules, allowing mass spectra to be recorded for large biomolecules such as proteins and carbohydrates. **Electrospray ionization (ESI)**, for example, forms ions by creating a fine spray of charged droplets in an electric field. Evaporation of the charged droplets forms gaseous ions that are then analyzed by their m/z ratio. ESI and related techniques have extended mass spectrometry into the analysis of nonvolatile compounds with molecular weights greater than 100,000 daltons (atomic mass units).

13.5 Electromagnetic Radiation

Infrared (IR) spectroscopy and **nuclear magnetic resonance (NMR)** spectroscopy (Chapter 14) both use a form of electromagnetic radiation as their energy source. To understand IR and NMR, therefore, you need to understand some of the properties of **electromagnetic radiation**—radiant energy having dual properties of both waves and particles.

The particles of electromagnetic radiation are called **photons**, each having a discrete amount of energy called a **quantum**. Because electromagnetic radiation also has wave properties, it can be characterized by its **wavelength** and **frequency**.

Length units used to report wavelength include:

Unit	Length
meter (m)	1 m
centimeter (cm)	10^{-2} m
micrometer (μm)	10^{-6} m
nanometer (nm)	10^{-9} m
Angstrom (\AA)	10^{-10} m

- **Wavelength (λ)** is the distance from one point on a wave (e.g., the peak or trough) to the same point on the adjacent wave. A variety of different length units are used for λ , depending on the type of radiation.

- Frequency (ν) is the number of waves passing a point per unit time. Frequency is reported in cycles per second (s^{-1}), which is also called hertz (Hz).

You come into contact with many different kinds of electromagnetic radiation in your daily life. For example, you use visible light to see the words on this page, you may cook with microwaves, and you should use sunscreen to protect your skin from the harmful effects of ultraviolet radiation.

The different forms of electromagnetic radiation make up the **electromagnetic spectrum**. The spectrum is arbitrarily divided into different regions, as shown in Figure 13.8. All electromagnetic radiation travels at the speed of light (c), 3.0×10^8 m/s.

The speed of electromagnetic radiation (c) is directly proportional to its wavelength and frequency:

$$c = \lambda\nu$$

The speed of light (c) is a constant, so wavelength and frequency are *inversely* related:

- $\lambda = c/\nu$: Wavelength increases as frequency decreases.
- $\nu = c/\lambda$: Frequency increases as wavelength decreases.

The energy (E) of a photon is directly proportional to its frequency:

$$E = h\nu \quad h = \text{Planck's constant } (6.63 \times 10^{-34} \text{ J}\cdot\text{s})$$


Frequency and wavelength are *inversely* proportional ($\nu = c/\lambda$), however, so energy and wavelength are *inversely* proportional:

$$E = h\nu = \frac{hc}{\lambda}$$

- E increases as ν increases.
- E decreases as λ increases.


When electromagnetic radiation strikes a molecule, some wavelengths—but not all—are absorbed. Only some wavelengths are absorbed because molecules have discrete energy levels. The energies of their electronic, vibrational, and nuclear spin states are *quantized*, not *continuous*.

Figure 13.8
The electromagnetic spectrum



- Visible light occupies only a small region of the electromagnetic spectrum.

- For absorption to occur, the energy of the photon must match the difference between two energy states in a molecule.



- The *larger* the energy difference between two states, the *higher* the energy of radiation needed for absorption, the *higher* the frequency, and the *shorter* the wavelength.

Problem 13.10 Which of the following has the higher frequency: (a) light having a wavelength of 10^2 or 10^4 nm; (b) light having a wavelength of 100 nm or 100 μm ; (c) red light or blue light?

Problem 13.11 Which of the following has the higher energy: (a) light having a ν of 10^4 Hz or 10^8 Hz; (b) light having a λ of 10 nm or 1000 nm; (c) red light or blue light?

Problem 13.12 The difference in energy between two electronic states is ~ 400 kJ/mol, whereas the difference in the energy between two vibrational states is ~ 20 kJ/mol. Which transition requires the higher ν of radiation?

13.6 Infrared Spectroscopy

Organic chemists use infrared (IR) spectroscopy to identify the functional groups in a compound.

13.6A Background

Using the wavenumber scale results in IR frequencies in a numerical range that is easier to report than the corresponding frequencies given in hertz ($4000\text{--}400\text{ cm}^{-1}$ compared to $1.2 \times 10^{14}\text{--}1.2 \times 10^{15}$ Hz).


Infrared radiation ($\lambda = 2.5\text{--}25\ \mu\text{m}$) is the energy source in infrared spectroscopy. These are somewhat longer wavelengths than visible light, so they are lower in frequency and lower in energy than visible light. Frequencies in IR spectroscopy are reported using a unit called the **wavenumber** ($\tilde{\nu}$):

$$\tilde{\nu} = \frac{1}{\lambda}$$


Wavenumber is *inversely* proportional to wavelength and reported in reciprocal centimeters (cm^{-1}). Wavenumber ($\tilde{\nu}$) is *proportional* to frequency (ν). **Frequency (and therefore energy) increases as the wavenumber increases.** Using the wavenumber scale, IR absorptions occur from $4000\text{ cm}^{-1}\text{--}400\text{ cm}^{-1}$.

- Absorption of IR light causes changes in the vibrational motions of a molecule.

Covalent bonds are not static. They are more like springs with weights on each end. When two atoms are bonded to each other, the bond stretches back and forth. When three or more atoms are joined together, bonds can also bend. These bond stretching and bending vibrations represent the different vibrational modes available to a molecule.



These vibrations are quantized, so they occur only at specific frequencies, which correspond to the frequency of IR light. When the frequency of IR light matches the frequency of a particular vibrational mode, the IR light is absorbed, causing the amplitude of the particular bond stretch or bond bend to increase.




- Different kinds of bonds vibrate at different frequencies, so they absorb different frequencies of IR light.
- IR spectroscopy distinguishes between the different kinds of bonds in a molecule, so it is possible to determine the functional groups present.

Problem 13.13 Which of the following has higher energy: (a) IR light of 3000 cm^{-1} or 1500 cm^{-1} in wavenumber; (b) IR light having a wavelength of $10\text{ }\mu\text{m}$ or $20\text{ }\mu\text{m}$?

13.6B Characteristics of an IR Spectrum

In an IR spectrometer, light passes through a sample. Frequencies that match vibrational frequencies are absorbed, and the remaining light is transmitted to a detector. A spectrum plots the amount of transmitted light versus its wavenumber. The IR spectrum of 1-propanol, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$, illustrates several important features of IR spectroscopy.



- An IR spectrum has broad lines.
- The absorption peaks go *down* on a page. The y axis measures **percent transmittance**: 100% transmittance means that all the light shone on a sample is transmitted and none is absorbed; 0% transmittance means that none of the light shone on a sample is transmitted and all is absorbed. Most absorptions lie between these two extremes.
- **Each peak corresponds to a particular kind of bond, and each bond type (such as O-H and C-H) occurs at a characteristic frequency.**




Figure 13.9

Comparing the functional group region and fingerprint region of two compounds

- **A** and **B** show peaks in the same regions for their C=O group and sp^3 hybridized C–H bonds.
- **A** and **B** are different compounds, so their fingerprint regions are quite different.

- IR spectra have both a wavelength and a wavenumber scale on the x axis. Wavelengths are recorded in μm (2.5–25). Wavenumber, frequency, and energy *decrease* from left to right. Where a peak occurs is reported in reciprocal centimeters (cm^{-1}).

Conceptually, the IR spectrum is divided into two regions:

- The **functional group region** occurs at $\geq 1500 \text{ cm}^{-1}$. Common functional groups give one or two peaks in this region, at a characteristic frequency.
- The **fingerprint region** occurs at $< 1500 \text{ cm}^{-1}$. This region often contains a complex set of peaks and is unique for every compound.

Compare, for example, the IR spectra of 5-methyl-2-hexanone (**A**) and ethyl propanoate (**B**) in Figure 13.9. The IR spectra look similar in their functional group regions because both compounds contain a carbonyl group (C=O) and several sp^3 hybridized C–H bonds. Since **A** and **B** are different compounds, however, their fingerprint regions look very different.

13.7 IR Absorptions

13.7A Where Particular Bonds Absorb in the IR

Where a particular bond absorbs in the IR depends on **bond strength** and **atom mass**.

- **Bond strength:** stronger bonds vibrate at higher frequency, so they absorb at higher $\tilde{\nu}$.
- **Atom mass:** bonds with lighter atoms vibrate at higher frequency, so they absorb at higher $\tilde{\nu}$.

Thinking of bonds as springs with weights on each end illustrates these trends. The strength of the spring is analogous to bond strength, and the mass of the weights is analogous to atomic mass. For two springs with the same weights on each end, the **stronger spring vibrates at a higher frequency**. For two springs of the same strength, **springs with lighter weights vibrate at higher frequency** than those with heavier weights. Hooke's law, as shown in Figure 13.10, describes the relationship of frequency to mass and bond strength.

Figure 13.10
Hooke's law: How the frequency of bond vibration depends on atom mass and bond strength

The frequency of bond vibration can be derived from Hooke's law, which describes the motion of a vibrating spring:

Hooke's law

$$\tilde{\nu} = k \sqrt{\frac{f}{m}}$$

f = force constant
 m = mass
 k = constant

stronger bond \dashrightarrow higher frequency

smaller mass \dashrightarrow higher frequency

- The force constant (f) is the strength of the bond (or spring). The larger the value of f , the stronger the bond, and the higher the $\tilde{\nu}$ of vibration.
- The mass (m) is the mass of atoms (or weights). The smaller the value of m , the higher the $\tilde{\nu}$ of vibration.

As a result, **bonds absorb in four predictable regions in an IR spectrum**. These four regions, and the bonds that absorb there, are summarized in Figure 13.11. Remembering the information in this figure will help you analyze the spectra of unknown compounds. To help you remember it, keep in mind the following two points:

- Absorptions for bonds to hydrogen always occur on the left side of the spectrum (the high wavenumber region). H has so little mass that H–Z bonds (where Z = C, O, and N) vibrate at high frequencies.
- Bond strength decreases in going from $\text{C}\equiv\text{C} \rightarrow \text{C}=\text{C} \rightarrow \text{C}-\text{C}$, so the frequency of vibration decreases—that is, the absorptions for these bonds move farther to the right side of the spectrum.

The functional group region consists of absorptions for single bonds to hydrogen (all H–Z bonds), as well as absorptions for all multiple bonds. Most absorptions in the functional group region are due to bond stretching (rather than bond bending). The fingerprint region consists of absorptions due to all other single bonds (except H–Z bonds), often making it a complex region that is very difficult to analyze.

Besides learning the general regions of the IR spectrum, it is also important to learn the specific absorption values for common bonds. Table 13.2 lists the most important IR absorptions in the functional group region. Other details of IR absorptions will be presented in later chapters when new functional groups are introduced. Appendix E contains a detailed list of the characteristic IR absorption frequencies for common bonds.

Figure 13.11
Summary: The four regions of the IR spectrum






Table 13.2 Important IR Absorptions


Bond type	Approximate $\tilde{\nu}$ (cm^{-1})	Intensity
O-H	3600–3200	strong, broad
N-H	3500–3200	medium
C-H	~3000	
• $\text{C}_{sp^3}\text{-H}$	3000–2850	strong
• $\text{C}_{sp^2}\text{-H}$	3150–3000	medium
• $\text{C}_{sp}\text{-H}$	3300	medium
$\text{C}\equiv\text{C}$	2250	medium
$\text{C}\equiv\text{N}$	2250	medium
$\text{C}=\text{O}$	1800–1650 (often ~1700)	strong
$\text{C}=\text{C}$	1650	medium
	1600, 1500	medium

Even subtle differences that affect bond strength affect the frequency of an IR absorption. Recall from Section 1.10 that the strength of a C–H bond increases as the percent *s*-character of the hybrid orbital on the carbon increases; thus:




- The higher the percent *s*-character, the stronger the bond and the higher the wavenumber of absorption.

Problem 13.14 Which bond in each pair absorbs at higher wavenumber?



Finally, almost all bonds in a molecule give rise to an absorption peak in an IR spectrum, but a few do not. **For a bond to absorb in the IR, there must be a change in dipole moment during the vibration.** Thus, symmetrical, nonpolar bonds do not absorb in the IR. The carbon–carbon triple bond of 2-butyne, for example, does not have an IR stretching absorption at 2250 cm^{-1} because the $\text{C}\equiv\text{C}$ bond is nonpolar and there is no change in dipole moment when the bond stretches along its axis. This type of vibration is said to be **IR inactive**.


Stretching along the bond axis
does not change the dipole moment.




13.7B IR Absorptions in Hydrocarbons

The IR spectra of hexane, 1-hexene, and 1-hexyne illustrate the important differences that characterize the IR spectra of hydrocarbons above 1500 cm^{-1} . Although all three compounds contain C–C bonds and sp^3 hybridized C–H bonds, the absorption peaks due to C=C and C≡C readily distinguish the alkene and alkyne.


Note, too, that the C–H absorptions in alkanes, alkenes, and alkynes have a characteristic appearance and position. The sp^3 hybridized C–H bonds are often seen as a broad, strong absorption at $< 3000\text{ cm}^{-1}$, whereas sp^2 and sp hybridized C–H bonds absorb at somewhat higher frequency.



- The **alkane** $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ has only C–C single bonds and sp^3 hybridized C atoms. Therefore, it has only one major absorption above 1500 cm^{-1} , its C_{sp^3} –H absorption at $3000\text{--}2850\text{ cm}^{-1}$.



- The **alkene** $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ has a C=C and C_{sp^2} –H, in addition to its sp^3 hybridized C atoms. Therefore, there are three major absorptions above 1500 cm^{-1} :
 - C_{sp^2} –H at $3150\text{--}3000\text{ cm}^{-1}$
 - C_{sp^3} –H at $3000\text{--}2850\text{ cm}^{-1}$
 - C=C at 1650 cm^{-1}




- The **alkyne** $\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ has a C≡C and C_{sp} –H, in addition to its sp^3 hybridized C atoms. Therefore, there are three major absorptions:
 - C_{sp} –H at 3300 cm^{-1}
 - C_{sp^3} –H at $3000\text{--}2850\text{ cm}^{-1}$
 - C≡C at $\sim 2250\text{ cm}^{-1}$


Problem 13.15 How do the IR spectra of the isomers cyclopentane and 1-pentene differ?

13.7C IR Absorptions in Oxygen-Containing Compounds


The most important IR absorptions for oxygen-containing compounds occur at **3600–3200 cm^{-1} for an OH group**, and at approximately **1700 cm^{-1} for a C=O**, as illustrated in the IR spectra of an **alcohol** (2-butanol), a **ketone** (2-butanone), and an **ether** (diethyl ether). The peak at $\sim 3000 \text{ cm}^{-1}$ in each spectrum is due to $\text{C}_{\text{sp}^3}\text{-H}$ bonds.



- The **OH** group in the alcohol $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$ shows a strong absorption at 3600–3200 cm^{-1} .




- The **C=O** group in the ketone $\text{CH}_3\text{COCH}_2\text{CH}_3$ shows a strong absorption at $\sim 1700 \text{ cm}^{-1}$.
- The exact location of the **C=O** absorption depends on the particular type of carbonyl group, whether the carbonyl carbon is part of a ring, and whether there are nearby double bonds. These details are discussed in Chapters 21 and 22.




- $(\text{CH}_3\text{CH}_2)_2\text{O}$ has neither an OH group nor a C=O, so its only absorption above 1500 cm^{-1} occurs at $\sim 3000 \text{ cm}^{-1}$, due to sp^3 hybridized C–H bonds. Compounds that contain an oxygen atom but do not show an OH or C=O absorption are **ethers**.

13.7D IR Absorptions in Nitrogen-Containing Compounds


Common functional groups that contain nitrogen atoms are also distinguishable by their IR absorptions above 1500 cm^{-1} , as illustrated by the IR spectra of an **amine** (octylamine), an **amide** (propanamide), and a **nitrile** (octanenitrile). Additional details on the IR spectra of these compounds are given in Chapters 22 and 25.



- The **N-H** bonds in the amine $\text{CH}_3(\text{CH}_2)_7\text{NH}_2$ give rise to two weak absorptions at 3300 and 3400 cm^{-1} .



- The **amide** $\text{CH}_3\text{CH}_2\text{CONH}_2$ exhibits absorptions above 1500 cm^{-1} for both its **N-H** and **C=O** groups:
 - N-H** (two peaks) at 3200 and 3400 cm^{-1}
 - C=O** at 1660 cm^{-1}




- The **C≡N** group of the **nitrile** $\text{CH}_3(\text{CH}_2)_6\text{CN}$ absorbs in the triple bond region at $\sim 2250\text{ cm}^{-1}$.

Sample Problem 13.6 How can the two isomers having molecular formula C_2H_6O be distinguished by IR spectroscopy?


Solution

First, draw the structures of the compounds and then locate the functional groups. One compound is an alcohol and one is an ether.





Although both compounds have sp^3 hybridized C-H bonds, ethanol has an OH group that gives a strong absorption at $3600-3200\text{ cm}^{-1}$, and dimethyl ether does not. This feature distinguishes the two isomers.

Problem 13.16 How do the three isomers of molecular formula C_3H_6O (**A**, **B**, and **C**) differ in their IR spectra?



Sample Problem 13.7 shows how the region above 1500 cm^{-1} in an IR spectrum can be used for functional group identification.


Sample Problem 13.7 What functional groups are responsible for the absorptions above 1500 cm^{-1} in compounds **A** and **B**?




Solution

- a. Compound **A** has two major absorptions above 1500 cm^{-1} : The absorption at $\sim 3000\text{ cm}^{-1}$ is due to C-H bonds and the absorption at $\sim 1700\text{ cm}^{-1}$ is due to a C=O group.
- b. Compound **B** has two major absorptions above 1500 cm^{-1} : The absorption at $\sim 3000\text{ cm}^{-1}$ is due to C-H bonds and the absorption at $\sim 2250\text{ cm}^{-1}$ is due to a triple bond, either a $C\equiv C$ or a $C\equiv N$. Because there is no absorption due to an sp hybridized C-H bond at 3300 cm^{-1} , this IR spectrum *cannot* be due to a terminal alkyne ($HC\equiv CR$) but may still be due to an internal alkyne.

Problem 13.17 What functional groups are responsible for the absorptions above 1500 cm^{-1} in the IR spectra for compounds **A** and **B**?



Problem 13.18 What are the major IR absorptions in the functional group region for each compound?




13.8 IR and Structure Determination


New instruments for determining blood alcohol concentration use IR spectroscopy for analyzing the C–H absorption of $\text{CH}_3\text{CH}_2\text{OH}$ in exhaled air. Figure 12.10 illustrated an earlier method based on oxidation chemistry.

Since its introduction, IR spectroscopy has proven to be a valuable tool for determining the functional groups in organic molecules.

In the 1940s, IR spectroscopy played a key role in elucidating the structure of the antibiotic penicillin G, the chapter-opening molecule. **β -Lactams**, four-membered rings that contain an amide, have a carbonyl group that absorbs at $\sim 1760\text{ cm}^{-1}$, a much higher frequency than that observed for most amides and many other carbonyl groups. Because penicillin G had an IR absorption at this frequency, **A** became the leading candidate for the structure of penicillin rather than **B**, a possibility originally considered more likely. Structure **A** was later confirmed by X-ray analysis.





IR spectroscopy is often used to determine the outcome of a chemical reaction. For example, oxidation of the hydroxy group in **C** to form the carbonyl group in periplanone **B** is accompanied by the disappearance of the OH absorption ($3600\text{--}3200\text{ cm}^{-1}$) and the appearance of a carbonyl absorption near 1700 cm^{-1} in the IR spectrum of the product. Periplanone **B** is the sex pheromone of the female American cockroach.



The combination of IR and mass spectral data provides key information on the structure of an unknown compound. The mass spectrum reveals the molecular weight of the unknown (and the molecular formula if an exact mass is available), and the IR spectrum helps to identify the important functional groups.

HOW TO Use MS and IR for Structure Determination


Example What information is obtained from the mass spectrum and IR spectrum of an unknown compound **X**? Assume **X** contains the elements C, H, and O.



Step [1] Use the molecular ion to determine possible molecular formulas. Use an exact mass (when available) to determine a molecular formula.

- Use the procedure outlined in Sample Problem 13.2 to calculate possible molecular formulas. For a molecular ion at $m/z = 88$:

HOW TO, continued . . .



- Discounting C_7H_4 (a hydrocarbon) and C_6O (because it contains no H's) gives three possible formulas for **X**.
- If high-resolution mass spectral data are available, the molecular formula can be determined directly. If the molecular ion had an exact mass of 88.0580, the molecular formula of **X** is $\text{C}_4\text{H}_8\text{O}_2$ (exact mass = 88.0524) rather than $\text{C}_5\text{H}_{12}\text{O}$ (exact mass = 88.0888) or $\text{C}_3\text{H}_4\text{O}_3$ (exact mass = 88.0160).

Step [2] Calculate the number of degrees of unsaturation (Section 10.2).

- For a compound of molecular formula $\text{C}_4\text{H}_8\text{O}_2$, the maximum number of H's = $2n + 2 = 2(4) + 2 = 10$.
- Because the compound contains only 8 H's, it has $10 - 8 = 2$ H's fewer than the maximum number.
- Because each degree of unsaturation removes 2 H's, **X** has one degree of unsaturation. **X** has one ring or one π bond.

Step [3] Determine what functional group is present from the IR spectrum.

- The two major absorptions in the IR spectrum above 1500 cm^{-1} are due to sp^3 hybridized C-H bonds ($\sim 3000\text{--}2850 \text{ cm}^{-1}$) and a C=O group (1740 cm^{-1}). Thus, the one degree of unsaturation in **X** is due to the presence of the **C=O**.

Mass spectrometry and IR spectroscopy give valuable but limited information on the identity of an unknown. Although the mass spectral and IR data reveal that **X** has a molecular formula of $\text{C}_4\text{H}_8\text{O}_2$ and contains a carbonyl group, more data are needed to determine its complete structure. In Chapter 14, we will learn how other spectroscopic data can be used for that purpose.

Problem 13.19 Which of the following possible structures for **X** can be excluded on the basis of its IR spectrum: (a) $\text{CH}_3\text{COOCH}_2\text{CH}_3$; (b) $\text{HOCH}_2\text{CH}_2\text{CH}_2\text{CHO}$; (c) $\text{CH}_3\text{CH}_2\text{COOCH}_3$; (d) $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$?

Problem 13.20 Propose structures consistent with each set of data: (a) a hydrocarbon with a molecular ion at $m/z = 68$ and IR absorptions at 3310 , $3000\text{--}2850$, and 2120 cm^{-1} ; (b) a compound containing C, H, and O with a molecular ion at $m/z = 60$ and IR absorptions at $3600\text{--}3200$ and $3000\text{--}2850 \text{ cm}^{-1}$.

KEY CONCEPTS

Mass Spectrometry and Infrared Spectroscopy

Mass Spectrometry (MS; 13.1–13.4)


- Mass spectrometry measures the molecular weight of a compound (13.1A).
- The mass of the molecular ion (**M**) = the molecular weight of a compound. Except for isotope peaks at $M + 1$ and $M + 2$, the molecular ion has the highest mass in a mass spectrum (13.1A).
- The base peak is the tallest peak in a mass spectrum (13.1A).
- A compound with an odd number of N atoms gives an odd molecular ion. A compound with an even number of N atoms (including zero) gives an even molecular ion (13.1B).
- Organic monochlorides show two peaks for the molecular ion (M and $M + 2$) in a 3:1 ratio (13.2).
- Organic monobromides show two peaks for the molecular ion (M and $M + 2$) in a 1:1 ratio (13.2).
- The fragmentation of radical cations formed in a mass spectrometer gives lower molecular weight fragments, often characteristic of a functional group (13.3).
- High-resolution mass spectrometry gives the molecular formula of a compound (13.4A).

Electromagnetic Radiation (13.5)


- The wavelength and frequency of electromagnetic radiation are *inversely* related by the following equations: $\lambda = c/\nu$ or $\nu = c/\lambda$ (13.5).
- The energy of a photon is proportional to its frequency; the higher the frequency, the higher the energy: $E = h\nu$ (13.5).

Infrared Spectroscopy (IR; 13.6 and 13.7)

- Infrared spectroscopy identifies functional groups.
- IR absorptions are reported in wavenumbers, $\tilde{\nu} = 1/\lambda$.
- The functional group region from **4000–1500 cm^{-1}** is the most useful region of an IR spectrum.
- C–H, O–H, and N–H bonds absorb at high frequency, $\geq 2500 \text{ cm}^{-1}$.
- As bond strength increases, the $\tilde{\nu}$ of absorption increases; thus, triple bonds absorb at higher $\tilde{\nu}$ than double bonds.








- The higher the percent *s*-character, the stronger the bond, and the higher the $\tilde{\nu}$ of an IR absorption.



PROBLEMS

Mass Spectrometry

13.21 What molecular ion is expected for each compound?




13.22 Which compound gives a molecular ion at $m/z = 122$: $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{C}_6\text{H}_5\text{COCH}_2\text{CH}_3$, or $\text{C}_6\text{H}_5\text{OCH}_2\text{CH}_3$?

13.23 Propose two molecular formulas for each molecular ion: (a) 102; (b) 98; (c) 119; (d) 74.


13.24 Propose four possible structures for a hydrocarbon with a molecular ion at $m/z = 112$.

13.25 Match each structure to its mass spectrum.



13.26 Propose two possible structures for a hydrocarbon having an exact mass of 96.0939 that forms ethylcyclopentane upon hydrogenation with H_2 and Pd-C.

13.27 What cations are formed in the mass spectrometer by α cleavage of each of the following compounds?



13.28 For each compound, assign likely structures to the fragments at each m/z value, and explain how each fragment is formed.

- $C_6H_5CH_2CH_2OH$: peaks at $m/z = 104, 91$
- $CH_2=C(CH_3)CH_2CH_2OH$: peaks at $m/z = 71, 68, 41, 31$

13.29 The mass spectrum of 3-ethyl-3-methylheptane $[(CH_3CH_2)_2C(CH_3)CH_2CH_2CH_2CH_3]$ shows fragments at $m/z = 127, 113,$ and 85 . Propose structures for the ions that give rise to these peaks.

13.30 Suppose you have two bottles, labeled ketone **A** and ketone **B**. You know that one bottle contains $CH_3CO(CH_2)_5CH_3$ and one contains $CH_3CH_2CO(CH_2)_4CH_3$, but you do not know which ketone is in which bottle. Ketone **A** gives a fragment at $m/z = 99$ and ketone **B** gives a fragment at $m/z = 113$. What are the likely structures of ketones **A** and **B** from these fragmentation data?

13.31 Propose a structure consistent with each set of data.

- A compound that contains a benzene ring and has a molecular ion at $m/z = 107$
- A hydrocarbon that contains only sp^3 hybridized carbons and a molecular ion at $m/z = 84$
- A compound that contains a carbonyl group and gives a molecular ion at $m/z = 114$
- A compound that contains C, H, N, and O and has an exact mass for the molecular ion at 101.0841


13.32 A low-resolution mass spectrum of the neurotransmitter dopamine gave a molecular ion at $m/z = 153$. Two possible molecular formulas for this molecular ion are $C_8H_{11}NO_2$ and $C_7H_{11}N_3O$. A high-resolution mass spectrum provided an exact mass at 153.0680. Which of the possible molecular formulas is the correct one?

13.33 Explain why compounds containing an odd number of nitrogen atoms have an odd molecular ion in their mass spectra.


- 13.34** Can the exact mass obtained in a high-resolution mass spectrum distinguish between two isomers such as $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ and $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{CH}_3$?
- 13.35** Primary (1°) alcohols often show a peak in their mass spectra at $m/z = 31$. Suggest a structure for this fragment.
- 13.36** Like alcohols, ethers undergo α cleavage by breaking a carbon-carbon bond between an alkyl group and the carbon bonded to the ether oxygen atom; that is, the red C-C bond in $\text{R}-\text{CH}_2\text{OR}'$ is broken. With this in mind, propose structures for the fragments formed by α cleavage of $(\text{CH}_3)_2\text{CHCH}_2\text{OCH}_2\text{CH}_3$. Suggest a reason why an ether fragments by α cleavage.

Infrared Spectroscopy


- 13.37** Which of the indicated bonds absorbs at higher $\tilde{\nu}$ in an IR spectrum?




- 13.38** What major IR absorptions are present above 1500 cm^{-1} for each compound?




- 13.39** How would each of the following pairs of compounds differ in their IR spectra?















- 13.40** Morphine, heroin, and oxycodone are three addicting analgesic narcotics. How could IR spectroscopy be used to distinguish these three compounds from each other?



- 13.41** Tell how IR spectroscopy could be used to determine when each reaction is complete.




13.42 Match each compound to its IR spectrum.



- 13.43 Draw the structures of the seven compounds of molecular formula C_3H_6O . For each compound tell what prominent IR absorptions it exhibits above 1500 cm^{-1} .


Combined Spectroscopy Problems

- 13.44 Propose possible structures consistent with each set of data. Assume each compound has an sp^3 hybridized C–H absorption in its IR spectrum, and that other major IR absorptions above 1500 cm^{-1} are listed.
- A compound having a molecular ion at 72 and an absorption in its IR spectrum at 1725 cm^{-1}
 - A compound having a molecular ion at 55 and an absorption in its IR spectrum at $\sim 2250\text{ cm}^{-1}$
 - A compound having a molecular ion of 74 and an absorption in its IR spectrum at $3600\text{--}3200\text{ cm}^{-1}$
- 13.45 A chiral hydrocarbon **X** exhibits a molecular ion at 82 in its mass spectrum. The IR spectrum of **X** shows peaks at 3300, 3000–2850, and 2250 cm^{-1} . Propose a structure for **X**.
- 13.46 A chiral compound **Y** has a strong absorption at $2970\text{--}2840\text{ cm}^{-1}$ in its IR spectrum and gives the following mass spectrum. Propose a structure for **Y**.




- 13.47 Treatment of benzaldehyde (C_6H_5CHO) with $Zn(Hg)$ in aqueous HCl forms a compound **Z** that has a molecular ion at 92 in its mass spectrum. **Z** shows absorptions at $3150\text{--}2950$, 1605, and 1496 cm^{-1} in its IR spectrum. Give a possible structure for **Z**.
- 13.48 Reaction of *tert*-butyl pentyl ether [$CH_3CH_2CH_2CH_2CH_2OC(CH_3)_3$] with HBr forms 1-bromopentane ($CH_3CH_2CH_2CH_2CH_2Br$) and compound **H**. **H** has a molecular ion in its mass spectrum at 56 and gives peaks in its IR spectrum at $3150\text{--}3000$, $3000\text{--}2850$, and 1650 cm^{-1} . Propose a structure for **H** and draw a stepwise mechanism that accounts for its formation.
- 13.49 Reaction of pentanoyl chloride ($CH_3CH_2CH_2CH_2COCl$) with lithium dimethyl cuprate [$LiCu(CH_3)_2$] forms a compound **J** that has a molecular ion in its mass spectrum at 100, as well as fragments at $m/z = 85$, 57, and 43 (base). The IR spectrum of **J** has strong peaks at 2962 and 1718 cm^{-1} . Propose a structure for **J**.
- 13.50 Benzonitrile (C_6H_5CN) is reduced to two different products depending on the reducing agent used. Treatment with lithium aluminum hydride followed by water forms **K**, which has a molecular ion in its mass spectrum at 107 and the following IR absorptions: 3373, 3290, 3062, 2920, and 1600 cm^{-1} . Treatment with a milder reducing agent forms **L**, which has a molecular ion in its mass spectrum at 106 and the following IR absorptions: 3086, 2850, 2820, 2736, 1703, and 1600 cm^{-1} . **L** shows fragments in its mass spectrum at $m/z = 105$ and 77. Propose structures for **K** and **L** and explain how you arrived at your conclusions.
- 13.51 Treatment of anisole ($CH_3OC_6H_5$) with Cl_2 and $FeCl_3$ forms **P**, which has peaks in its mass spectrum at $m/z = 142$ (M), 144 (M + 2), 129, and 127. **P** has absorptions in its IR spectrum at $3096\text{--}2837$ (several peaks), 1582, and 1494 cm^{-1} . Propose possible structures for **P**.

- 13.52 Reaction of $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ with NaH forms compound **W**, which gives the IR and mass spectra shown below. Propose a structure for **W** and draw a stepwise mechanism that accounts for its formation.




Challenge Problems

- 13.53 Explain why a carbonyl absorption shifts to lower frequency in an α,β -unsaturated carbonyl compound—a compound having a carbonyl group bonded directly to a carbon-carbon double bond. For example, the carbonyl absorption occurs at 1720 cm^{-1} for cyclohexanone, and at 1685 cm^{-1} for 2-cyclohexenone.




cyclohexanone



2-cyclohexenone
(an α,β -unsaturated carbonyl compound)

- 13.54 Oxidation of citronellol, a constituent of rose and geranium oils, with PCC in the presence of added NaOCOCH_3 forms compound **A**. **A** has a molecular ion in its mass spectrum at 154 and a strong peak in its IR spectrum at 1730 cm^{-1} , in addition to C-H stretching absorptions. Without added NaOCOCH_3 , oxidation of citronellol with PCC yields isopulegone, which is then converted to **B** with aqueous base. **B** has a molecular ion at 152, and a peak in its IR spectrum at 1680 cm^{-1} in addition to C-H stretching absorptions.



- Identify the structures of **A** and **B**.
- Draw a mechanism for the conversion of citronellol to isopulegone.
- Draw a mechanism for the conversion of isopulegone to **B**.

14

Nuclear Magnetic Resonance Spectroscopy

- 14.1 An introduction to NMR spectroscopy
- 14.2 ^1H NMR: Number of signals
- 14.3 ^1H NMR: Position of signals
- 14.4 The chemical shift of protons on sp^2 and sp hybridized carbons
- 14.5 ^1H NMR: Intensity of signals
- 14.6 ^1H NMR: Spin-spin splitting
- 14.7 More complex examples of splitting
- 14.8 Spin-spin splitting in alkenes
- 14.9 Other facts about ^1H NMR spectroscopy
- 14.10 Using ^1H NMR to identify an unknown
- 14.11 ^{13}C NMR spectroscopy
- 14.12 Magnetic resonance imaging (MRI)



Melatonin, a hormone synthesized by the pineal gland, is thought to induce sleep. Because melatonin synthesis is inhibited by light, melatonin levels in the body rise as less light falls upon the eye, and drop quickly at dawn. For this reason, melatonin has become a popular dietary supplement for travelers suffering from jetlag and individuals with mild sleep disorders. Modern spectroscopic techniques have been used to characterize the structure of melatonin. In Chapter 14, we learn how nuclear magnetic resonance spectroscopy plays a key role in organic structure determination.

In Chapter 14 we continue our study of organic structure determination by learning about **nuclear magnetic resonance (NMR)** spectroscopy. NMR spectroscopy is the most powerful tool for characterizing organic molecules, because it can be used to identify the carbon–hydrogen framework in a compound.

14.1 An Introduction to NMR Spectroscopy

Two common types of NMR spectroscopy are used to characterize organic structure:


- ^1H NMR (**proton NMR**) is used to determine the number and type of hydrogen atoms in a molecule; and
- ^{13}C NMR (**carbon NMR**) is used to determine the type of carbon atoms in a molecule.

Before you can learn how to use NMR spectroscopy to determine the structure of a compound, you need to understand a bit about the physics behind it. Keep in mind, though, that NMR stems from the same basic principle as all other forms of spectroscopy. Energy interacts with a molecule, and absorptions occur only when the incident energy matches the energy difference between two states.

14.1A The Basis of NMR Spectroscopy

The source of energy in NMR is radio waves. Radiation in the radiofrequency region of the electromagnetic spectrum (so-called **RF** radiation) has very long wavelengths, so its corresponding frequency and energy are both low. **When these low-energy radio waves interact with a molecule, they can change the nuclear spins of some elements, including ^1H and ^{13}C .**


When a charged particle such as a proton spins on its axis, it creates a magnetic field. For the purpose of this discussion, therefore, a nucleus is a tiny bar magnet, symbolized by \uparrow . Normally these nuclear magnets are randomly oriented in space, but in the presence of an external magnetic field, B_0 , they are oriented with or against this applied field. More nuclei are oriented *with* the applied field because this arrangement is lower in energy, but the **energy difference between these two states is very small** (< 0.4 J/mol).



In a magnetic field, there are now two different energy states for a proton:

- A **lower energy state with the nucleus aligned in the same direction as B_0**
- A **higher energy state with the nucleus aligned opposed to B_0**

When an external energy source ($h\nu$) that matches the energy difference (ΔE) between these two states is applied, energy is absorbed, causing the **nucleus to “spin flip” from one orientation to another**. The energy difference between these two nuclear spin states corresponds to the low-frequency radiation in the RF region of the electromagnetic spectrum.



- A nucleus is in *resonance* when it absorbs RF radiation and “spin flips” to a higher energy state.

Thus, two variables characterize NMR:

- **An applied magnetic field, B_0 .** Magnetic field strength is measured in tesla (T).
- **The frequency ν of radiation used for resonance,** measured in hertz (Hz) or megahertz (MHz); (1 MHz = 10^6 Hz)

The frequency needed for resonance and the applied magnetic field strength are proportionally related:

$$\nu \propto B_0$$


- **The stronger the magnetic field, the larger the energy difference between the two nuclear spin states, and the higher the ν needed for resonance.**

NMR spectrometers are referred to as 300 MHz instruments, 500 MHz instruments, and so forth, depending on the frequency of RF radiation used for resonance.

Early NMR spectrometers used a magnetic field strength of ~ 1.4 T, which required RF radiation of 60 MHz for resonance. Modern NMR spectrometers use stronger magnets, thus requiring higher frequencies of RF radiation for resonance. For example, a magnetic field strength of 7.05 T requires a frequency of 300 MHz for a proton to be in resonance. These spectrometers use very powerful magnetic fields to create a small, but measurable energy difference between the two possible spin states. A schematic of an NMR spectrometer is shown in Figure 14.1.

If all protons absorbed at the same frequency in a given magnetic field, the spectra of all compounds would consist of a single absorption, rendering NMR useless for structure determination. Fortunately, however, this is not the case.

Figure 14.1 Schematic of an NMR spectrometer



An NMR spectrometer. The sample is dissolved in a solvent, usually CDCl_3 (deuteriochloroform), and placed in a magnetic field. A radiofrequency generator then irradiates the sample with a short pulse of radiation, causing resonance. When the nuclei fall back to their lower energy state, the detector measures the energy released, and a spectrum is recorded. The superconducting magnets in modern NMR spectrometers have coils that are cooled in liquid helium and conduct electricity with essentially no resistance.


- All protons do *not* absorb at the same frequency. Protons in different environments absorb at slightly different frequencies, and so they are distinguishable by NMR.

The frequency at which a particular proton absorbs is determined by its electronic environment, as discussed in Section 14.3. Because electrons are moving charged particles, they create a magnetic field opposed to the applied field B_0 , and the size of the magnetic field generated by the electrons around a proton determines where it absorbs. Modern NMR spectrometers use a constant magnetic field strength B_0 , and then a narrow range of frequencies is applied to achieve the resonance of all protons.

Only nuclei that contain odd mass numbers (such as ^1H , ^{13}C , ^{19}F , and ^{31}P) or odd atomic numbers (such as ^2H and ^{14}N) give rise to NMR signals. Because both ^1H and ^{13}C , the less abundant isotope of carbon, are NMR active, NMR allows us to map the carbon and hydrogen framework of an organic molecule.

14.1B A ^1H NMR Spectrum

An NMR spectrum plots the **intensity of a signal** against its **chemical shift** measured in **parts per million (ppm)**. The common scale of chemical shifts is called the δ (delta) scale. The proton NMR spectrum of *tert*-butyl methyl ether [$\text{CH}_3\text{OC}(\text{CH}_3)_3$] illustrates several important features:



tert-Butyl methyl ether (MTBE) is the high-octane gasoline additive that has contaminated the water supply in some areas (Section 3.4).

$(\text{CH}_3)_4\text{Si}$
tetramethylsilane
TMS

- NMR absorptions generally appear as sharp signals. The ^1H NMR spectrum of $\text{CH}_3\text{OC}(\text{CH}_3)_3$ consists of two signals: a tall peak at 1.2 ppm due to the $(\text{CH}_3)_3\text{C}-$ group, and a smaller peak at 3.2 ppm due to the $\text{CH}_3\text{O}-$ group.
- **Increasing chemical shift is plotted from right to left.** Most protons absorb somewhere from 0–12 ppm.
- The terms **upfield** and **downfield** describe the relative location of signals. Upfield means to the *right*. The $(\text{CH}_3)_3\text{C}-$ peak is upfield from the $\text{CH}_3\text{O}-$ peak. Downfield means to the *left*. The $\text{CH}_3\text{O}-$ peak is downfield from the $(\text{CH}_3)_3\text{C}-$ peak.

NMR absorptions are measured relative to the position of a reference signal at 0 ppm on the δ scale due to **tetramethylsilane (TMS)**. TMS is a volatile and inert compound that gives a single peak upfield from other typical NMR absorptions.

Although chemical shifts are measured relative to the TMS signal at 0 ppm, this reference is often not plotted on a spectrum.

The *positive* direction of the δ scale is *downfield* from TMS. A very small number of absorptions occur upfield from the TMS signal, which is defined as the negative direction of the δ scale. (See Problem 14.72.)

The **chemical shift** on the x axis gives the position of an NMR signal, measured in ppm, according to the following equation:

$$\text{chemical shift (in ppm on the } \delta \text{ scale)} = \frac{\text{observed chemical shift (in Hz) downfield from TMS}}{\nu \text{ of the NMR spectrometer (in MHz)}}$$

A chemical shift gives absorptions as a fraction of the NMR operating frequency, making it independent of the spectrometer used to record a spectrum. Because the frequency of the radiation required for resonance is proportional to the strength of the applied magnetic field, B_0 , reporting NMR absorptions in frequency is meaningless unless the value of B_0 is also reported. By reporting the absorption as a fraction of the NMR operating frequency, though, we get units—ppm—that are independent of the spectrometer.

Sample Problem 14.1

Calculate the chemical shift of an absorption that occurs at 1500 Hz downfield from TMS using a 300 MHz NMR spectrometer.

Solution

Use the equation that defines the chemical shift in ppm:

$$\text{chemical shift} = \frac{1500 \text{ Hz downfield from TMS}}{300 \text{ MHz operating frequency}} = 5 \text{ ppm}$$

Problem 14.1

The ^1H NMR spectrum of CH_3OH recorded on a 500 MHz NMR spectrometer consists of two signals, one due to the CH_3 protons at 1715 Hz and one due to the OH proton at 1830 Hz, both measured downfield from TMS. (a) Calculate the chemical shift of each absorption. (b) Do the CH_3 protons absorb upfield or downfield from the OH proton?

Problem 14.2

The ^1H NMR spectrum of 1,2-dimethoxyethane ($\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_3$) recorded on a 300 MHz NMR spectrometer consists of signals at 1017 Hz and 1065 Hz downfield from TMS. (a) Calculate the chemical shift of each absorption. (b) At what frequency would each absorption occur if the spectrum were recorded on a 500 MHz NMR spectrometer?

Four different features of a ^1H NMR spectrum provide information about a compound's structure:

- [1] **Number of signals** (Section 14.2)
- [2] **Position of signals** (Sections 14.3 and 14.4)
- [3] **Intensity of signals** (Section 14.5)
- [4] **Spin-spin splitting of signals** (Sections 14.6–14.8)

14.2 ^1H NMR: Number of Signals


How many ^1H NMR signals does a compound exhibit? The number of NMR signals *equals* the number of different types of protons in a compound.

14.2A General Principles


- Protons in different environments give different NMR signals. Equivalent protons give the same NMR signal.

Any CH_3 group is different from any CH_2 group, which is different from any CH group in a molecule. Two CH_3 groups may be identical (as in CH_3OCH_3) or different (as in $\text{CH}_3\text{OCH}_2\text{CH}_3$), depending on what each CH_3 group is bonded to.


In many compounds, deciding whether two protons are in identical or different environments is intuitive.



All equivalent H's
1 NMR signal



2 types of H's
2 NMR signals



3 types of H's
3 NMR signals

tert-Butyl methyl ether $[\text{CH}_3\text{OC}(\text{CH}_3)_3]$ (Section 14.1) exhibits two NMR signals because it contains two different kinds of protons: one CH_3 group is bonded to $-\text{OC}(\text{CH}_3)_3$, whereas the other three CH_3 groups are each bonded to the same group, $[-\text{C}(\text{CH}_3)_2]\text{OCH}_3$.

- **CH_3OCH_3** : Each CH_3 group is bonded to the same group ($-\text{OCH}_3$), making both CH_3 groups equivalent.
- **$\text{CH}_3\text{CH}_2\text{Cl}$** : The protons of the CH_3 group are different from those of the CH_2 group.
- **$\text{CH}_3\text{OCH}_2\text{CH}_3$** : The protons of the CH_2 group are different from those in each CH_3 group. The two CH_3 groups are also different from each other; one CH_3 group is bonded to $-\text{OCH}_2\text{CH}_3$ and the other is bonded to $-\text{CH}_2\text{OCH}_3$.


In some cases, it is less obvious by inspection if two protons are equivalent or different. To rigorously determine whether two protons are in identical environments (and therefore give rise to one NMR signal), replace each H atom in question by another atom Z (for example, $Z = \text{Cl}$). If substitution by Z yields the same compound or enantiomers, the two protons are equivalent, as shown in Sample Problem 14.2.

Sample Problem 14.2

How many different kinds of H atoms does $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ contain?

Solution

In comparing two H atoms, replace each H by Z (for example, $Z = \text{Cl}$), and examine the substitution products that result. The two CH_3 groups are identical because substitution of one H by Cl gives $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$ (1-chloropentane). There are two different types of CH_2 groups, because substitution of one H by Cl gives two different products:



Thus, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ has **three** different types of protons and gives **three** NMR signals.

Figure 14.2 gives the number of NMR signals exhibited by four additional molecules. All protons—not just protons bonded to carbon atoms—give rise to NMR signals. Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), for example, gives three NMR signals, one of which is due to its OH proton.

Problem 14.3

How many ^1H NMR signals does each compound show?

- | | | | |
|--|---|--|--|
| a. CH_3CH_3 | c. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ | e. $\text{CH}_3\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$ | g. $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ |
| b. $\text{CH}_3\text{CH}_2\text{CH}_3$ | d. $(\text{CH}_3)_2\text{CHCH}(\text{CH}_3)_2$ | f. $\text{CH}_3\text{OCH}_2\text{CH}(\text{CH}_3)_2$ | h. $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ |

Problem 14.4

How many different types of protons does $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$ contain?


14.2B Determining Equivalent Protons in Alkenes and Cycloalkanes

To determine equivalent protons in cycloalkanes and alkenes that have restricted bond rotation, always draw in all bonds to hydrogen.




Figure 14.2

The number of ^1H NMR signals of some representative organic compounds




Then, in comparing two H atoms on a ring or double bond, **two protons are equivalent only if they are cis (or trans) to the same groups**, as illustrated with 1,1-dichloroethylene, 1-bromo-1-chloroethylene, and chloroethylene.




- **1,1-Dichloroethylene:** The two H atoms on the C=C are both cis to a Cl atom. Thus, both H atoms are equivalent.
- **1-Bromo-1-chloroethylene:** H_a is cis to a Cl atom and H_b is cis to a Br atom. Thus, H_a and H_b are different, giving rise to two NMR signals.
- **Chloroethylene:** H_a is bonded to the carbon with the Cl atom, making it different from H_b and H_c. Of the remaining two H atoms, H_b is cis to a Cl atom and H_c is cis to a H atom, making them different. All three H atoms in this compound are different.

Proton equivalency in cycloalkanes can be determined similarly.



- **Cyclopropane:** All H atoms are equivalent, so there is only one NMR signal.
- **Chlorocyclopropane:** There are now three kinds of H atoms: H_a is bonded to a carbon bonded to a Cl; both H_b protons are cis to the Cl whereas both H_c protons are cis to another H.


Problem 14.5 How many ¹H NMR signals does each dimethylcyclopropane show?



14.2C Enantiotopic and Diastereotopic Protons


Let's look more closely at the protons of a single sp^3 hybridized CH₂ group to determine whether these two protons are always equivalent to *each other*. Two examples illustrate different outcomes.

CH₃CH₂Br has two different types of protons—those of the CH₃ group and those of the CH₂ group—meaning that the two H atoms of the CH₂ group are *equivalent to each other*. To confirm this fact, we replace each H of the CH₂ group by an atom Z and examine the products of substitution. In this case, substitution of each H by Z creates a new stereogenic center, forming two products that are **enantiomers**.



- When substitution of two H atoms by Z forms enantiomers, the two H atoms are equivalent and give a single NMR signal. These two H atoms are called *enantiotopic* protons.


In contrast, the two H atoms of the CH_2 group in (2*R*)-2-chlorobutane, which contains one stereogenic center, are *not* equivalent to each other. Substitution of each H by Z forms two **diastereomers**, and thus, these two H atoms give *different* NMR signals.



- When substitution of two H atoms by Z forms diastereomers, the two H atoms are not equivalent, and give two NMR signals. These two H atoms are called *diastereotopic* protons.

Sample Problem 14.3


Label the protons in each indicated CH_2 group as enantiotopic, diastereotopic, or neither.




Solution

To determine equivalency in these cases, look for whether the compound has a stereogenic center to begin with and whether a new stereogenic is formed when H is replaced by Z.


- a. The compound is achiral and has no stereogenic center. Since no new stereogenic center is formed on substitution of H by Z, the protons are **neither** enantiotopic nor diastereotopic. The H's within the CH_2 group are *equivalent* to each other and give *one* NMR signal.




- b. The compound is achiral and has no stereogenic center. Since a new stereogenic center is formed on substitution of H by Z, the protons are **enantiotopic**. The H's within the CH_2 group are *equivalent* to each other and give *one* NMR signal.



- c. The compound has one stereogenic center to begin with. Since a new stereogenic center is formed on substitution of H by Z, the protons are **diastereotopic**. The H's within the CH_2 group are *different* from each other and give *different* NMR signals.



Problem 14.6 Label the protons in each indicated CH₂ group as enantiotopic, diastereotopic, or neither.



Problem 14.7 How many ¹H NMR signals would you expect for each compound: (a) CH₃CH(Cl)CH₂CH₃; (b) ClCH₂CH(CH₃)OCH₃; (c) CH₃CH(Br)CH₂CH₂CH₃?


14.3 ¹H NMR: Position of Signals

In the NMR spectrum of *tert*-butyl methyl ether in Section 14.1B, why does the CH₃O– group absorb downfield from the –C(CH₃)₃ group?

- Where a particular proton absorbs depends on its electronic environment.

14.3A Shielding and Deshielding Effects

To understand how the electronic environment around a nucleus affects its chemical shift, recall that in a magnetic field, an electron creates a small magnetic field that opposes the applied magnetic field, B_0 . **Electrons are said to shield the nucleus from B_0 .**




In the vicinity of the nucleus, therefore, the magnetic field generated by the circulating electron *decreases* the external magnetic field that the proton “feels.” Because the proton experiences a lower magnetic field strength, it needs a lower frequency to achieve resonance. Lower frequency is to the right in an NMR spectrum, toward lower chemical shift, so **shielding shifts an absorption upfield**, as shown in Figure 14.3a.

Figure 14.3 How chemical shift is affected by electron density around a nucleus


a. Shielding effects

- An electron shields the nucleus.
- The absorption shifts *upfield*.



b. Deshielding effects


- Decreased electron density deshields a nucleus.
- The absorption shifts *downfield*.



← Increasing chemical shift
Increasing ν

← Increasing chemical shift
Increasing ν

Figure 14.4
Shielding and deshielding effects



- As the electron density around the nucleus increases, the nucleus feels a smaller resultant magnetic field, so a lower frequency is needed to achieve resonance.
 - **The absorption shifts upfield.**
- As the electron density around the nucleus decreases, the nucleus feels a larger resultant magnetic field, so a higher frequency is needed to achieve resonance.
 - **The absorption shifts downfield.**

What happens if the electron density around a nucleus is *decreased*, instead? For example, how do the chemical shifts of the protons in CH_4 and CH_3Cl compare?


The less shielded the nucleus becomes, the more of the applied magnetic field (B_0) it feels. This *deshielded* nucleus experiences a higher magnetic field strength, so it needs a higher frequency to achieve resonance. Higher frequency is to the *left* in an NMR spectrum, toward higher chemical shift, so **deshielding shifts an absorption downfield**, as shown in Figure 14.3b for CH_3Cl versus CH_4 . The electronegative Cl atom withdraws electron density from the carbon and hydrogen atoms in CH_3Cl , thus deshielding them relative to those in CH_4 .

Remember the trend:
Decreased electron density deshields a nucleus and an absorption moves downfield.


- **Protons near electronegative atoms are deshielded, so they absorb downfield.**

Figure 14.4 summarizes the effects of shielding and deshielding.


These electron density arguments explain the relative position of NMR signals in many compounds.



- The H_b protons are **deshielded** because they are closer to the electronegative Cl atom, so they absorb **downfield** from H_a .



- Because F is more electronegative than Br, the H_b protons are more **deshielded** than the H_a protons and absorb farther **downfield**.



- The larger number of electronegative Cl atoms (two versus one) **deshields** H_b more than H_a , so it absorbs **downfield** from H_a .

Sample Problem 14.4

Which of the underlined protons in each pair absorbs farther downfield: (a) $\text{CH}_3\text{CH}_2\text{CH}_3$ or CH_3OCH_3 ; (b) CH_3OCH_3 or CH_3SCH_3 ?

Solution

- The CH_3 group in CH_3OCH_3 is deshielded by the electronegative O atom. **Deshielding shifts the absorption downfield.**
- Because oxygen is more electronegative than sulfur, the CH_3 group in CH_3OCH_3 is more **deshielded** and absorbs **downfield**.








Problem 14.8

For each compound, which of the underlined protons absorbs farther downfield: (a) $\text{FCH}_2\text{CH}_2\text{CH}_2\text{Cl}$; (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$; (c) $\text{CH}_3\text{OC}(\text{CH}_3)_3$?

14.3B Chemical Shift Values

Not only is the *relative* position of NMR absorptions predictable, but it is also possible to predict the approximate chemical shift value for a given type of proton.

Table 14.1 Characteristic Chemical Shifts of Common Types of Protons

Type of proton	Chemical shift (ppm)	Type of proton	Chemical shift (ppm)
 sp^3 <ul style="list-style-type: none"> • RCH_3 ~0.9 • R_2CH_2 ~1.3 • R_3CH ~1.7 	0.9–2	 sp^2	4.5–6
 $Z = C, O, N$	1.5–2.5		6.5–8
$-C\equiv C-H$	~2.5		9–10
 $Z = N, O, X$	2.5–4		10–12
		$RO-H$ or $R-N-H$	1–5


- Protons in a given environment absorb in a predictable region in an NMR spectrum.

A more detailed list of characteristic chemical shift values is found in Appendix F.

Table 14.1 lists the typical chemical shift values for the most common bonds encountered in organic molecules.

Table 14.1 illustrates that absorptions for a given type of C–H bond occur in a narrow range of chemical shift values, usually 1–2 ppm. For example, all sp^3 hybridized C–H bonds in alkanes and cycloalkanes absorb between 0.9 and 2.0 ppm. By contrast, absorptions due to N–H and O–H protons can occur over a broader range. For example, the OH proton of an alcohol is found anywhere in the 1–5 ppm range. The position of these absorptions is affected by the extent of hydrogen bonding, making it more variable.

The chemical shift of a particular type of C–H bond is also affected by the number of R groups bonded to the carbon atom.





Increasing alkyl substitution
Increasing chemical shift

- The chemical shift of a C–H bond increases with increasing alkyl substitution.


Problem 14.9

For each compound, first label each different type of proton and then rank the protons in order of increasing chemical shift.



14.4 The Chemical Shift of Protons on sp^2 and sp Hybridized Carbons


The chemical shift of protons bonded to benzene rings, C–C double bonds, and C–C triple bonds merits additional comment.



Each of these functional groups contains π bonds with **loosely held π electrons**. When placed in a magnetic field, these π electrons move in a circular path, inducing a new magnetic field. How this induced magnetic field affects the chemical shift of a proton depends on the direction of the induced field *in the vicinity of the absorbing proton*.


Protons on Benzene Rings

In a magnetic field, the six π electrons in **benzene** circulate around the ring, creating a ring current. The magnetic field induced by these moving electrons reinforces the applied magnetic field in the vicinity of the protons. The protons thus feel a stronger magnetic field and a higher frequency is needed for resonance, so the **protons are deshielded and the absorption is downfield**.



Protons on Carbon–Carbon Double Bonds

A similar phenomenon occurs with protons on carbon–carbon double bonds. In a magnetic field, the loosely held π electrons create a magnetic field that reinforces the applied field in the vicinity of the protons. Because the protons now feel a stronger magnetic field, they require a higher frequency for resonance. **The protons are deshielded and the absorption is downfield**.



Protons on Carbon–Carbon Triple Bonds

In a magnetic field, the π electrons of a carbon–carbon triple bond are induced to circulate, but in this case the induced magnetic field *opposes* the applied magnetic field (B_0). The proton thus feels a weaker magnetic field, so a lower frequency is needed for resonance. **The nucleus is shielded and the absorption is upfield**.





Table 14.2 summarizes the shielding and deshielding effects due to circulating π electrons.

To remember the chemical shifts of some common bond types, it is helpful to think of a ^1H NMR spectrum as being divided into six different regions (Figure 14.5).

Table 14.2 Effect of π Electrons on Chemical Shift Values


Proton type	Effect	Chemical shift (ppm)
	highly deshielded	6.5–8
	deshielded	4.5–6
$-\text{C}\equiv\text{C}-\text{H}$	shielded	~2.5

Figure 14.5
Regions in the ^1H NMR spectrum



- Shielded protons absorb at lower chemical shift (to the right).
- Deshielded protons absorb at higher chemical shift (to the left).
- Note: The drawn chemical shift scale is not linear.


Sample Problem 14.5 Rank H_a , H_b , and H_c in order of increasing chemical shift.



Solution

The H_a protons are bonded to an sp^3 hybridized carbon, so they are shielded and absorb upfield compared to H_b and H_c . Because the H_b protons are deshielded by the electronegative oxygen atom on the C to which they are bonded, they absorb downfield from H_a . The H_c proton is deshielded by two factors. The electronegative O atom withdraws electron density from H_c . Moreover, because H_c is bonded directly to a $\text{C}=\text{C}$, the magnetic field induced by the π electrons causes further deshielding. Thus, in order of increasing chemical shift, $\text{H}_a < \text{H}_b < \text{H}_c$.

Problem 14.10 Rank each group of protons in order of increasing chemical shift.




14.5 ^1H NMR: Intensity of Signals

The relative intensity of ^1H NMR signals also provides information about a compound's structure.

- The area under an NMR signal is proportional to the number of absorbing protons.

For example, in the ^1H NMR spectrum of $\text{CH}_3\text{OC}(\text{CH}_3)_3$, the ratio of the area under the downfield peak (due to the $\text{CH}_3\text{O}-$ group) to the upfield peak [due to the $-\text{C}(\text{CH}_3)_3$ group] is 1:3. An NMR spectrometer automatically integrates the area under the peaks, and prints out a stepped curve (an **integral**) on the spectrum. **The height of each step is proportional to the area under the peak, which is in turn proportional to the number of absorbing protons.**



Integrals can be manually measured, but modern NMR spectrometers automatically calculate and plot the value of each integral in arbitrary units. If the heights of two integrals are 20 units and 60 units, the ratio of absorbing protons is 20:60, or 1:3, or 2:6, or 3:9, and so forth. This tells the *ratio*, not the absolute number of protons. Integration ratios are approximate, and often values must be rounded to the nearest whole number.


Problem 14.11 Which compounds give a ^1H NMR spectrum with two signals in a ratio of 2:3?

- a. $\text{CH}_3\text{CH}_2\text{Cl}$ b. $\text{CH}_3\text{CH}_2\text{CH}_3$ c. $\text{CH}_3\text{CH}_2\text{OCH}_2\text{CH}_3$ d. $\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_3$

Knowing the molecular formula of a compound and integration values from its ^1H NMR spectrum gives the *actual number* of protons responsible for a particular signal.

HOW TO Determine the Number of Protons Giving Rise to an NMR Signal

Example A compound of molecular formula $\text{C}_9\text{H}_{10}\text{O}_2$ gives the following integrated ^1H NMR spectrum. How many protons give rise to each signal?



Step [1] Determine the number of integration units per proton by dividing the total number of integration units by the total number of protons.

- Total number of integration units: $54 + 23 + 33 = 110$ units
- Total number of protons = 10
- Divide: $110 \text{ units}/10 \text{ protons} = \mathbf{11 \text{ units per proton}}$

Step [2] Determine the number of protons giving rise to each signal.

- To determine the number of H atoms giving rise to each signal, divide each integration value by the answer of Step [1] and round to the nearest whole number.

$$\text{Answer: } \begin{array}{l} \text{Signal [A]:} \\ \frac{54}{11} = 4.9 \approx \mathbf{5 \text{ H}} \end{array} \quad \begin{array}{l} \text{Signal [B]:} \\ \frac{23}{11} = 2.1 \approx \mathbf{2 \text{ H}} \end{array} \quad \begin{array}{l} \text{Signal [C]:} \\ \frac{33}{11} = \mathbf{3 \text{ H}} \end{array}$$


Problem 14.12 A compound of molecular formula $\text{C}_8\text{H}_{14}\text{O}_2$ gives three NMR signals having the indicated integration values: signal [A] 14 units, signal [B] 12 units, and signal [C] 44 units. How many protons give rise to each signal?

Problem 14.13 Compound **A** exhibits two signals in its ^1H NMR spectrum at 2.64 and 3.69 ppm and the ratio of the absorbing signals is 2:3. Compound **B** exhibits two signals in its ^1H NMR spectrum at 2.09 and 4.27 ppm and the ratio of the absorbing signals is 3:2. Which compound corresponds to $\text{CH}_3\text{O}_2\text{CCH}_2\text{CH}_2\text{CO}_2\text{CH}_3$ (dimethyl succinate) and which compound corresponds to $\text{CH}_3\text{CO}_2\text{CH}_2\text{CH}_2\text{O}_2\text{CCH}_3$ (ethylene diacetate)?

14.6 ^1H NMR: Spin–Spin Splitting

The ^1H NMR spectra you have seen up to this point have been limited to one or more single absorptions called **singlets**. In the ^1H NMR spectrum of $\text{BrCH}_2\text{CHBr}_2$, however, the two signals for the two different kinds of protons are each split into more than one peak. The splitting pat-

terns, the result of **spin-spin splitting**, can be used to determine how many protons reside on the carbon atoms near the absorbing proton.




To understand spin-spin splitting, we must distinguish between the **absorbing protons** that give rise to an NMR signal, and the **adjacent protons** that cause the signal to split. **The number of adjacent protons determines the observed splitting pattern.**

- The CH_2 signal appears as **two peaks**, called a **doublet**. The relative area under the peaks of a doublet is 1:1.
- The CH signal appears as **three peaks**, called a **triplet**. The relative area under the peaks of a triplet is 1:2:1.


Spin-spin splitting occurs only between nonequivalent protons on the same carbon or adjacent carbons. To illustrate how spin-spin splitting arises, we'll examine nonequivalent protons on adjacent carbons, the more common example. Spin-spin splitting arises because protons are little magnets that can be aligned with or against an applied magnetic field, and this affects the magnetic field that a nearby proton feels.

14.6A Splitting: How a Doublet Arises

First, let's examine how the doublet due to the CH_2 group in $\text{BrCH}_2\text{CHBr}_2$ arises. The CH_2 group contains the absorbing protons and the CH group contains the adjacent proton that causes the splitting.



When placed in an applied magnetic field (B_0), the adjacent proton (CHBr_2) can be aligned with (\uparrow) or against (\downarrow) B_0 . As a result, the absorbing protons (CH_2Br) feel two slightly different magnetic fields—one slightly larger than B_0 and one slightly smaller than B_0 . Because the absorbing protons feel two different magnetic fields, they absorb at two different frequencies in the NMR spectrum, thus splitting a single absorption into a doublet.



Keep in mind the difference between an **NMR signal** and an **NMR peak**. An NMR signal is the entire absorption due to a particular kind of proton. NMR peaks are contained within a signal. A doublet constitutes one signal that is split into two peaks.

Table 14.3 Names for a Given Number of Peaks in an NMR Signal

Number of peaks	Name	Number of peaks	Name
1	singlet	5	quintet
2	doublet	6	sextet
3	triplet	7	septet
4	quartet	> 7	multiplet


Rule [1] Equivalent protons don't split each other's signals.

Rule [2] A set of n nonequivalent protons splits the signal of a nearby proton into $n + 1$ peaks.

- In $\text{BrCH}_2\text{CHBr}_2$, for example, *one* adjacent CH proton splits an NMR signal into *two* peaks (a doublet), and *two* adjacent CH_2 protons split an NMR signal into *three* peaks (a triplet). Names for split NMR signals containing two to seven peaks are given in Table 14.3. An NMR signal having more than seven peaks is called a **multiplet**.
- The inside peaks of a split NMR signal are always most intense, with the area under the peaks decreasing from the inner to the outer peaks in a given splitting pattern.

Rule [3] Splitting is observed for nonequivalent protons on the same carbon or adjacent carbons.

If H_a and H_b are not equivalent, splitting is observed when:




H_a and H_b are on the **same** carbon.

H_a and H_b are on **adjacent** carbons.

The splitting of an NMR signal reveals the number of nearby nonequivalent protons. It tells nothing about the absorbing proton itself.


Splitting is not generally observed between protons separated by more than three σ bonds. Although H_a and H_b are not equivalent to each other in 2-butanone and ethyl methyl ether, H_a and H_b are separated by four σ bonds and so they are too far away to split each other's NMR signals.



2-butanone

H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b



ethyl methyl ether

H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b











Table 14.4 illustrates common splitting patterns observed for adjacent nonequivalent protons.

Predicting splitting is always a two-step process:


- **Determine if two protons are equivalent or different.** Only nonequivalent protons split each other.
- **Determine if two nonequivalent protons are close enough to split each other's signals.** Splitting is observed only for nonequivalent protons on the *same* carbon or *adjacent* carbons.

Several examples of spin-spin splitting in specific compounds illustrate the result of this two-step strategy.


Table 14.4 Common Splitting Patterns Observed in ^1H NMR

Example	Pattern	Analysis (H_a and H_b are not equivalent.)
[1] 		<ul style="list-style-type: none"> H_a: one adjacent H_b proton \rightarrow two peaks \rightarrow a doublet H_b: one adjacent H_a proton \rightarrow two peaks \rightarrow a doublet
[2] 		<ul style="list-style-type: none"> H_a: two adjacent H_b protons \rightarrow three peaks \rightarrow a triplet H_b: one adjacent H_a proton \rightarrow two peaks \rightarrow a doublet
[3] 		<ul style="list-style-type: none"> H_a: two adjacent H_b protons \rightarrow three peaks \rightarrow a triplet H_b: two adjacent H_a protons \rightarrow three peaks \rightarrow a triplet
[4] 		<ul style="list-style-type: none"> H_a: three adjacent H_b protons \rightarrow four peaks \rightarrow a quartet* H_b: two adjacent H_a protons \rightarrow three peaks \rightarrow a triplet
[5] 		<ul style="list-style-type: none"> H_a: three adjacent H_b protons \rightarrow four peaks \rightarrow a quartet* H_b: one adjacent H_a proton \rightarrow two peaks \rightarrow a doublet


*The relative area under the peaks of a quartet is 1:3:3:1.




- All protons are equivalent (H_a), so there is no splitting and the NMR signal is one singlet.



- There are two NMR signals. H_a and H_b are nonequivalent protons bonded to adjacent C atoms, so they are close enough to split each other's NMR signals. The H_a signal is split into a triplet by the two H_b protons. The H_b signal is split into a triplet by the two H_a protons.









- There are three NMR signals. H_a has no adjacent nonequivalent protons, so its signal is a singlet. The H_b signal is split into a quartet by the three H_c protons. The H_c signal is split into a triplet by the two H_b protons.




- There are two NMR signals. H_a and H_b are nonequivalent protons on the same carbon, so they are close enough to split each other's NMR signals. The H_a signal is split into a doublet by H_b . The H_b signal is split into a doublet by H_a .

Problem 14.14 Into how many peaks will each indicated proton be split?



Problem 14.15 Although $\text{Cl}_2\text{CHCHCl}_2$ and $\text{Br}_2\text{CHCHCl}_2$ each have only two hydrogens, these compounds have very different ^1H NMR spectra. For each compound, give the number of ^1H NMR signals and indicate into how many peaks each signal is split.

Problem 14.16 For each compound give the number of ^1H NMR signals, and then determine how many peaks are present for each NMR signal.



Problem 14.17 Sketch the NMR spectrum of $\text{CH}_3\text{CH}_2\text{Cl}$, giving the approximate location of each NMR signal.

14.7 More Complex Examples of Splitting

Up to now you have studied examples of spin–spin splitting where the absorbing proton has nearby protons on *one* adjacent carbon only. What happens when the absorbing proton has nonequivalent protons on *two* adjacent carbons? Different outcomes are possible, depending on whether the adjacent nonequivalent protons are *equivalent to* or *different from* each other.

For example, 2-bromopropane $[(\text{CH}_3)_2\text{CHBr}]$ has two types of protons— H_a and H_b —so it exhibits two NMR signals, as shown in Figure 14.6.


- The H_a protons have only one adjacent nonequivalent proton (H_b), so they are split into two peaks, a **doublet**.
- H_b has three H_a protons on each side. Because the six H_a protons are *equivalent to each other*, the $n + 1$ rule can be used to determine splitting: $6 + 1 = 7$ peaks, a **septet**.

This is a specific example of a general rule:


- Whenever two (or three) sets of adjacent protons are equivalent to each other, use the $n + 1$ rule to determine the splitting pattern.

A different outcome results when an absorbing proton is flanked by adjacent protons that are *not equivalent to each other*. Consider the splitting pattern expected for the H_b protons in the

Figure 14.6
The ^1H NMR spectrum of 2-bromopropane, $[(\text{CH}_3)_2\text{CHBr}]$



^1H NMR spectrum of $\text{CH}_3\text{CH}_2\text{CH}_2\text{Z}$. H_b has protons on both adjacent carbons, but since H_a and H_c are *not equivalent to each other*, we cannot merely add them together and use the $n + 1$ rule.




Instead, to determine the splitting of H_b , we must consider the effect of the H_a protons and the H_c protons *separately*. The three H_a protons split the H_b signal into four peaks, and the two H_c protons split each of these four peaks into three peaks—that is, the NMR signal due to H_b consists of $4 \times 3 = 12$ peaks. Figure 14.7 shows a splitting diagram illustrating how these 12 peaks arise.

- When two sets of adjacent protons are *different from each other* (n protons on one adjacent carbon and m protons on the other), the number of peaks in an NMR signal = $(n + 1)(m + 1)$.

It is only possible to see 12 peaks in an NMR spectrum when the coupling constants between each set of nonequivalent protons—that is, J_{ab} and J_{bc} in this example—are different; in other words, $J_{ab} \neq J_{bc}$. Such is the case with the nonequivalent protons on carbon-carbon double bonds, which is discussed in Section 14.8. **In practice, with flexible alkyl chains it is more common for J_{ab} and J_{bc} to be very similar or identical.** In this case, peaks overlap and many fewer than 12 peaks are observed.


The ^1H NMR spectrum of 1-bromopropane ($\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$) illustrates the result of peak overlap (Figure 14.8).



$\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ has three different types of protons— H_a , H_b , and H_c —so it exhibits three NMR signals. H_a and H_c are each triplets because they are adjacent to two H_b protons. H_b has protons on both adjacent carbons, and H_a and H_c are *not equivalent to each other*. The three H_a protons should split the H_b signal into four peaks, and the two H_c protons should split each of these four peaks into three peaks—that is, the NMR signal due to H_b should once again consist of $4 \times 3 = 12$ peaks. However, since $J_{ab} = J_{bc}$ in this case, peak overlap occurs and a multiplet of only six peaks is observed.


Figure 14.7

A splitting diagram for the H_b protons in $\text{CH}_3\text{CH}_2\text{CH}_2\text{Z}$



- The H_b signal is split into 12 peaks, a quartet of triplets. The number of peaks actually seen for the signal depends on the relative size of the coupling constants, J_{ab} and J_{bc} . When $J_{ab} \gg J_{bc}$, as drawn in this diagram, all 12 lines of the pattern are visible. When J_{ab} and J_{bc} are similar in magnitude, peaks overlap and fewer lines are observed.

Figure 14.8
The ^1H NMR spectrum
of 1-bromopropane,
 $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$



- H_a and H_c are both triplets.
- The signal for H_b appears as a multiplet of six peaks (a sextet), due to peak overlap.


In $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$, the n protons on one adjacent carbon and the m protons on the other adjacent carbon split the observed signal into $n + m + 1$ peaks. In other words, the 3 H_a protons and 2 H_c protons split the NMR signal into $3 + 2 + 1 = 6$ peaks, as shown in the sextet in Figure 14.8.

Sample Problem 14.6


How many peaks are present in the NMR signal of each indicated proton?

- a. $\text{ClCH}_2\text{CH}_2\text{CH}_2\text{Cl}$ b. $\text{ClCH}_2\text{CH}_2\text{CH}_2\text{Br}$

Solution



- H_b has two H_a protons on each adjacent C. Because the four H_a protons are equivalent to each other, the $n + 1$ rule can be used to determine splitting: $4 + 1 = 5$ **peaks**, a quintet.



- H_b has two H_a protons on one adjacent C and two H_c protons on the other. Because H_a and H_c are not equivalent to each other, the maximum number of peaks for $\text{H}_b = (n + 1)(m + 1) = (2 + 1)(2 + 1) = 9$ **peaks**. However, since this molecule has a flexible alkyl chain, it is likely that J_{ab} and J_{bc} are very similar, so that peak overlap occurs. In this case, the number of peaks for $\text{H}_b = n + m + 1 = 2 + 2 + 1 = 5$ **peaks**.

Problem 14.18



How many peaks are present in the NMR signal of each indicated proton?

- a. $(\text{CH}_3)_2\text{CHCO}_2\text{CH}_3$ b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ c. $\begin{array}{c} \text{Cl} \quad \text{CH}_2\text{Br} \\ | \quad | \\ \text{C} = \text{C} \\ | \quad | \\ \text{H} \quad \text{H} \end{array}$ d. $\begin{array}{c} \text{H} \quad \text{H} \\ | \quad | \\ \text{C} = \text{C} \\ | \quad | \\ \text{Br} \quad \text{H} \end{array}$ (all H atoms)

Problem 14.19


Describe the ^1H NMR spectrum of each compound. State how many NMR signals are present, the splitting pattern for each signal, and the approximate chemical shift.

- a. $\text{CH}_3\text{OCH}_2\text{CH}_3$ c. $\text{CH}_3\text{OCH}_2\text{CH}_2\text{CH}_2\text{OCH}_3$



14.8 Spin–Spin Splitting in Alkenes

Protons on carbon–carbon double bonds often give characteristic splitting patterns. A disubstituted double bond can have two **geminal protons** (on the same carbon atom), two **cis protons**, or two **trans protons**. When these protons are different, each proton splits the NMR signal of the other, so that each proton appears as a doublet. **The magnitude of the coupling constant J for these doublets depends on the arrangement of hydrogen atoms.**




Thus, the *E* and *Z* isomers of 3-chloropropenoic acid both exhibit two doublets for the two alkenyl protons, but the coupling constant is larger when the protons are trans compared to when the protons are cis, as shown in Figure 14.9.

When a double bond is monosubstituted, there are three nonequivalent protons, and the pattern is more complicated because all three protons are coupled to each other. For example, vinyl acetate ($\text{CH}_2=\text{CHOCOCH}_3$) has four different types of protons, three of which are bonded to the double bond. Besides the singlet for the CH_3 group, each proton on the double bond is coupled to two other different protons on the double bond, giving the spectrum in Figure 14.10.

- H_b has two nearby nonequivalent protons that split its signal, the geminal proton H_c and the trans proton H_d . H_d splits the H_b signal into a doublet, and the H_c proton splits the doublet into two doublets. This pattern of four peaks is called a **doublet of doublets**.
- H_c has two nearby nonequivalent protons that split its signal, the geminal proton H_b and the cis proton H_d . H_d splits the H_c signal into a doublet, and the H_b proton splits the doublet into two doublets, forming another **doublet of doublets**.
- H_d has two nearby nonequivalent protons that split its signal, the trans proton H_b and the cis proton H_c . H_b splits the H_d signal into a doublet, and the H_c proton splits the doublet into two doublets, forming another **doublet of doublets**.

Splitting diagrams for the three alkenyl protons in vinyl acetate are drawn in Figure 14.11. Note that each pattern is different in appearance because the magnitude of the coupling constants forming them is different.

Figure 14.9
 ^1H NMR spectra for the alkenyl protons of (*E*)- and (*Z*)-3-chloropropenoic acid




- Although both (*E*)- and (*Z*)-3-chloropropenoic acid show two doublets in their ^1H NMR spectra for their alkenyl protons, $J_{\text{trans}} > J_{\text{cis}}$.


Figure 14.10

The ^1H NMR spectrum of vinyl acetate ($\text{CH}_2=\text{CHOCOCH}_3$)


Vinyl acetate is polymerized to poly(vinyl acetate) (Problem 15.30), a polymer used in paints, glues, and adhesives.

**Figure 14.11**

Splitting diagram for the alkenyl protons in vinyl acetate ($\text{CH}_2=\text{CHOCOCH}_3$)



Problem 14.20 Draw a splitting diagram for H_b in *trans*-1,3-dichloropropene, given that $J_{ab} = 13.1$ Hz and $J_{bc} = 7.2$ Hz.

*trans*-1,3-dichloropropene


Problem 14.21 Identify **A** and **B**, isomers of molecular formula $\text{C}_3\text{H}_4\text{Cl}_2$, from the given ^1H NMR data: Compound **A** exhibits signals at 1.75 (doublet, 3 H, $J = 6.9$ Hz) and 5.89 (quartet, 1 H, $J = 6.9$ Hz) ppm. Compound **B** exhibits signals at 4.16 (singlet, 2 H), 5.42 (doublet, 1 H, $J = 1.9$ Hz), and 5.59 (doublet, 1 H, $J = 1.9$ Hz) ppm.

14.9 Other Facts About ^1H NMR Spectroscopy

14.9A OH Protons

- Under usual conditions, an OH proton does not split the NMR signal of adjacent protons.
- The signal due to an OH proton is not split by adjacent protons.

Figure 14.12
The ^1H NMR spectrum of ethanol ($\text{CH}_3\text{CH}_2\text{OH}$)



Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$), for example, has three different types of protons, so there are three signals in its ^1H NMR spectrum, as shown in Figure 14.12.

- The H_a signal is split by the two H_b protons into three peaks, a **triplet**.
- The H_b signal is split by only the three H_a protons into four peaks, a **quartet**. The adjacent OH proton does *not* split the signal due to H_b .
- H_c is a **singlet** because OH protons are *not* split by adjacent protons.


Why is a proton bonded to an oxygen atom a singlet in a ^1H NMR spectrum? Protons on electronegative elements rapidly **exchange** between molecules in the presence of trace amounts of acid or base. It is as if the CH_2 group in ethanol never “feels” the presence of the OH proton, because the OH proton is rapidly moving from one molecule to another. We therefore see a peak due to the OH proton, but it is a single peak with no splitting. This phenomenon usually occurs with NH and OH protons.

Problem 14.22 How many signals are present in the ^1H NMR spectrum for each molecule? What splitting is observed in each signal: (a) $(\text{CH}_3)_3\text{CCH}_2\text{OH}$; (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$; (c) $(\text{CH}_3)_2\text{CHNH}_2$?

14.9B Cyclohexane Conformations

How does the rotation around carbon–carbon σ bonds and the ring flip of cyclohexane rings affect an NMR spectrum? Because these processes are rapid at room temperature, an NMR spectrum records an **average** of all conformations that interconvert.

Thus, even though each cyclohexane carbon has two different types of hydrogens—one axial and one equatorial—the two chair forms of cyclohexane rapidly interconvert them, and an **NMR spectrum shows a single signal for the average environment** that it “sees.”



Axial and equatorial H's rapidly interconvert. NMR sees an average environment and shows one signal.


14.9C Protons on Benzene Rings

Benzene has six equivalent, deshielded protons and exhibits a single peak in its ^1H NMR spectrum at 7.27 ppm. Monosubstituted benzene derivatives—that is, benzene rings with one H atom





Figure 14.13

The 6.5–8 ppm region of the ^1H NMR spectrum of three benzene derivatives

A benzene ring with one substituent Z



A monosubstituted benzene ring has three different types of H atoms: H_a, H_b, and H_c.





- The appearance of the signals in the 6.5–8 ppm region of the ^1H NMR spectrum depends on the identity of Z in $\text{C}_6\text{H}_5\text{Z}$.

We will learn more about the spectroscopic absorptions of benzene derivatives in Chapter 17.

replaced by another substituent Z—contain five deshielded protons that are no longer all equivalent to each other. The identity of Z determines the appearance of this region of a ^1H NMR spectrum (6.5–8 ppm), as shown in Figure 14.13. We will not analyze the splitting patterns observed for the ring protons of monosubstituted benzenes.

Problem 14.23


What protons in alcohol **A** give rise to each signal in its ^1H NMR spectrum? Explain all splitting patterns observed for absorptions between 0–7 ppm.

**14.10 Using ^1H NMR to Identify an Unknown**

Once we know a compound's molecular formula from its mass spectral data and the identity of its functional group from its IR spectrum, we can then use its ^1H NMR spectrum to determine its structure. A suggested procedure is illustrated for compound **X**, whose molecular formula ($\text{C}_4\text{H}_8\text{O}_2$) and functional group ($\text{C}=\text{O}$) were determined in Section 13.8.

HOW TO Use ^1H NMR Data to Determine a Structure

Example Using its ^1H NMR spectrum, determine the structure of an unknown compound **X** that has molecular formula $\text{C}_4\text{H}_8\text{O}_2$ and contains a $\text{C}=\text{O}$ absorption in its IR spectrum.




Step [1] Determine the number of different kinds of protons.

- The number of NMR signals equals the number of different types of protons.
- This molecule has three NMR signals ([A], [B], and [C]) and therefore **three** types of protons (H_a , H_b , and H_c).


Step [2] Use the integration data to determine the number of H atoms giving rise to each signal (Section 14.5).

- Total number of integration units: $14 + 11 + 15 = 40$ units
- Total number of protons = 8
- Divide: $40 \text{ units} / 8 \text{ protons} = 5 \text{ units per proton}$
- Then, divide each integration value by this answer (5 units per proton) and round to the nearest whole number.




Step [3] Use individual splitting patterns to determine what carbon atoms are bonded to each other.

- Start with the singlets. Signal [C] is due to a CH_3 group with no adjacent nonequivalent H atoms. Possible structures include:



- Because signal [A] is a **triplet**, there must be **2 H's** (CH_2 group) on the adjacent carbon.
- Because signal [B] is a **quartet**, there must be **3 H's** (CH_3 group) on the adjacent carbon.
- This information suggests that **X** has an **ethyl** group CH_3CH_2- .


HOW TO, continued . . .



To summarize, **X** contains CH_3- , CH_3CH_2- , and $\text{C}=\text{O}$ (from the IR). Comparing these atoms with the molecular formula shows that one O atom is missing. Because O atoms do not absorb in a ^1H NMR spectrum, their presence can only be inferred by examining the chemical shift of protons near them. O atoms are more electronegative than C, thus deshielding nearby protons, and shifting their absorption downfield.

Step [4] Use chemical shift data to complete the structure.

- Put the structure together in a manner that preserves the splitting data and is consistent with the reported chemical shifts.
- In this example, two isomeric structures (**A** and **B**) are possible for **X** considering the splitting data only:




- Chemical shift information distinguishes the two possibilities. The electronegative O atom deshields adjacent H's, shifting them downfield between 3 and 4 ppm. If **A** is the correct structure, the singlet due to the CH_3 group (H_c) should occur downfield, whereas if **B** is the correct structure, the quartet due to the CH_2 group (H_b) should occur downfield.
- Because the NMR of **X** has a singlet (not a quartet) at 3.7, **A is the correct structure**.


Problem 14.24 Propose a structure for a compound of molecular formula $\text{C}_7\text{H}_{14}\text{O}_2$ with an IR absorption at 1740 cm^{-1} and the following ^1H NMR data:

Absorption	ppm	Integration value
singlet	1.2	26
triplet	1.3	10
quartet	4.1	6

Problem 14.25 Propose a structure for a compound of molecular formula C_3H_8O with an IR absorption at $3600\text{--}3200\text{ cm}^{-1}$ and the following NMR spectrum:



Problem 14.26 The 1H NMR spectrum of melatonin, the chapter-opening molecule, is more complex than other examples we have encountered, but the chemical shift and splitting patterns observed for several peaks can be explained by what we have learned about 1H NMR thus far. (a) Which protons in melatonin give rise to signals [A]–[D]? (b) Explain the splitting pattern observed in signal [C].



Problem 14.27 Identify products **A** and **B** from the given 1H NMR data.


- Treatment of $CH_2=CHCOCH_3$ with one equivalent of HCl forms compound **A**. **A** exhibits the following absorptions in its 1H NMR spectrum: 2.2 (singlet, 3 H), 3.05 (triplet, 2 H), and 3.6 (triplet, 2 H) ppm. What is the structure of **A**?
- Treatment of acetone $[(CH_3)_2C=O]$ with dilute aqueous base forms **B**. Compound **B** exhibits four singlets in its 1H NMR spectrum at 1.3 (6 H), 2.2 (3 H), 2.5 (2 H), and 3.8 (1 H) ppm. What is the structure of **B**?

14.11 ^{13}C NMR Spectroscopy

^{13}C NMR spectroscopy is also an important tool for organic structure analysis. The physical basis for ^{13}C NMR is the same as for 1H NMR. When placed in a magnetic field, B_0 , ^{13}C nuclei can align themselves with or against B_0 . More nuclei are aligned with B_0 because this arrange-

ment is lower in energy, but these nuclei can be made to spin flip against the applied field by applying RF radiation of the appropriate frequency.

^{13}C NMR spectra, like ^1H NMR spectra, plot peak intensity versus chemical shift, using TMS as the reference signal at 0 ppm. ^{13}C occurs in only 1.1% natural abundance, however, so ^{13}C NMR signals are much weaker than ^1H NMR signals. To overcome this limitation, modern spectrometers irradiate samples with many pulses of RF radiation and use mathematical tools to increase signal sensitivity and decrease background noise. The spectrum of acetic acid (CH_3COOH) illustrates the general features of a ^{13}C NMR spectrum.



^{13}C NMR spectra are easier to analyze than ^1H spectra because signals are not split. **Each type of carbon atom appears as a single peak.**

Why aren't ^{13}C signals split by nearby carbon atoms? Recall from Section 14.6 that splitting occurs when two NMR active nuclei—like two protons—are close to each other. Because of the low natural abundance of ^{13}C nuclei (1.1%), the chance of two ^{13}C nuclei being bonded to each other is very small (0.01%), and so no carbon–carbon splitting is observed.

A ^{13}C NMR signal can also be split by nearby protons. This ^1H – ^{13}C splitting is usually eliminated from a spectrum, however, by using an instrumental technique that decouples the proton–carbon interactions, so that every peak in a ^{13}C NMR spectrum is a singlet.

Two features of ^{13}C NMR spectra provide the most structural information: the **number of signals** observed and the **chemical shifts** of those signals.


14.11A ^{13}C NMR: Number of Signals

- The number of signals in a ^{13}C spectrum gives the number of different types of carbon atoms in a molecule.

Carbon atoms in the same environment give the same NMR signal, whereas carbons in different environments give different NMR signals. The ^{13}C NMR spectrum of CH_3COOH has two signals because there are two different types of carbon atoms—the C of the CH_3 group and the C of the carbonyl ($\text{C}=\text{O}$).

- Because ^{13}C NMR signals are not split, the number of signals equals the number of lines in the ^{13}C NMR spectrum.


Thus, the ^{13}C NMR spectra of dimethyl ether, chloroethane, and methyl acetate exhibit one, two, and three lines, respectively, because these compounds contain one, two, and three different types of carbon atoms.



In contrast to what occurs in proton NMR, peak intensity is not proportional to the number of absorbing carbons, so ^{13}C NMR signals are not integrated.


Sample Problem 14.7

How many lines are observed in the ^{13}C NMR spectrum of each compound?




Solution

The number of different types of carbons equals the number of lines in a ^{13}C NMR spectrum.



Problem 14.28

How many lines are observed in the ^{13}C NMR spectrum of each compound?



Problem 14.29

Draw all constitutional isomers of molecular formula $\text{C}_3\text{H}_6\text{Cl}_2$.

- How many signals does each isomer exhibit in its ^1H NMR spectrum?
- How many lines does each isomer exhibit in its ^{13}C NMR spectrum?
- When only the number of signals in both ^1H and ^{13}C NMR spectroscopy is considered, is it possible to distinguish all of these constitutional isomers?







14.11B ^{13}C NMR: Position of Signals

In contrast to the small range of chemical shifts in ^1H NMR (0–12 ppm usually), ^{13}C NMR absorptions occur over a much broader range, 0–220 ppm. The chemical shifts of carbon atoms in ^{13}C NMR depend on the same effects as the chemical shifts of protons in ^1H NMR:


- The sp^3 hybridized C atoms of alkyl groups are shielded and absorb upfield.
- Electronegative elements like halogen, nitrogen, and oxygen shift absorptions downfield.
- The sp^2 hybridized C atoms of alkenes and benzene rings absorb downfield.
- Carbonyl carbons are highly deshielded, and absorb farther downfield than other carbon types.

Table 14.5 lists common ^{13}C chemical shift values. The ^{13}C NMR spectra of 1-propanol ($\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$) and methyl acetate ($\text{CH}_3\text{CO}_2\text{CH}_3$) in Figure 14.14 illustrate these principles.

Table 14.5 Common ^{13}C Chemical Shift Values

Type of carbon	Chemical shift (ppm)	Type of carbon	Chemical shift (ppm)
	5–45		100–140
 Z = N, O, X	30–80		120–150
	65–100		160–210

Problem 14.30 Which of the indicated carbon atoms in each molecule absorbs farther downfield?



Problem 14.31 Identify the carbon atoms that give rise to each NMR signal.







Figure 14.14
Representative ^{13}C NMR
spectra

a. 1-Propanol



- The three types of C's in 1-propanol—identified as C_a , C_b , and C_c —give rise to three ^{13}C NMR signals.
- Deshielding increases with increasing proximity to the electronegative O atom, and the absorption shifts downfield; thus, in order of increasing chemical shift: $C_a < C_b < C_c$.

b. Methyl acetate



- The three types of C's in methyl acetate—identified as C_a , C_b , and C_c —give rise to three ^{13}C NMR signals.
- The carbonyl carbon (C_b) is highly deshielded, so it absorbs farthest downfield.
- C_a , an sp^3 hybridized C that is not bonded to an O atom, is the most shielded, and so it absorbs farthest upfield.
- Thus, in order of increasing chemical shift: $C_a < C_c < C_b$.


Problem 14.32

A compound of molecular formula $\text{C}_4\text{H}_8\text{O}_2$ shows no IR peaks at $3600\text{--}3200$ or 1700 cm^{-1} . It exhibits one singlet in its ^1H NMR spectrum at 3.69 ppm, and one line in its ^{13}C NMR spectrum at 67 ppm. What is the structure of this unknown?

Problem 14.33

Draw the structure of a compound of molecular formula $\text{C}_4\text{H}_8\text{O}$ that has a signal in its ^{13}C NMR spectrum at > 160 ppm. Then draw the structure of an isomer of molecular formula $\text{C}_4\text{H}_8\text{O}$ that has all of its ^{13}C NMR signals at < 160 ppm.

Figure 14.15
Magnetic resonance imaging



- An MRI instrument: An MRI instrument is especially useful for visualizing soft tissue. In 2002, 60 million MRI procedures were performed. The 2003 Nobel Prize in Physiology or Medicine was awarded to chemist Paul C. Lauterbur and physicist Sir Peter Mansfield for their contributions in developing magnetic resonance imaging.
- An MRI image of the lower back: **A** labels spinal cord compression from a herniated disc. **B** labels the spinal cord, which would not be visualized with conventional X-rays.

14.12 Magnetic Resonance Imaging (MRI)

Magnetic resonance imaging (MRI)—NMR spectroscopy in medicine—is a powerful diagnostic technique (Figure 14.15a). The “sample” is the patient, who is placed in a large cavity in a magnetic field, and then irradiated with RF energy. Because RF energy has very low frequency and low energy, the method is safer than X-rays or computed tomography (CT) scans that employ high-frequency, high-energy radiation that is known to damage living cells.


Living tissue contains protons (especially the H atoms in H_2O) in different concentrations and environments. When irradiated with RF energy, these protons are excited to a higher energy spin state, and then fall back to the lower energy spin state. These data are analyzed by a computer that generates a plot that delineates tissues of different proton density (Figure 14.15b). MRIs can be recorded in any plane. Moreover, because the calcium present in bones is not NMR active, an MRI instrument can “see through” bones such as the skull and visualize the soft tissue underneath.

KEY CONCEPTS


Nuclear Magnetic Resonance Spectroscopy

1H NMR Spectroscopy

- The **number of signals** equals the number of different types of protons (14.2).
- The **position of a signal** (its chemical shift) is determined by shielding and deshielding effects.
 - Shielding shifts an absorption upfield; deshielding shifts an absorption downfield.
 - Electronegative atoms withdraw electron density, deshield a nucleus, and shift an absorption downfield (14.3).



- Loosely held π electrons can either shield or deshield a nucleus. Protons on benzene rings and double bonds are deshielded and absorb downfield, whereas protons on triple bonds are shielded and absorb upfield (14.4).



[3] The **area under an NMR signal** is proportional to the number of absorbing protons (14.5).

[4] **Spin-spin splitting** tells about nearby nonequivalent protons (14.6–14.8).

- Equivalent protons do not split each other's signals.
- A set of n nonequivalent protons on the same carbon or adjacent carbons splits an NMR signal into $n + 1$ peaks.
- OH and NH protons do not cause splitting (14.9).
- When an absorbing proton has two sets of nearby nonequivalent protons that are equivalent to each other, use the $n + 1$ rule to determine splitting.
- When an absorbing proton has two sets of nearby nonequivalent protons that are not equivalent to each other, the number of peaks in the NMR signal = $(n + 1)(m + 1)$. In flexible alkyl chains, peak overlap often occurs, resulting in $n + m + 1$ peaks in an NMR signal.

^{13}C NMR Spectroscopy (14.11)

[1] **The number of signals equals the number of different types of carbon atoms.** All signals are **single peaks**.










[2] The relative position of ^{13}C signals is determined by shielding and deshielding effects.

- Carbons that are sp^3 hybridized are shielded and absorb upfield.
- Electronegative elements (N, O, and halogen) shift absorptions downfield.
- The carbons of alkenes and benzene rings absorb downfield.
- Carbonyl carbons are highly deshielded, and absorb farther downfield than other carbon types.





PROBLEMS

^1H NMR Spectroscopy—Determining Equivalent Protons





14.34 How many different types of protons are present in each compound?



14.35 How many ^1H NMR signals does each compound give?



14.36 How many ^1H NMR signals does each natural product exhibit?



^1H NMR—Chemical Shift and Integration

14.37 Using a 300 MHz NMR instrument:

- How many Hz downfield from TMS is a signal at 2.5 ppm?
- If a signal comes at 1200 Hz downfield from TMS, at what ppm does it occur?
- If two peaks are separated by 2 ppm, how many Hz does this correspond to?



14.38 Acetone exhibits a singlet in its ^1H NMR spectrum at 2.16 ppm. If CH_2Cl_2 exhibits a singlet 1570 Hz downfield from acetone on a 500 MHz NMR spectrometer, what is the chemical shift of the singlet due to CH_2Cl_2 ?

14.39 Which of the indicated protons in each pair absorbs farther downfield?

- $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{OCH}_3$
- $\text{CH}_3\text{CH}_2\text{CH}_2\text{I}$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$
- $\text{CH}_3\text{OCH}_2\text{CH}_3$ or $\text{CH}_3\text{OCH}_2\text{CH}_2\text{CH}_3$
- $\text{CH}_3\text{CH}_2\text{CHBr}_2$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$


14.40 A compound of molecular formula C_6H_{10} gives three signals in its ^1H NMR spectrum with the following integration units: 13, 33, 73 units. How many protons are responsible for each signal?

14.41 How could you use chemical shift and integration data in ^1H NMR spectroscopy to distinguish between each pair of compounds? The ^1H NMR spectrum of each compound contains only singlets.

- $\text{CH}_3\text{CO}_2\text{C}(\text{CH}_3)_3$ and $\text{CH}_3\text{CO}_2\text{CH}_3$
- $\text{CH}_3\text{OCH}_2\text{CH}_2\text{OCH}_3$ and $\text{CH}_3\text{OCH}_2\text{OCH}_3$
-  and 













^1H NMR—Splitting

14.42 Which compounds give one singlet in the ^1H NMR spectrum?




14.43 For the five isomeric alkanes of molecular formula C_6H_{14} , label each type of proton and indicate how many peaks each will exhibit in its ^1H NMR signal.

14.44 Into how many peaks will the signal for each of the indicated protons be split?



14.45 How can you use ^1H NMR spectroscopy to distinguish between $\text{CH}_2=\text{C}(\text{Br})\text{CO}_2\text{CH}_3$ and methyl (2E)-3-bromo-2-propenoate, $\text{BrCH}=\text{CHCO}_2\text{CH}_3$?

14.46 Label the signals due to H_a , H_b , and H_c in the ^1H NMR spectrum of acrylonitrile ($\text{CH}_2=\text{CHCN}$). Draw a splitting diagram for the absorption due to the H_a proton.









^{13}C NMR

14.47 Draw the four constitutional isomers having molecular formula $\text{C}_4\text{H}_9\text{Br}$ and indicate how many different kinds of carbon atoms each has.


14.48 Which compounds in Problem 14.42 give one signal in their ^{13}C NMR spectra?

14.49 Explain why the carbonyl carbon of an aldehyde or ketone absorbs farther downfield than the carbonyl carbon of an ester in a ^{13}C NMR spectrum.

14.50 How many ^{13}C NMR signals does each compound exhibit?



14.51 Rank the indicated carbon atoms in each compound in order of increasing chemical shift.



14.52 Identify the carbon atoms that give rise to the signals in the ^{13}C NMR spectrum of each compound.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$; ^{13}C NMR: 14, 19, 35, and 62 ppm
 b. $(\text{CH}_3)_2\text{CHCHO}$; ^{13}C NMR: 16, 41, and 205 ppm
 c. $\text{CH}_2=\text{CHCH}(\text{OH})\text{CH}_3$; ^{13}C NMR: 23, 69, 113, and 143 ppm

- 14.53 a. How many signals does dimethyl fumarate ($\text{CH}_3\text{O}_2\text{CCH}=\text{CHCO}_2\text{CH}_3$, with a trans C=C) exhibit in its ^{13}C NMR spectrum?
 b. Draw the structure of an isomer of dimethyl fumarate that has each of the following number of signals in its ^{13}C NMR spectrum: [1] three; [2] four; [5] five.

Combined Spectroscopy Problems

Additional spectroscopy problems are located at the end of Chapters 15–23 and 25.

14.54 Propose a structure consistent with each set of spectral data:

- a. $\text{C}_4\text{H}_8\text{Br}_2$: IR peak at $3000\text{--}2850\text{ cm}^{-1}$; NMR (ppm):
 1.87 (singlet, 6 H)
 3.86 (singlet, 2 H)
 b. $\text{C}_3\text{H}_6\text{Br}_2$: IR peak at $3000\text{--}2850\text{ cm}^{-1}$; NMR (ppm):
 2.4 (quintet)
 3.5 (triplet)
 c. $\text{C}_5\text{H}_{10}\text{O}_2$: IR peak at 1740 cm^{-1} ; NMR (ppm):
 1.15 (triplet, 3 H) 2.30 (quartet, 2 H)
 1.25 (triplet, 3 H) 4.72 (quartet, 2 H)
 d. $\text{C}_6\text{H}_{14}\text{O}$: IR peak at $3600\text{--}3200\text{ cm}^{-1}$; NMR (ppm):
 0.8 (triplet, 6 H) 1.5 (quartet, 4 H)
 1.0 (singlet, 3 H) 1.6 (singlet, 1 H)
 e. $\text{C}_6\text{H}_{14}\text{O}$: IR peak at $3000\text{--}2850\text{ cm}^{-1}$; NMR (ppm):
 1.10 (doublet, 30 units)
 3.60 (septet, 5 units)
 f. $\text{C}_3\text{H}_6\text{O}$: IR peak at 1730 cm^{-1} ; NMR (ppm):
 1.11 (triplet)
 2.46 (multiplet)
 9.79 (triplet)

14.55 Identify the structures of isomers **A** and **B** (molecular formula $\text{C}_9\text{H}_{10}\text{O}$).

Compound **A**: IR peak at 1742 cm^{-1} ; ^1H NMR data (ppm) at 2.15 (singlet, 3 H), 3.70 (singlet, 2 H), and 7.20 (broad singlet, 5 H).


Compound **B**: IR peak at 1688 cm^{-1} ; ^1H NMR data (ppm) at 1.22 (triplet, 3 H), 2.98 (quartet, 2 H), and 7.28–7.95 (multiplet, 5 H).

14.56 Compound **C** has a molecular ion in its mass spectrum at 146 and a prominent absorption in its IR spectrum at 1762 cm^{-1} . **C** shows the following ^1H NMR spectral data: 1.47 (doublet, 3 H), 2.07 (singlet, 6 H), and 6.84 (quartet, 1 H) ppm. What is the structure of **C**?


14.57 As we will learn in Chapter 20, reaction of $(\text{CH}_3)_2\text{CO}$ with $\text{LiC}\equiv\text{CH}$ followed by H_2O affords compound **D**, which has a molecular ion in its mass spectrum at 84 and prominent absorptions in its IR spectrum at $3600\text{--}3200$, 3303, 2938, and 2120 cm^{-1} . **D** shows the following ^1H NMR spectral data: 1.53 (singlet, 6 H), 2.37 (singlet, 1 H), and 2.43 (singlet, 1 H) ppm. What is the structure of **D**?

14.58 Identify the structures of isomers **E** and **F** (molecular formula $C_4H_8O_2$).

Compound E: IR absorption at 1743 cm^{-1}




Compound F: IR absorption at 1730 cm^{-1}




14.59 Identify the structures of isomers **H** and **I** (molecular formula $C_8H_{11}N$).

a. **Compound H:** IR absorptions at 3365 , 3284 , 3026 , 2932 , 1603 , and 1497 cm^{-1}




b. **Compound I:** IR absorptions at 3367 , 3286 , 3027 , 2962 , 1604 , and 1492 cm^{-1}




14.60 Propose a structure consistent with each set of data.

a. $C_9H_{10}O_2$: IR absorption at 1718 cm^{-1}



b. C_9H_{12} : IR absorption at $2850\text{--}3150\text{ cm}^{-1}$




14.61 Propose a structure consistent with each set of data.



a. Compound **J**: molecular ion at 72; IR peak at 1710 cm^{-1} ; ^1H NMR data (ppm) at 1.0 (triplet, 3 H), 2.1 (singlet, 3 H), and 2.4 (quartet, 2 H)

b. Compound **K**: molecular ion at 88; IR peak at $3600\text{--}3200\text{ cm}^{-1}$; ^1H NMR data (ppm) at 0.9 (triplet, 3 H), 1.2 (singlet, 6 H), 1.5 (quartet, 2 H), and 1.6 (singlet, 1 H)


14.62 In the presence of a small amount of acid, a solution of acetaldehyde (CH_3CHO) in methanol (CH_3OH) was allowed to stand and a new compound **L** was formed. **L** has a molecular ion in its mass spectrum at 90 and IR absorptions at 2992 and 2941 cm^{-1} . **L** shows three signals in its ^{13}C NMR at 19, 52, and 101 ppm. The ^1H NMR spectrum of **L** is given below. What is the structure of **L**?




- 14.63 Treatment of $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_2\text{CH}_3$ with TsOH affords two products (**M** and **N**) with molecular formula C_6H_{12} . The ^1H NMR spectra of **M** and **N** are given below. Propose structures for **M** and **N** and draw a mechanism to explain their formation.




- 14.64 Compound **O** has molecular formula $\text{C}_{10}\text{H}_{12}\text{O}$ and shows an IR absorption at 1687 cm^{-1} . The ^1H NMR spectrum of **O** is given below. What is the structure of **O**?



14.65 Compound **P** has molecular formula $C_5H_9ClO_2$. Deduce the structure of **P** from its 1H and ^{13}C NMR spectra.



14.66 Treatment of 2-butanone ($CH_3COCH_2CH_3$) with strong base followed by CH_3I forms a compound **Q**, which gives a molecular ion in its mass spectrum at 86. The IR ($> 1500\text{ cm}^{-1}$ only) and 1H NMR spectrum of **Q** are given below. What is the structure of **Q**?



14.67 Low molecular weight esters (RCO_2R) often have characteristic odors. Using its molecular formula and 1H NMR spectral data, identify each ester.


- Compound **R**, the odor of banana: $C_7H_{14}O_2$; 1H NMR: 0.93 (doublet, 6 H), 1.52 (multiplet, 2 H), 1.69 (multiplet, 1 H), 2.04 (singlet, 3 H), and 4.10 (triplet, 2 H) ppm
- Compound **S**, the odor of rum: $C_7H_{14}O_2$; 1H NMR: 0.94 (doublet, 6 H), 1.15 (triplet, 3 H), 1.91 (multiplet, 1 H), 2.33 (quartet, 2 H), and 3.86 (doublet, 2 H) ppm

- 14.68** When 2-bromo-3,3-dimethylbutane is treated with $\text{K}^+ \text{OC}(\text{CH}_3)_3$, a single product **T** having molecular formula C_6H_{12} is formed. When 3,3-dimethyl-2-butanol is treated with H_2SO_4 , the major product **U** has the same molecular formula. Given the following ^1H NMR data, what are the structures of **T** and **U**? Explain in detail the splitting patterns observed for the three split signals in **T**.

^1H NMR of **T**: 1.01 (singlet, 9 H), 4.82 (doublet of doublets, 1 H, $J = 10, 1.7$ Hz), 4.93 (doublet of doublets, 1 H, $J = 18, 1.7$ Hz), and 5.83 (doublet of doublets, 1 H, $J = 18, 10$ Hz) ppm


^1H NMR of **U**: 1.60 (singlet) ppm

- 14.69** In a Baeyer–Villiger reaction, ketones ($\text{R}_2\text{C}=\text{O}$) are converted to esters (RCO_2R) by using peroxy acids. With an unsymmetrical ketone, two possible esters can be formed, as shown for 3,3-dimethyl-2-butanone as starting material. How could you use spectroscopic techniques— ^1H NMR, IR, and MS—to determine which ester (**A** or **B**) is formed?




- 14.70** Propose a structure consistent with each set of data.

- a. A compound **X** (molecular formula $\text{C}_6\text{H}_{12}\text{O}_2$) gives a strong peak in its IR spectrum at 1740 cm^{-1} . The ^1H NMR spectrum of **X** shows only two singlets, including one at 3.5 ppm. The ^{13}C NMR spectrum is given below. Propose a structure for **X**.




- b. A compound **Y** (molecular formula C_6H_{10}) gives four lines in its ^{13}C NMR spectrum (27, 30, 67, and 93 ppm), and the IR spectrum given here. Propose a structure for **Y**.




Challenge Problems

- 14.71 The ^1H NMR spectrum of *N,N*-dimethylformamide shows three singlets at 2.9, 3.0, and 8.0 ppm. Explain why the two CH_3 groups are not equivalent to each other, thus giving rise to two NMR signals.



- 14.72 18-Annulene shows two signals in its ^1H NMR spectrum, one at 8.9 (12 H) and one at -1.8 (6 H) ppm. Using a similar argument to that offered for the chemical shift of benzene protons, explain why both shielded and deshielded values are observed for 18-annulene.




- 14.73 Explain why the ^{13}C NMR spectrum of 3-methyl-2-butanol shows five signals.
- 14.74 Since ^{31}P has an odd mass number, ^{31}P nuclei absorb in the NMR and, in many ways, these nuclei behave similarly to protons in NMR spectroscopy. With this in mind, explain why the ^1H NMR spectrum of methyl dimethylphosphonate, $\text{CH}_3\text{PO}(\text{OCH}_3)_2$, consists of two doublets at 1.5 and 3.7 ppm.

15

Radical Reactions

- 15.1 Introduction
- 15.2 General features of radical reactions
- 15.3 Halogenation of alkanes
- 15.4 The mechanism of halogenation
- 15.5 Chlorination of other alkanes
- 15.6 Chlorination versus bromination
- 15.7 Halogenation as a tool in organic synthesis
- 15.8 The stereochemistry of halogenation reactions
- 15.9 Application: The ozone layer and CFCs
- 15.10 Radical halogenation at an allylic carbon
- 15.11 Application: Oxidation of unsaturated lipids
- 15.12 Application: Antioxidants
- 15.13 Radical addition reactions to double bonds
- 15.14 Polymers and polymerization



Polystyrene, an inexpensive polymer synthesized from the monomer styrene, $C_6H_5CH=CH_2$, is one of the six compounds—called the “Big Six”—that account for three-quarters of the synthetic polymers produced in the United States. The polystyrene foam used in packaging materials and drinking cups for hot beverages is called Styrofoam, a trademark of the Dow Chemical Company. Polystyrene is also used to form the housings of small kitchen appliances, televisions, computers, and CD cases. Although recycled polystyrene can be molded into trays and trash cans, the polystyrene used in food packaging and beverage cups is contaminated with food, making it difficult to clean and recycle. In Chapter 15, we learn about the synthesis of polymers like polystyrene.


A **small but significant group of reactions** involves the homolysis of nonpolar bonds to form highly reactive **radical intermediates**. Although they are unlike other organic reactions, radical transformations are important in many biological and industrial processes. The gases O_2 and NO (nitric oxide) are both radicals. This means that many oxidation reactions with O_2 involve radical intermediates, and biological processes mediated by NO such as blood clotting and neurotransmission may involve radicals. Many useful industrial products such as Styrofoam and polyethylene are prepared by radical processes.

In Chapter 15 we examine the cleavage of nonpolar bonds by radical reactions.

15.1 Introduction

Radicals were first discussed in Section 6.3.

- A **radical** is a reactive intermediate with a single unpaired electron, formed by homolysis of a covalent bond.




A **radical contains an atom that does not have an octet of electrons**, making it reactive and unstable. Radical processes involve single electrons, so half-headed arrows are used to show the movement of electrons. **One half-headed arrow is used for each electron**.

Carbon radicals are classified as **primary (1°)**, **secondary (2°)**, or **tertiary (3°)** by the number of R groups bonded to the carbon with the unpaired electron. A carbon radical is sp^2 hybridized and **trigonal planar**, like sp^2 hybridized carbocations. The unhybridized p orbital contains the unpaired electron and extends above and below the trigonal planar carbon.



Classification of carbon radicals



The trigonal planar geometry of a carbon radical



Bond dissociation energies for the cleavage of C–H bonds are used as a measure of radical stability. For example, two different radicals can be formed by cleavage of the C–H bonds in $CH_3CH_2CH_3$.





Cleavage of the stronger 1° C–H bond to form the 1° radical ($CH_3CH_2CH_2\cdot$) requires more energy than cleavage of the weaker 2° C–H bond to form the 2° radical [$(CH_3)_2CH\cdot$] $—410$ versus 397 kJ/mol. This makes the 2° radical more stable, because less energy is required for its formation, as illustrated in Figure 15.1. Thus, **cleavage of the weaker bond forms the more stable radical**. This is a specific example of a general trend.

- The stability of a radical increases as the number of alkyl groups bonded to the radical carbon increases.

Figure 15.1

The relative stability of 1° and 2° carbon radicals







The **lower** the bond dissociation energy for a C–H bond, the **more stable** the resulting carbon radical.

Thus, a 3° radical is more stable than a 2° radical, and a 2° radical is more stable than a 1° radical. Increasing alkyl substitution increases radical stability in the same way it increases carbocation stability. Alkyl groups are more polarizable than hydrogen atoms, so they can more easily donate electron density to the electron-deficient carbon radical, thus increasing stability.



Unlike carbocations, however, less stable radicals generally do *not* rearrange to more stable radicals. This difference can be used to distinguish between reactions involving radical intermediates and those involving carbocations.

Problem 15.1

Classify each radical as 1°, 2°, or 3°.

**Problem 15.2**

Draw the most stable radical that can result from cleavage of a C–H bond in each molecule.

**15.2 General Features of Radical Reactions**


Radicals are formed from covalent bonds by adding energy in the form of **heat** (Δ) or **light** ($h\nu$). Some radical reactions are carried out in the presence of a **radical initiator**, a compound that contains an especially weak bond that serves as a source of radicals. **Peroxides**, compounds with the general structure **RO–OR**, are the most commonly used radical initiators. Heating a peroxide readily causes homolysis of the weak O–O bond, forming two $\text{RO}\cdot$ radicals.

15.2A Two Common Reactions of Radicals

Radicals undergo two main types of reactions: **they react with σ bonds**, and **they add to π bonds**, in both cases achieving an octet of electrons.

[1] Reaction of a Radical $\text{X}\cdot$ with a C–H Bond


A radical $\text{X}\cdot$ **abstracts a hydrogen atom from a C–H σ bond to form H–X and a carbon radical**. One electron from the C–H bond is used to form the new H–X bond, and the other electron in the C–H bond remains on carbon. The result is that the original radical $\text{X}\cdot$ is now surrounded by an octet of electrons, and a new radical is formed.



This radical reaction is typically seen with the nonpolar C–H bonds of **alkanes**, which cannot react with polar or ionic electrophiles and nucleophiles.

[2] Reaction of a Radical X• with a C=C

A radical X• also adds to the π bond of a carbon–carbon double bond. One electron from the double bond is used to form a new C–X bond, and the other electron remains on the other carbon originally part of the double bond.




Whenever a radical reacts with a stable single or double bond, a new radical is formed in the products.


Although the electron-rich double bond of an **alkene** reacts with electrophiles by ionic addition mechanisms, it also reacts with radicals because these reactive intermediates are also electron deficient.

15.2B Two Radicals Reacting with Each Other

A radical, once formed, rapidly reacts with whatever is available. Usually that means a stable σ or π bond. Occasionally, however, two radicals come into contact with each other, and they react to form a σ bond.



The reaction of a radical with oxygen, a diradical in its ground state electronic configuration, is another example of two radicals reacting with each other. In this case, the reaction of O₂ with X• forms a new radical, thus preventing X• from reacting with an organic substrate.




Compounds that prevent radical reactions from occurring are called *radical inhibitors* or *radical scavengers*. Besides O₂, vitamin E and related compounds, discussed in Section 15.12, are radical scavengers, too. The fact that these compounds inhibit a reaction often suggests that the reaction occurs via radical intermediates.

Problem 15.3 Draw the products formed when a chlorine atom (Cl•) reacts with each species.

- a. CH₃–CH₃ b. CH₂=CH₂ c. $\cdot\ddot{\text{O}}\cdot$ d. O₂


15.3 Halogenation of Alkanes

In the presence of light or heat, alkanes react with halogens to form alkyl halides. Halogenation is a **radical substitution reaction**, because a halogen atom X replaces a hydrogen via a mechanism that involves radical intermediates.






Halogenation of alkanes is useful only with Cl₂ and Br₂. Reaction with F₂ is too violent and reaction with I₂ is too slow to be useful. With an alkane that has more than one type of hydrogen atom, a mixture of alkyl halides may result (Reaction [3]).

Figure 15.2

Complete halogenation of CH_4 using excess Cl_2 

When asked to draw the products of halogenation of an alkane, **draw the products of monohalogenation only**, unless specifically directed to do otherwise.

Examples




In these examples of halogenation, a halogen has replaced a single hydrogen atom on the alkane. Can the other hydrogen atoms be replaced, too? Figure 15.2 shows that when CH_4 is treated with excess Cl_2 , all four hydrogen atoms can be successively replaced by Cl to form CCl_4 . **Monohalogenation**—the substitution of a single H by X—can be achieved experimentally by adding halogen X_2 to an excess of alkane.

Sample Problem 15.1

Draw all the constitutional isomers formed by monohalogenation of $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_3$ with Cl_2 and $h\nu$.




Solution

Substitute Cl for H on every carbon, and then check to see if any products are identical. The starting material has five C's, but replacement of one H atom on two C's gives the same product. Thus, **$(\text{CH}_3)_2\text{CHCH}_2\text{CH}_3$ affords four monochloro substitution products.**



Problem 15.4

Draw all constitutional isomers formed by monochlorination of each alkane.



Problem 15.5

Compounds **A** and **B** are isomers having molecular formula C_5H_{12} . Heating **A** with Cl_2 gives a single product of monohalogenation, whereas heating **B** under the same conditions forms three constitutional isomers. What are the structures of **A** and **B**?

15.4 The Mechanism of Halogenation


Unlike nucleophilic substitution, which proceeds by two different mechanisms depending on the starting material and reagent, all halogenation reactions of alkanes—regardless of the halogen and alkane used—proceed by the *same* mechanism. Three facts about halogenation suggest that the mechanism involves radical, not ionic, intermediates.

Fact	Explanation
[1] Light, heat, or added peroxide is necessary for the reaction.	<ul style="list-style-type: none"> Light or heat provides the energy needed for homolytic bond cleavage to form radicals. Breaking the weak O–O bond of peroxides initiates radical reactions as well.
[2] O ₂ inhibits the reaction.	<ul style="list-style-type: none"> The diradical O₂ removes radicals from a reaction mixture, thus preventing reaction.
[3] No rearrangements are observed.	<ul style="list-style-type: none"> Radicals do not rearrange.

15.4A The Steps of Radical Halogenation

The chlorination of ethane illustrates the **three distinct parts of radical halogenation** (Mechanism 15.1):

Overall reaction




- Initiation:** Two radicals are formed by homolysis of a σ bond and this begins the reaction.
- Propagation:** A radical reacts with another reactant to form a new σ bond and another radical.
- Termination:** Two radicals combine to form a stable bond. Removing radicals from the reaction mixture without generating any new radicals stops the reaction.



Mechanism 15.1 Radical Halogenation of Alkanes

Initiation



Step [1] Bond cleavage forms two radicals.



- Homolysis of the weakest bond in the starting materials requires energy from light or heat.
- Thus, the Cl–Cl bond ($\Delta H^\circ = 242$ kJ/mol), which is weaker than either the C–C or C–H bond in ethane ($\Delta H^\circ = 368$ and 410 kJ/mol, respectively), is broken to form two chlorine radicals.

Propagation

Steps [2] and [3] One radical reacts and a new radical is formed.




Repeat Steps [2], [3], [2], [3], again and again.

- The Cl \cdot radicals abstract a hydrogen atom from ethane (Step [2]). This forms H–Cl and leaves one unpaired electron on carbon, generating the ethyl radical (CH₃CH₂ \cdot).
- CH₃CH₂ \cdot abstracts a chlorine atom from Cl₂ (Step [3]), forming CH₃CH₂Cl and a new chlorine radical (Cl \cdot).
- The Cl \cdot radical formed in Step [3] is a reactant in Step [2], so Steps [2] and [3] can occur repeatedly without an additional initiation reaction (Step [1]).
- In each propagation step, one radical is consumed and one radical is formed. The two products—CH₃CH₂Cl and HCl—are formed during propagation.

Termination

Step [4] Two radicals react to form a σ bond.



- To terminate the chain, two radicals react with each other in one of three ways (Steps [4a, b, and c]) to form stable bonds.

Although initiation generates the $\text{Cl}\cdot$ radicals needed to begin the reaction, the propagation steps ([2] and [3]) form the two reaction products— $\text{CH}_3\text{CH}_2\text{Cl}$ and HCl . Once the process has begun, propagation occurs over and over without the need for Step [1] to occur. **A mechanism such as radical halogenation that involves two or more repeating steps is called a chain mechanism.** Each propagation step involves a reactive radical abstracting an atom from a stable bond to form a new bond and another radical that continues the chain.

Usually a radical reacts with a stable bond to propagate the chain, but occasionally two radicals combine, and this reaction terminates the chain. Depending on the reaction and the reaction conditions, some radical chain mechanisms can repeat thousands of times before termination occurs.

Termination Step [4a] forms Cl_2 , a reactant, whereas Step [4c] forms $\text{CH}_3\text{CH}_2\text{Cl}$, one of the reaction products. Termination Step [4b] forms $\text{CH}_3\text{CH}_2\text{-CH}_2\text{CH}_3$, which is neither a reactant nor a desired product. The formation of a small quantity of $\text{CH}_3\text{CH}_2\text{-CH}_2\text{CH}_3$, however, is evidence that ethyl radicals are formed in the reaction.

The most important steps of radical halogenation are those that lead to product formation—the propagation steps—so subsequent discussion of this reaction concentrates on these steps only.

Problem 15.6 Using Mechanism 15.1 as a guide, write the mechanism for the reaction of CH_4 with Br_2 to form CH_3Br and HBr . Classify each step as initiation, propagation, or termination.

15.4B Energy Changes During the Chlorination of Ethane

Figure 15.3 shows how bond dissociation energies (Section 6.4) can be used to calculate ΔH° for the two propagation steps in the chlorination of ethane. Because the overall ΔH° is negative, the reaction is **exothermic**. Moreover, because the transition state for the first propagation step is higher in energy than the transition state for the second propagation step, the **first step is rate-determining**. Both of these facts are illustrated in the energy diagram in Figure 15.4.

Problem 15.7 Calculate ΔH° for the two propagation steps in the reaction of CH_4 with Br_2 to form CH_3Br and HBr (Problem 15.6).

Problem 15.8 Calculate ΔH° for the rate-determining step of the reaction of CH_4 with I_2 . Explain why this result illustrates that this reaction is extremely slow.

Figure 15.3
Energy changes in the propagation steps during the chlorination of ethane






Figure 15.4
Energy diagram for the propagation steps in the chlorination of ethane



- Because radical halogenation consists of two propagation steps, the energy diagram has two energy barriers.
- The first step is rate-determining because its transition state is at higher energy.
- The reaction is exothermic because $\Delta H^\circ_{\text{overall}}$ is negative.


15.5 Chlorination of Other Alkanes

Recall from Section 15.3 that the chlorination of $\text{CH}_3\text{CH}_2\text{CH}_3$ affords a 1:1 mixture of $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ (formed by removal of a 1° hydrogen) and $(\text{CH}_3)_2\text{CHCl}$ (formed by removal of a 2° hydrogen).






$\text{CH}_3\text{CH}_2\text{CH}_3$ has six 1° hydrogen atoms and only two 2° hydrogens, so the expected product ratio of $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$ to $(\text{CH}_3)_2\text{CHCl}$ (assuming all hydrogens are *equally* reactive) is 3:1. Because the observed ratio is 1:1, however, the 2° C–H bonds must be more reactive; that is, **it must be easier to homolytically cleave a 2° C–H bond than a 1° C–H bond**. Recall from Section 15.2 that 2° C–H bonds are weaker than 1° C–H bonds. Thus,

- The weaker the C–H bond, the more readily the hydrogen atom is removed in radical halogenation.



When alkanes react with Cl_2 , a mixture of products results, with more product formed by cleavage of the weaker C–H bond than you would expect on statistical grounds.

Problem 15.9 Which C–H bond in each compound is most readily broken during radical halogenation?




15.6 Chlorination versus Bromination

Although alkanes undergo radical substitution reactions with both Cl_2 and Br_2 , chlorination and bromination exhibit two important differences:

- Chlorination is *faster* than bromination.
- Although chlorination is *unselective*, yielding a mixture of products, bromination is often *selective*, yielding one major product.

For example, propane reacts rapidly with Cl_2 to form a 1:1 mixture of 1° and 2° alkyl chlorides. On the other hand, propane reacts with Br_2 much more slowly and forms 99% $(\text{CH}_3)_2\text{CHBr}$.




This is a specific example of the **reactivity–selectivity principle: less reactive reagents are more selective. In bromination, the major (and sometimes exclusive) product results from cleavage of the weakest C–H bond.**

Sample Problem 15.2

Draw the major product formed when 3-ethylpentane is heated with Br_2 .





Solution

Keep in mind: **the more substituted the carbon atom, the weaker the C–H bond.** The major bromination product in 3-ethylpentane is formed by cleavage of the sole 3° C–H bond, its weakest C–H bond.



Problem 15.10

Draw the major product formed when each cycloalkane is heated with Br_2 .




To explain the difference between chlorination and bromination, we return to the Hammond postulate (Section 7.15) to estimate the relative energy of the transition states of the rate-determining steps of these reactions. The **rate-determining step is the abstraction of a hydrogen atom by the halogen radical**, so we must compare these steps for bromination and chlorination. Keep in mind:

- Transition states in endothermic reactions resemble the products. The more stable product is formed faster.
- Transition states in exothermic reactions resemble the starting materials. The relative stability of the products does not greatly affect the relative energy of the transition states, so a mixture of products often results.

Bromination: $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{Br}_2$

A bromine radical can abstract either a 1° or a 2° hydrogen from propane, generating either a 1° radical or a 2° radical. Calculating ΔH° using bond dissociation energies reveals that both reactions are **endothermic**, but **it takes less energy to form the more stable 2° radical**.




According to the Hammond postulate, the transition state of an endothermic reaction resembles the products, so the energy of activation to form the more stable 2° radical is lower and it is formed faster, as shown in the energy diagram in Figure 15.5. Because the 2° radical $[(\text{CH}_3)_2\text{CH}\cdot]$ is converted to 2-bromopropane $[(\text{CH}_3)_2\text{CHBr}]$ in the second propagation step, this **2° alkyl halide is the major product of bromination**.

- **Conclusion:** Because the rate-determining step in bromination is endothermic, the more stable radical is formed faster, and often a single radical halogenation product predominates.

Chlorination: $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{Cl}_2$

A chlorine radical can also abstract either a 1° or a 2° hydrogen from propane, generating either a 1° radical or a 2° radical. Calculating ΔH° using bond dissociation energies reveals that both reactions are **exothermic**.



Figure 15.5
Energy diagram for a selective endothermic reaction: $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{Br}\cdot \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\cdot$ or $(\text{CH}_3)_2\text{CH}\cdot + \text{HBr}$



- The transition state to form the less stable 1° radical ($\text{CH}_3\text{CH}_2\text{CH}_2\cdot$) is higher in energy than the transition state to form the more stable 2° radical $[(\text{CH}_3)_2\text{CH}\cdot]$. Thus, **the 2° radical is formed faster**.

Figure 15.6

Energy diagram for a nonselective exothermic reaction: $\text{CH}_3\text{CH}_2\text{CH}_3 + \text{Cl}\cdot \rightarrow \text{CH}_3\text{CH}_2\dot{\text{C}}\text{H}_2$ or $(\text{CH}_3)_2\dot{\text{C}}\text{H} + \text{HCl}$



Because chlorination has an *exothermic* rate-determining step, the transition state to form both radicals resembles the same starting material, $\text{CH}_3\text{CH}_2\text{CH}_3$. As a result, the relative stability of the two radicals is much less important and both radicals are formed. An energy diagram for these processes is drawn in Figure 15.6. Because the 1° and 2° radicals are converted to 1-chloropropane ($\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$) and 2-chloropropane [$(\text{CH}_3)_2\text{CHCl}$], respectively, in the second propagation step, **both alkyl halides are formed in chlorination.**

- **Conclusion:** Because the rate-determining step in chlorination is exothermic, the transition state resembles the starting material, both radicals are formed, and a mixture of products results.


Problem 15.11 Why is the reaction of methylcyclohexane with Cl_2 not a useful method to prepare 1-chloro-1-methylcyclohexane? What other constitutional isomers are formed in the reaction mixture?

Problem 15.12 Reaction of $(\text{CH}_3)_3\text{CH}$ with Cl_2 forms two products: $(\text{CH}_3)_2\text{CHCH}_2\text{Cl}$ (63%) and $(\text{CH}_3)_3\text{CCl}$ (37%). Why is the major product formed by cleavage of the stronger 1° C-H bond?

15.7 Halogenation as a Tool in Organic Synthesis

Halogenation is a useful tool because it adds a functional group to a previously unfunctionalized molecule, making an **alkyl halide**. These alkyl halides can then be converted to alkenes by elimination, and to alcohols and ethers by nucleophilic substitution.


Sample Problem 15.3 Show how cyclohexane can be converted to cyclohexene by a stepwise sequence.




Solution

There is no one-step method to convert an alkane to an alkene. A two-step method is needed:

[1] **Radical halogenation** produces an alkyl halide.



[2] **Elimination of HCl** with a strong base produces cyclohexene.



Problem 15.13 Synthesize each compound from $(CH_3)_3CH$.

- a. $(CH_3)_3CBr$ b. $(CH_3)_2C=CH_2$ c. $(CH_3)_3COH$ d. $(CH_3)_2C(Cl)CH_2Cl$

Problem 15.14 Show all steps and reagents needed to convert cyclohexane into each compound: (a) the two enantiomers of *trans*-1,2-dibromocyclohexane; and (b) 1,2-epoxycyclohexane.

15.8 The Stereochemistry of Halogenation Reactions


The stereochemistry of a reaction product depends on whether the reaction occurs at a stereogenic center or at another atom, and whether a new stereogenic center is formed. The rules predicting the stereochemistry of reaction products are summarized in Table 15.1.

Table 15.1 Rules for Predicting the Stereochemistry of Reaction Products

Starting material	Result
Achiral	<ul style="list-style-type: none"> An achiral starting material always gives either an achiral or a racemic product.
Chiral	<ul style="list-style-type: none"> If a reaction does not occur at a stereogenic center, the configuration at a stereogenic center is retained in the product. If a reaction occurs at a stereogenic center, we must know the mechanism to predict the stereochemistry of the product.


15.8A Halogenation of an Achiral Starting Material

Halogenation of the **achiral starting material** $CH_3CH_2CH_2CH_3$ forms two constitutional isomers by replacement of either a 1° or 2° hydrogen.



- 1-Chlorobutane ($CH_3CH_2CH_2CH_2Cl$) has no stereogenic center and thus it is an **achiral** compound.
- 2-Chlorobutane [$CH_3CH(Cl)CH_2CH_3$] has a new stereogenic center, and so an **equal amount of two enantiomers** must form—a **racemic mixture**.


A racemic mixture results when a new stereogenic center is formed because the first propagation step generates a **planar, sp^2 hybridized radical**. Cl_2 then reacts with the planar radical from either the front or back side to form an equal amount of two enantiomers.




Thus, the achiral starting material butane forms an achiral product (1-chlorobutane) and a racemic mixture of two enantiomers [($2R$)- and ($2S$)-2-chlorobutane].

15.8B Halogenation of a Chiral Starting Material

Let's now examine chlorination of the chiral starting material ($2R$)-2-bromobutane at C2 and C3.




Chlorination at C2 occurs at the stereogenic center. Abstraction of a hydrogen atom at C2 forms a trigonal planar sp^2 hybridized radical that is now achiral. This achiral radical then reacts with Cl_2 from either side to form a new stereogenic center, resulting in an **equal amount of two enantiomers—a racemic mixture**.



- Radical halogenation reactions occur with racemization at a stereogenic center.





Chlorination at C3 does *not* occur at the stereogenic center, but it forms a new stereogenic center. Because no bond is broken to the stereogenic center at C2, **its configuration is retained** during the reaction. Abstraction of a hydrogen atom at C3 forms a trigonal planar sp^2 hybridized radical that still contains this stereogenic center. Reaction of the radical with Cl_2 from either side forms a new stereogenic center, so the products have two stereogenic centers: the configuration at C2 is the same in both compounds, but the configuration at C3 is different, making them **diastereomers**.



Thus, four isomers are formed by chlorination of (2*R*)-2-bromobutane at C2 and C3. Attack at the stereogenic center (C2) gives a product with one stereogenic center, resulting in a mixture of enantiomers. Attack at C3 forms a new stereogenic center, giving a mixture of diastereomers.

Problem 15.15 What products are formed from monochlorination of (2*R*)-2-bromobutane at C1 and C4? Assign *R* and *S* designations to each stereogenic center.

Problem 15.16 Draw the monochlorination products formed when each compound is heated with Cl₂. Include the stereochemistry at any stereogenic center.




15.9 Application: The Ozone Layer and CFCs

The 1995 Nobel Prize in Chemistry was awarded to Mario Molina, Paul Crutzen, and F. Sherwood Rowland for their work in elucidating the interaction of ozone with CFCs. What began as a fundamental research project turned out to have important implications in the practical world.




Propane and butane are now used as propellants in spray cans in place of CFCs.



O₃ destruction is most severe in the region of the South Pole, where a large ozone hole is visible with satellite imaging.


Ozone is formed in the upper atmosphere by reaction of oxygen molecules with oxygen atoms. Ozone is also decomposed with sunlight back to these same two species. The overall result of these reactions is to convert high-energy ultraviolet light into heat.

The synthesis and decomposition of O₃ in the upper atmosphere




Ozone is vital to life; it acts like a shield, protecting the earth's surface from destructive ultraviolet radiation. A decrease in ozone concentration in this protective layer would have some immediate consequences, including an increase in the incidence of skin cancer and eye cataracts. Other long-term effects include a reduced immune response, interference with photosynthesis in plants, and harmful effects on the growth of plankton, the mainstay of the ocean food chain.

Current research suggests that **chlorofluorocarbons (CFCs)** are responsible for destroying ozone in the upper atmosphere. CFCs are simple halogen-containing organic compounds manufactured under the trade name Freons.



CFCl₃
trichlorofluoromethane
CFC 11
Freon 11




CF₂Cl₂
dichlorodifluoromethane
CFC 12
Freon 12

CFCs are inert, odorless, and nontoxic, and they have been used as refrigerants, solvents, and aerosol propellants. Because CFCs are volatile and water insoluble, they readily escape into the upper atmosphere, where they are decomposed by high-energy sunlight to form radicals that destroy ozone by the radical chain mechanism shown in Figure 15.7.


The overall result is that O₃ is consumed as a reactant and O₂ molecules are formed. In this way, a small amount of CFC can destroy a large amount of O₃. These findings led to a ban on the use of CFCs in aerosol propellants in the United States in 1978 and to the phasing out of their use in refrigeration systems.

Figure 15.7

CFCs and the destruction of the ozone layer



Initiation: CFCs are decomposed by sunlight to form chlorine radicals.




Propagation: Ozone is destroyed by a chain reaction with radical intermediates.



- The chain reaction is initiated by homolysis of a C–Cl bond in CFCl_3 .
- Propagation consists of two steps. Reaction of $\text{Cl}\cdot$ with O_3 forms chlorine monoxide ($\text{ClO}\cdot$), which reacts with oxygen atoms to form O_2 and $\text{Cl}\cdot$.

Newer alternatives to CFCs are **hydrochlorofluorocarbons (HCFCs)** and **hydrofluorocarbons (HFCs)** such as CH_2FCF_3 . These compounds have many properties in common with CFCs, but they are largely decomposed by $\text{HO}\cdot$ before they reach the stratosphere and therefore they have little impact on stratospheric O_3 .



CH_2FCF_3
HFC-134a




This HFC is decomposed before it reaches the stratosphere.

Problem 15.17


Nitric oxide, $\text{NO}\cdot$, is another radical also thought to cause ozone destruction by a similar mechanism. One source of $\text{NO}\cdot$ in the stratosphere is supersonic aircraft whose jet engines convert small amounts of N_2 and O_2 to $\text{NO}\cdot$. Write the propagation steps for the reaction of O_3 with $\text{NO}\cdot$.

15.10 Radical Halogenation at an Allylic Carbon

Now let's examine radical halogenation at an **allylic carbon**—the carbon adjacent to a **double bond**. Homolysis of the allylic C–H bond of propene generates the **allyl radical**, which has an unpaired electron on the carbon adjacent to the double bond.




The bond dissociation energy for this process (364 kJ/mol) is even less than that for a 3° C–H bond (381 kJ/mol). Because the weaker the C–H bond, the more stable the resulting radical, an **allyl radical is more stable than a 3° radical**, and the following order of radical stability results:



The position of the atoms and the σ bonds stays the same in drawing resonance structures. Resonance structures differ in the location of only π bonds and nonbonded electrons.


The allyl radical is more stable than other radicals because two resonance structures can be drawn for it.



- The “true” structure of the allyl radical is a hybrid of the two resonance structures. In the hybrid, the π bond and the unpaired electron are delocalized.
- Delocalizing electron density lowers the energy of the hybrid, thus stabilizing the allyl radical.


Problem 15.18

Draw a second resonance structure for each radical. Then draw the hybrid.



15.10A Selective Bromination at Allylic C–H Bonds

Because allylic C–H bonds are weaker than other sp^3 hybridized C–H bonds, the **allylic carbon can be selectively halogenated** by using *N*-bromosuccinimide (**NBS**, Section 10.15) in the presence of light or peroxides. Under these conditions only the allylic C–H bond in cyclohexene reacts to form an allylic halide.




NBS contains a weak N–Br bond that is homolytically cleaved with light to generate a bromine radical, initiating an allylic halogenation reaction. Propagation then consists of the usual two steps of radical halogenation as shown in Mechanism 15.2.



Mechanism 15.2 Allylic Bromination with NBS

Initiation



Step [1] Cleavage of the N–Br bond forms two radicals.



- The reaction begins with homolysis of the weak N–Br bond in NBS using light energy. This generates a Br· radical that begins the radical halogenation process.


Propagation

Steps [2] and [3] One radical reacts and a new radical is formed in each step.




- The Br· radical abstracts an allylic hydrogen atom to afford an allylic radical in Step [2]. (Only one Lewis structure of the allylic radical is drawn.)
- The allylic radical reacts with Br₂ in the second propagation step to form the product of allylic halogenation. Because the Br· radical formed in Step [3] is also a reactant in Step [2], Steps [2] and [3] repeatedly occur without the need for Step [1].

Besides acting as a source of Br· to initiate the reaction, NBS generates a low concentration of Br₂ needed in the second chain propagation step (Step [3] of the mechanism). The HBr formed in Step [2] reacts with NBS to form Br₂, which is then used for halogenation in Step [3] of the mechanism.



A low concentration of Br₂ (from NBS) favors allylic substitution (over addition) in part because bromine is needed for only one step of the mechanism. When Br₂ adds to a double bond, a low Br₂ concentration would first form a low concentration of bridged bromonium ion (Section 10.13), which must then react with more bromine (in the form of Br[–]) in a second step to form a dibromide. If concentrations of both intermediates—bromonium ion and Br[–]—are low, the overall rate of addition is very slow.

Thus, an alkene with allylic C–H bonds undergoes two different reactions depending on the reaction conditions.



- Treatment of cyclohexene with Br₂ (in an organic solvent like CCl₄) leads to **addition** via **ionic intermediates** (Section 10.13).
- Treatment of cyclohexene with NBS (+ *hν* or ROOR) leads to **allylic substitution**, via **radical intermediates**.

Problem 15.19

Draw the products of each reaction.





Figure 15.8
The oxidation of unsaturated lipids with O_2



- Oxidation is shown at one allylic carbon only. Reaction at the other labeled allylic carbon is also possible.

15.11 Application: Oxidation of Unsaturated Lipids

Oils—triacylglycerols having one or more sites of unsaturation in their long carbon chains—are susceptible to oxidation at their allylic carbon atoms. Oxidation occurs by way of a radical chain mechanism, as shown in Figure 15.8.


- **Step [1]** Oxygen in the air abstracts an allylic hydrogen atom to form an allylic radical because the allylic C-H bond is weaker than the other C-H bonds.
- **Step [2]** The allylic radical reacts with another molecule of O_2 to form a peroxy radical.
- **Step [3]** The peroxy radical abstracts an allylic hydrogen from another lipid molecule to form a hydroperoxide and another allylic radical that continues the chain. Steps [2] and [3] can repeat again and again until some other radical terminates the chain.

The hydroperoxides formed by this process are unstable and decompose to other oxidation products, many of which have a disagreeable odor and taste. **This process turns an oil rancid. Unsaturated lipids are more easily oxidized than saturated ones** because they contain weak allylic C-H bonds that are readily cleaved in Step [1] of this reaction, forming resonance-stabilized allylic radicals. Because saturated fats have no double bonds and thus no weak allylic C-H bonds, they are much less susceptible to air oxidation, resulting in increased shelf life of products containing them.

Problem 15.23

Draw a second resonance structure for the allylic radical formed as a product of Step [1] in Figure 15.8. What hydroperoxide is formed using this Lewis structure?


Problem 15.24 Which C–H bond is most readily cleaved in linoleic acid? Draw all possible resonance structures for the resulting radical. Draw all the hydroperoxides formed by reaction of this resonance-stabilized radical with O_2 .





15.12 Application: Antioxidants

An *antioxidant* is a compound that stops an oxidation reaction from occurring.

- Naturally occurring antioxidants such as **vitamin E** prevent radical reactions that can cause cell damage.
- Synthetic antioxidants such as **BHT**—butylated hydroxy toluene—are added to packaged and prepared foods to prevent oxidation and spoilage.



The purported health benefits of antioxidants have made them a popular component in anti-aging formulations.




Vitamin E and BHT are radical inhibitors, so they terminate radical chain mechanisms by reacting with radicals. How do they trap radicals? Both vitamin E and BHT use a hydroxy group bonded to a benzene ring—a general structure called a **phenol**.

Radicals ($R\cdot$) abstract a hydrogen atom from the OH group of an antioxidant, forming a new resonance-stabilized radical. **This new radical does not participate in chain propagation**, but rather terminates the chain and halts the oxidation process. All phenols (including vitamin E and BHT) inhibit oxidation by this radical process.



Hazelnuts, almonds, and many other types of nuts are an excellent source of the natural antioxidant vitamin E.




The many nonpolar C–C and C–H bonds of vitamin E make it fat soluble, and thus it dissolves in the nonpolar interior of the cell membrane, where it is thought to inhibit the oxidation of the unsaturated fatty acid residues in the phospholipids. Oxidative damage to lipids in cells via radical mechanisms is thought to play an important role in the aging process. For this reason, many anti-aging formulas with antioxidants like vitamin E are now popular consumer products.

Problem 15.25 Draw all resonance structures for the radical that results from hydrogen atom abstraction from BHT.

15.13 Radical Addition Reactions to Double Bonds


We now turn our attention to the second common reaction of radicals, addition to double bonds. Because an alkene contains an electron-rich, easily broken π bond, it reacts with an electron-deficient radical.




Radicals react with alkenes via a radical chain mechanism that consists of initiation, propagation, and termination steps analogous to those discussed previously for radical substitution.

15.13A Addition of HBr

HBr adds to alkenes to form alkyl bromides in the presence of light, heat, or peroxides.



The regioselectivity of addition to an unsymmetrical alkene is *different* from the addition of HBr without added light, heat, or peroxides.



- HBr addition to propene *without* added light, heat, or peroxides gives 2-bromopropane: the **H atom is added to the less substituted carbon**. This reaction occurs via **carbocation** intermediates (Section 10.10).
- HBr addition to propene *with* added light, heat, or peroxides gives 1-bromopropane: the **Br atom is added to the less substituted carbon**. This reaction occurs via **radical** intermediates.

Problem 15.26 Draw the product(s) formed when each alkene is treated with either [1] HBr alone; or [2] HBr in the presence of peroxides.

- a. $\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ b. c. $\text{CH}_3\text{CH}=\text{CHCH}_2\text{CH}_2\text{CH}_3$

15.13B The Mechanism of the Radical Addition of HBr to an Alkene

In the presence of added light, heat, or peroxides, HBr addition to an alkene forms radical intermediates, and like other radical reactions, proceeds by a mechanism with three distinct parts: initiation, propagation, and termination. Mechanism 15.3 is written for the reaction of $\text{CH}_3\text{CH}=\text{CH}_2$ with HBr and ROOR to form $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$.


The first propagation step (Step [3] of the mechanism, the addition of $\text{Br}\cdot$ to the double bond) is worthy of note. With propene there are two possible paths for this step, depending on which



Mechanism 15.3 Radical Addition of HBr to an Alkene

Initiation


Steps [1] and [2] Abstraction of H from HBr occurs by a two-step process.




- With ROOR to initiate the reaction, two steps are needed to form Br \cdot . Homolysis of the weak O–O bond of the peroxide forms RO \cdot , which abstracts a hydrogen atom from HBr to form Br \cdot .

Propagation

Steps [3] and [4] The π bond is broken and the C–H and C–Br σ bonds are formed.



- The first step of propagation forms the C–Br bond when the Br \cdot radical adds to the terminal carbon, leading to a 2 $^\circ$ carbon radical.




- The 2 $^\circ$ radical abstracts a H atom from HBr, forming the new C–H bond and completing the addition reaction. Because a new Br \cdot radical is also formed in this step, Steps [3] and [4] occur repeatedly.

Repeat Steps [3], [4], [3], [4], and so forth.


Termination

Step [5] Two radicals react to form a bond.




- To terminate the chain, two radicals (for example two Br \cdot radicals) react with each other to form a stable bond, preventing further propagation via Steps [3] and [4].

carbon atom of the double bond forms the new bond to bromine. Path [A] forms a less stable 1 $^\circ$ radical whereas Path [B] forms a more stable 2 $^\circ$ radical. **The more stable 2 $^\circ$ radical forms faster**, so Path [B] is preferred.



The mechanism also illustrates why the regioselectivity of HBr addition is different depending on the reaction conditions. In both reactions, H and Br add to the double bond, but the *order* of addition depends on the mechanism.



- In radical addition (HBr with added light, heat, or ROOR), $Br\cdot$ adds first to generate the more stable radical.
- In ionic addition (HBr alone), H^+ adds first to generate the more stable carbocation.

Problem 15.27 When HBr adds to $(CH_3)_2C=CH_2$ under radical conditions, two radicals are possible products in the first step of chain propagation. Draw the structure of both radicals and indicate which one is formed. Then draw the preferred product from HBr addition under radical conditions.

Problem 15.28 What reagents are needed to convert 1-ethylcyclohexene into (a) 1-bromo-2-ethylcyclohexane; (b) 1-bromo-1-ethylcyclohexane; (c) 1,2-dibromo-1-ethylcyclohexane?

15.13C Energy Changes in the Radical Addition of HBr

The energy changes during propagation in the radical addition of HBr to $CH_2=CH_2$ can be calculated from bond dissociation energies, as shown in Figure 15.9.

Both propagation steps for the addition of HBr are exothermic, so propagation is exothermic (energetically favorable) overall. For the addition of HCl or HI, however, one of the chain-propagating steps is quite endothermic, and thus too difficult to be part of a repeating chain mechanism. Thus, **HBr adds to alkenes under radical conditions, but HCl and HI do not.**

Problem 15.29 Draw an energy diagram for the two propagation steps in the radical addition of HBr to propene. Draw the transition state for each step.

15.14 Polymers and Polymerization

Polymers—large molecules made up of repeating units of smaller molecules called **monomers**—include such biologically important compounds as proteins and carbohydrates. They also include such industrially important plastics as polyethylene, poly(vinyl chloride) (PVC), and polystyrene.


15.14A Synthetic Polymers

Many synthetic polymers—that is, those synthesized in the lab—are among the most widely used organic compounds in modern society. Although some synthetic polymers resemble natural substances, many have different and unusual properties that make them more useful than naturally occurring materials. Soft drink bottles, plastic bags, food wrap, compact discs, Teflon, and Styrofoam are all made of synthetic polymers. In this section we examine polymers derived from alkene monomers. Chapter 30 is devoted to a detailed discussion of the synthesis and properties of several different types of synthetic polymers.

HDPE (high-density polyethylene) and **LDPE** (low-density polyethylene) are two common types of polyethylene prepared under different reaction conditions and having different physical properties. HDPE is opaque and rigid, and is used in milk containers and water jugs. LDPE is less opaque and more flexible, and is used in plastic bags and electrical insulation. Products containing HDPE and LDPE (and other plastics) are often labeled with a symbol indicating recycling ease: the lower the number, the easier to recycle.




Figure 15.9
Energy changes during
the propagation steps:
 $\text{CH}_2=\text{CH}_2 + \text{HBr} \rightarrow \text{CH}_3\text{CH}_2\text{Br}$




- **Polymerization** is the joining together of monomers to make polymers.

For example, joining **ethylene monomers** together forms the polymer **polyethylene**, a plastic used in milk containers and sandwich bags.



Many ethylene derivatives having the general structure $\text{CH}_2=\text{CHZ}$ are also used as monomers for polymerization. The identity of Z affects the physical properties of the resulting polymer, making some polymers more suitable for one consumer product (e.g., plastic bags or food wrap) than another (e.g., soft drink bottles or compact discs). Polymerization of $\text{CH}_2=\text{CHZ}$ usually affords polymers with the Z groups on every other carbon atom in the chain. Table 15.2 lists some common monomers and polymers used in medicine or dentistry.



Sample Problem 15.5

What polymer is formed when $\text{CH}_2=\text{CHCO}_2\text{H}$ (acrylic acid) is polymerized? The resulting polymer, poly(acrylic acid), is used in disposable diapers because it absorbs 30 times its weight in water.

Solution

Draw three or more alkene monomers, break one bond of each double bond, and join the alkenes together with single bonds. With unsymmetrical alkenes, substituents are bonded to every other carbon.











Table 15.2 Common Monomers and Polymers Used in Medicine and Dentistry

Monomer	→	Polymer	Consumer product
$\text{CH}_2=\text{CHCl}$ vinyl chloride	→	 poly(vinyl chloride) PVC	 PVC blood bags and tubing
$\text{CH}_2=\text{CHCH}_3$ propene	→	 polypropylene	 polypropylene syringes
$\text{CF}_2=\text{CF}_2$ tetrafluoroethylene	→	 polytetrafluoroethylene Teflon	 dental floss

Problem 15.30

(a) Draw the structure of polystyrene, the chapter-opening molecule, which is formed by polymerizing the monomer styrene, $\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$. (b) What monomer is used to form poly(vinyl acetate), a polymer used in paints and adhesives?

**15.14B Radical Polymerization**

The alkene monomers used in polymerization are prepared from petroleum.


The polymers described in Section 15.14A are prepared by polymerization of alkene monomers by **adding a radical to a π bond**. The mechanism resembles the radical addition of HBr to an alkene, except that a **carbon radical rather than a bromine atom is added to the double bond**. Mechanism 15.4 is written with the general monomer $\text{CH}_2=\text{CHZ}$, and again has three parts: initiation, propagation, and termination.



Mechanism 15.4 Radical Polymerization of $\text{CH}_2=\text{CHZ}$

Initiation


Steps [1] and [2] A carbon radical is formed by a two-step process.



- Chain initiation begins with homolysis of the weak O–O bond of the peroxide to form $\text{RO}\cdot$, which then adds to a molecule of monomer to form a carbon radical.

Propagation


Step [3] The polymer chain grows.



- Chain propagation consists of a single step that joins monomer units together.
- In Step [3], the carbon radical formed during initiation adds to another alkene molecule to form a new C–C bond and another carbon radical. Addition always forms the more substituted carbon radical—that is, the **unpaired electron is always located on the carbon atom having the Z substituent**.
- This carbon radical reacts with more monomer, so that Step [3] occurs repeatedly, and the polymer chain grows.

Termination


Step [4] Two radicals combine to form a bond.



- To terminate the chain, two radicals combine to form a stable bond, thus ending the polymerization process.

In radical polymerization, the more substituted radical always adds to the less substituted end of the monomer, a process called **head-to-tail polymerization**.

The **more substituted radical** adds to the **less substituted end** of the double bond.



Problem 15.31 Draw the steps of the mechanism that converts vinyl chloride ($\text{CH}_2=\text{CHCl}$) into poly(vinyl chloride).

KEY CONCEPTS


Radical Reactions

General Features of Radicals

- A radical is a reactive intermediate with a single unpaired electron (15.1).
- A carbon radical is sp^2 hybridized and trigonal planar (15.1).
- The stability of a radical increases as the number of C atoms bonded to the radical carbon increases (15.1).
- Allylic radicals are stabilized by resonance, making them more stable than 3° radicals (15.10).


Radical Reactions

[1] Halogenation of alkanes (15.4)




- The reaction follows a radical chain mechanism.
- The weaker the C–H bond, the more readily the hydrogen is replaced by X.
- Chlorination is faster and less selective than bromination (15.6).
- Radical substitution at a stereogenic center results in racemization (15.8).

[2] Allylic halogenation (15.10)




- The reaction follows a radical chain mechanism.

[3] Radical addition of HBr to an alkene (15.13)



- A radical addition mechanism is followed.
- Br bonds to the less substituted carbon atom to form the more substituted, more stable radical.

[4] Radical polymerization of alkenes (15.14)




- A radical addition mechanism is followed.

PROBLEMS


Radicals and Bond Strength

15.32 With reference to the indicated C–H bonds in 2-methylbutane:




- Rank the C–H bonds in order of increasing bond strength.
- Draw the radical resulting from cleavage of each C–H bond, and classify it as 1°, 2°, or 3°.
- Rank the radicals in order of increasing stability.
- Rank the C–H bonds in order of increasing ease of H abstraction in a radical halogenation reaction.

15.33 Rank each group of radicals in order of increasing stability.




15.34 Why is a benzylic C–H bond unusually weak?






Halogenation of Alkanes




15.35 Rank the indicated hydrogen atoms in order of increasing ease of abstraction in a radical halogenation reaction.



15.36 Draw all constitutional isomers formed by monochlorination of each alkane with Cl_2 and $h\nu$.







15.37 What is the major monobromination product formed by heating each alkane with Br_2 ?







15.38 Five isomeric alkanes (**A–E**) having the molecular formula C_6H_{14} are each treated with $\text{Cl}_2 + h\nu$ to give alkyl halides having molecular formula $\text{C}_6\text{H}_{13}\text{Cl}$. **A** yields five constitutional isomers. **B** yields four constitutional isomers. **C** yields two constitutional isomers. **D** yields three constitutional isomers, two of which possess stereogenic centers. **E** yields three constitutional isomers, only one of which possesses a stereogenic center. Identify the structures of **A–E**.

15.39 What alkane is needed to make each alkyl halide by radical halogenation?




15.40 Which alkyl halides can be prepared in good yield by radical halogenation of an alkane?





15.41 Explain why chlorination of cyclohexane with two equivalents of Cl_2 in the presence of light is a poor method to prepare 1,2-dichlorocyclohexane.

15.42 Explain why radical bromination of *p*-xylene forms **C** rather than **D**.






- 15.43 a. What product(s) (excluding stereoisomers) are formed when **Y** is heated with Cl_2 ?
 b. What product(s) (excluding stereoisomers) are formed when **Y** is heated with Br_2 ?
 c. What steps are needed to convert **Y** to the alkene **Z**?






Resonance

15.44 Draw resonance structures for each radical.




Allylic Halogenation

15.45 Draw the products formed when each alkene is treated with NBS + $h\nu$.












15.46 Is it possible to prepare 5-bromo-1-methylcyclopentene in good yield by allylic bromination of 1-methylcyclopentene? Explain.

15.47 Draw all constitutional isomers formed when **X** is treated with NBS + $h\nu$.





Reactions

15.48 Draw the organic products formed in each reaction.









15.49 What reagents are needed to convert cyclopentene into (a) bromocyclopentane; (b) *trans*-1,2-dibromocyclopentane; (c) 3-bromocyclopentene?

15.50 Treatment of a hydrocarbon **A** (molecular formula C_9H_{18}) with Br_2 in the presence of light forms alkyl halides **B** and **C**, both having molecular formula $C_9H_{17}Br$. Reaction of either **B** or **C** with $KOC(CH_3)_3$ forms compound **D** (C_9H_{16}) as the major product. Ozonolysis of **D** forms cyclohexanone and acetone. Identify the structures of **A–D**.



Stereochemistry and Reactions


15.51 Draw the products formed in each reaction and include the stereochemistry around any stereogenic centers.



15.52 (a) Draw all stereoisomers of molecular formula $C_5H_{10}Cl_2$ formed when (2*R*)-2-chloropentane is heated with Cl_2 . (b) Assuming that products having different physical properties can be separated into fractions by some physical method (such as fractional distillation), how many different fractions would be obtained? (c) Which of these fractions would be optically active?

15.53 (a) Draw all stereoisomers of molecular formula $C_7H_{15}Cl$ formed when (3*S*)-3-methylhexane is heated with Cl_2 . (b) Assuming that products having different physical properties can be separated by fractional distillation, how many different fractions would be obtained? (c) How many fractions would be optically active?

15.54 Draw the six products (including stereoisomers) formed when **A** is treated with NBS + $h\nu$.




Mechanisms

15.55 Consider the following bromination: $(\text{CH}_3)_3\text{CH} + \text{Br}_2 \xrightarrow{\Delta} (\text{CH}_3)_3\text{CBr} + \text{HBr}$.


- Calculate ΔH° for this reaction by using the bond dissociation energies in Table 6.2.
- Draw out a stepwise mechanism for the reaction, including the initiation, propagation, and termination steps.
- Calculate ΔH° for each propagation step.
- Draw an energy diagram for the propagation steps.
- Draw the structure of the transition state of each propagation step.

15.56 Draw a stepwise mechanism for the following reaction.



15.57 Although CH_4 reacts with Cl_2 to form CH_3Cl and HCl , the corresponding reaction of CH_4 with I_2 does not occur at an appreciable rate, even though the $\text{I}-\text{I}$ bond is much weaker than the $\text{Cl}-\text{Cl}$ bond. Explain why this is so.

15.58 An alternative mechanism for the propagation steps in the radical chlorination of CH_4 is drawn below. Calculate ΔH° for these steps and explain why this pathway is unlikely.




15.59 When 3,3-dimethyl-1-butene is treated with HBr alone, the major product is 2-bromo-2,3-dimethylbutane. When the same alkene is treated with HBr and peroxide, the sole product is 1-bromo-3,3-dimethylbutane. Explain these results by referring to the mechanisms.

15.60 Write a stepwise mechanism that shows how a very small amount of $\text{CH}_3\text{CH}_2\text{Cl}$ could form during the chlorination of CH_4 .


15.61 Write out the two propagation steps for the addition of HCl to propene and calculate ΔH° for each step. Which step prohibits chain propagation from repeatedly occurring?

Synthesis


15.62 Devise a synthesis of each compound from cyclopentane and any other required organic or inorganic reagents.




15.63 Devise a synthesis of each target compound from methylcyclohexane. You may use any other required organic or inorganic reagents.



15.64 Devise a synthesis of each target compound from the indicated starting material. You may use any other required organic or inorganic reagents.




15.65 Devise a synthesis of each compound using CH_3CH_3 as the only source of carbon atoms. You may use any other required organic or inorganic reagents.




15.66 Devise a synthesis of $\text{OHC}(\text{CH}_2)_4\text{CHO}$ from cyclohexane using any required organic or inorganic reagents.

Radical Oxidation Reactions

- 15.67** As described in Section 9.16, the leukotrienes, important components in the asthmatic response, are synthesized from arachidonic acid via the hydroperoxide 5-HPETE. Write a stepwise mechanism for the conversion of arachidonic acid to 5-HPETE with O_2 .




- 15.68** Ethers are oxidized with O_2 to form hydroperoxides that decompose violently when heated. Draw a stepwise mechanism for this reaction.




- 15.69** (a) Ignoring stereoisomers, what two allylic hydroperoxides are formed by the oxidation of 1-hexene with O_2 ? (b) Draw a stepwise mechanism that shows how these hydroperoxides are formed.

Antioxidants

- 15.70** Draw all resonance structures of the radical resulting from abstraction of a hydrogen atom from the antioxidant BHA (butylated hydroxy anisole).




- 15.71** In cells, vitamin C exists largely as its conjugate base **X**. **X** is an antioxidant because radicals formed in oxidation processes abstract the indicated H atom, forming a new radical that halts oxidation. Draw the structure of the radical formed by H abstraction, and explain why this H atom is most easily removed.




Polymers and Polymerization

- 15.72** What monomer is needed to form each polymer?




- 15.73** (a) Hard contact lenses, which first became popular in the 1960s, were made by polymerizing methyl methacrylate [$CH_2=C(CH_3)CO_2CH_3$] to form poly(methyl methacrylate) (PMMA). Draw the structure of PMMA. (b) More comfortable softer contact lenses introduced in the 1970s were made by polymerizing hydroxyethyl methacrylate [$CH_2=C(CH_3)CO_2CH_2CH_2OH$] to form poly(hydroxyethyl methacrylate) (poly-HEMA). Draw the structure of poly-HEMA. Since neither polymer allows oxygen from the air to pass through to the retina, newer contact lenses that are both comfortable and oxygen-permeable have now been developed.

15.74 Draw a stepwise mechanism for the following polymerization reaction.



15.75 As we will learn in Section 30.2C, styrene derivatives such as **A** can be polymerized by way of cationic rather than radical intermediates. Cationic polymerization is an example of electrophilic addition to an alkene involving carbocations.




- Draw a short segment of the polymer formed by the polymerization of **A**.
- Why does **A** react faster than styrene ($\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$) in a cationic polymerization?

Spectroscopy

15.76 **A** and **B**, isomers of molecular formula $\text{C}_3\text{H}_5\text{Cl}_3$, are formed by the radical chlorination of a dihalide **C** of molecular formula $\text{C}_3\text{H}_6\text{Cl}_2$.

- Identify the structures of **A** and **B** from the following ^1H NMR data:
 Compound **A**: singlet at 2.23 and singlet at 4.04 ppm
 Compound **B**: doublet at 1.69, multiplet at 4.34, and doublet at 5.85 ppm
- What is the structure of **C**?

15.77 Identify the structure of a minor product formed from the radical chlorination of propane, which has molecular formula $\text{C}_3\text{H}_6\text{Cl}_2$ and exhibits the given ^1H NMR spectrum.




15.78 Radical chlorination of CH_3CH_3 forms two minor products **X** and **Y** of molecular formula $\text{C}_2\text{H}_4\text{Cl}_2$.


- Identify the structures of **X** and **Y** from the following ^1H NMR data:
 Compound **X**: singlet at 3.7 ppm
 Compound **Y**: doublet at 2.1 and quartet at 5.9 ppm
- Draw a stepwise mechanism that shows how each product is formed from CH_3CH_3 .

Challenge Problems


15.79 Draw a stepwise mechanism for the following addition reaction to an alkene.




- 15.80** In the presence of a radical initiator ($Z\cdot$), tributyltin hydride (R_3SnH , $R = CH_3CH_2CH_2CH_2$) reduces alkyl halides to alkanes: $R'X + R_3SnH \rightarrow R'H + R_3SnX$. The mechanism consists of a radical chain process with an intermediate tin radical:



This reaction has been employed in many radical cyclization reactions. Draw a stepwise mechanism for the following reaction.




- 15.81** $PGF_{2\alpha}$ (Sections 4.15 and 29.6) is synthesized in cells from arachidonic acid ($C_{20}H_{32}O_2$) using a cyclooxygenase enzyme that catalyzes a multistep radical pathway. Part of this process involves the conversion of radical **A** to PGG_2 , an unstable intermediate, which is then transformed to $PGF_{2\alpha}$ and other prostaglandins. Draw a stepwise mechanism for the conversion of **A** to PGG_2 . (Hint: The mechanism begins with radical addition to a carbon-carbon double bond to form a resonance-stabilized radical.)



Conjugation, Resonance, and Dienes

16



- 16.1 Conjugation
- 16.2 Resonance and allylic carbocations
- 16.3 Common examples of resonance
- 16.4 The resonance hybrid
- 16.5 Electron delocalization, hybridization, and geometry
- 16.6 Conjugated dienes
- 16.7 Interesting dienes and polyenes
- 16.8 The carbon–carbon σ bond length in 1,3-butadiene
- 16.9 Stability of conjugated dienes
- 16.10 Electrophilic addition: 1,2- versus 1,4-addition
- 16.11 Kinetic versus thermodynamic products
- 16.12 The Diels–Alder reaction
- 16.13 Specific rules governing the Diels–Alder reaction
- 16.14 Other facts about the Diels–Alder reaction
- 16.15 Conjugated dienes and ultraviolet light



Lycopene is a red pigment found in tomatoes, watermelon, papaya, guava, and pink grapefruit. An antioxidant like vitamin E, lycopene contains many conjugated double bonds—double bonds separated by only one single bond—that allow π electron density to delocalize and give the molecule added stability. In Chapter 16 we learn about such conjugated unsaturated systems.


Chapter 16 is the first of three chapters that discuss the chemistry of conjugated molecules—molecules with overlapping p orbitals on three or more adjacent atoms. Chapter 16 focuses mainly on acyclic conjugated compounds, whereas Chapters 17 and 18 discuss the chemistry of benzene and related compounds that have a p orbital on every atom in a ring.

Much of Chapter 16 is devoted to the properties and reactions of 1,3-dienes. To understand these compounds, however, we must first learn about the consequences of having p orbitals on three or more adjacent atoms. Because the ability to draw resonance structures is also central to mastering this material, the key aspects of resonance theory are presented in detail.

16.1 Conjugation


The word *conjugation* is derived from the Latin *conjugatus*, meaning “to join.”

Conjugation occurs whenever p orbitals can overlap on three or more adjacent atoms. Two common conjugated systems are 1,3-dienes and allylic carbocations.




16.1A 1,3-Dienes

1,3-Dienes such as 1,3-butadiene contain two carbon–carbon double bonds joined by a single σ bond. Each carbon atom of a 1,3-diene is bonded to three other atoms and has no nonbonded electron pairs, so each carbon atom is sp^2 hybridized and has one p orbital containing an electron. **The four p orbitals on adjacent atoms make a 1,3-diene a conjugated system.**




What is special about conjugation? Having three or more p orbitals on adjacent atoms allows p orbitals to overlap and electrons to delocalize.




- When p orbitals overlap, the electron density in each of the π bonds is spread out over a larger volume, thus lowering the energy of the molecule and making it more stable.

Conjugation makes 1,3-butadiene inherently different from 1,4-pentadiene, a compound having two double bonds separated by more than one σ bond. The π bonds in 1,4-pentadiene are too far apart to be conjugated.




1,4-Pentadiene is an **isolated diene**. The electron density in each π bond of an isolated diene is *localized* between two carbon atoms. In 1,3-butadiene, however, the electron density of both π

Figure 16.1
Electrostatic potential plots
for a conjugated and an
isolated diene




bonds is *delocalized* over the four atoms of the diene. Electrostatic potential maps in Figure 16.1 clearly indicate the difference between these localized and delocalized π bonds.

Problem 16.1 Classify each diene as isolated or conjugated.




Problem 16.2 Label each double bond in 5-HPETE (5-hydroperoxyeicosatetraenoic acid) as isolated or conjugated. 5-HPETE is an intermediate in the biological conversion of arachidonic acid to leukotrienes, potent molecules that contribute to the asthmatic response (Section 9.16).




16.1B Allylic Carbocations

The **allyl carbocation** is another example of a conjugated system. The three carbon atoms of the allyl carbocation—the positively charged carbon atom and the two that form the double bond—are sp^2 hybridized with a p orbital. The p orbitals for the double bond carbons each contain an electron, whereas the p orbital for the carbocation is empty.




- Three p orbitals on three adjacent atoms, even if one of the p orbitals is empty, make the allyl carbocation conjugated.

Conjugation stabilizes the allyl carbocation because overlap of three adjacent p orbitals delocalizes the electron density of the π bond over three atoms.



Problem 16.3 Which of the following species are conjugated?




16.2 Resonance and Allylic Carbocations

The word *resonance* is used in two different contexts. In NMR spectroscopy, a nucleus is *in resonance* when it absorbs energy, promoting it to a higher energy state. In drawing molecules, there is *resonance* when two different Lewis structures can be drawn for the same arrangement of atoms.


Recall from Section 1.5 that resonance structures are two or more different Lewis structures for the same arrangement of atoms. Being able to draw correct resonance structures is crucial to understanding conjugation and the reactions of conjugated dienes.

- Two resonance structures differ in the placement of π bonds and nonbonded electrons. The placement of atoms and σ bonds stays the same.

We have already drawn resonance structures for the acetate anion (Section 2.5C) and the allyl radical (Section 15.10). The **conjugated allyl carbocation** is another example of a species for which two resonance structures can be drawn. Drawing resonance structures for the allyl carbocation is a way to use Lewis structures to illustrate how conjugation delocalizes electrons.




The true structure of the allyl carbocation is a hybrid of the two resonance structures. In the hybrid, the π bond is delocalized over all three atoms. As a result, the positive charge is also delocalized over the two terminal carbons. Delocalizing electron density lowers the energy of the hybrid, thus stabilizing the allyl carbocation and making it more stable than a normal 1° carbocation. Experimental data show that its stability is comparable to a more highly substituted 2° carbocation.




The electrostatic potential maps in Figure 16.2 compare the resonance-stabilized allyl carbocation with $\text{CH}_3\text{CH}_2\text{CH}_2^+$, a localized 1° carbocation. The electron-deficient region—the site of the positive charge—is concentrated on a single carbon atom in the 1° carbocation $\text{CH}_3\text{CH}_2\text{CH}_2^+$. In the allyl carbocation, however, the electron-poor region is spread out on both terminal carbons.

Figure 16.2
Electrostatic potential maps for a localized and a delocalized carbocation



Problem 16.4 Draw a second resonance structure for each carbocation. Then draw the hybrid.



How “real” is the delocalization of charge and electron density in the allyl carbocation? Recall from the discussion of NMR spectroscopy in Chapter 14 that an NMR absorption shifts downfield (to higher chemical shift) as the electron density around the nucleus decreases (is deshielded). Thus, a positively charged carbocation, such as **A** in Figure 16.3, is highly deshielded, so its ^{13}C NMR absorption is far downfield at 330 ppm. A resonance-stabilized carbocation, however, such as **B** in Figure 16.3, has less positive charge concentrated on any given carbon atom (because the charge is delocalized), so its ^{13}C NMR absorption is farther upfield at 224 ppm.

Problem 16.5 How many ^{13}C NMR signals are predicted for carbocations **A** and **B** in Figure 16.3?


Problem 16.6 Use resonance theory and the Hammond postulate to explain why 3-chloro-1-propene ($\text{CH}_2=\text{CHCH}_2\text{Cl}$) is more reactive than 1-chloropropane ($\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$) in $\text{S}_{\text{N}}1$ reactions.

16.3 Common Examples of Resonance


When are resonance structures drawn for a molecule or reactive intermediate? Because resonance involves delocalizing π bonds and nonbonded electrons, one or both of these structural features must be present to draw additional resonance forms. There are four common bonding patterns for which more than one Lewis structure can be drawn.

Type [1] The Three Atom “Allyl” System, $\text{X}=\text{Y}-\text{Z}^*$

- For any group of three atoms having a double bond $\text{X}=\text{Y}$ and an atom Z that contains a p orbital with zero, one, or two electrons, two resonance structures are possible:




The asterisk [*] corresponds to a charge, a radical, or a lone pair.



This is called **allyl** type resonance because it can be drawn for allylic carbocations, allylic carbanions, and allylic radicals.


X , Y , and Z may all be carbon atoms, as in the case of an allylic carbocation (resonance structures **A** and **B**), or they may be heteroatoms, as in the case of the acetate anion (resonance structures **C** and **D**). The atom Z bonded to the multiple bond can be charged (a net positive or negative charge) or neutral (having zero, one, or two nonbonded electrons). **The two resonance**

Figure 16.3
 ^{13}C chemical shifts for a localized and a resonance-stabilized carbocation




- The absorption shifts *upfield* as the amount of positive charge *decreases*.

structures differ in the location of the double bond, and either the charge, the radical, or the lone pair, generalized by [*].




Type [2] Conjugated Double Bonds

Cyclic, completely conjugated rings like benzene have two resonance structures, drawn by moving the electrons in a cyclic manner around the ring. Three resonance structures can be drawn for conjugated dienes, two of which involve charge separation.



Type [3] Cations Having a Positive Charge Adjacent to a Lone Pair


- When a lone pair and a positive charge are located on adjacent atoms, two resonance structures can be drawn.



The overall charge is the same in both resonance structures. Based on formal charge, a neutral X in one structure must bear a (+) charge in the other.

Type [4] Double Bonds Having One Atom More Electronegative Than the Other


- For a double bond $\text{X}=\text{Y}$ in which the electronegativity of $\text{Y} > \text{X}$, a second resonance structure can be drawn by moving the π electrons onto Y.



Sample Problem 16.1 illustrates how to apply these different types of resonance to actual molecules.

Sample Problem 16.1


Draw two more resonance structures for each species.




Solution

Mentally breaking a molecule into two- or three-atom units can make it easier to draw additional resonance structures.

- a. Think of the top three atoms of the six-membered ring in **A** as an “allyl” unit. Moving the π bond forms a new “allyl” unit in **B**, and moving the π bond in **B** generates a third resonance structure **C**. No new valid resonance structures are generated by moving electrons in **C**.




- b. Compound **D** contains a carbonyl group, so moving the electron pair in the double bond to the more electronegative oxygen atom separates the charge and generates structure **E**. **E** now has a three-atom “allyl” unit, so the remaining π bond can be moved to form structure **F**.



Problem 16.7

Draw additional resonance structures for each ion.




16.4 The Resonance Hybrid

The lower its energy, the more a resonance structure contributes to the overall structure of the hybrid.


Although the resonance hybrid is some combination of all of its valid resonance structures, the **hybrid more closely resembles the most stable resonance structure**. Recall from Section 1.5C that the most stable resonance structure is called the **major contributor** to the hybrid, and the less stable resonance structures are called the **minor contributors**. Two identical resonance structures are equal contributors to the hybrid.

Use the following three rules to evaluate the relative stabilities of two or more valid resonance structures.


Rule [1] Resonance structures with more bonds and fewer charges are more stable.



Rule [2] Resonance structures in which every atom has an octet are more stable.




Rule [3] Resonance structures that place a negative charge on a more electronegative atom are more stable.



Sample Problem 16.2 illustrates how to determine the relative energy of contributing resonance structures and the hybrid.


Sample Problem 16.2

Draw a second resonance structure for carbocation **A**, as well as the hybrid of both resonance structures. Then use Rules [1]–[3] to rank the relative stability of both resonance structures and the hybrid.



Solution


Because **A** contains a positive charge and a lone pair on adjacent atoms, a second resonance structure **B** can be drawn. Because **B** has more bonds and all second-row atoms have octets, **B** is more stable than **A**, making it the major contributor to the hybrid **C**. Because the hybrid is more stable than either resonance contributor, the order of stability is:



Problem 16.8 Draw a second resonance structure and the hybrid for each species, and then rank the two resonance structures and the hybrid in order of increasing stability.



Problem 16.9 Draw all possible resonance structures for the following cation and indicate which structure makes the largest contribution to the resonance hybrid.




Problem 16.10 For acetic acid ($\text{CH}_3\text{CO}_2\text{H}$): (a) Draw three resonance structures; (b) draw a structure for the resonance hybrid; (c) rank the three resonance structures and the resonance hybrid in order of increasing energy.

16.5 Electron Delocalization, Hybridization, and Geometry


To delocalize nonbonded electrons or electrons in π bonds, there must be p orbitals that can overlap. This may mean that the hybridization of an atom is different than would have been predicted using the rules first outlined in Chapter 1.

For example, there are two Lewis structures (**A** and **B**) for the resonance-stabilized anion ($\text{CH}_3\text{COCH}_2^-$).



Based on structure **A**, the indicated carbon is sp^3 hybridized, with the lone pair of electrons in an sp^3 hybrid orbital. Based on structure **B**, though, it is sp^2 hybridized with the unhybridized p orbital forming the π portion of the double bond.


Delocalizing electrons stabilizes a molecule. The electron pair on the carbon atom adjacent to the $\text{C}=\text{O}$ can only be delocalized, though, if it has a p orbital that can overlap with two other p orbitals on two adjacent atoms. Thus, the terminal carbon atom is sp^2 hybridized with trigonal planar geometry. **Three adjacent p orbitals make the anion conjugated.**



- In any system $\text{X}=\text{Y}-\text{Z}^-$, Z is sp^2 hybridized, and the nonbonded electron pair occupies a p orbital to make the system conjugated.


Sample Problem 16.3

Determine the hybridization around the indicated carbon atom in the following anion.




Solution

Because this is an example of an allyl-type system ($\text{X}=\text{Y}-\text{Z}^-$), a second resonance structure can be drawn that “moves” the lone pair and the π bond. To delocalize the lone pair and make the system conjugated, the indicated carbon atom must be sp^2 hybridized with the lone pair occupying a p orbital.



Problem 16.11

Determine the hybridization of the indicated atom in each species.




16.6 Conjugated Dienes


Compounds with many π bonds are called **polyenes**.

In the remainder of Chapter 16 we examine **conjugated dienes**, compounds having two double bonds joined by one σ bond. Conjugated dienes are also called **1,3-dienes**. 1,3-Butadiene ($\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$) is the simplest conjugated diene.

Three stereoisomers are possible for 1,3-dienes with alkyl groups bonded to each end carbon of the diene ($\text{RCH}=\text{CH}-\text{CH}=\text{CHR}$).




Two possible conformations result from rotation around the C–C bond that joins the two double bonds.



- The *s-cis* conformation has two double bonds on the *same* side of the single bond.
- The *s-trans* conformation has two double bonds on *opposite* sides of the single bond.


Keep in mind that stereoisomers are discrete molecules, whereas conformations interconvert. Three structures drawn for 2,4-hexadiene illustrate the differences between stereoisomers and conformations in a 1,3-diene:



Problem 16.12 Draw the structure consistent with each description.

- (*2E,4E*)-2,4-octadiene in the *s-trans* conformation
- (*3E,5Z*)-3,5-nonadiene in the *s-cis* conformation
- (*3Z,5Z*)-4,5-dimethyl-3,5-decadiene. Draw both the *s-cis* and *s-trans* conformations.


Problem 16.13 Neuroprotectin D1 (NPD1) is synthesized in the body from highly unsaturated essential fatty acids. NPD1 is a potent natural anti-inflammatory agent.



- Label each double bond as conjugated or isolated.
- Label each double bond as *E* or *Z*.
- For each conjugated system, label the given conformation as *s-cis* or *s-trans*.



Figure 16.4

Biologically active organic compounds that contain conjugated double bonds



16.7 Interesting Dienes and Polyenes

Isoprene and **lycopene** are two naturally occurring compounds with conjugated double bonds.



The Blue Ridge Mountains

Isoprene, the common name for 2-methyl-1,3-butadiene, is given off by plants as the temperature rises, a process thought to increase a plant's tolerance for heat stress. Isoprene is a component of the blue haze seen above forested hillsides, such as Virginia's Blue Ridge Mountains.

Unlike most of the organic compounds encountered up to this point, **lycopene**, the chapter-opening molecule, is colored. The 11 conjugated double bonds of lycopene cause its red color, a phenomenon discussed in Section 16.15.


Simvastatin and calcitriol are two drugs that contain conjugated double bonds in addition to other functional groups (Figure 16.4). Simvastatin is the generic name of the widely used cholesterol-lowering medicine Zocor. Calcitriol, a biologically active hormone formed from vitamin D₃ obtained in the diet, is responsible for regulating calcium and phosphorus metabolism. Sold under the trade name of Rocaltrol, calcitriol is used to treat patients who are unable to convert vitamin D₃ to the active hormone. Since calcitriol promotes the absorption of calcium ions, it is also used to treat hypocalcemia, the presence of low calcium levels in the blood.

16.8 The Carbon–Carbon σ Bond Length in 1,3-Butadiene

Four features distinguish conjugated dienes from isolated dienes.


- [1] **The C–C single bond joining the two double bonds is unusually short.**
- [2] **Conjugated dienes are more stable than similar isolated dienes.**
- [3] **Some reactions of conjugated dienes are different than reactions of isolated double bonds.**
- [4] **Conjugated dienes absorb longer wavelengths of ultraviolet light.**

The bond length of the carbon–carbon double bonds in 1,3-butadiene is similar to an isolated double bond, but the central carbon–carbon single bond is shorter than the C–C bond in ethane.



The C–C σ bond is **shorter** than the C–C σ bond in ethane.


The observed bond distances can be explained by looking at hybridization. Each carbon atom in 1,3-butadiene is sp^2 hybridized, so the central C–C single bond is formed by the overlap of two sp^2 hybridized orbitals, rather than the sp^3 hybridized orbitals used to form the C–C bond in CH_3CH_3 .



Recall from Section 1.10B that increasing percent s-character decreases bond length.

- Based on hybridization, a $\text{C}_{sp^2}-\text{C}_{sp^2}$ bond should be shorter than a $\text{C}_{sp^3}-\text{C}_{sp^3}$ bond because it is formed from orbitals having a higher percent s-character.

A resonance argument can also be used to explain the shorter C–C σ bond length of 1,3-butadiene. 1,3-Butadiene can be represented by three resonance structures:




Structures **B** and **C** have charge separation and fewer bonds than **A**, making them less stable resonance structures and only minor contributors to the resonance hybrid. **B** and **C** both contain a double bond between the central carbon atoms, however, so the hybrid must have a partial double bond there. This makes the central C–C bond shorter than a C–C single bond in an alkane.

- Based on resonance, the central C–C bond in 1,3-butadiene is shorter because it has partial double bond character.


Finally, 1,3-butadiene is a conjugated molecule with four overlapping p orbitals on adjacent atoms. As a result, the π electrons are not localized between the carbon atoms of the double bonds, but rather delocalized over four atoms. This places more electron density between the central two carbon atoms of 1,3-butadiene than would normally be present. This *shortens* the bond. Drawing resonance structures illustrates this delocalization.

The overlap of adjacent p orbitals increases the electron density in the C–C σ bond.




Problem 16.14 Using hybridization, predict how the bond length of the C–C σ bond in $\text{HC}\equiv\text{C}-\text{C}\equiv\text{CH}$ should compare with the C–C σ bonds in CH_3CH_3 and $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$.

Problem 16.15 Use resonance theory to explain why both C–O bond lengths are equal in the acetate anion.



16.9 Stability of Conjugated Dienes


In Section 12.3 we learned that hydrogen adds to alkenes to form alkanes, and that the heat released in this reaction, the **heat of hydrogenation**, can be used as a measure of alkene stability.



The relative stability of conjugated and isolated dienes can also be determined by comparing their heats of hydrogenation.

- When hydrogenation gives the same alkane from two dienes, the more stable diene has the smaller heat of hydrogenation.

For example, both 1,4-pentadiene (an isolated diene) and (3*E*)-1,3-pentadiene (a conjugated diene) are hydrogenated to pentane with two equivalents of H₂. Because less energy is released in converting the conjugated diene to pentane, it must be lower in energy (more stable) to begin with. The relative energies of these isomeric pentadienes are illustrated in Figure 16.5.



- A conjugated diene has a smaller heat of hydrogenation and is more stable than a similar isolated diene.

Why is a conjugated diene more stable than an isolated diene? Because a conjugated diene has overlapping *p* orbitals on four adjacent atoms, its π electrons are delocalized over four atoms. This delocalization, which cannot occur in an isolated diene, is illustrated by drawing resonance structures.

No resonance structures can be drawn for 1,4-pentadiene, but three can be drawn for (3*E*)-1,3-pentadiene (or any other conjugated diene). The hybrid of these resonance structures illustrates that the two adjacent π bonds are delocalized in a conjugated diene, making it lower in energy than an isolated diene.






Figure 16.5


Relative energies of an isolated and conjugated diene



Problem 16.16 Which diene in each pair has the larger heat of hydrogenation?



Problem 16.17 Rank the following compounds in order of increasing stability.




16.10 Electrophilic Addition: 1,2- Versus 1,4-Addition


Recall from Chapters 10 and 11 that the characteristic reaction of compounds with π bonds is **addition**. The π bonds in conjugated dienes undergo addition reactions, too, but they differ in two ways from the addition reactions to isolated double bonds.

- **Electrophilic addition in conjugated dienes gives a mixture of products.**
- **Conjugated dienes undergo a unique addition reaction not seen in alkenes or isolated dienes.**


We learned in Chapter 10 that HX adds to the π bond of alkenes to form alkyl halides.



With an **isolated diene**, electrophilic addition of one equivalent of HBr yields *one* product and Markovnikov's rule is followed. The H atom bonds to the less substituted carbon—that is, the carbon atom of the double bond that had more H atoms to begin with.



With a **conjugated diene**, electrophilic addition of one equivalent of HBr affords *two* products.



The ends of the 1,3-diene are called C1 and C4 arbitrarily, without regard to IUPAC numbering.


- The **1,2-addition product** results from Markovnikov addition of HBr across two adjacent carbon atoms (C1 and C2) of the diene.
- The **1,4-addition product** results from addition of HBr to the two end carbons (C1 and C4) of the diene. 1,4-Addition is also called **conjugate addition**.

The mechanism of electrophilic addition of HX involves **two steps**: addition of H^+ (from HX) to form a resonance-stabilized carbocation, followed by nucleophilic attack of X^- at either electrophilic end of the carbocation to form two products. Mechanism 16.1 illustrates the reaction of 1,3-butadiene with HBr.




Mechanism 16.1 Electrophilic Addition of HBr to a 1,3-Diene—1,2- and 1,4-Addition

Step [1] Addition of the electrophile (H^+) to the π bond



- H^+ (from HBr) always adds to a terminal C of the 1,3-diene to form a resonance-stabilized carbocation. This is the slow step of the mechanism because two bonds are broken and only one is formed.

Step [2] Nucleophilic attack of Br^-




- Nucleophilic attack of Br^-** can occur at either site of the resonance-stabilized carbocation that bears a (+) charge, forming either the 1,2-addition product or the 1,4-addition product.

Like the electrophilic addition of HX to an alkene, the addition of HBr to a conjugated diene forms the more stable carbocation in Step [1], the rate-determining step. In this case, however, the carbocation is both 2° and **allylic**, and thus two Lewis structures can be drawn for it. In the second step, nucleophilic attack of Br^- can then occur at two different electrophilic sites, forming two different products.

- Addition of HX to a conjugated diene forms 1,2- and 1,4-products because of the resonance-stabilized allylic carbocation intermediate.


Sample Problem 16.4

Draw the products of the following reaction.




Solution


Write the steps of the mechanism to determine the structure of the products. Addition of H^+ forms the more stable 2° allylic carbocation, for which two resonance structures can be drawn. Nucleophilic attack of Br^- at either end of the allylic carbocation gives two constitutional isomers, formed by 1,2-addition and 1,4-addition to the diene.



Problem 16.18 Draw the products formed when each diene is treated with one equivalent of HCl.




Problem 16.19 Draw a stepwise mechanism for the following reaction.




16.11 Kinetic Versus Thermodynamic Products

The amount of 1,2- and 1,4-addition products formed in the electrophilic addition reactions of 1,3-butadiene, a conjugated diene, depends greatly on the reaction conditions.



- At low temperature the major product is formed by 1,2-addition.
- At higher temperature the major product is formed by 1,4-addition.

Moreover, when a mixture containing predominately the 1,2-product is heated, the 1,4-addition product becomes the major product at equilibrium.




- The 1,2-product is formed faster because it predominates at low temperature. The product that is formed faster is called the *kinetic product*.
- The 1,4-product must be more stable because it predominates at equilibrium. The product that predominates at equilibrium is called the *thermodynamic product*.

In many of the reactions we have learned thus far, the more stable product is formed faster—that is, the kinetic and thermodynamic products are the same. The electrophilic addition of HBr to 1,3-butadiene is different, in that **the more stable product is formed more slowly**—that is, the kinetic and thermodynamic products are *different*. Why is the more stable product formed more slowly?

To answer this question, recall that the rate of a reaction is determined by its energy of activation (E_a), whereas the amount of product present at equilibrium is determined by its stability (Figure 16.6). When a single starting material **A** forms two different products (**B** and **C**) by two exothermic pathways, the relative height of the energy barriers determines how fast **B** and **C** are formed, whereas the relative energies of **B** and **C** determine the amount of each at equilibrium. In an exothermic reaction, the relative energies of **B** and **C** do not determine the relative energies of activation to form **B** and **C**.

Figure 16.7


Energy diagram for the two-step mechanism:
 $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + \text{HBr} \rightarrow$
 $\text{CH}_3\text{CH}(\text{Br})\text{CH}=\text{CH}_2 +$
 $\text{CH}_3\text{CH}=\text{CHCH}_2\text{Br}$



- At low temperature, the energy of activation is the more important factor. Because most molecules do not have enough kinetic energy to overcome the higher energy barrier at lower temperature, they react by the faster pathway, forming the kinetic product.
- At higher temperature, most molecules have enough kinetic energy to reach either transition state. The two products are in equilibrium with each other, and the more stable compound—which is lower in energy—becomes the major product.

Problem 16.20

Label each product in the following reaction as a 1,2-product or 1,4-product, and decide which is the kinetic product and which is the thermodynamic product.




16.12 The Diels–Alder Reaction


Diels and Alder shared the 1950 Nobel Prize in Chemistry for unraveling the intricate details of this remarkable reaction.

The arrows may be drawn in a clockwise or counterclockwise direction to show the flow of electrons in a Diels–Alder reaction.

The **Diels–Alder reaction**, named for German chemists Otto Diels and Kurt Alder, is an addition reaction between a **1,3-diene** and an alkene called a **dienophile**, to form a new six-membered ring.



Three curved arrows are needed to show the cyclic movement of electron pairs because three π bonds break and two σ bonds and one π bond form. Because each new σ bond is ~ 100 kJ/mol stronger than a π bond that is broken, a typical Diels–Alder reaction releases ~ 200 kJ/mol of energy. The following equations illustrate three examples of the Diels–Alder reaction:




All Diels–Alder reactions have the following features in common:

- [1] They are initiated by heat; that is, the Diels–Alder reaction is a *thermal* reaction.
- [2] They form new six-membered rings.
- [3] Three π bonds break, and two new C–C σ bonds and one new C–C π bond form.
- [4] They are concerted; that is, all old bonds are broken and all new bonds are formed in a single step.

The Diels–Alder reaction forms new carbon–carbon bonds, so it can be used to synthesize larger, more complex molecules from smaller ones. For example, Figure 16.8 illustrates a Diels–Alder reaction used in the synthesis of tetrodotoxin, a toxin isolated from many different types of puffer fish.

Diels–Alder reactions may seem complicated at first, but they are really less complicated than many of the reactions you have already learned, especially those with multistep mechanisms and carbocation intermediates. The key is to learn how to arrange the starting materials to more easily visualize the structure of the product.


Figure 16.8
Synthesis of a natural product
using the Diels–Alder reaction



- Tetrodotoxin, a complex natural product containing several six-membered rings joined together, is a poison isolated from the ovaries and liver of the puffer fish, so named because the fish inflates itself into a ball when alarmed. Eating fish tainted with trace amounts of this potent toxin results in weakness, paralysis, and eventually death. One step in the synthesis of tetrodotoxin involves forming a six-membered ring by a Diels–Alder reaction.


HOW TO Draw the Product of a Diels–Alder Reaction

Example Draw the product of the following Diels–Alder reaction:




Step [1] Arrange the 1,3-diene and the dienophile next to each other, with the diene drawn in the *s-cis* conformation.


- This step is key: **Rotate the diene** so that it is drawn in the *s-cis* conformation, and place the end C's of the diene close to the double bond of the dienophile.



Step [2] Cleave the three π bonds and use arrows to show where the new bonds will be formed.



Problem 16.21 Draw the product formed when each diene and dienophile react in a Diels–Alder reaction.




16.13 Specific Rules Governing the Diels–Alder Reaction

Several rules govern the course of the Diels–Alder reaction.

16.13A Diene Reactivity


Rule [1] The diene can react only when it adopts the *s-cis* conformation.

Both ends of the conjugated diene must be close to the π bond of the dienophile for reaction to occur. Thus, an acyclic diene in the *s-trans* conformation must rotate about the central C–C σ bond to form the *s-cis* conformation before reaction can take place.



This rotation is prevented in cyclic dienes. As a result:

- When the two double bonds are constrained in the *s-cis* conformation, the diene is unusually *reactive*.
- When the two double bonds are constrained in the *s-trans* conformation, the diene is *unreactive*.



Problem 16.22 Rank the following dienes in order of increasing reactivity in a Diels–Alder reaction.




Problem 16.23 Why is (2*Z*,4*Z*)-2,4-hexadiene considerably less reactive than (2*E*,4*E*)-2,4-hexadiene in a Diels–Alder reaction?


16.13B Dienophile Reactivity

Rule [2] Electron-withdrawing substituents in the dienophile increase the reaction rate.

In a Diels–Alder reaction, the conjugated diene acts as a nucleophile and the dienophile acts as an electrophile. As a result, electron-withdrawing groups make the dienophile more electrophilic (and, thus, more reactive) by withdrawing electron density from the carbon–carbon double bond. If *Z* is an electron-withdrawing group, then the reactivity of the dienophile increases as follows:



A carbonyl group is an effective electron-withdrawing group because the carbonyl carbon bears a partial positive charge (δ^+), which withdraws electron density from the carbon–carbon double bond of the dienophile. Common dienophiles that contain a carbonyl group are shown in Figure 16.9.



Problem 16.24 Rank the following dienophiles in order of increasing reactivity.





Figure 16.9
Common dienophiles in the Diels–Alder reaction




16.13C Stereospecificity


Rule [3] The stereochemistry of the dienophile is retained in the product.

- A **cis dienophile** forms a **cis-substituted cyclohexene**.
- A **trans dienophile** forms a **trans-substituted cyclohexene**.


The two **cis** COOH groups of maleic acid become two **cis** substituents in a Diels–Alder adduct. The COOH groups can be drawn both above or both below the plane to afford a single achiral **meso** compound. The **trans dienophile** fumaric acid yields two enantiomers with **trans** COOH groups.



A **cyclic dienophile** forms a **bicyclic product**. A bicyclic system in which the two rings share a common C–C bond is called a **fused ring system**. The two H atoms at the ring fusion must be **cis**, because they were **cis** in the starting dienophile. A bicyclic system of this sort is said to be **cis-fused**.




Problem 16.25 Draw the products of each Diels–Alder reaction, and indicate the stereochemistry.



16.13D The Rule of Endo Addition

Rule [4] When endo and exo products are possible, the endo product is preferred.

To understand the rule of endo addition, we must first examine Diels–Alder products that result from cyclic 1,3-dienes. When cyclopentadiene reacts with a dienophile such as ethylene, a new six-membered ring forms, and above the ring there is a one atom “bridge.” This carbon atom originated as the sp^3 hybridized carbon of the diene that was not involved in the reaction.




The product of the Diels–Alder reaction of a cyclic 1,3-diene is bicyclic, but the carbon atoms shared by both rings are *non-adjacent*. Thus, this bicyclic product differs from the fused ring system obtained when the dienophile is cyclic.

- A bicyclic ring system in which the two rings share non-adjacent carbon atoms is called a *bridged* ring system.

Fused and bridged bicyclic ring systems are compared in Figure 16.10.

When cyclopentadiene reacts with a substituted alkene as the dienophile ($\text{CH}_2=\text{CHZ}$), the substituent Z can be oriented in one of two ways in the product. The terms **endo** and **exo** are used to indicate the position of Z.



- A substituent on one bridge is *endo* if it is closer to the *longer* bridge that joins the two carbons common to both rings.
- A substituent is *exo* if it is closer to the *shorter* bridge that joins the carbons together.

Figure 16.10
Fused and bridged bicyclic ring systems compared






Figure 16.11

How endo and exo products are formed in the Diels–Alder reaction


Pathway [1] With Z oriented under the diene, the endo product is formed.




Pathway [2] With Z oriented away from the diene, the exo product is formed.



To help you distinguish endo and exo, remember that **endo** is under the newly formed six-membered ring.




In a Diels–Alder reaction, the **endo** product is preferred, as shown in two examples.



The Diels–Alder reaction is concerted, and the reaction occurs with the diene and the dienophile arranged one above the other, as shown in Figure 16.11, not side-by-side. In theory, the substituent Z can be oriented either directly under the diene to form the endo product (Pathway [1] in Figure 16.11) or away from the diene to form the exo product (Pathway [2] in Figure 16.11). In practice, though, the **endo product is the major product**. The transition state leading to the endo product allows more interaction between the electron-rich diene and the electron-withdrawing substituent Z on the dienophile, an energetically favorable arrangement.

Problem 16.26 Draw the product of each Diels–Alder reaction.



16.14 Other Facts About the Diels–Alder Reaction


16.14A Retrosynthetic Analysis of a Diels–Alder Product

The Diels–Alder reaction is used widely in organic synthesis, so you must be able to look at a compound and determine what conjugated diene and what dienophile were used to make it. To draw the starting materials from a given Diels–Alder adduct:

- **Locate the six-membered ring that contains the C=C.**
- **Draw three arrows around the cyclohexene ring, beginning with the π bond. Each arrow moves two electrons to the adjacent bond, cleaving one π bond and two σ bonds, and forming three π bonds.**
- **Retain the stereochemistry of substituents on the C=C of the dienophile. Cis substituents on the six-membered ring give a cis dienophile.**

This stepwise retrosynthetic analysis gives the 1,3-diene and dienophile needed for any Diels–Alder reaction, as shown in the two examples in Figure 16.12.

Problem 16.27 What diene and dienophile are needed to prepare each product?



16.14B Retro Diels–Alder Reaction

A reactive diene like 1,3-cyclopentadiene readily undergoes a Diels–Alder reaction with *itself*; that is, **1,3-cyclopentadiene dimerizes because one molecule acts as the diene and another acts as the dienophile.**






Figure 16.12

Finding the diene and dienophile needed for a Diels–Alder reaction



The formation of dicyclopentadiene is so rapid that it takes only a few hours at room temperature for cyclopentadiene to completely dimerize. How, then, can cyclopentadiene be used in a Diels–Alder reaction if it really exists as a dimer?

When heated, dicyclopentadiene undergoes a **retro Diels–Alder reaction**, and two molecules of cyclopentadiene are re-formed. If cyclopentadiene is immediately treated with a different dienophile, it reacts to form a new Diels–Alder adduct with this dienophile.




16.14C Application: Diels–Alder Reaction in the Synthesis of Steroids


Recall from Section 4.15 that lipids are water-insoluble biomolecules that have diverse structures.

Steroids are tetracyclic lipids containing three six-membered rings and one five-membered ring. The four rings are designated as **A**, **B**, **C**, and **D**.


The steroid skeleton




Steroids exhibit a wide range of biological properties, depending on the substitution pattern of functional groups on the rings. They include **cholesterol** (a component of cell membranes that is implicated in cardiovascular disease), **estrone** (a female sex hormone responsible for the regulation of the menstrual cycle), and **cortisone** (a hormone responsible for the control of inflammation and the regulation of carbohydrate metabolism).



Diels–Alder reactions have been used widely in the laboratory syntheses of steroids. The key Diels–Alder reactions used to prepare the C ring of estrone and the B ring of cortisone are as follows:




Problem 16.28 Draw the product (**A**) of the following Diels-Alder reaction. **A** was a key intermediate in the synthesis of the addicting pain reliever morphine, isolated from the opium poppy.




16.15 Conjugated Dienes and Ultraviolet Light

Recall from Chapter 13 that the absorption of infrared energy can promote a molecule from a lower vibrational state to a higher one. In a similar fashion, the absorption of ultraviolet (UV) light can promote an electron from a lower electronic state to a higher one. Ultraviolet light has a slightly shorter wavelength (and, thus, higher frequency) than visible light. The most useful region of UV light for this purpose is **200–400 nm**.




16.15A General Principles

When electrons in a lower energy state (the **ground state**) absorb light having the appropriate energy, an electron is promoted to a higher electronic state (the **excited state**).




The energy difference between the two states depends on the location of the electron. The promotion of electrons in σ bonds and unconjugated π bonds requires light having a wavelength of < 200 nm; that is, it has a shorter wavelength and higher energy than light in the UV region of the electromagnetic spectrum. With conjugated dienes, however, the energy difference between the ground and excited states decreases, so longer wavelengths of light can be used to promote electrons. The wavelength of UV light absorbed by a compound is often referred to as its λ_{max} . 1,3-Butadiene, for example, absorbs UV light at $\lambda_{\text{max}} = 217$ nm and 1,3-cyclohexadiene has a λ_{max} of 256 nm.



- Conjugated dienes and polyenes absorb light in the UV region of the electromagnetic spectrum (200–400 nm).


As the number of conjugated π bonds increases, the energy difference between the ground and excited state decreases, shifting the absorption to longer wavelengths.




With molecules having eight or more conjugated π bonds, the absorption shifts from the UV to the visible region and the compound takes on the color of those wavelengths of visible light it does *not* absorb. For example, lycopene absorbs visible light at $\lambda_{\text{max}} = 470$ nm, in the blue-green region of the visible spectrum. Because it does not absorb light in the red region, lycopene appears bright red (Figure 16.13).

Figure 16.13

Why lycopene appears red



Problem 16.29 Which compound in each pair absorbs UV light at longer wavelength?




16.15B Sunscreens



Commercial sunscreens are given an **SPF** rating (sun protection factor), according to the amount of sunscreen present. The higher the number, the greater the protection.


Ultraviolet radiation from the sun is high enough in energy to cleave bonds, forming radicals that can prematurely age skin and cause skin cancers. The ultraviolet region is often subdivided, based on the wavelength of UV light: UV-A (320–400 nm), UV-B (290–320 nm), and UV-C (< 290 nm). Fortunately, much of the highest energy UV light (UV-C) is filtered out by the ozone layer, so that only UV light having wavelengths > 290 nm reaches the skin's surface. Much of this UV light is absorbed by **melanin**, the highly conjugated colored pigment in the skin that serves as the body's natural protection against the harmful effects of UV radiation.

Prolonged exposure to the sun can allow more UV radiation to reach your skin than melanin can absorb. A commercial sunscreen can offer added protection, however, because it contains conjugated compounds that absorb UV light, thus shielding your skin (for a time) from the harmful effects of UV radiation. Two sunscreens that have been used for this purpose are *para*-aminobenzoic acid (PABA) and padimate O.



Many sunscreens contain more than one component to filter out different regions of the UV spectrum. Conjugated compounds generally shield the skin from UV-B radiation, but often have little effect on longer-wavelength UV-A radiation, which does not burn the skin, but can still cause long-term damage to skin cells.

Problem 16.30 Which of the following compounds might be an ingredient in a commercial sunscreen? Explain why or why not.



KEY CONCEPTS


Conjugation, Resonance, and Dienes

Conjugation and Delocalization of Electron Density


- The overlap of *p* orbitals on three or more adjacent atoms allows electron density to delocalize, thus adding stability (16.1).
- An allyl carbocation ($\text{CH}_2=\text{CHCH}_2^+$) is more stable than a 1° carbocation because of *p* orbital overlap (16.2).
- In any system $\text{X}=\text{Y}-\text{Z}$, Z is sp^2 hybridized to allow the lone pair to occupy a *p* orbital, making the system conjugated (16.5).

Four Common Examples of Resonance (16.3)

[1] The three-atom “allyl” system:



[2] Conjugated double bonds:



[3] Cations having a positive charge adjacent to a lone pair:



[4] Double bonds involving one atom more electronegative than the other:



Rules on Evaluating the Relative “Stability” of Resonance Structures (16.4)


- Structures with more bonds and fewer charges are more stable.
- Structures in which every atom has an octet are more stable.
- Structures that place a negative charge on a more electronegative atom are more stable.

The Unusual Properties of Conjugated Dienes

- The C–C σ bond joining the two double bonds is unusually short (16.8).
- Conjugated dienes are more stable than the corresponding isolated dienes. ΔH° of hydrogenation is smaller for a conjugated diene than for an isolated diene converted to the same product (16.9).
- The reactions are unusual:
 - Electrophilic addition affords products of 1,2-addition and 1,4-addition (16.10, 16.11).
 - Conjugated dienes undergo the Diels–Alder reaction, a reaction that does not occur with isolated dienes (16.12–16.14).
- Conjugated dienes absorb UV light in the 200–400 nm region. As the number of conjugated π bonds increases, the absorption shifts to longer wavelength (16.15).


Reactions of Conjugated Dienes

[1] Electrophilic addition of HX (X = halogen) (16.10–16.11)



- The mechanism has two steps.
- Markovnikov’s rule is followed. Addition of H^+ forms the more stable allylic carbocation.
- The 1,2-product is the kinetic product. When H^+ adds to the double bond, X^- adds to the end of the allylic carbocation to which it is closer (C2 not C4). The kinetic product is formed faster at low temperature.
- The thermodynamic product has the more substituted, more stable double bond. The thermodynamic product predominates at equilibrium. With 1,3-butadiene, the thermodynamic product is the 1,4-product.

[2] Diels–Alder reaction (16.12–16.14)




- The reaction forms two σ and one π bond in a six-membered ring.
- The reaction is initiated by heat.
- The mechanism is concerted: All bonds are broken and formed in a single step.
- The diene must react in the *s*-cis conformation (16.13A).
- Electron-withdrawing groups in the dienophile increase the reaction rate (16.13B).
- The stereochemistry of the dienophile is retained in the product (16.13C).
- Endo products are preferred (16.13D).


PROBLEMS

Conjugation

16.31 Which of the following systems are conjugated?




16.32 Label each double bond in the following natural products as isolated or conjugated.




16.33 Explain why 2,3-di-*tert*-butyl-1,3-butadiene does not behave like a conjugated diene in a Diels–Alder reaction.

Resonance and Hybridization


16.34 Draw all reasonable resonance structures for each species.



16.35 Draw all reasonable resonance structures for each species.




16.36 Explain why the cyclopentadienide anion **A** gives only one signal in its ^{13}C NMR spectrum.




16.37 Explain why the N atoms in $\text{C}_6\text{H}_5\text{NH}_2$ and $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$ are hybridized differently.

16.38 Explain each statement using resonance theory.

a. The indicated C–H bond in propene is more acidic than the indicated C–H bond in propane.



b. The bond dissociation energy for the C–C bond in ethane is much higher than the bond dissociation energy for the indicated C–C bond in 1-butene.



Nomenclature and Stereoisomers in Conjugated Dienes


16.39 Draw the structure of each compound.

- (3Z)-1,3-pentadiene in the *s*-trans conformation
- (2E,4Z)-1-bromo-3-methyl-2,4-hexadiene
- (2E,4E,6E)-2,4,6-octatriene
- (2E,4E)-3-methyl-2,4-hexadiene in the *s*-cis conformation

16.40 Draw and name all dienes of molecular formula C_5H_8 . When *E* and *Z* isomers are possible, draw and name each stereoisomer.

16.41 Draw all possible stereoisomers of 2,4-heptadiene and label each double bond as *E* or *Z*.

16.42 Label each pair of compounds as stereoisomers or conformations.




16.43 Rank the following dienes in order of increasing heat of hydrogenation.




Electrophilic Addition


16.44 Draw the products formed when each compound is treated with one equivalent of HBr.




16.45 Treatment of alkenes **A** and **B** with HBr gives the same alkyl halide **C**. Draw a mechanism for each reaction, including all reasonable resonance structures for any intermediate.



16.46 Draw a stepwise mechanism for the following reaction.




16.47 Addition of HCl to alkene **X** forms two alkyl halides **Y** and **Z**.



- Label **Y** and **Z** as a 1,2-addition product or a 1,4-addition product.
- Label **Y** and **Z** as the kinetic or thermodynamic product and explain why.
- Explain why addition of HCl occurs at the indicated C=C (called an exocyclic double bond), rather than the other C=C (called an endocyclic double bond).

16.48 Explain, with reference to the mechanism, why addition of one equivalent of HCl to diene **A** forms only two products of electrophilic addition, even though four constitutional isomers are possible.




16.49 The major product formed by addition of HBr to $(CH_3)_2C=CH-CH=C(CH_3)_2$ is the same at low and high temperature. Draw the structure of the major product and explain why the kinetic and thermodynamic products are the same in this reaction.


Diels–Alder Reaction

16.50 Explain why methyl vinyl ether ($\text{CH}_2=\text{CHOCH}_3$) is not a reactive dienophile in the Diels–Alder reaction.


16.51 Draw the products of the following Diels–Alder reactions. Indicate stereochemistry where appropriate.




16.52 What diene and dienophile are needed to prepare each Diels–Alder product?




16.53 Give two different ways to prepare the following compound by the Diels–Alder reaction. Explain which method is preferred.




16.54 Compounds containing triple bonds are also Diels–Alder dienophiles. With this in mind, draw the products of each reaction.




16.55 Diels–Alder reaction of a monosubstituted diene (such as $\text{CH}_2=\text{CH}-\text{CH}=\text{CHOCH}_3$) with a monosubstituted dienophile (such as $\text{CH}_2=\text{CHCHO}$) gives a mixture of products, but the 1,2-disubstituted product often predominates. Draw the resonance hybrid for each reactant and use the charge distribution of the hybrids to explain why the 1,2-disubstituted product is the major product.




16.56 What is the structure of the product formed when **A** is heated in the presence of maleic acid? Explain why only one product is formed even though **A** has four double bonds.




- 16.57 The following reactions have been used to synthesize dieldrin and aldrin (named for Diels and Alder), two pesticides having a similar story to DDT (Section 7.4). Identify the lettered compounds in this reaction scheme.




- 16.58 Devise a stepwise synthesis of each compound from dicyclopentadiene using a Diels–Alder reaction as one step. You may also use organic compounds having ≤ 4 C's, and any required organic or inorganic reagents.




- 16.59 Intramolecular Diels–Alder reactions are possible when a substrate contains both a 1,3-diene and a dienophile, as shown in the following general reaction.



With this in mind, draw the product of each intramolecular Diels–Alder reaction.




- 16.60 A transannular Diels–Alder reaction is an intramolecular reaction that occurs when the diene and dienophile are contained in one ring, resulting in the formation of a tricyclic ring system. Draw the product formed when the following triene undergoes a transannular Diels–Alder reaction.




General Reactions

- 16.61 Draw a stepwise mechanism for the following reaction.




- 16.62 Draw the products of each reaction. Indicate the stereochemistry of Diels–Alder products.





- 16.63** Like alkenes, conjugated dienes can be prepared by elimination reactions. Draw a stepwise mechanism for the acid-catalyzed dehydration of 3-methyl-2-buten-1-ol $[(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{OH}]$ to isoprene $[\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}=\text{CH}_2]$.
- 16.64** (a) Draw the two isomeric dienes formed when $\text{CH}_2=\text{CHCH}_2\text{CH}(\text{Cl})\text{CH}(\text{CH}_3)_2$ is treated with an alkoxide base. (b) Explain why the major product formed in this reaction does not contain the more highly substituted alkene.

Spectroscopy

- 16.65** The treatment of isoprene $[\text{CH}_2=\text{C}(\text{CH}_3)\text{CH}=\text{CH}_2]$ with one equivalent of mCPBA forms **A** as the major product. **A** gives a molecular ion at 84 in its mass spectrum, and peaks at $2850\text{--}3150\text{ cm}^{-1}$ in its IR spectrum. The $^1\text{H NMR}$ spectrum of **A** is given below. What is the structure of **A**?




- 16.66** The treatment of $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{Br}$ with H_2O forms **B** (molecular formula $\text{C}_5\text{H}_{10}\text{O}$) as one of the products. Determine the structure of **B** from its $^1\text{H NMR}$ and IR spectra.




UV Absorption

- 16.67** Rank the following compounds in the order of increasing λ_{max} .



- 16.68** Explain why ferulic acid, a natural product found in rice, oats, and other plants, is both an antioxidant and a sunscreen.



Benzene and Aromatic Compounds

17

- 17.1 Background
- 17.2 The structure of benzene
- 17.3 Nomenclature of benzene derivatives
- 17.4 Spectroscopic properties
- 17.5 Interesting aromatic compounds
- 17.6 Benzene's unusual stability
- 17.7 The criteria for aromaticity—Hückel's rule
- 17.8 Examples of aromatic compounds
- 17.9 What is the basis of Hückel's rule?
- 17.10 The inscribed polygon method for predicting aromaticity
- 17.11 Buckminsterfullerene—Is it aromatic?



Capsaicin is responsible for the characteristic spicy flavor of jalapeño and habañero peppers. Although it first produces a burning sensation on contact with the mouth or skin, repeated application desensitizes the area to pain. This property has made it the active ingredient in several topical creams for the treatment of chronic pain. Capsaicin has also been used as an animal deterrent in pepper sprays, and as an additive to make birdseed squirrel-proof. Capsaicin is an aromatic compound because it contains a benzene ring. In this chapter, we learn about the characteristics of aromatic compounds like capsaicin.


The hydrocarbons we have examined thus far—including the alkanes, alkenes, and alkynes, as well as the conjugated dienes and polyenes of Chapter 16—have been aliphatic hydrocarbons. In Chapter 17, we continue our study of conjugated systems with **aromatic hydrocarbons**.

We begin with **benzene** and then examine other cyclic, planar, and conjugated ring systems to learn the modern definition of what it means to be aromatic. Then, in Chapter 18, we will learn about the reactions of aromatic compounds, highly unsaturated hydrocarbons that do not undergo addition reactions like other unsaturated compounds. An explanation of this behavior relies on an understanding of the structure of aromatic compounds presented in Chapter 17.


17.1 Background

For 6 C's, the maximum number of H's = $2n + 2 = 2(6) + 2 = 14$. Because benzene contains only 6 H's, it has $14 - 6 = 8$ H's fewer than the maximum number. This corresponds to 8 H's/2 H's for each degree of unsaturation = **four degrees of unsaturation in benzene**.

Benzene (C₆H₆) is the simplest aromatic hydrocarbon (or arene). Since its isolation by Michael Faraday from the oily residue remaining in the illuminating gas lines in London in 1825, it has been recognized as an unusual compound. Based on the calculation introduced in Section 10.2, **benzene has four degrees of unsaturation, making it a highly unsaturated hydrocarbon**. But, whereas unsaturated hydrocarbons such as alkenes, alkynes, and dienes readily undergo addition reactions, *benzene does not*. For example, bromine adds to ethylene to form a dibromide, but benzene is inert under similar conditions.




Benzene *does* react with bromine, but only in the presence of FeBr₃ (a Lewis acid), and the reaction is a **substitution**, *not* an addition.



Thus, any structure proposed for benzene must account for its high degree of unsaturation and its lack of reactivity towards electrophilic addition.

In the last half of the nineteenth century August Kekulé proposed structures that were close to the modern description of benzene. In the Kekulé model, benzene was thought to be a rapidly equilibrating mixture of two compounds, each containing a six-membered ring with three alternating π bonds. These structures are now called **Kekulé structures**. In the Kekulé description, the bond between any two carbon atoms is sometimes a single bond and sometimes a double bond.



Although benzene is still drawn as a six-membered ring with three alternating π bonds, in reality **there is no equilibrium between two different kinds of benzene molecules**. Instead, current descriptions of benzene are based on resonance and electron delocalization due to orbital overlap, as detailed in Section 17.2.

In the nineteenth century, many other compounds having properties similar to those of benzene were isolated from natural sources. Because these compounds possessed strong and characteristic odors, they were called **aromatic compounds**. It is their chemical properties, though, not their odor that make these compounds special.


- Aromatic compounds resemble benzene—they are unsaturated compounds that do not undergo the addition reactions characteristic of alkenes.

17.2 The Structure of Benzene

Any structure for benzene must account for the following:


- It contains a six-membered ring and three additional degrees of unsaturation.
- It is planar.
- All C–C bond lengths are equal.

Although the Kekulé structures satisfy the first two criteria, they break down with the third, because having three alternating π bonds means that benzene should have three short double bonds alternating with three longer single bonds.



Resonance

Benzene is conjugated, so we must use resonance and orbitals to describe its structure. The resonance description of benzene consists of two equivalent Lewis structures, each with three double bonds that alternate with three single bonds.



The electrons in the π bonds are **delocalized** around the ring.

The resonance description of benzene matches the Kekulé description with one important exception. **The two Kekulé representations are not in equilibrium with each other.** Instead, the true structure of benzene is a resonance **hybrid** of the two Lewis structures, with the dashed lines of the hybrid indicating the position of the π bonds.

Some texts draw benzene as a hexagon with an inner circle:



The circle represents the **six π electrons**, distributed over the six atoms of the ring.

We will use one of the two Lewis structures and not the hybrid in drawing benzene, because it is easier to keep track of the electron pairs in the π bonds (the π electrons).


- Because each π bond has two electrons, benzene has six π electrons.

The resonance hybrid of benzene explains why all C–C bond lengths are the same. Each C–C bond is single in one resonance structure and double in the other, so the actual bond length (139 pm) is intermediate between a carbon–carbon single bond (153 pm) and a carbon–carbon double bond (134 pm).



Hybridization and Orbitals

Each carbon atom in a benzene ring is surrounded by three atoms and no lone pairs of electrons, making it **sp^2 hybridized and trigonal planar with all bond angles 120°** . Each carbon also has a p orbital with one electron that extends above and below the plane of the molecule.




The six adjacent p orbitals overlap, delocalizing the six electrons over the six atoms of the ring and making benzene a conjugated molecule. Because each p orbital has two lobes, one above and one below the plane of the benzene ring, the overlap of the p orbitals creates two “doughnuts” of electron density, as shown in Figure 17.1a. The electrostatic potential plot in Figure 17.1b also shows that the electron-rich region is concentrated above and below the plane of the molecule, where the six π electrons are located.


- Benzene's six π electrons make it electron rich and so it readily reacts with electrophiles.

Problem 17.1

Draw all possible resonance structures for the antihistamine diphenhydramine, the active ingredient in Benadryl.

**Problem 17.2**

What orbitals are used to form the bonds indicated in each molecule? Of the indicated C–C bonds, which is the shortest?

**17.3 Nomenclature of Benzene Derivatives**

Many organic molecules contain a benzene ring with one or more substituents, so we must learn how to name them. Many common names are recognized by the IUPAC system, however, so this complicates the nomenclature of benzene derivatives somewhat.

Figure 17.1


Two views of the electron density in a benzene ring

a. View of the p orbital overlap



- Overlap of six adjacent p orbitals creates two rings of electron density, one above and one below the plane of the benzene ring.


b. Electrostatic potential plot




- The electron-rich region (in red) is concentrated above and below the ring carbons, where the six π electrons are located. (The electron-rich region below the plane is hidden from view.)

17.3A Monosubstituted Benzenes

To name a benzene ring with one substituent, **name the substituent and add the word *benzene***. Carbon substituents are named as alkyl groups.




Many monosubstituted benzenes, such as those with methyl (CH_3 -), hydroxy ($-\text{OH}$), and amino ($-\text{NH}_2$) groups, have common names that you must learn, too.




17.3B Disubstituted Benzenes

There are three different ways that two groups can be attached to a benzene ring, so a prefix—**ortho**, **meta**, or **para**—can be used to designate the relative position of the two substituents. Ortho, meta, and para are also abbreviated as *o*, *m*, and *p*, respectively.



If the two groups on the benzene ring are different, **alphabetize the names of the substituents** preceding the word *benzene*. If one of the substituents is part of a **common root**, name the **molecule as a derivative of that monosubstituted benzene**.




17.3C Polysubstituted Benzenes

For three or more substituents on a benzene ring:

- [1] **Number to give the lowest possible numbers around the ring.**
- [2] **Alphabetize the substituent names.**


- [3] When substituents are part of common roots, name the molecule as a derivative of that monosubstituted benzene. The substituent that comprises the common root is located at C1.

Examples of naming polysubstituted benzenes



- Assign the lowest set of numbers.
- Alphabetize the names of all the substituents.

4-chloro-1-ethyl-2-propylbenzene




- Name the molecule as a derivative of the common root **aniline**.
- Designate the position of the NH_2 group as "1," and then assign the lowest possible set of numbers to the other substituents.

2,5-dichloroaniline


17.3D Naming Aromatic Rings as Substituents

A benzene substituent (C_6H_5-) is called a **phenyl group**, and it can be abbreviated in a structure as **Ph-**.




- A phenyl group (C_6H_5-) is formed by removing one hydrogen from benzene (C_6H_6).

Benzene, therefore, can be represented as PhH , and phenol would be PhOH .




The **benzyl** group, another common substituent that contains a benzene ring, differs from a phenyl group.




Finally, substituents derived from other substituted aromatic rings are collectively called **aryl groups**.

Examples of aryl groups



Problem 17.3 Give the IUPAC name for each compound.



Problem 17.4 Draw the structure corresponding to each name:

- a. isobutylbenzene d. *m*-bromoaniline
 b. *o*-dichlorobenzene e. 4-chloro-1,2-diethylbenzene
 c. *cis*-1,2-diphenylcyclohexane f. 3-*tert*-butyl-2-ethyltoluene

Problem 17.5 Draw and name all the isomeric trichlorobenzenes (molecular formula $\text{C}_6\text{H}_3\text{Cl}_3$).

17.4 Spectroscopic Properties

The important IR and NMR absorptions of aromatic compounds are summarized in Table 17.1.

The absorption at 6.5–8.0 ppm in the ^1H NMR spectrum is particularly characteristic of compounds containing benzene rings. **All aromatic compounds have highly deshielded protons due to the ring current effect of the circulating π electrons**, as discussed in Section 14.4. Observing whether a new compound absorbs in this region of a ^1H NMR spectrum is one piece of data used to determine if it is aromatic.

^{13}C NMR spectroscopy is used to determine the substitution patterns in disubstituted benzenes, because each line in a spectrum corresponds to a different kind of carbon atom. For example, *o*-, *m*-, and *p*-dibromobenzene each exhibit a different number of lines in its ^{13}C NMR spectrum, as shown in Figure 17.2.

Table 17.1 Characteristic Spectroscopic Absorptions of Benzene Derivatives




Type of spectroscopy	Type of C, H	Absorption
IR absorptions	$\text{C}_{\text{sp}^2}\text{-H}$	3150–3000 cm^{-1}
	$\text{C}=\text{C}$ (arene)	1600, 1500 cm^{-1}
^1H NMR absorptions	 (aryl H)	6.5–8 ppm (highly deshielded protons)
	 (benzylic H)	1.5–2.5 ppm (somewhat deshielded $\text{C}_{\text{sp}^3}\text{-H}$)
^{13}C NMR absorption	C_{sp^2} of arenes	120–150 ppm


Figure 17.2
 ^{13}C NMR absorptions
 of the three isomeric
 dibromobenzenes



- The number of signals (lines) in the ^{13}C NMR spectrum of a disubstituted benzene with two identical groups indicates whether they are ortho, meta, or para to each other.

Problem 17.6 What is the structure of a compound of molecular formula $C_{10}H_{14}O_2$ that shows a strong IR absorption at $3150\text{--}2850\text{ cm}^{-1}$ and gives the following ^1H NMR absorptions: 1.4 (triplet, 6 H), 4.0 (quartet, 4 H), and 6.8 (singlet, 4 H) ppm?

Problem 17.7 How many ^{13}C NMR signals does each compound exhibit?



Problem 17.8 How do the isomeric trichlorobenzenes ($C_6H_3Cl_3$) drawn in Problem 17.5 differ in their ^{13}C NMR spectra?

17.5 Interesting Aromatic Compounds


BTX contains **benzene**, **toluene**, and **xylene** (the common name for dimethylbenzene).

Benzene and **toluene**, the simplest aromatic hydrocarbons obtained from petroleum refining, are useful starting materials for synthetic polymers. They are two components of the **BTX** mixture added to gasoline to boost octane ratings.


The components of the gasoline additive BTX




benzene



toluene



p-xylene




naphthalene
(used in mothballs)

Compounds containing two or more benzene rings that share carbon–carbon bonds are called **polycyclic aromatic hydrocarbons (PAHs)**. Naphthalene, the simplest PAH, is the active ingredient in mothballs.


Benzo[*a*]pyrene, a more complicated PAH shown in Figure 17.3, is formed by the incomplete combustion of organic materials. It is found in cigarette smoke, automobile exhaust, and the fumes from charcoal grills. When ingested or inhaled, benzo[*a*]pyrene and other similar PAHs are oxidized to carcinogenic products, as discussed in Section 9.17.

Helicene and **twistoflex** are two synthetic PAHs whose unusual shapes are shown in Figure 17.4. Helicene consists of six benzene rings. Because the rings at both ends are not bonded to each other, all of the rings twist slightly, creating a rigid helical shape that prevents the hydrogen atoms on both ends from crashing into each other. Similarly, to reduce steric hindrance between the hydrogen atoms on nearby benzene rings, twistoflex is also nonplanar.

Figure 17.3
Benzo[*a*]pyrene, a
common PAH



- Benzo[*a*]pyrene, produced by the incomplete oxidation of organic compounds in tobacco, is found in cigarette smoke.


Figure 17.4 Helicene and twistoflex—Two synthetic polycyclic aromatic hydrocarbons

Both helicene and twistoflex are chiral molecules—that is, they are not superimposable on their mirror images, even though neither of them contains a stereogenic center. It's their shape that makes them chiral, not the presence of carbon atoms bonded to four different groups. Each ring system is twisted into a shape that lacks a mirror plane, and each structure is rigid, thus creating the chirality.


Many widely used drugs contain a benzene ring. Six examples are shown in Figure 17.5.

17.6 Benzene's Unusual Stability


Considering benzene as the hybrid of two resonance structures adequately explains its equal C–C bond lengths, but does not account for its unusual stability and lack of reactivity towards addition.

Figure 17.5
Selected drugs that contain a benzene ring


- Trade name: **Zoloft**
- Generic name: **sertraline**
- Use: a psychotherapeutic drug for depression and panic disorders




- Trade name: **Valium**
- Generic name: **diazepam**
- Use: a sedative




- Trade name: **Novocain**
- Generic name: **procaine**
- Use: a local anesthetic



- Trade name: **Viracept**
- Generic name: **nelfinavir**
- Use: an antiviral drug used to treat HIV




- Trade name: **Viagra**
- Generic name: **sildenafil**
- Use: a drug used to treat erectile dysfunction



- Trade name: **Claritin**
- Generic name: **loratadine**
- Use: an antihistamine for seasonal allergies

Heats of hydrogenation, which were used in Section 16.9 to show that conjugated dienes are more stable than isolated dienes, can also be used to estimate the stability of benzene. Equations [1]–[3] compare the heats of hydrogenation of cyclohexene, 1,3-cyclohexadiene, and benzene, all of which give cyclohexane when treated with excess hydrogen in the presence of a metal catalyst.



The relative stability of conjugated dienes versus isolated dienes was first discussed in Section 16.9.

The addition of one mole of H_2 to cyclohexene releases -120 kJ/mol of energy (Equation [1]). If each double bond is worth -120 kJ/mol of energy, then the addition of two moles of H_2 to 1,3-cyclohexadiene (Equation [2]) should release 2×-120 kJ/mol = -240 kJ/mol of energy. The observed value, however, is -232 kJ/mol. This is slightly smaller than expected because 1,3-cyclohexadiene is a conjugated diene, and conjugated dienes are more stable than two isolated carbon-carbon double bonds.


The hydrogenations of cyclohexene and 1,3-cyclohexadiene occur readily at room temperature, but benzene can be hydrogenated only under forcing conditions, and even then the reaction is extremely slow. If each double bond is worth -120 kJ/mol of energy, then the addition of three moles of H_2 to benzene should release 3×-120 kJ/mol = -360 kJ/mol of energy. In fact, the observed heat of hydrogenation is only -208 kJ/mol, which is 152 kJ/mol less than predicted and even lower than the observed value for 1,3-cyclohexadiene. Figure 17.6 compares the hypothetical and observed heats of hydrogenation for benzene.

The huge difference between the hypothetical and observed heats of hydrogenation for benzene cannot be explained solely on the basis of resonance and conjugation.


- The low heat of hydrogenation of benzene means that benzene is especially stable, even more so than the conjugated compounds introduced in Chapter 16. This unusual stability is characteristic of aromatic compounds.

Figure 17.6

A comparison between the observed and hypothetical heats of hydrogenation for benzene




Benzene's unusual behavior in chemical reactions is not limited to hydrogenation. As mentioned in Section 17.1, **benzene does not undergo addition reactions typical of other highly unsaturated compounds, including conjugated dienes.** Benzene does not react with Br_2 to yield an addition product. Instead, in the presence of a Lewis acid, bromine *substitutes* for a hydrogen atom, thus yielding a product that retains the benzene ring.



This behavior is characteristic of aromatic compounds. The structural features that distinguish aromatic compounds from the rest are discussed in Section 17.7.

Problem 17.9

Compounds **A** and **B** are both hydrogenated to methylcyclohexane. Which compound has the larger heat of hydrogenation? Which compound is more stable?



17.7 The Criteria for Aromaticity—Hückel's Rule


Four structural criteria must be satisfied for a compound to be aromatic:

- A molecule must be cyclic, planar, completely conjugated, and contain a particular number of π electrons.

[1] A molecule must be cyclic.

- To be aromatic, each p orbital must overlap with p orbitals on two adjacent atoms.

The p orbitals on all six carbons of benzene continuously overlap, so benzene is aromatic. 1,3,5-Hexatriene has six p orbitals, too, but the two on the terminal carbons cannot overlap with each other, so **1,3,5-hexatriene is not aromatic.**



[2] A molecule must be planar.


- All adjacent p orbitals must be aligned so that the π electron density can be delocalized.



cyclooctatetraene
not aromatic




a tub-shaped,
eight-membered ring



Adjacent p orbitals cannot overlap.
Electrons cannot delocalize.

For example, cyclooctatetraene resembles benzene in that it is a cyclic molecule with alternating double and single bonds. Cyclooctatetraene is tub shaped, however, **not planar**, so overlap between adjacent π bonds is impossible. **Cyclooctatetraene, therefore, is not aromatic**, so it undergoes addition reactions like those of other alkenes.



[3] A molecule must be completely conjugated.


- Aromatic compounds must have a p orbital on every atom.

A completely conjugated ring




benzene
a p orbital on **every C**
aromatic

These rings are not completely conjugated.



1,3-cyclohexadiene
not aromatic



1,3,5-cycloheptatriene
not aromatic

Both 1,3-cyclohexadiene and 1,3,5-cycloheptatriene contain at least one carbon atom that does not have a p orbital, and so they are not completely conjugated and therefore **not aromatic**.

[4] A molecule must satisfy Hückel's rule, and contain a particular number of π electrons.

Some compounds satisfy the first three criteria for aromaticity, but still they show none of the stability typical of aromatic compounds. For example, **cyclobutadiene** is so highly reactive that it can only be prepared at extremely low temperatures.



a planar, cyclic, completely conjugated molecule
that is **not aromatic**

It turns out that in addition to being cyclic, planar, and completely conjugated, a compound needs a particular number of π electrons to be aromatic. Erich Hückel first recognized in 1931 that the following criterion, expressed in two parts and now known as **Hückel's rule**, had to be satisfied, as well:

- An aromatic compound must contain $4n + 2$ π electrons ($n = 0, 1, 2$, and so forth).
- Cyclic, planar, and completely conjugated compounds that contain $4n$ π electrons are especially unstable, and are said to be **antiaromatic**.

Hückel's rule refers to the number of π electrons, not the number of atoms in a particular ring.

Thus, compounds that contain 2, 6, 10, 14, 18, and so forth π electrons are aromatic, as shown in Table 17.2. **Benzene is aromatic and especially stable because it contains 6 π electrons. Cyclobutadiene is antiaromatic and especially unstable because it contains 4 π electrons.**




Table 17.2 The Number of π Electrons That Satisfy Hückel's Rule


n	$4n + 2$
0	2
1	6
2	10
3	14
4, etc.	18

Considering aromaticity, all compounds can be classified in one of three ways:


- | | |
|--|--|
| [1] Aromatic | • A cyclic, planar, completely conjugated compound with $4n + 2$ π electrons. |
| [2] Antiaromatic | • A cyclic, planar, completely conjugated compound with $4n$ π electrons. |
| [3] Not aromatic (or nonaromatic) | • A compound that lacks one (or more) of the four requirements to be aromatic or antiaromatic. |

Note, too, the relationship between each compound type and a similar open-chained molecule having the same number of π electrons.

- An aromatic compound is *more* stable than a similar acyclic compound having the same number of π electrons. Benzene is more stable than 1,3,5-hexatriene.
- An antiaromatic compound is *less* stable than an acyclic compound having the same number of π electrons. Cyclobutadiene is less stable than 1,3-butadiene.
- A compound that is not aromatic is *similar* in stability to an acyclic compound having the same number of π electrons. 1,3-Cyclohexadiene is similar in stability to *cis,cis*-2,4-hexadiene, so it is not aromatic.




^1H NMR spectroscopy readily indicates whether a compound is aromatic. The protons on sp^2 hybridized carbons in aromatic hydrocarbons are highly deshielded and absorb at 6.5–8 ppm, whereas hydrocarbons that are not aromatic absorb at 4.5–6 ppm, typical of protons bonded to the C=C of an alkene. Thus, benzene absorbs at 7.3 ppm, whereas cyclooctatetraene, which is not aromatic, absorbs farther upfield, at 5.8 ppm for the protons on its sp^2 hybridized carbons.



Many compounds in addition to benzene are aromatic. Several examples are presented in Section 17.8.

Problem 17.10 Estimate where the protons bonded to the sp^2 hybridized carbons will absorb in the ^1H NMR spectrum of each compound.



17.8 Examples of Aromatic Compounds


In Section 17.8 we look at many different types of aromatic compounds.

17.8A Aromatic Compounds with a Single Ring


Benzene is the most common aromatic compound having a single ring. **Completely conjugated rings larger than benzene are also aromatic if they are planar and have $4n + 2 \pi$ electrons.**

- Hydrocarbons containing a single ring with alternating double and single bonds are called *annulenes*.

To name an annulene, indicate the number of atoms in the ring in brackets and add the word *annulene*. Thus, benzene is [6]-annulene. Both **[14]-annulene** and **[18]-annulene** are cyclic, planar, completely conjugated molecules that follow Hückel's rule, and so they are aromatic.




$$\begin{aligned} & \text{[14]-annulene} \\ & 4n + 2 = 4(3) + 2 = \\ & 14 \pi \text{ electrons} \\ & \text{aromatic} \end{aligned}$$



$$\begin{aligned} & \text{[18]-annulene} \\ & 4n + 2 = 4(4) + 2 = \\ & 18 \pi \text{ electrons} \\ & \text{aromatic} \end{aligned}$$

[10]-Annulene has 10π electrons, which satisfies Hückel's rule, but a planar molecule would place the two H atoms inside the ring too close to each other, so the ring puckers to relieve this strain. Because **[10]-annulene is not planar**, the 10π electrons can't delocalize over the entire ring and it is **not aromatic**.


[10]-Annulene fits Hückel's rule, but it's **not planar**.



[10]-annulene
 10π electrons
not aromatic

=

The molecule puckers to keep these H's farther away from each other.




3-D representation

Problem 17.11 Would [16]-, [20]- or [22]-annulene be aromatic if each ring is planar?

Problem 17.12 Explain why an annulene cannot have an odd number of carbon atoms in the ring.


17.8B Aromatic Compounds with More Than One Ring

Hückel's rule for determining aromaticity can be applied only to monocyclic systems, but many aromatic compounds containing several benzene rings joined together are also known. Two or more six-membered rings with alternating double and single bonds can be fused together to form **polycyclic aromatic hydrocarbons (PAHs)**. Joining two benzene rings together forms **naphthalene**. There are two different ways to join three rings together, forming **anthracene** and **phenanthrene**, and many more complex hydrocarbons are known.



As the number of fused benzene rings increases, the number of resonance structures increases as well. Although two resonance structures can be drawn for benzene, naphthalene is a hybrid of three resonance structures.

Three resonance structures for naphthalene



Problem 17.13 Draw the four resonance structures for anthracene.


17.8C Aromatic Heterocycles

Recall from Section 9.3 that a **heterocycle** is a ring that contains at least one heteroatom.

Heterocycles containing oxygen, nitrogen, or sulfur—atoms that also have at least one lone pair of electrons—can also be aromatic. With heteroatoms, we must always determine whether the lone pair is localized on the heteroatom or part of the delocalized π system. Two examples, **pyridine** and **pyrrole**, illustrate these different possibilities.

Pyridine


Pyridine is a heterocycle containing a six-membered ring with three π bonds and one nitrogen atom. Like benzene, two resonance structures (with all neutral atoms) can be drawn.



two resonance structures for pyridine
6 π electrons


Pyridine is cyclic, planar, and completely conjugated, because the three single and double bonds alternate around the ring. **Pyridine has six π electrons, two from each π bond, thus satisfying Hückel's rule and making pyridine aromatic.** The nitrogen atom of pyridine also has a nonbonded electron pair, which is localized on the N atom, so it is *not* part of the delocalized π electron system of the aromatic ring.

How is the nitrogen atom of the pyridine ring hybridized? The N atom is surrounded by three groups (two atoms and a lone electron pair), making it **sp^2 hybridized**, and leaving one unhybridized p orbital with one electron that overlaps with adjacent p orbitals. The lone pair on N resides in an sp^2 hybrid orbital that is perpendicular to the delocalized π electrons.



Pyrrole

Pyrrole contains a five-membered ring with two π bonds and one nitrogen atom. The N atom also has a lone pair of electrons.




Pyrrole is cyclic and planar, with a total of four π electrons from the two π bonds. Is the non-bonded electron pair localized on N or part of a delocalized π electron system? The lone pair on N is *adjacent* to a double bond. Recall the following general rule from Section 16.5:

- In any system $X=Y-Z$, Z is sp^2 hybridized and the lone pair occupies a p orbital to make the system conjugated.

If the lone pair on the N atom occupies a p orbital:

- Pyrrole has a p orbital on every adjacent atom, so it is completely conjugated.
- Pyrrole has six π electrons—four from the π bonds and two from the lone pair.



Scombroid fish poisoning, associated with facial flushing, hives, and general itching, is caused by the ingestion of inadequately refrigerated fish, typically mahimahi (pictured) and tuna. Bacteria convert the amino acid histidine (Chapter 28) to histamine, which, when consumed in large amounts, results in this clinical syndrome.

Because pyrrole is cyclic, planar, completely conjugated, and has $4n + 2$ π electrons, **pyrrole is aromatic.** The number of electrons—not the size of the ring—determines whether a compound is aromatic.

Electrostatic potential maps, shown in Figure 17.7 for pyridine and pyrrole, confirm that the **lone pair in pyridine is localized on N**, whereas the **lone pair in pyrrole is part of the delocalized π system**. Thus, a fundamental difference exists between the N atoms in pyridine and pyrrole.

- When a heteroatom is already part of a double bond (as in the N of pyridine), its lone pair *cannot* occupy a p orbital and so it cannot be delocalized over the ring.
- When a heteroatom is *not* part of a double bond (as in the N of pyrrole), its lone pair can be located in a p orbital and *delocalized* over a ring to make it aromatic.


Histamine

Histamine, a biologically active amine formed in many tissues, has an aromatic heterocycle with two N atoms, one of which is similar to the N atom of pyridine and one of which is similar to the N atom of pyrrole.


Figure 17.7
Electrostatic potential maps of
pyridine and pyrrole



- In pyridine, the nonbonded electron pair is localized on the N atom in an sp^2 hybridized orbital, as shown by the region of high electron density (in red) on N.
- In pyrrole, the nonbonded electron pair is in a p orbital and is delocalized over the ring, so the entire ring is electron rich (red).




Histamine has a five-membered ring with two π bonds and two nitrogen atoms, each of which contains a lone pair of electrons. The heterocycle has four π electrons from the two double bonds. The lone pair on N1 also occupies a p orbital, making the heterocycle completely conjugated, and giving it a total of six π electrons. The lone pair on N1 is thus delocalized over the five-membered ring and the heterocycle is aromatic. The lone pair on N2 occupies an sp^2 hybrid orbital perpendicular to the delocalized π electrons.




- N1 resembles the N atom of pyrrole.
- N2 resembles the N atom of pyridine.

Histamine produces a wide range of physiological effects in the body. Excess histamine is responsible for the runny nose and watery eyes symptomatic of hay fever. It also stimulates the overproduction of stomach acid, and contributes to the formation of hives. These effects result from the interaction of histamine with two different cellular receptors. We will learn more about antihistamines and antiulcer drugs, compounds that block the effects of histamine, in Section 25.6.

Problem 17.14 Which heterocycles are aromatic?




Problem 17.15




Quinine is isolated from the bark of the cinchona tree native to the Andes Mountains.

- (a) How is each N atom in quinine, an effective antimalarial drug that reduces fever, hybridized?
 (b) In what type of orbital does the lone pair on each N reside?



Problem 17.16

- Januvia is the trade name for sitagliptin, a drug that increases the body's ability to lower blood sugar levels, and thus it was introduced in 2006 for the treatment of type 2 diabetes. (a) Explain why the five-membered ring in sitagliptin is aromatic. (b) Determine the hybridization of each N atom. (c) In what type of orbital does the lone pair on each N atom reside?




17.8D Charged Aromatic Compounds

Both negatively and positively charged ions can also be aromatic if they possess all the necessary elements.


Cyclopentadienyl Anion

The **cyclopentadienyl anion** is a cyclic and planar anion with two double bonds and a nonbonded electron pair. In this way it resembles pyrrole. The two π bonds contribute four electrons and the lone pair contributes two more, for a total of six. By Hückel's rule, having **six π electrons confers aromaticity**. Like the N atom in pyrrole, the **negatively charged carbon atom must be sp^2 hybridized**, and the **nonbonded electron pair must occupy a p orbital** for the ring to be completely conjugated.

The cyclopentadienyl anion




The ring is completely conjugated with **6 π electrons**.




The lone pair resides in a p orbital.


- The cyclopentadienyl anion is aromatic because it is cyclic, planar, completely conjugated, and has six π electrons.

We can draw **five equivalent resonance structures for the cyclopentadienyl anion**, delocalizing the negative charge over every carbon atom of the ring.



The tropylium cation

The ring is completely conjugated with **6 π electrons**.




One p orbital is vacant.


- The tropylium cation is aromatic because it is cyclic, planar, completely conjugated, and has six π electrons delocalized over the seven atoms of the ring.

Problem 17.20 Draw the seven resonance structures for the tropylium cation.

Problem 17.21 Assuming the rings are planar, which ions are aromatic?




Problem 17.22 Compound **A** exhibits a peak in its ^1H NMR spectrum at 7.6 ppm, indicating that it is aromatic. How are the carbon atoms of the triple bonds hybridized? In what type of orbitals are the π electrons of the triple bonds contained? How many π electrons are delocalized around the ring in **A**?



17.9 What Is the Basis of Hückel's Rule?

Why does the number of π electrons determine whether a compound is aromatic? Cyclobutadiene is cyclic, planar, and completely conjugated, just like benzene, but why is benzene aromatic and cyclobutadiene antiaromatic?



A complete explanation is beyond the scope of an introductory organic chemistry text, but nevertheless, you can better understand the basis of aromaticity by learning more about orbitals and bonding.

17.9A Bonding and Antibonding Orbitals

So far we have used the following basic concepts to describe how bonds are formed:

- Hydrogen uses its 1s orbital to form σ bonds with other elements.
- Second-row elements use hybrid orbitals (sp , sp^2 , or sp^3) to form σ bonds.
- Second-row elements use p orbitals to form π bonds.

This description of bonding is called **valence bond theory**. In valence bond theory, a covalent bond is formed by the overlap of two atomic orbitals, and the electron pair in the resulting bond is shared by both atoms. Thus, a carbon–carbon double bond consists of a σ bond, formed by overlap of two sp^2 hybrid orbitals, each containing one electron, and a π bond, formed by overlap of two p orbitals, each containing one electron.


This description of bonding works well for most of the organic molecules we have encountered thus far. Unfortunately, it is inadequate for describing systems with many adjacent p orbitals that overlap, as there are in aromatic compounds. To more fully explain the bonding in these systems, we must utilize **molecular orbital (MO) theory**.

MO theory describes bonds as the mathematical combination of atomic orbitals that form a new set of orbitals called **molecular orbitals (MOs)**. A molecular orbital occupies a region of space *in a molecule* where electrons are likely to be found. When forming molecular orbitals from atomic orbitals, keep in mind:

- A set of n atomic orbitals forms n molecular orbitals.

If *two* atomic orbitals combine, *two* molecular orbitals are formed. This is fundamentally different than valence bond theory. Because aromaticity is based on p orbital overlap, what does MO theory predict will happen when two p (atomic) orbitals combine?

The two lobes of each p orbital are opposite in phase, with a node of electron density at the nucleus. When two p orbitals combine, two molecular orbitals should form. The two p orbitals can add together constructively—that is, with like phases interacting—or destructively—that is, with opposite phases interacting.




- When two p orbitals of similar phase overlap side-by-side, a π bonding molecular orbital results.
- When two p orbitals of opposite phase overlap side-by-side, a π^* antibonding molecular orbital results.

A π bonding MO is lower in energy than the two atomic p orbitals from which it is formed because a stable bonding interaction results when orbitals of similar phase combine. A bonding interaction holds nuclei together. Similarly, a π^* antibonding MO is higher in energy because a destabilizing node results when orbitals of opposite phase combine. A destabilizing interaction pushes nuclei apart.

If two atomic p orbitals each have one electron and then combine to form MOs, the two electrons will occupy the lower energy π bonding MO, as shown in Figure 17.8.

Figure 17.8

Combination of two p orbitals to form π and π^* molecular orbitals




- Two atomic p orbitals combine to form two molecular orbitals. The bonding π MO is lower in energy than the two p orbitals from which it was formed, and the antibonding π^* MO is higher in energy than the two p orbitals from which it was formed.
- Two electrons fill the lower energy bonding MO first.

17.9B Molecular Orbitals Formed When More Than Two p Orbitals Combine

The molecular orbital description of benzene is much more complex than the two MOs formed in Figure 17.8. Because each of the six carbon atoms of benzene has a p orbital, six atomic p orbitals combine to form six π molecular orbitals, as shown in Figure 17.9. A description of the exact appearance and energies of these six MOs requires more sophisticated mathematics and

Figure 17.9

How the six p orbitals of benzene overlap to form six molecular orbitals



- Depicted in this diagram are the interactions of the six atomic p orbitals of benzene, which form six molecular orbitals. When orbitals of like phase combine, a bonding interaction results. When orbitals of opposite phase combine, a destabilizing node results.

understanding of MO theory than is presented in this text. Nevertheless, note that the six MOs are labeled ψ_1 – ψ_6 , with ψ_1 being the lowest in energy and ψ_6 the highest.

The most important features of the six benzene MOs are as follows:

- **The larger the number of bonding interactions, the lower in energy the MO.** The lowest energy molecular orbital (ψ_1) has all bonding interactions between the p orbitals.
- **The larger the number of nodes, the higher in energy the MO.** The highest energy MO (ψ_6^*) has all nodes between the p orbitals.
- Three MOs are lower in energy than the starting p orbitals, making them bonding MOs (ψ_1 , ψ_2 , ψ_3), whereas three MOs are higher in energy than the starting p orbitals, making them antibonding MOs (ψ_4^* , ψ_5^* , ψ_6^*).
- The two pairs of MOs (ψ_2 and ψ_3 ; ψ_4^* and ψ_5^*) with the same energy are called **degenerate orbitals**.
- **The highest energy orbital that contains electrons is called the *highest occupied molecular orbital* (HOMO).** For benzene, the degenerate orbitals ψ_2 and ψ_3 are the HOMOs.
- **The lowest energy orbital that does not contain electrons is called the *lowest unoccupied molecular orbital* (LUMO).** For benzene, the degenerate orbitals ψ_4^* and ψ_5^* are the LUMOs.

To fill the MOs, the six electrons are added, two to an orbital, beginning with the lowest energy orbital. As a result, the six electrons completely fill the bonding MOs, leaving the antibonding MOs empty. This is what gives benzene and other aromatic compounds their special stability and this is why six π electrons satisfies Hückel's $4n + 2$ rule.

- All bonding MOs (and HOMOs) are completely filled in aromatic compounds. No π electrons occupy antibonding MOs.

17.10 The Inscribed Polygon Method for Predicting Aromaticity

An inscribed polygon is also called a **Frost circle**.


To predict whether a compound has π electrons completely filling bonding MOs, we must know how many bonding molecular orbitals and how many π electrons it has. It is possible to predict the relative energies of cyclic, completely conjugated compounds, without sophisticated math (or knowing what the resulting MOs look like) by using the **inscribed polygon method**.

HOW TO Use the Inscribed Polygon Method to Determine the Relative Energies of MOs for Cyclic, Completely Conjugated Compounds

Example Plot the relative energies of the MOs of benzene

Step [1] Draw the polygon in question inside a circle with its vertices touching the circle and one of the vertices pointing down. Mark the points at which the polygon intersects the circle.

- Inscribe a hexagon inside a circle for benzene. The six vertices of the hexagon form six points of intersection, corresponding to the six MOs of benzene. The pattern—a single MO having the lowest energy, two degenerate pairs of MOs, and a single highest energy MO—matches that found in Figure 17.9.



Step [2] Draw a line horizontally through the center of the circle and label MOs as bonding, nonbonding, or antibonding.

- **MOs below this line are bonding**, and lower in energy than the p orbitals from which they were formed. Benzene has three bonding MOs.


—Continued

HOW TO, continued . . .

- **MOs at this line are nonbonding**, and equal in energy to the p orbitals from which they were formed. Benzene has no nonbonding MOs.
- **MOs above this line are antibonding**, and higher in energy than the p orbitals from which they were formed. Benzene has three antibonding MOs.

Step [3] Add the electrons, beginning with the lowest energy MO.

- **All the bonding MOs (and the HOMOs) are completely filled in aromatic compounds. No π electrons occupy antibonding MOs.**
- Benzene is aromatic because it has six π electrons that completely fill the bonding MOs.



This method works for all monocyclic, completely conjugated hydrocarbons regardless of ring size. Figure 17.10 illustrates MOs for completely conjugated five- and seven-membered rings using this method. The total number of MOs always equals the number of vertices of the polygon. Because both systems have three bonding MOs, each needs six π electrons to fully occupy them, making the cyclopentadienyl anion and the tropylium cation aromatic, as we learned in Section 17.8D.

The inscribed polygon method is consistent with Hückel's $4n + 2$ rule; that is, there is always one lowest energy bonding MO that can hold two π electrons and the other bonding MOs come in degenerate pairs that can hold a total of four π electrons. For the compound to be aromatic, these MOs must be completely filled with electrons, so the “magic numbers” for aromaticity fit Hückel's $4n + 2$ rule (Figure 17.11).

Figure 17.10

Using the inscribed polygon method for five- and seven-membered rings






Figure 17.11
MO patterns for cyclic,
completely conjugated
systems




Sample Problem 17.1 Use the inscribed polygon method to show why cyclobutadiene is not aromatic.



Solution


Cyclobutadiene has four MOs (formed from its four p orbitals), to which its four π electrons must be added.

Step [1] Inscribe a square with a vertex down and mark its four points of intersection with the circle.



- The four points of intersection correspond to the four MOs of cyclobutadiene.

Steps [2] and [3] Draw a line through the center of the circle, label the MOs, and add the electrons.



- Cyclobutadiene has four MOs—one bonding, two nonbonding, and one antibonding.
- Adding cyclobutadiene's four π electrons to these orbitals places two in the lowest energy bonding MO and one each in the two nonbonding MOs.
- Separating electrons in two degenerate MOs keeps like charges farther away from each other.

Conclusion: Cyclobutadiene is not aromatic because its HOMOs, two degenerate nonbonding MOs, are not completely filled.

The procedure followed in Sample Problem 17.1 also illustrates why cyclobutadiene is antiaromatic. Having the two unpaired electrons in nonbonding MOs suggests that cyclobutadiene should be a highly unstable diradical. In fact, antiaromatic compounds resemble cyclobutadiene because their HOMOs contain two unpaired electrons, making them especially unstable.

Problem 17.23 Use the inscribed polygon method to show why the following cation is aromatic:




Problem 17.24 Use the inscribed polygon method to show why the cyclopentadienyl cation and radical are not aromatic.


17.11 Buckminsterfullerene—Is It Aromatic?

The two most common elemental forms of carbon are diamond and graphite. Diamond, one of the hardest substances known, is used for industrial cutting tools, whereas graphite, a slippery black substance, is used as a lubricant. Their physical characteristics are so different because their molecular structures are very different.

The structure of diamond consists of a continuous tetrahedral network of sp^3 hybridized carbon atoms, thus creating an infinite array of chair cyclohexane rings (without the hydrogen atoms). The structure of graphite, on the other hand, consists of parallel sheets of sp^2 hybridized carbon atoms, thus creating an infinite array of benzene rings. The parallel sheets are then held together by weak intermolecular interactions.



diamond
an “infinite” array of six-membered rings,
covalently bonded in three dimensions



graphite
an “infinite” array of benzene rings,
covalently bonded in two dimensions

Three sheets of graphite,
viewed edge-on



Graphite exists in planar sheets
of benzene rings, held together
by weak intermolecular forces.



Buckminsterfullerene (or buckyball) was discovered by Smalley, Curl, and Kroto, who shared the 1996 Nobel Prize in Chemistry for their work. Its unusual name stems from its shape, which resembles the geodesic dome invented by R. Buckminster Fuller. The pattern of five- and six-membered rings also resembles the pattern of rings on a soccer ball.

Buckminsterfullerene (C_{60}) is a third elemental form of carbon. Its structure consists of 20 hexagons and 12 pentagons of sp^2 hybridized carbon atoms joined in a spherical arrangement. It is completely conjugated because each carbon atom has a p orbital with an electron in it.



buckminsterfullerene, C_{60}


20 hexagons + 12 pentagons
of carbon atoms joined together

[The 60 C's of buckminsterfullerene are drawn. Each C also
contains a p orbital with one electron, which is not drawn.]

Is C_{60} aromatic? Although it is completely conjugated, it is not planar. Because of its curvature, it is not as stable as benzene. In fact, it undergoes addition reactions with electrophiles in much the same way as ordinary alkenes. Benzene, on the other hand, undergoes substitution reactions with electrophiles, which preserves the unusually stable benzene ring intact. These reactions are the subject of Chapter 18.

Problem 17.25 How many ^{13}C NMR signals does C_{60} exhibit?

Diamond and graphite are two
elemental forms of carbon.



KEY CONCEPTS

Benzene and Aromatic Compounds


Comparing Aromatic, Antiaromatic, and Nonaromatic Compounds (17.7)

- **Aromatic compound**
 - A cyclic, planar, completely conjugated compound that contains $4n + 2 \pi$ electrons ($n = 0, 1, 2, 3,$ and so forth).
 - An aromatic compound is more stable than a similar acyclic compound having the same number of π electrons.
- **Antiaromatic compound**
 - A cyclic, planar, completely conjugated compound that contains $4n \pi$ electrons ($n = 0, 1, 2, 3,$ and so forth).
 - An antiaromatic compound is less stable than a similar acyclic compound having the same number of π electrons.
- **Nonaromatic compound**
 - A compound that lacks one (or more) of the four requirements to be aromatic or antiaromatic.

Properties of Aromatic Compounds

- Every atom in the ring has a p orbital to delocalize electron density (17.2).
- They are unusually stable. ΔH° for hydrogenation is much less than expected, given the number of degrees of unsaturation (17.6).
- They do not undergo the usual addition reactions of alkenes (17.6).
- ^1H NMR spectra show highly deshielded protons because of ring currents that reinforce the applied magnetic field (17.4).
- All bonding MOs and HOMOs are completely filled and no electrons occupy antibonding orbitals (17.9).

Examples of Aromatic Compounds with Six π Electrons (17.8)



Examples of Compounds That Are Not Aromatic (17.8)











PROBLEMS

Benzene Structure and Nomenclature

- 17.26** Early structural studies on benzene had to explain the following experimental evidence. When benzene was treated with Br_2 (plus a Lewis acid), a single substitution product of molecular formula $\text{C}_6\text{H}_5\text{Br}$ was formed. When this product was treated with another equivalent of Br_2 , three different compounds of molecular formula $\text{C}_6\text{H}_4\text{Br}_2$ were formed.
- Explain why a single Kekulé structure is consistent with the first result, but does not explain the second result.
 - Then explain why a resonance description of benzene is consistent with the results of both reactions.
- 17.27** Draw all aromatic hydrocarbons that have molecular formula C_9H_{12} . Give the IUPAC name for each compound.
- 17.28** Draw all aromatic hydrocarbons that have molecular formula C_8H_{10} . For each compound, determine how many isomers of molecular formula $\text{C}_8\text{H}_9\text{Br}$ would be formed if one H atom on the benzene ring were replaced by a Br atom.

17.29 Give the IUPAC name for each compound.



17.30 Draw a structure corresponding to each name.






- a. *p*-dichlorobenzene
 b. *m*-chlorophenol
 c. *p*-iodoaniline
 d. *o*-bromonitrobenzene

- e. 2,6-dimethoxytoluene
 f. 2-phenyl-1-butene
 g. 2-phenyl-2-propen-1-ol
 h. *trans*-1-benzyl-3-phenylcyclopentane






17.31 a. Draw the 14 constitutional isomers of molecular formula C_8H_9Cl that contain a benzene ring.
 b. Name all compounds that contain a trisubstituted benzene ring.
 c. For which compound(s) are stereoisomers possible? Draw all possible stereoisomers.

Aromaticity









17.32 How many π electrons are contained in each molecule?






17.33 Which compounds are aromatic? For any compound that is not aromatic, state why this is so.




17.34 Which of the following heterocycles are aromatic?




17.35 Label each compound as aromatic, antiaromatic, or not aromatic. Assume all completely conjugated rings are planar.




- 17.36 Hydrocarbon **A** possesses a significant dipole, even though it is composed of only C–C and C–H bonds. Explain why the dipole arises and use resonance structures to illustrate the direction of the dipole. Which ring is more electron rich?

**A**


- 17.37 Pentalene, azulene, and heptalene are conjugated hydrocarbons that do not contain a benzene ring. Which hydrocarbons are especially stable or unstable based on the number of π electrons they contain? Explain your choices.



pentalene




azulene




heptalene

- 17.38 Rank the indicated C–C bonds in order of increasing bond length, and explain why you chose this order.




- 17.39 The purine heterocycle occurs commonly in the structure of DNA.



purine


- How is each N atom hybridized?
- In what type of orbital does each lone pair on a N atom reside?
- How many π electrons does purine contain?
- Why is purine aromatic?

- 17.40


**C**

- How many π electrons does **C** contain?
- How many π electrons are delocalized in the ring?
- Explain why **C** is aromatic.


- 17.41 Explain the observed rate of reactivity of the following 2° alkyl halides in an S_N1 reaction.



- 17.42 Draw a stepwise mechanism for the following reaction.




- 17.43 Explain why α -pyrone reacts with Br_2 to yield a substitution product (like benzene does), rather than an addition product to one of its C=C bonds.


 α -pyrone



Resonance

17.44 Draw additional resonance structures for each species.



17.45 The carbon–carbon bond lengths in naphthalene are not equal. Use a resonance argument to explain why bond (a) is shorter than bond (b).



17.46  pyrrole  furan

a. Draw all reasonable resonance structures for pyrrole and explain why pyrrole is less resonance stabilized than benzene.


b. Draw all reasonable resonance structures for furan and explain why furan is less resonance stabilized than pyrrole.

Acidity

17.47 Which compound in each pair is the stronger acid?



17.48 Treatment of indene with NaNH_2 forms its conjugate base in a Brønsted–Lowry acid–base reaction. Draw all reasonable resonance structures for indene's conjugate base, and explain why the $\text{p}K_a$ of indene is lower than the $\text{p}K_a$ of most hydrocarbons.




17.49 Considering both 5-methyl-1,3-cyclopentadiene (**A**) and 7-methyl-1,3,5-cycloheptatriene (**B**), which labeled H atom is most acidic? Which labeled H atom is least acidic? Explain your choices.




17.50 Draw the conjugate bases of pyrrole and cyclopentadiene. Explain why the sp^3 hybridized C–H bond of cyclopentadiene is more acidic than the N–H bond of pyrrole.

17.51 a. Explain why protonation of pyrrole occurs at C2 to form **A**, rather than on the N atom to form **B**.
b. Explain why **A** is more acidic than **C**, the conjugate acid of pyridine.




Inscribed Polygon Method

17.52 Use the inscribed polygon method to show the pattern of molecular orbitals in cyclooctatetraene.




- Label the MOs as bonding, antibonding, or nonbonding.
 - Indicate the arrangement of electrons in these orbitals for cyclooctatetraene, and explain why cyclooctatetraene is not aromatic.
 - Treatment of cyclooctatetraene with potassium forms a dianion. How many π electrons does this dianion contain?
 - How are the π electrons in this dianion arranged in the molecular orbitals?
 - Classify the dianion of cyclooctatetraene as aromatic, antiaromatic, or not aromatic, and explain why this is so.
- 17.53 Use the inscribed polygon method to show the pattern of molecular orbitals in 1,3,5,7-cyclononatetraene and use it to label its cation, radical, and anion as aromatic, antiaromatic, or not aromatic.



Spectroscopy

17.54 How many ^{13}C NMR signals does each compound exhibit?



17.55 Which of the diethylbenzene isomers (ortho, meta, or para) corresponds to each set of ^{13}C NMR spectral data?


[A] ^{13}C NMR signals: 16, 29, 125, 127.5, 128.4, and 144 ppm

[B] ^{13}C NMR signals: 15, 26, 126, 128, and 142 ppm


[C] ^{13}C NMR signals: 16, 29, 128, and 141 ppm

17.56 Propose a structure consistent with each set of data.


a. $\text{C}_{10}\text{H}_{14}$: IR absorptions at 3150–2850, 1600, and 1500 cm^{-1}



b. C_9H_{12} : ^{13}C NMR signals at 21, 127, and 138 ppm



c. C_8H_{10} : IR absorptions at 3108–2875, 1606, and 1496 cm^{-1}



17.57 Propose a structure consistent with each set of data.

a. Compound **A**:

Molecular formula: $C_8H_{10}O$

IR absorption at 3150–2850 cm^{-1}

1H NMR data: 1.4 (triplet, 3 H), 3.95 (quartet, 2 H), and 6.8–7.3 (multiplet, 5 H) ppm


b. Compound **B**:

Molecular formula: $C_9H_{10}O_2$


IR absorption at 1669 cm^{-1}

1H NMR data: 2.5 (singlet, 3 H), 3.8 (singlet, 3 H), 6.9 (doublet, 2 H), and 7.9 (doublet, 2 H) ppm


17.58 Thymol (molecular formula $C_{10}H_{14}O$) is the major component of the oil of thyme. Thymol shows IR absorptions at 3500–3200, 3150–2850, 1621, and 1585 cm^{-1} . The 1H NMR spectrum of thymol is given below. Propose a possible structure for thymol.



- 17.59** [7]-Paracyclophane is an unusual aromatic compound with a bridge connecting two para carbons. Explain why the labeled protons absorb in different regions of the ^1H NMR spectrum, even though both are bonded to sp^3 hybridized C atoms.




- 17.60** You have a sample of a compound of molecular formula $\text{C}_{11}\text{H}_{15}\text{NO}_2$, which has a benzene ring substituted by two groups, $(\text{CH}_3)_2\text{N}-$ and $-\text{CO}_2\text{CH}_2\text{CH}_3$, and exhibits the given ^{13}C NMR. What disubstituted benzene isomer corresponds to these ^{13}C data?




General Problems

- 17.61** Answer the following questions about curcumin, a yellow pigment isolated from turmeric, a tropical perennial in the ginger family and a principal ingredient in curry powder.




- In Chapter 11 we learned that most enols, compounds that contain a hydroxy group bonded to a $\text{C}=\text{C}$, are unstable and tautomerize to carbonyl groups. Draw the keto form of the enol of curcumin, and explain why the enol is more stable than many other enols.
 - Explain why the enol $\text{O}-\text{H}$ proton is more acidic than an alcohol $\text{O}-\text{H}$ proton.
 - Why is curcumin colored?
 - Explain why curcumin is an antioxidant.
- 17.62** Stanozolol is an anabolic steroid that promotes muscle growth. Although stanozolol has been used by athletes and body builders, many physical and psychological problems result from prolonged use and it is banned in competitive sports.




- Explain why the nitrogen heterocycle—a pyrazole ring—is aromatic.
- In what type of orbital is the lone pair on each N atom contained?
- Draw all reasonable resonance structures for stanozolol.
- Explain why the pK_a of the $\text{N}-\text{H}$ bond in the pyrazole ring is comparable to the pK_a of the $\text{O}-\text{H}$ bond, making it considerably more acidic than amines such as CH_3NH_2 ($pK_a = 40$).

Challenge Problems


17.63 Explain why compound **A** is much more stable than compound **B**.



17.64 (*R*)-Carvone, the major component of the oil of spearmint, undergoes acid-catalyzed isomerization to carvacrol, a major component of the oil of thyme. Draw a stepwise mechanism and explain why this isomerization occurs.



17.65 Explain why triphenylene resembles benzene in that it does not undergo addition reactions with Br_2 , but phenanthrene reacts with Br_2 to yield the addition product drawn. (Hint: Draw resonance structures for both triphenylene and phenanthrene, and use them to determine how delocalized each π bond is.)



17.66 Although benzene itself absorbs at 128 ppm in its ^{13}C NMR spectrum, the carbons of substituted benzenes absorb either upfield or downfield from this value depending on the substituent. Explain the observed values for the carbon ortho to the given substituent in the monosubstituted benzene derivatives **X** and **Y**.



Electrophilic Aromatic Substitution

18

- 18.1 Electrophilic aromatic substitution
- 18.2 The general mechanism
- 18.3 Halogenation
- 18.4 Nitration and sulfonation
- 18.5 Friedel–Crafts alkylation and Friedel–Crafts acylation
- 18.6 Substituted benzenes
- 18.7 Electrophilic aromatic substitution of substituted benzenes
- 18.8 Why substituents activate or deactivate a benzene ring
- 18.9 Orientation effects in substituted benzenes
- 18.10 Limitations on electrophilic substitution reactions with substituted benzenes
- 18.11 Disubstituted benzenes
- 18.12 Synthesis of benzene derivatives
- 18.13 Halogenation of alkyl benzenes
- 18.14 Oxidation and reduction of substituted benzenes
- 18.15 Multistep synthesis



LSD, commonly referred to as “acid,” is a powerful hallucinogen prepared from lysergic acid, the principal organic compound derived from one of the ergot fungi. Immortalized in the 1967 Beatles’ song, “Lucy in the Sky with Diamonds,” LSD produces sensory illusions, making it difficult for the user to distinguish between reality and fantasy. Given its potent biological properties, LSD has been the target of several different laboratory syntheses. A key step in one of them involves carbon–carbon bond formation using electrophilic aromatic substitution, the most common reaction of aromatic compounds and the subject of Chapter 18.

Chapter 18 discusses the chemical reactions of benzene and other aromatic compounds. Although aromatic rings are unusually stable, making benzene unreactive in most of the reactions discussed so far, benzene acts as a nucleophile with certain electrophiles, yielding substitution products with an intact aromatic ring.


We begin with the basic features and mechanism of electrophilic aromatic substitution (Sections 18.1–18.5), the basic reaction of benzene. Next, we discuss the electrophilic aromatic substitution of substituted benzenes (Sections 18.6–18.12), and conclude with other useful reactions of benzene derivatives (Sections 18.13–18.14). The ability to interconvert resonance structures and evaluate their relative stabilities is crucial to understanding this material.

18.1 Electrophilic Aromatic Substitution


Based on its structure and properties, what kinds of reactions should benzene undergo? Are any of its bonds particularly weak? Does it have electron-rich or electron-deficient atoms?

- Benzene has six π electrons delocalized in six p orbitals that overlap above and below the plane of the ring. These loosely held π electrons make the benzene ring electron rich, and so it reacts with electrophiles.
- Because benzene's six π electrons satisfy Hückel's rule, benzene is especially stable. Reactions that keep the aromatic ring intact are therefore favored.

As a result, **the characteristic reaction of benzene is *electrophilic aromatic substitution*—a hydrogen atom is replaced by an electrophile.**



Benzene does *not* undergo addition reactions like other unsaturated hydrocarbons, because addition would yield a product that is not aromatic. Substitution of a hydrogen, on the other hand, keeps the aromatic ring intact.



Five specific examples of electrophilic aromatic substitution are shown in Figure 18.1. The basic mechanism, discussed in Section 18.2, is the same in all five cases. The reactions differ only in the identity of the electrophile, E^+ .


Problem 18.1 Why is benzene less reactive towards electrophiles than an alkene, even though it has more π electrons than an alkene (six versus two)?

18.2 The General Mechanism

No matter what electrophile is used, all electrophilic aromatic substitution reactions occur via a **two-step mechanism**: addition of the electrophile E^+ to form a resonance-stabilized carbocation, followed by deprotonation with base, as shown in Mechanism 18.1.

Figure 18.1

Five examples of electrophilic aromatic substitution




Friedel–Crafts alkylation and acylation, named for Charles Friedel and James Crafts who discovered the reactions in the nineteenth century, form new carbon–carbon bonds.




Mechanism 18.1 General Mechanism—Electrophilic Aromatic Substitution

Step [1] Addition of the electrophile (E^+) to form a carbocation



- Addition of the electrophile (E^+) forms a new C–E bond using two π electrons from the benzene ring, and generating a carbocation. This carbocation intermediate is not aromatic, but it is resonance stabilized—**three resonance structures can be drawn**.
- Step [1] is rate-determining because the aromaticity of the benzene ring is lost.


Step [2] Loss of a proton to re-form the aromatic ring



- In Step [2], a base ($B:$) removes the proton from the carbon bearing the electrophile, thus re-forming the aromatic ring. This step is fast because the aromaticity of the benzene ring is restored.
- Any of the three resonance structures of the carbocation intermediate can be used to draw the product. The choice of resonance structure affects how curved arrows are drawn, but not the identity of the product.

The first step in electrophilic aromatic substitution forms a carbocation, for which three resonance structures can be drawn. To help keep track of the location of the positive charge:

- Always draw in the H atom on the carbon bonded to E. This serves as a reminder that it is the only sp^3 hybridized carbon in the carbocation intermediate.
- Notice that the positive charge in a given resonance structure is always located ortho or para to the new C–E bond. In the hybrid, therefore, the charge is delocalized over three atoms of the ring.



This two-step mechanism for electrophilic aromatic substitution applies to all of the electrophiles in Figure 18.1. **The net result of addition of an electrophile (E^+) followed by elimination of a proton (H^+) is substitution of E for H.**

The energy changes in electrophilic aromatic substitution are shown in Figure 18.2. The mechanism consists of two steps, so the energy diagram has two energy barriers. Because the first step is rate-determining, its transition state is higher in energy.


Problem 18.2

In Step [2] of Mechanism 18.1, loss of a proton to form the substitution product was drawn using one resonance structure only. Use curved arrows to show how the other two resonance structures can be converted to the substitution product (PhE) by removal of a proton with $:B$.

18.3 Halogenation

The general mechanism outlined in Mechanism 18.1 can now be applied to each of the five specific examples of electrophilic aromatic substitution shown in Figure 18.1. For each mechanism we must learn how to generate a specific electrophile. This step is *different* with each electro-


Figure 18.2
Energy diagram for electrophilic aromatic substitution:
 $PhH + E^+ \rightarrow PhE + H^+$




- The mechanism has two steps, so there are two energy barriers.
- Step [1] is rate-determining; its transition state is at higher energy.

Figure 18.3


Examples of biologically active aryl chlorides




Generic name: **bupropion**
Trade names: **Wellbutrin, Zyban**
antidepressant,
also used to reduce nicotine cravings



chlorpheniramine
antihistamine



2,4-D
2,4-dichlorophenoxy-
acetic acid
herbicide



2,4,5-T
2,4,5-trichlorophenoxy-
acetic acid
herbicide


the active components in **Agent Orange**,
a defoliant used in the Vietnam War

Herbicides were used extensively during the Vietnam War to defoliate dense jungle areas. The concentration of certain herbicide by-products in the soil remains high today.



18.4 Nitration and Sulfonation


Nitration and **sulfonation** of benzene introduce two different functional groups on an aromatic ring. Nitration is an especially useful reaction because a nitro group can then be reduced to an NH_2 group, a common benzene substituent, in a reaction discussed in Section 18.14.



Generation of the electrophile in both nitration and sulfonation requires strong acid. In **nitration**, the electrophile is $^+\text{NO}_2$ (the **nitronium ion**), formed by protonation of HNO_3 followed by loss of water (Mechanism 18.3).




Mechanism 18.3 Formation of the Nitronium Ion ($^+\text{NO}_2$) for Nitration



In **sulfonation**, protonation of sulfur trioxide, SO_3 , forms a positively charged sulfur species ($^+\text{SO}_3\text{H}$) that acts as an electrophile (Mechanism 18.4).




Mechanism 18.4 Formation of the Electrophile $^+\text{SO}_3\text{H}$ for Sulfonation



These steps illustrate how to generate the electrophile E^+ for nitration and sulfonation, the process that begins any mechanism for electrophilic aromatic substitution. To complete either of these mechanisms, you must replace the electrophile E^+ by either $^+\text{NO}_2$ or $^+\text{SO}_3\text{H}$ in the general mechanism (Mechanism 18.1). Thus, **the two-step sequence that replaces H by E is the same regardless of E^+** . This is shown in Sample Problem 18.1 using the reaction of benzene with the nitronium ion.


Sample Problem 18.1 Draw a stepwise mechanism for the nitration of a benzene ring.




Solution

We must first generate the electrophile and then write the two-step mechanism for electrophilic aromatic substitution using it.

Generation of the electrophile $^+\text{NO}_2$




Two-step mechanism for substitution



Any species with a lone pair of electrons can be used to remove the proton in the last step. In this case, the mechanism is drawn with HSO_4^- , formed when $^+\text{NO}_2$ is generated as the electrophile.

Problem 18.4

Draw a stepwise mechanism for the sulfonation of an alkyl benzene such as **A** to form a substituted benzenesulfonic acid **B**. Treatment of **B** with base forms a sodium salt **C** that can be used as a synthetic detergent to clean away dirt (see Problem 3.15).






18.5 Friedel–Crafts Alkylation and Friedel–Crafts Acylation



Friedel–Crafts alkylation and **Friedel–Crafts acylation** form new carbon–carbon bonds.

18.5A General Features

In **Friedel–Crafts alkylation**, treatment of benzene with an alkyl halide and a Lewis acid (AlCl_3) forms an alkyl benzene. This reaction is an **alkylation** because it results in transfer of an alkyl group from one atom to another (from Cl to benzene).



**Friedel–Crafts alkylation—
General reaction****Examples**

In **Friedel–Crafts acylation**, a benzene ring is treated with an **acid chloride** (RCOCl) and AlCl_3 to form a ketone. Because the new group bonded to the benzene ring is called an **acyl group**, the transfer of an acyl group from one atom to another is an **acylation**.




**Friedel–Crafts acylation—
General reaction****Example**

Acid chlorides are also called **acyl chlorides**.

Problem 18.5 What product is formed when benzene is treated with each organic halide in the presence of AlCl_3 ?



Problem 18.6 What acid chloride would be needed to prepare each of the following ketones from benzene using a Friedel–Crafts acylation?


**18.5B Mechanism**

The mechanisms of alkylation and acylation proceed in a manner analogous to those for halogenation, nitration, and sulfonation. The unique feature in each reaction is how the electrophile is generated.

In **Friedel–Crafts alkylation**, the Lewis acid AlCl_3 reacts with the alkyl chloride to form a Lewis acid–base complex, illustrated with $\text{CH}_3\text{CH}_2\text{Cl}$ and $(\text{CH}_3)_3\text{CCl}$ as alkyl chlorides. The identity of the alkyl chloride determines the exact course of the reaction as shown in Mechanism 18.5.



Mechanism 18.5 Formation of the Electrophile in Friedel–Crafts Alkylation—Two Possibilities




- For CH_3Cl and 1°RCl , the Lewis acid–base complex itself serves as the electrophile for electrophilic aromatic substitution.
- With 2° and 3°RCl , the Lewis acid–base complex reacts further to give a 2° or 3° carbocation, which serves as the electrophile. Carbocation formation occurs only with 2° and 3° alkyl chlorides, because they afford more stable carbocations.

In either case, the electrophile goes on to react with benzene in the two-step mechanism characteristic of electrophilic aromatic substitution, illustrated in Mechanism 18.6 using the 3° carbocation, $(\text{CH}_3)_3\text{C}^+$.



Mechanism 18.6 Friedel–Crafts Alkylation Using a 3° Carbocation




- Addition of the electrophile (a 3° carbocation) forms a new carbon–carbon bond in Step [1].
- AlCl_4^- removes a proton on the carbon bearing the new substituent, thus re-forming the aromatic ring in Step [2].

In **Friedel–Crafts acylation**, the Lewis acid AlCl_3 ionizes the carbon–halogen bond of the acid chloride, thus forming a positively charged carbon electrophile called an **acylium ion**, which is resonance stabilized (Mechanism 18.7). The positively charged carbon atom of the acylium ion then goes on to react with benzene in the two-step mechanism of electrophilic aromatic substitution.




Mechanism 18.7 Formation of the Electrophile in Friedel–Crafts Acylation




To complete the mechanism for acylation, insert the electrophile into the general mechanism and draw the last two steps, as illustrated in Sample Problem 18.2.

Sample Problem 18.2 Draw a stepwise mechanism for the following Friedel–Crafts acylation.



Solution

First generate the acylium ion, and then write the two-step mechanism for electrophilic aromatic substitution using it for the electrophile.



Problem 18.7 Draw a stepwise mechanism for the Friedel–Crafts alkylation of benzene with $\text{CH}_3\text{CH}_2\text{Cl}$ and AlCl_3 .


18.5C Other Facts About Friedel–Crafts Alkylation

Three additional facts about Friedel–Crafts alkylations must be kept in mind.


[1] Vinyl halides and aryl halides do *not* react in Friedel–Crafts alkylation.

Most Friedel–Crafts reactions involve carbocation electrophiles. Because the carbocations derived from vinyl halides and aryl halides are highly unstable and do not readily form, these organic halides do *not* undergo Friedel–Crafts alkylation.

Unreactive halides in the Friedel–Crafts alkylation







vinyl halide



aryl halide


Problem 18.8 Which halides are unreactive in a Friedel–Crafts alkylation reaction?



[2] Rearrangements can occur.

The Friedel–Crafts reaction can yield products having rearranged carbon skeletons when 1° and 2° alkyl halides are used as starting materials, as shown in Equations [1] and [2]. In both reactions, the carbon atom bonded to the halogen in the starting material (labeled in red) is not bonded to the benzene ring in the product, thus indicating that a rearrangement has occurred.

Recall from Section 9.9 that a 1,2-shift converts a less stable carbocation to a more stable carbocation by shift of a hydrogen atom or an alkyl group.




The result in Equation [1] is explained by a carbocation rearrangement involving a 1,2-hydride shift: **the less stable 2° carbocation (formed from the 2° halide) rearranges to a more stable 3° carbocation**, as illustrated in Mechanism 18.8.




Mechanism 18.8 Friedel–Crafts Alkylation Involving Carbocation Rearrangement

Steps [1] and [2] Formation of a 2° carbocation




- Reaction of the alkyl chloride with AlCl₃ forms a complex that decomposes in Step [2] to form a **2° carbocation**.

Step [3] Carbocation rearrangement



- **1,2-Hydride shift** converts the less stable 2° carbocation to a more stable 3° carbocation.

Steps [4] and [5] Addition of the carbocation and loss of a proton




- Friedel–Crafts alkylation occurs by the usual two-step process: **addition of the carbocation** followed by **loss of a proton** to form the alkylated product.


Rearrangements can occur even when no free carbocation is formed initially. For example, the 1° alkyl chloride in Equation [2] forms a complex with AlCl₃, which does *not* decompose to an unstable 1° carbocation, as shown in Mechanism 18.9. Instead, a **1,2-hydride shift** forms a 2° carbocation, which then serves as the electrophile in the two-step mechanism for electrophilic aromatic substitution.



Mechanism 18.9 A Rearrangement Reaction Beginning with a 1° Alkyl Chloride



Problem 18.9 Draw a stepwise mechanism for the following reaction.





Problem 18.10 Offer an explanation as to why rearrangements do not occur with the acylium ion formed in a Friedel–Crafts acylation reaction.

[3] Other functional groups that form carbocations can also be used as starting materials.


Although Friedel–Crafts alkylation works well with alkyl halides, any compound that readily forms a carbocation can be used instead. The two most common alternatives are alkenes and alcohols, both of which afford carbocations in the presence of strong acid.

- Protonation of an alkene forms a carbocation, which can then serve as an electrophile in a Friedel–Crafts alkylation.


- Protonation of an alcohol, followed by loss of water, likewise forms a carbocation.



Each carbocation can then go on to react with benzene to form a product of electrophilic aromatic substitution. For example:



Problem 18.11 Draw the product of each reaction.




18.5D Intramolecular Friedel–Crafts Reactions

All of the Friedel–Crafts reactions discussed thus far have resulted from intermolecular reaction of a benzene ring with an electrophile. Starting materials that contain both units are capable of **intramolecular reaction**, and this forms a new ring. For example, treatment of compound **A**,

Figure 18.4


Intramolecular Friedel–Crafts acylation in the synthesis of LSD



Sertraline (Problem 18.12) is an SSRI—selective serotonin reuptake inhibitor. Antidepressants of this type act by increasing the concentration of serotonin (Section 25.6C), a compound that plays a key role in mood, sleep, perception, and temperature regulation.

- Intramolecular Friedel–Crafts acylation formed a product containing a new six-membered ring (in red), which was converted to LSD in several steps.
- LSD was first prepared by Swiss chemist Albert Hoffman in 1938 from a related organic compound isolated from the ergot fungus that attacks rye and other grains. Ergot has a long history as a dreaded poison, affecting individuals who become ill from eating ergot-contaminated bread. The hallucinogenic effects of LSD were first discovered when Hoffman accidentally absorbed a small amount of the drug through his fingertips.


which contains both a benzene ring and an acid chloride, with AlCl_3 , forms α -tetralone by an intramolecular Friedel–Crafts acylation reaction.




Such an intramolecular Friedel–Crafts acylation was a key step in the synthesis of LSD, the molecule that introduced Chapter 18, as shown in Figure 18.4.

Problem 18.12

Draw a stepwise mechanism for the intramolecular Friedel–Crafts acylation of compound **A** to form **B**. **B** can be converted in one step to the antidepressant sertraline (trade name Zoloft).

**Problem 18.13**

Intramolecular reactions are also observed in Friedel–Crafts alkylation. Draw the intramolecular alkylation product formed from each of the following reactants. (Watch out for rearrangements!)



18.6 Substituted Benzenes

Many substituted benzene rings undergo electrophilic aromatic substitution. Common substituents include halogens, OH, NH₂, alkyl, and many functional groups that contain a carbonyl. Each substituent either increases or decreases the electron density in the benzene ring, and this affects the course of electrophilic aromatic substitution, as we will learn in Section 18.7.

What makes a substituent on a benzene ring electron donating or electron withdrawing? The answer is **inductive effects** and **resonance effects**, both of which can add or remove electron density.


Inductive Effects

Inductive effects stem from the **electronegativity** of the atoms in the substituent and the **polarizability** of the substituent group.

- Atoms more electronegative than carbon—including N, O, and X—pull electron density away from carbon and thus exhibit an electron-withdrawing inductive effect.
- Polarizable alkyl groups donate electron density, and thus exhibit an electron-donating inductive effect.


Considering inductive effects *only*, an NH₂ group withdraws electron density and CH₃ donates electron density.

Electron-withdrawing inductive effect



- N is **more electronegative** than C.
- N inductively withdraws electron density.

Electron-donating inductive effect



- Alkyl groups are **polarizable**, making them electron-donating groups.

Inductive and resonance effects were first discussed in Sections 2.5B and 2.5C, respectively.

Problem 18.14


Which substituents have an electron-withdrawing and which have an electron-donating inductive effect: (a) CH₃CH₂CH₂CH₂- ; (b) Br- ; (c) CH₃CH₂O-

Resonance Effects

Resonance effects can either donate or withdraw electron density, depending on whether they place a positive or negative charge on the benzene ring.


- A resonance effect is electron donating when resonance structures place a negative charge on carbons of the benzene ring.
- A resonance effect is electron withdrawing when resonance structures place a positive charge on carbons of the benzene ring.

An electron-donating resonance effect is observed whenever an atom Z having a lone pair of electrons is directly bonded to a benzene ring (general structure—C₆H₅—Z:). Common examples of Z include N, O, and halogen. For example, five resonance structures can be drawn for aniline (C₆H₅NH₂). Because three of them place a negative charge on a carbon atom of the benzene ring, an NH₂ group donates electron density to a benzene ring by a resonance effect.




In contrast, an electron-withdrawing resonance effect is observed in substituted benzenes having the general structure C₆H₅—Y=Z, where Z is more electronegative than Y. For exam-

ple, seven resonance structures can be drawn for benzaldehyde (C_6H_5CHO). Because three of them place a positive charge on a carbon atom of the benzene ring, a CHO group withdraws electron density from a benzene ring by a resonance effect.



Problem 18.15

Draw all resonance structures for each compound and use the resonance structures to determine if the substituent has an electron-donating or electron-withdrawing resonance effect.




Considering Both Inductive and Resonance Effects

To predict whether a substituted benzene is more or less electron rich than benzene itself, we must consider the **net balance of both the inductive and the resonance effects**. Alkyl groups, for instance, donate electrons by an inductive effect, but they have no resonance effect because they lack nonbonded electron pairs or π bonds. As a result,

- An alkyl group is an electron-donating group and an alkyl benzene is more electron rich than benzene.

When electronegative atoms, such as N, O, or halogen, are bonded to the benzene ring, they inductively *withdraw* electron density from the ring. All of these groups also have a nonbonded pair of electrons, so they *donate* electron density to the ring by resonance. **The identity of the element determines the net balance of these opposing effects.**



- When a neutral O or N atom is bonded directly to a benzene ring, the resonance effect dominates and the net effect is electron donation.
- When a halogen X is bonded to a benzene ring, the inductive effect dominates and the net effect is electron withdrawal.


Thus, NH_2 and OH are electron-donating groups because the resonance effect predominates, whereas Cl and Br are electron-withdrawing groups because the inductive effect predominates.

Finally, the inductive and resonance effects in compounds having the general structure $C_6H_5-Y=Z$ (with Z more electronegative than Y) are **both electron withdrawing**; in other words, the two effects *reinforce* each other. This is true for benzaldehyde (C_6H_5CHO) and all other compounds that contain a carbonyl group bonded directly to the benzene ring.

Thus, on balance, an **NH_2 group is electron donating**, so the benzene ring of aniline ($C_6H_5NH_2$) has more electron density than benzene. An **aldehyde group (CHO), on the other hand, is**


Figure 18.5

The effect of substituents on the electron density in substituted benzenes



- The NH_2 group donates electron density, making the benzene ring more electron rich (redder), whereas the CHO group withdraws electron density, making the benzene ring less electron rich (greener).

electron withdrawing, so the benzene ring of benzaldehyde (C_6H_5CHO) has less electron density than benzene. These effects are illustrated in the electrostatic potential maps in Figure 18.5. These compounds represent examples of the general structural features in electron-donating and electron-withdrawing substituents:




- Common electron-donating groups are alkyl groups or groups with an N or O atom (with a lone pair) bonded to the benzene ring.
- Common electron-withdrawing groups are halogens or groups with an atom Y bearing a full or partial positive charge (+ or δ^+) bonded to the benzene ring.

The net effect of electron donation and withdrawal on the reactions of substituted aromatics is discussed in Sections 18.7–18.9.


Sample Problem 18.3

Classify each substituent as electron donating or electron withdrawing.




Solution

Draw out the atoms and bonds of the substituent to clearly see lone pairs and multiple bonds. Always look at the atom bonded directly to the benzene ring to determine electron-donating or electron-withdrawing effects. An O or N atom with a lone pair of electrons makes a substituent electron donating. A halogen or an atom with a partial positive charge makes a substituent electron withdrawing.



Problem 18.16 Classify each substituent as electron donating or electron withdrawing.



18.7 Electrophilic Aromatic Substitution of Substituted Benzenes


Electrophilic aromatic substitution is a general reaction of *all* aromatic compounds, including polycyclic aromatic hydrocarbons, heterocycles, and substituted benzene derivatives. A substituent affects two aspects of electrophilic aromatic substitution:

- **The rate of reaction:** A substituted benzene reacts faster or slower than benzene itself.
- **The orientation:** The new group is located either ortho, meta, or para to the existing substituent. The identity of the first substituent determines the position of the second substituent.

Toluene ($C_6H_5CH_3$) and nitrobenzene ($C_6H_5NO_2$) illustrate two possible outcomes.


[1] Toluene

Toluene reacts **faster** than benzene in all substitution reactions. Thus, its **electron-donating CH_3 group activates the benzene ring** to electrophilic attack. Although three products are possible, compounds with the new group ortho or para to the CH_3 group predominate. The CH_3 group is therefore called an **ortho, para director**.



[2] Nitrobenzene


Nitrobenzene reacts **more slowly** than benzene in all substitution reactions. Thus, its **electron-withdrawing NO_2 group deactivates the benzene ring** to electrophilic attack. Although three products are possible, the compound with the new group meta to the NO_2 group predominates. The NO_2 group is called a **meta director**.



Substituents either activate or deactivate a benzene ring towards electrophiles, and direct selective substitution at specific sites on the ring. **All substituents can be divided into three general types.**

[1] Ortho, para directors and activators

- Substituents that *activate* a benzene ring and direct substitution ortho and para.




[2] Ortho, para deactivators

- Substituents that *deactivate* a benzene ring and direct substitution ortho and para.




[3] Meta directors

- Substituents that direct substitution meta.
- All meta directors *deactivate* the ring.




To learn these lists: **Keep in mind that the halogens are in a class by themselves.** Then learn the general structures for each type of substituent.

- All ortho, para directors are R groups or have a nonbonded electron pair on the atom bonded to the benzene ring.




- All meta directors have a full or partial positive charge on the atom bonded to the benzene ring.



Sample Problem 18.4 shows how this information can be used to predict the products of electrophilic aromatic substitution reactions.

Sample Problem 18.4


Draw the products of each reaction and state whether the reaction is faster or slower than a similar reaction with benzene.




Solution

To draw the products:

- Draw the Lewis structure for the substituent to see if it has a lone pair or partial positive charge on the atom bonded to the benzene ring.
- Classify the substituent—ortho, para activating, ortho, para deactivating, or meta deactivating—and draw the products.




The lone pair on N makes this group an **ortho, para activator**.
This compound reacts faster than benzene.




The δ^+ on this C makes the group a **meta deactivator**.
This compound reacts more slowly than benzene.

Problem 18.17 Draw the products of each reaction.



Problem 18.18 Draw the products formed when each compound is treated with HNO_3 and H_2SO_4 . State whether the reaction occurs faster or slower than a similar reaction with benzene.




18.8 Why Substituents Activate or Deactivate a Benzene Ring

- **Why do substituents activate or deactivate a benzene ring?**
- **Why are particular orientation effects observed?** Why are some groups ortho, para directors and some groups meta directors?

To understand why some substituents make a benzene ring react faster than benzene itself (activators), whereas others make it react slower (deactivators), we must evaluate the rate-determining step (the first step) of the mechanism. Recall from Section 18.2 that the first step in electrophilic aromatic substitution is the addition of an electrophile (E^+) to form a resonance-stabilized carbocation. The Hammond postulate (Section 7.15) makes it possible to predict the relative rate of the reaction by looking at the stability of the carbocation intermediate.

- The more stable the carbocation, the lower in energy the transition state that forms it, and the faster the reaction.



The principles of inductive effects and resonance effects, first introduced in Section 18.6, can now be used to predict carbocation stability.

- Electron-donating groups stabilize the carbocation and activate a benzene ring towards electrophilic attack.
- Electron-withdrawing groups destabilize the carbocation and deactivate a benzene ring towards electrophilic attack.

The energy diagrams in Figure 18.6 illustrate the effect of electron-donating and electron-withdrawing groups on the energy of the transition state of the rate-determining step in electrophilic aromatic substitution. From Section 18.6, we now know which groups increase or decrease the rate of reaction of substituted benzenes with electrophiles.

- All activators are either R groups or they have an N or O atom with a lone pair bonded directly to the benzene ring. These are the electron-donor groups of Section 18.6.






Figure 18.6 Energy diagrams comparing the rate of electrophilic aromatic substitution of substituted benzenes




- Electron-donor groups **D** stabilize the carbocation intermediate, lower the energy of the transition state, and increase the rate of reaction.
- Electron-withdrawing groups **W** destabilize the carbocation intermediate, raise the energy of the transition state, and decrease the rate of reaction.

- All deactivators are either halogens or they have an atom with a partial or full positive charge bonded directly to the benzene ring. These are the electron-withdrawing groups of Section 18.6.




Deactivating groups:			
$-\ddot{\text{F}}:$	$-\ddot{\text{Cl}}:$	$-\ddot{\text{Br}}:$	$-\ddot{\text{I}}:$
$-\text{CHO}$	$-\text{COR}$		
$-\text{COOR}$	$-\text{COOH}$		
$-\text{CN}$	$-\text{SO}_3\text{H}$	$-\text{NO}_2$	$-\text{NR}_3^+$

Problem 18.19 Label each compound as more or less reactive than benzene in electrophilic aromatic substitution.



Problem 18.20 Rank the compounds in each group in order of increasing reactivity in electrophilic aromatic substitution.



18.9 Orientation Effects in Substituted Benzenes


To understand why particular orientation effects arise, you must keep in mind the general structures for ortho, para directors and for meta directors already given in Section 18.7. There are two general types of ortho, para directors and one general type of meta director:

- All ortho, para directors are R groups or have a nonbonded electron pair on the atom bonded to the benzene ring.
- All meta directors have a full or partial positive charge on the atom bonded to the benzene ring.

To evaluate the directing effects of a given substituent, we can follow a stepwise procedure.

HOW TO Determine the Directing Effects of a Particular Substituent

Step [1] Draw all resonance structures for the carbocation formed from attack of an electrophile E^+ at the ortho, meta, and para positions of a substituted benzene ($\text{C}_6\text{H}_5 - \text{A}$).



- There are at least three resonance structures for each site of reaction.
- Each resonance structure places a positive charge ortho or para to the new C–E bond.


Step [2] Evaluate the stability of the intermediate resonance structures. The electrophile attacks at those positions that give the most stable carbocation.

Sections 18.9A–C show how this two-step procedure can be used to determine the directing effects of the CH_3 group in toluene, the NH_2 group in aniline, and the NO_2 group in nitrobenzene, respectively.

18.9A The CH₃ Group—An ortho, para Director

To determine why a **CH₃ group directs electrophilic aromatic substitution to the ortho and para positions**, first draw all resonance structures that result from electrophilic attack at the ortho, meta, and para positions to the CH₃ group.

Always draw in the H atom at the site of electrophilic attack. This will help you keep track of where the charges go.




Note that the positive charge in all resonance structures is always ortho or para to the new C–E bond. It is *not* necessarily ortho or para to the CH₃ group.

To evaluate the stability of the resonance structures, determine whether any are especially stable or unstable. In this example, **attack ortho or para to CH₃ generates a resonance structure that places a positive charge on a carbon atom with the CH₃ group**. The electron-donating CH₃ group *stabilizes* the adjacent positive charge. In contrast, attack meta to the CH₃ group does *not* generate any resonance structure stabilized by electron donation. Other alkyl groups are ortho, para directors for the same reason.

- **Conclusion:** The CH₃ group directs electrophilic attack ortho and para to itself because an electron-donating inductive effect stabilizes the carbocation intermediate.

18.9B The NH₂ Group—An ortho, para Director

To determine why an **amino group (NH₂) directs electrophilic aromatic substitution to the ortho and para positions**, follow the same procedure.




Attack at the meta position generates the usual three resonance structures. Because of the lone pair on the N atom, attack at the ortho and para positions generates a fourth resonance structure, which is stabilized because **every atom has an octet of electrons. This additional resonance structure can be drawn for all substituents that have an N, O, or halogen atom bonded directly to the benzene ring.**

- **Conclusion:** The NH_2 group directs electrophilic attack ortho and para to itself because the carbocation intermediate has additional resonance stabilization.

18.9C The NO_2 Group—A meta Director




To determine why a **nitro group (NO_2)** directs electrophilic aromatic substitution to the **meta position**, follow the same procedure.



Attack at each position generates three resonance structures. One resonance structure resulting from attack at the ortho and para positions is especially *destabilized*, because it contains a positive charge on two adjacent atoms. Attack at the meta position does not generate any particularly unstable resonance structures.

- **Conclusion:** With the NO_2 group (and all meta directors), meta attack occurs because attack at the ortho or para position gives a destabilized carbocation intermediate.


Problem 18.21 Draw all resonance structures for the carbocation formed by ortho attack of the electrophile $^+\text{NO}_2$ on each starting material. Label any resonance structures that are especially stable or unstable.



Problem 18.22 Use the procedure illustrated in Sections 18.9A–C to show why chlorine is an ortho, para director.

Figure 18.7 summarizes the reactivity and directing effects of the common substituents on benzene rings. You do not need to memorize this list. Instead, follow the general procedure outlined in Sections 18.9A–C to predict particular substituent effects.

Figure 18.7
The reactivity and directing effects of common substituted benzenes



In summary:


- [1] All ortho, para directors except the halogens activate the benzene ring.
- [2] All meta directors deactivate the benzene ring.
- [3] The halogens deactivate the benzene ring.

18.10 Limitations on Electrophilic Substitution Reactions with Substituted Benzenes


Although electrophilic aromatic substitution works well with most substituted benzenes, halogenation and the Friedel–Crafts reactions have some additional limitations that must be kept in mind.

18.10A Halogenation of Activated Benzenes


Considering all electrophilic aromatic substitution reactions, halogenation occurs the most readily. As a result, benzene rings activated by strong electron-donating groups—OH, NH₂, and their alkyl derivatives (OR, NHR, and NR₂)—undergo **polyhalogenation** when treated with X₂ and FeX₃. For example, aniline (C₆H₅NH₂) and phenol (C₆H₅OH) both give a tribromo derivative when treated with Br₂ and FeBr₃. Substitution occurs at all hydrogen atoms ortho and para to the NH₂ and OH groups.



Monosubstitution of H by Br occurs with Br₂ *alone* without added catalyst to form a mixture of ortho and para products.




Problem 18.23 Draw the products of each reaction.



18.10B Limitations in Friedel–Crafts Reactions


Friedel–Crafts reactions are the most difficult electrophilic aromatic substitution reactions to carry out in the laboratory. For example, they do not occur when the benzene ring is substituted with NO₂ (a strong deactivator) or with NH₂, NHR, or NR₂ (strong activators).

A benzene ring deactivated by a strong electron-withdrawing group—that is, any of the meta directors—is not electron rich enough to undergo Friedel–Crafts reactions.




Friedel–Crafts reactions also do not occur with NH₂ groups, which are strong activating groups. NH₂ groups are strong Lewis bases (due to the nonbonded electron pair on N), so they react with

AlCl_3 , the Lewis acid needed for alkylation or acylation. The resulting product contains a positive charge adjacent to the benzene ring, so the ring is now strongly deactivated and therefore unreactive in Friedel–Crafts reactions.




Problem 18.24 Which of the following compounds undergo Friedel–Crafts alkylation with CH_3Cl and AlCl_3 ? Draw the products formed when a reaction occurs.




Another limitation of the Friedel–Crafts alkylation arises because of **polyalkylation**. Treatment of benzene with an alkyl halide and AlCl_3 places an electron-donor R group on the ring. Because R groups activate a ring, the alkylated product ($\text{C}_6\text{H}_5\text{R}$) is now *more reactive* than benzene itself towards further substitution, and it reacts again with RCl to give products of polyalkylation.

To minimize polyalkylation a large excess of benzene is used relative to the amount of alkyl halide.



Polysubstitution does not occur with Friedel–Crafts acylation, because the product now has an electron-withdrawing group that deactivates the ring towards another electrophilic substitution.




18.11 Disubstituted Benzenes

What happens in electrophilic aromatic substitution when a disubstituted benzene ring is used as starting material? **To predict the products, look at the directing effects of both substituents and then determine the net result**, using the following three guidelines.

Rule [1] When the directing effects of two groups reinforce, the new substituent is located on the position directed by both groups.

For example, the CH_3 group in *p*-nitrotoluene is an ortho, para director and the NO_2 group is a meta director. These two effects reinforce each other so that one product is formed on treatment with Br_2 and FeBr_3 . Notice that the position para to the CH_3 group is “blocked” by a nitro group so no substitution can occur on that carbon.

ortho, para director




meta director

p-nitrotoluene

Rule [2] If the directing effects of two groups oppose each other, the more powerful activator “wins out.”

In compound **A**, the NHCOCH_3 group activates its two ortho positions, and the CH_3 group activates its two ortho positions to reaction with electrophiles. Because the NHCOCH_3 is a stronger activator, substitution occurs ortho to it.

stronger ortho, para director




weaker ortho, para director



A

Rule [3] No substitution occurs between two meta substituents because of crowding.

For example, no substitution occurs at the carbon atom between the two CH_3 groups in *m*-xylene, even though two CH_3 groups activate that position.




Sample Problem 18.5 Draw the products formed from nitration of each compound.




Solution


a. Both the OH and CH_3 groups are ortho, para directors. Because the OH group is a stronger activator, substitution occurs ortho to it.



- b. Both the OH and CH₃ groups are ortho, para directors whose directing effects reinforce each other in this case. No substitution occurs between the two meta substituents, however, so two products result.




Problem 18.25 Draw the products formed when each compound is treated with HNO₃ and H₂SO₄.



18.12 Synthesis of Benzene Derivatives


To synthesize benzene derivatives with more than one substituent, we must always take into account the directing effects of each substituent. In a disubstituted benzene, for example, **the directing effects indicate which substituent must be added to the ring first.**

For example, the Br group in *p*-bromonitrobenzene is an ortho, para director and the NO₂ group is a meta director. Because the two substituents are para to each other, the ortho, para director must be introduced *first* when synthesizing this compound from benzene.




Thus, Pathway [1], in which bromination precedes nitration, yields the desired para product, whereas Pathway [2], in which nitration precedes bromination, yields the undesired meta isomer.

Pathway [1]: Bromination before nitration



This pathway gives the desired product.


Pathway [2]: Nitration before bromination




This pathway does NOT form the desired product.

Pathway [1] yields both the desired para product as well as the undesired ortho isomer. Because these compounds are constitutional isomers, they are separable. Obtaining such a mixture of ortho and para isomers is often unavoidable.


Sample Problem 18.6 Devise a synthesis of *o*-nitrotoluene from benzene.

**Solution**

The CH_3 group in *o*-nitrotoluene is an ortho, para director and the NO_2 group is a meta director. Because the two substituents are ortho to each other, the **ortho, para director must be introduced first**. The synthesis thus involves two steps: Friedel–Crafts alkylation followed by nitration.



Problem 18.26 Devise a synthesis of each compound from the indicated starting material.




18.13 Halogenation of Alkyl Benzenes

Radical halogenation of alkanes was discussed in Chapter 15. The mechanism of radical halogenation at an allylic carbon was given in Section 15.10.


We finish Chapter 18 by learning some additional reactions of substituted benzenes that greatly expand the ability to synthesize benzene derivatives. These reactions do not involve the benzene ring itself, so they are not further examples of electrophilic aromatic substitution. In Section 18.13 we return to radical halogenation, and in Section 18.14 we examine useful oxidation and reduction reactions.

Benzylic C–H bonds are weaker than most other sp^3 hybridized C–H bonds, because homolysis forms a resonance-stabilized benzylic radical.



The bond dissociation energy for a benzylic C–H bond (356 kJ/mol) is even less than the bond dissociation energy for a 3° C–H bond (381 kJ/mol).

As a result, an alkyl benzene undergoes selective bromination at the weak benzylic C–H bond under radical conditions to form a **benzylic halide**. For example, radical bromination of ethylbenzene using either Br₂ (in the presence of light or heat) or *N*-bromosuccinimide (NBS, in the presence of light or peroxides) forms a benzylic bromide as the sole product.




The mechanism for halogenation at the benzylic position resembles other radical halogenation reactions, and so it involves initiation, propagation, and termination. Mechanism 18.10 illustrates the radical bromination of ethylbenzene using Br₂ (*hν* or Δ).



Mechanism 18.10 Benzylic Bromination

Initiation


Step [1] Bond cleavage forms two radicals.



- The reaction begins with homolysis of the Br–Br bond using energy from light or heat to form two Br• radicals.

Propagation

Steps [2] and [3] One radical reacts and a new radical is formed.



- Abstraction of a benzylic hydrogen by a Br• radical forms the resonance-stabilized benzylic radical in Step [2], which reacts with Br₂ in Step [3] to form the bromination product.
- Because the Br• radical formed in Step [3] is a reactant in Step [2], Steps [2] and [3] can occur repeatedly without additional initiation.


Termination

Step [4] Two radicals react to form a bond.



- To terminate the reaction, two radicals, for example two Br• radicals, react with each other to form a stable bond.

Thus, an alkyl benzene undergoes two different reactions with Br₂, depending on the reaction conditions.




- With Br₂ and FeBr₃ (**ionic conditions**), electrophilic aromatic substitution occurs, resulting in replacement of H by Br on the aromatic ring to form ortho and para isomers.
- With Br₂ and light or heat (**radical conditions**), substitution of H by Br occurs at the *benzylic* carbon of the alkyl group.

Problem 18.27 Explain why $C_6H_5CH_2CH_2Br$ is not formed during the radical bromination of $C_6H_5CH_2CH_3$.

Problem 18.28 Draw the products formed when isopropylbenzene [$C_6H_5CH(CH_3)_2$] is treated with each reagent: (a) $Br_2, FeBr_3$; (b) $Br_2, h\nu$; (c) $Cl_2, FeCl_3$.


The radical bromination of alkyl benzenes is a useful reaction because the resulting benzylic halide can serve as starting material for a variety of substitution and elimination reactions, thus making it possible to form many new substituted benzenes. Sample Problem 18.7 illustrates one possibility.

Sample Problem 18.7 Design a synthesis of styrene from ethylbenzene.




Solution

The double bond can be introduced by a two-step reaction sequence: bromination at the benzylic position under radical conditions, followed by elimination of HBr with strong base to form the π bond.



Problem 18.29 How could you use ethylbenzene to prepare each compound? More than one step is required.




18.14 Oxidation and Reduction of Substituted Benzenes


Oxidation and reduction reactions are valuable tools for preparing many other benzene derivatives. Because the mechanisms are complex and do not have general applicability, reagents and reactions are presented only, without reference to the detailed mechanism.

18.14A Oxidation of Alkyl Benzenes

Arenes containing at least one benzylic C–H bond are oxidized with $KMnO_4$ to benzoic acid, a carboxylic acid with the carboxy group ($COOH$) bonded directly to the benzene ring. With some alkyl benzenes, this also results in the cleavage of carbon–carbon bonds, so the product has fewer carbon atoms than the starting material.




Substrates with more than one alkyl group are oxidized to dicarboxylic acids. Compounds without a benzylic C–H bond are inert to oxidation.




18.14B Reduction of Aryl Ketones to Alkyl Benzenes

Ketones formed as products in Friedel–Crafts acylation can be reduced to alkyl benzenes by two different methods.



- The **Clemmensen reduction** uses zinc and mercury in the presence of strong acid.
- The **Wolff–Kishner reduction** uses hydrazine (NH_2NH_2) and strong base (KOH).

Because both C–O bonds in the starting material are converted to C–H bonds in the product, the reduction is difficult and the reaction conditions must be harsh.




We now know two different ways to introduce an alkyl group on a benzene ring (Figure 18.8):

- A **one-step method using Friedel–Crafts alkylation**
- A **two-step method using Friedel–Crafts acylation to form a ketone, followed by reduction**


Figure 18.8

Two methods to prepare an alkyl benzene



Although the two-step method seems more roundabout, it must be used to synthesize certain alkyl benzenes that cannot be prepared by the one-step Friedel–Crafts alkylation because of rearrangements.


Recall from Section 18.5C that propylbenzene cannot be prepared by a Friedel–Crafts alkylation. Instead, when benzene is treated with 1-chloropropane and AlCl_3 , isopropylbenzene is formed by a rearrangement reaction. Propylbenzene can be made, however, by a two-step procedure using Friedel–Crafts acylation followed by reduction.



Problem 18.30 Write out the two-step sequence that converts benzene to each compound:

(a) $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$; (b) $\text{C}_6\text{H}_5\text{CH}_2\text{C}(\text{CH}_3)_3$.


Problem 18.31 What steps are needed to convert benzene into *p*-isobutylacetophenone, a synthetic intermediate used in the synthesis of the anti-inflammatory agent ibuprofen.




Problem 18.32 Only one alkyl benzene with the general structure $\text{C}_6\text{H}_5\text{CH}_2\text{R}$ can be made by both Friedel–Crafts alkylation and Friedel–Crafts acylation followed by reduction. What is the identity of R in this compound?

18.14C Reduction of Nitro Groups

A nitro group (NO_2) is easily introduced on a benzene ring by nitration with strong acid (Section 18.4). This process is useful because the nitro group is readily reduced to an amino group (NH_2) under a variety of conditions. The most common methods use H_2 and a catalyst, or a metal (such as Fe or Sn) and a strong acid like HCl.




For example, reduction of ethyl *p*-nitrobenzoate with H_2 and a palladium catalyst forms ethyl *p*-aminobenzoate, a local anesthetic commonly called benzocaine.



Benzocaine is the active ingredient in the over-the-counter topical anesthetic Orajel.

Sample Problem 18.8 illustrates the utility of this process in a short synthesis.

Sample Problem 18.8 Design a synthesis of *m*-bromoaniline from benzene.




Solution

To devise a retrosynthetic plan, keep in mind:

- The NH_2 group cannot be introduced directly on the ring by electrophilic aromatic substitution. It must be added by a two-step process: nitration followed by reduction.
- Both the Br and NH_2 groups are ortho, para directors, but they are located meta to each other on the ring. However, an NO_2 group (from which an NH_2 group is made) is a meta director, and we can use this fact to our advantage.

Retrosynthetic Analysis


Working backwards gives the following **three-step retrosynthetic analysis**:






- [1] Form the NH_2 group by reduction of NO_2 .
- [2] Introduce the Br group meta to the NO_2 group by halogenation.
- [3] Add the NO_2 group by nitration.

Synthesis

The synthesis then involves three steps, and the order is crucial for success. Halogenation (Step [2] of the synthesis) must occur *before* reduction (Step [3]) in order to form the meta substitution product.




Problem 18.33 Synthesize each compound from benzene.



18.15 Multistep Synthesis

The reactions learned in Chapter 18 make it possible to synthesize a wide variety of substituted benzenes, as shown in Sample Problems 18.9–18.11.


Sample Problem 18.9 Synthesize *p*-nitrobenzoic acid from benzene.



Solution

Both groups on the ring (NO_2 and COOH) are meta directors. To place these two groups para to each other, remember that the COOH group is prepared by oxidizing an alkyl group, which is an ortho, para director.


Retrosynthetic Analysis



Working backwards:


- [1] Form the COOH group by oxidation of an alkyl group.
- [2] Introduce the NO_2 group para to the CH_3 group (an ortho, para director) by nitration.
- [3] Add the CH_3 group by Friedel–Crafts alkylation.

Synthesis



- Friedel–Crafts alkylation with CH_3Cl and AlCl_3 forms toluene in Step [1]. Because CH_3 is an ortho, para director, nitration yields the desired para product, which can be separated from its ortho isomer (Step [2]).
- Oxidation with KMnO_4 converts the CH_3 group into a COOH group, giving the desired product in Step [3].


Sample Problem 18.10 Synthesize *p*-chlorostyrene from benzene.



Solution

Both groups on the ring are ortho, para directors located para to each other. To introduce the double bond in the side chain, we must follow the two-step sequence in Sample Problem 18.7.


Retrosynthetic Analysis



Working backwards:


- [1] Form the double bond by two steps: benzylic halogenation followed by elimination.
- [2] Introduce the CH_3CH_2 group by Friedel-Crafts alkylation.
- [3] Add the Cl atom by chlorination.

Synthesis



- Chlorination in Step [1] followed by Friedel-Crafts alkylation in Step [2] forms the desired para product, which can be separated from its ortho isomer.
- Benzylic bromination followed by elimination with strong base $[\text{KOC}(\text{CH}_3)_3]$ (Steps [3] and [4]) forms the double bond of the target compound, *p*-chlorostyrene.

Sample Problem 18.11 Synthesize the trisubstituted benzene **A** from benzene.




Solution


Two groups (CH_3CO and NO_2) in **A** are meta directors located meta to each other, and the third substituent, an alkyl group, is an ortho, para director.

Retrosynthetic Analysis

With three groups on the benzene ring, begin by determining the possible disubstituted benzenes that are immediate precursors of the target compound, and then eliminate any that cannot be converted to the desired product. For example, three different disubstituted benzenes (**B–D**) can theoretically be precursors to **A**. However, conversion of compounds **B** or **D** to **A** would require a Friedel-Crafts reaction on a deactivated benzene ring, a reaction that does not occur. Thus, only **C** is a feasible precursor of **A**.




To complete the retrosynthetic analysis, prepare **C** from benzene:




- [1] Add the ketone by Friedel-Crafts acylation.
- [2] Add the alkyl group by the two-step process—Friedel-Crafts acylation followed by reduction. It is not possible to prepare butylbenzene by a one-step Friedel-Crafts alkylation because of a rearrangement reaction (Section 18.14B).

Synthesis



- Friedel-Crafts acylation followed by reduction with Zn(Hg), HCl yields butylbenzene (Steps [1]–[2]).
- Friedel-Crafts acylation gives the para product **C**, which can be separated from its ortho isomer (Step [3]).
- Nitration in Step [4] introduces the NO₂ group ortho to the alkyl group (an ortho, para director) and meta to the CH₃CO group (a meta director).

Problem 18.34 Synthesize each compound from benzene.




KEY CONCEPTS

Electrophilic Aromatic Substitution

Mechanism of Electrophilic Aromatic Substitution (18.2)





- Electrophilic aromatic substitution follows a two-step mechanism. Reaction of the aromatic ring with an electrophile forms a carbocation, and loss of a proton regenerates the aromatic ring.
- The first step is rate-determining.
- The intermediate carbocation is stabilized by resonance; a minimum of three resonance structures can be drawn. The positive charge is always located ortho or para to the new C–E bond.



Three Rules Describing the Reactivity and Directing Effects of Common Substituents (18.7–18.9)


- [1] All ortho, para directors except the halogens activate the benzene ring.
 [2] All meta directors deactivate the benzene ring.
 [3] The halogens deactivate the benzene ring and direct ortho, para.

Summary of Substituent Effects in Electrophilic Aromatic Substitution (18.6–18.9)

	Substituent	Inductive effect	Resonance effect	Reactivity	Directing effect
[1]	 R = alkyl	donating	none	activating	ortho, para
[2]	 Z = N or O	withdrawing	donating	activating	ortho, para
[3]	 X = halogen	withdrawing	donating	deactivating	ortho, para
[4]	 Y (δ^+ or +)	withdrawing	withdrawing	deactivating	meta


Five Examples of Electrophilic Aromatic Substitution

[1] Halogenation—Replacement of H by Cl or Br (18.3)




- Polyhalogenation occurs on benzene rings substituted by OH and NH₂ (and related substituents) (18.10A).


[2] Nitration—Replacement of H by NO₂ (18.4)





[3] Sulfonation—Replacement of H by SO₃H (18.4)



[4] Friedel–Crafts alkylation—Replacement of H by R (18.5)




Variations:



- Rearrangements can occur.
- Vinyl halides and aryl halides are unreactive.
- The reaction does not occur on benzene rings substituted by meta deactivating groups or NH₂ groups (18.10B).
- Polyalkylation can occur.


[5] Friedel–Crafts acylation—Replacement of H by RCO (18.5)




- The reaction does not occur on benzene rings substituted by meta deactivating groups or NH₂ groups (18.10B).

Other Reactions of Benzene Derivatives

[1] Benzylic halogenation (18.13)




[2] Oxidation of alkyl benzenes (18.14A)




- A benzylic C–H bond is needed for reaction.

[3] Reduction of ketones to alkyl benzenes (18.14B)



[4] Reduction of nitro groups to amino groups (18.14C)



PROBLEMS

Reactions


18.35 Draw the products formed when phenol (C_6H_5OH) is treated with each reagent.

- | | |
|---------------------------------|--|
| a. HNO_3, H_2SO_4 | g. $Cl_2, FeCl_3$ |
| b. SO_3, H_2SO_4 | h. product in (a), then Sn, HCl |
| c. $CH_3CH_2Cl, AlCl_3$ | i. product in (d), then $Zn(Hg), HCl$ |
| d. $(CH_3CH_2)_2CHCOCl, AlCl_3$ | j. product in (d), then $NH_2NH_2, ^-OH$ |
| e. $Br_2, FeBr_3$ | k. product in (c), then $Br_2, h\nu$ |
| f. Br_2 | l. product in (c), then $KMnO_4$ |


18.36 Draw the products formed when benzonitrile (C_6H_5CN) is treated with each reagent.

- | | | | | |
|-------------------|---------------------|--------------------|-----------------------------|-----------------------|
| a. $Br_2, FeBr_3$ | b. HNO_3, H_2SO_4 | c. SO_3, H_2SO_4 | d. $CH_3CH_2CH_2Cl, AlCl_3$ | e. $CH_3COCl, AlCl_3$ |
|-------------------|---------------------|--------------------|-----------------------------|-----------------------|


18.37 Draw the products formed when each compound is treated with $CH_3CH_2COCl, AlCl_3$.




18.38 Draw the products of each reaction.









18.39 What products are formed when benzene is treated with each alkyl chloride and $AlCl_3$?







18.40 Write out two different routes to ketone **A** using a Friedel-Crafts acylation.





18.41 Draw the products of each reaction.



18.42 You have learned two ways to make an alkyl benzene: Friedel–Crafts alkylation, and Friedel–Crafts acylation followed by reduction. Although some alkyl benzenes can be prepared by both methods, it is often true that only one method can be used to prepare a given alkyl benzene. Which method(s) can be used to prepare each of the following compounds from benzene? Show the steps that would be used.







18.43 Explain why each of the following reactions will not form the given product. Then, design a synthesis of **A** from benzene and **B** from phenol ($\text{C}_6\text{H}_5\text{OH}$).



Substituent Effects

18.44 Rank the compounds in each group in order of increasing reactivity in electrophilic aromatic substitution.



18.45 Draw all resonance structures for each compound, and explain why a particular substituent has an electron-donating or electron-withdrawing resonance effect: (a) $\text{C}_6\text{H}_5\text{NO}_2$; (b) $\text{C}_6\text{H}_5\text{F}$.

18.46 For each of the following substituted benzenes: [1] $\text{C}_6\text{H}_5\text{Br}$; [2] $\text{C}_6\text{H}_5\text{CN}$; [3] $\text{C}_6\text{H}_5\text{OCOCH}_3$:




a. Does the substituent donate or withdraw electron density by an inductive effect?

b. Does the substituent donate or withdraw electron density by a resonance effect?


c. On balance, does the substituent make a benzene ring more or less electron rich than benzene itself?

d. Does the substituent activate or deactivate the benzene ring in electrophilic aromatic substitution?

18.47 Which benzene ring in each compound is more reactive in electrophilic aromatic substitution?



18.48 For each N-substituted benzene, predict whether the compound reacts faster than, slower than, or at a similar rate to benzene in electrophilic aromatic substitution. Then draw the major product(s) formed when each compound reacts with a general electrophile E^+ .




18.49 Explain each statement in detail using resonance structures.

- A phenyl group (C_6H_5-) is an ortho, para director that activates a benzene ring towards electrophilic attack.
- A nitroso group ($-NO$) is an ortho, para director that deactivates a benzene ring towards electrophilic attack.


18.50 Explain the following observation. Ethyl 3-phenylpropanoate ($C_6H_5CH_2CH_2CO_2CH_2CH_3$) reacts with electrophiles to afford ortho- and para-disubstituted arenes, but ethyl 3-phenyl-2-propenoate ($C_6H_5CH=CHCO_2CH_2CH_3$) reacts with electrophiles to afford meta-disubstituted arenes.

18.51 Explain why the meta product is formed in the following reaction despite the fact that $-N(CH_3)_2$ is usually an ortho, para director.




Mechanisms


18.52 Draw a stepwise mechanism for each reaction.



18.53 Draw a stepwise, detailed mechanism for the following intramolecular reaction.




18.54 Draw a stepwise, detailed mechanism for the following reaction.




18.55 Friedel-Crafts alkylation of benzene with (2*R*)-2-chlorobutane and $AlCl_3$ affords *sec*-butylbenzene.

- How many stereogenic centers are present in the product?
- Would you expect the product to exhibit optical activity? Explain, with reference to the mechanism.


- 18.56** Although two products (**A** and **B**) are possible when naphthalene undergoes electrophilic aromatic substitution, only **A** is formed. Draw resonance structures for the intermediate carbocation to explain why this is observed.



- 18.57** Draw a stepwise mechanism for the following reaction, which is used to prepare the pesticide DDT.




- 18.58** Benzene undergoes electrophilic aromatic substitution with anhydrides, compounds having the general structure $(RCO)_2O$, in a reaction that resembles Friedel-Crafts acylation. Draw a stepwise mechanism for the reaction of benzene with glutaric anhydride in the presence of $AlCl_3$.



- 18.59** Benzyl bromide ($C_6H_5CH_2Br$) reacts rapidly with CH_3OH to afford benzyl methyl ether ($C_6H_5CH_2OCH_3$). Draw a stepwise mechanism for the reaction, and explain why this 1° alkyl halide reacts rapidly with a weak nucleophile under conditions that favor an S_N1 mechanism. Would you expect the para-substituted benzylic halides $CH_3OC_6H_4CH_2Br$ and $O_2NC_6H_4CH_2Br$ to each be more or less reactive than benzyl bromide in this reaction? Explain your reasoning.

- 18.60** Explain why HBr addition to $C_6H_5CH=CHCH_3$ forms only one alkyl halide, $C_6H_5CH(Br)CH_2CH_3$.







- 18.61** Draw a stepwise mechanism for the following reaction, which is used to prepare bisphenol A (BPA), a widely used monomer in polymer synthesis. Although BPA is not acutely toxic, safety concerns over low-dose exposure, especially in infants, have led to a re-examination of its use in baby bottles and infant formula cans.










Synthesis

- 18.62** Synthesize each compound from benzene and any other organic or inorganic reagents.

- isopropylbenzene
- butylbenzene
- o*-butylchlorobenzene
- m*-bromonitrobenzene
- o*-bromonitrobenzene










18.63 Synthesize each compound from benzene and any other organic or inorganic reagents.






18.64 Synthesize each compound from toluene ($C_6H_5CH_3$) and any other organic or inorganic reagents.

- a. $C_6H_5CH_2Br$
b. $C_6H_5CH_2OH$
c. $C_6H_5CH_2OC(CH_3)_3$
d. C_6H_5CHO







18.65 Devise a synthesis of each compound from phenol (C_6H_5OH) and any other organic or inorganic reagents.



18.66 Use the reactions in this chapter along with those learned in Chapters 11 and 12 to synthesize each compound. You may use benzene, acetylene ($HC\equiv CH$), two-carbon alcohols, ethylene oxide, and any inorganic reagents.


- a. $C_6H_5C\equiv CH$
b. $C_6H_5CH_2CH_2Br$
c. $C_6H_5C\equiv CCH_2CH_2OH$



18.67 Devise a synthesis of 1-phenyl-1-propyne ($C_6H_5C\equiv CCH_3$) from benzene and organic compounds having ≤ 3 C's. You may use any required inorganic or organic reagents.



18.68 Ibuprofen, a para-disubstituted arene with the structure $HO_2CCH_2C_6H_4CH_2CH(CH_3)_2$, is a much more potent analgesic than aspirin, but it was never sold commercially because it caused liver toxicity in some clinical trials. Devise a synthesis of ibuprofen from benzene and organic halides having fewer than five carbons.

18.69 Carboxylic acid **X** is an intermediate in the multistep synthesis of proparacaine, a local anesthetic. Devise a synthesis of **X** from phenol and any needed organic or inorganic reagents.




Spectroscopy

18.70 Identify the structures of isomers **A** and **B** (molecular formula C_8H_9Br).





18.71 Propose a structure of compound **C** (molecular formula $C_{10}H_{12}O$) consistent with the following data. **C** is partly responsible for the odor and flavor of raspberries.


Compound **C**: IR absorption at 1717 cm^{-1}




- 18.72 Compound **X** (molecular formula $C_{10}H_{12}O$) was treated with $NH_2NH_2, ^-OH$ to yield compound **Y** (molecular formula $C_{10}H_{14}$). Based on the 1H NMR spectra of **X** and **Y** given below, what are the structures of **X** and **Y**?



- 18.73 Reaction of *p*-cresol with two equivalents of 2-methyl-1-propene affords BHT, a preservative with molecular formula $C_{15}H_{24}O$. BHT gives the following 1H NMR spectral data: 1.4 (singlet, 18 H), 2.27 (singlet, 3 H), 5.0 (singlet, 1 H), and 7.0 (singlet, 2 H) ppm. What is the structure of BHT? Draw a stepwise mechanism illustrating how it is formed.




- 18.74 Compound **Z** (molecular formula C_9H_9ClO) can be converted to the antidepressant bupropion (Figure 18.3) by a series of reactions. **Z** shows a strong peak in its IR spectrum at 1683 cm^{-1} . The 1H NMR spectrum of **Z** shows peaks at 1.2 (triplet, 3 H), 2.9 (quartet, 2 H), and 7.2–8.0 (multiplet, 4 H) ppm. Propose a structure for **Z**.



Challenge Problems


18.75 The ^1H NMR spectrum of phenol ($\text{C}_6\text{H}_5\text{OH}$) shows three absorptions in the aromatic region: 6.70 (2 ortho H's), 7.14 (2 meta H's), and 6.80 (1 para H) ppm. Explain why the ortho and para absorptions occur at lower chemical shift than the meta absorption.

18.76 Explain the reactivity and orientation effects observed in each heterocycle.




- Pyridine is less reactive than benzene in electrophilic aromatic substitution and yields 3-substituted products.
- Pyrrole is more reactive than benzene in electrophilic aromatic substitution and yields 2-substituted products.

18.77 Draw a stepwise, detailed mechanism for the dienone-phenol rearrangement, a reaction that forms alkyl-substituted phenols from cyclohexadienes.



18.78 Draw a stepwise mechanism for the following intramolecular reaction, which is used in the synthesis of the female sex hormone estrone.



18.79 Although aryl halides are generally inert to nucleophilic substitution, aryl halides that also contain a nitro group ortho or para to the halogen undergo **nucleophilic aromatic substitution**, as shown in the following example.




- Keeping in mind that the reaction cannot follow an $\text{S}_{\text{N}}1$ or $\text{S}_{\text{N}}2$ mechanism, suggest a mechanism for this process.
- Explain why an electron-withdrawing NO_2 group is needed for this nucleophilic substitution to occur.
- Explain why *m*-chloronitrobenzene does not undergo this reaction.

19

Carboxylic Acids and the Acidity of the O-H Bond

- 19.1 Structure and bonding
- 19.2 Nomenclature
- 19.3 Physical properties
- 19.4 Spectroscopic properties
- 19.5 Interesting carboxylic acids
- 19.6 Aspirin, arachidonic acid, and prostaglandins
- 19.7 Preparation of carboxylic acids
- 19.8 Reactions of carboxylic acids—General features
- 19.9 Carboxylic acids—Strong organic Brønsted–Lowry acids
- 19.10 Inductive effects in aliphatic carboxylic acids
- 19.11 Substituted benzoic acids
- 19.12 Extraction
- 19.13 Sulfonic acids
- 19.14 Amino acids



Hexanoic acid is a low molecular weight carboxylic acid with the foul odor associated with dirty socks and locker rooms. Its common name, caproic acid, is derived from the Latin word *caper*, meaning “goat.” The fleshy coat of ginkgo seeds contains hexanoic acid, giving the seeds an unpleasant and even repulsive odor. It is likely that this foul odor served as an attractant for seed dispersal at some time during the 280 million years that *Ginkgo biloba* has existed on earth. Since only female ginkgo trees produce seeds, male trees, which are propagated by cuttings and grafts, are generally planted by landscapers in the United States. In Chapter 19 we learn about the properties of hexanoic acid and other carboxylic acids.


Chapter 19 serves as a transition between the preceding discussion of resonance and aromaticity, and the subsequent treatment of carbonyl chemistry. We pause to study the chemistry of the OH group by examining **carboxylic acids (RCOOH)**, and to a lesser extent, **phenols (PhOH)** and **alcohols (ROH)**.

In Chapter 19 we concentrate on the acidity of carboxylic acids, and revisit some of the factors that determine acidity, a topic first discussed in Chapter 2. Then, in Chapters 20 and 22 we will learn other reactions of carboxylic acids that occur at the carbonyl group.


19.1 Structure and Bonding

Carboxylic acids are organic compounds containing a carboxy group (COOH). Although the structure of a carboxylic acid is often abbreviated as **RCOOH** or **RCO₂H**, keep in mind that the central carbon atom of the functional group is doubly bonded to one oxygen atom and singly bonded to another.


The word **carboxy** (for a COOH group) is derived from **carbonyl** (C=O) + **hydroxy** (OH).



The carbon atom of a carboxy group is surrounded by three groups, making it ***sp*² hybridized** and **trigonal planar**, with bond angles of approximately 120°. The C=O of a carboxylic acid is shorter than its C–O.




The C–O single bond of a carboxylic acid is shorter than the C–O single bond of an alcohol. This can be explained by looking at the hybridization of the respective carbon atoms. In the alcohol, the carbon is *sp*³ hybridized, whereas in the carboxylic acid the carbon is *sp*² hybridized. As a result, the higher percent *s*-character in the *sp*² hybrid orbital shortens the C–O bond in the carboxylic acid.



Because oxygen is more electronegative than either carbon or hydrogen, **the C–O and O–H bonds are polar**. The electrostatic potential plot of acetic acid in Figure 19.1 shows that the carbon and hydrogen atoms are electron poor and the oxygen atoms are electron rich.

Figure 19.1

Electrostatic potential plot of acetic acid (CH₃COOH)



Acetic acid contains two electron-rich oxygen atoms (in red). Its carbonyl carbon and hydroxy hydrogen are both electron deficient.

19.2 Nomenclature

Both IUPAC and common names are used for carboxylic acids.

19.2A IUPAC System



In IUPAC nomenclature, carboxylic acids are identified by a suffix added to the parent name of the longest chain, and two different endings are used depending on whether the carboxy group is bonded to a chain or ring.

To name a carboxylic acid using the IUPAC system:

- [1] If the COOH is bonded to a chain of carbons, find the longest chain containing the COOH group, and change the *-e* ending of the parent alkane to the suffix *-oic acid*. If the COOH group is bonded to a ring, name the ring and add the words *carboxylic acid*.
- [2] Number the carbon chain or ring to put the **COOH group at C1**, but omit this number from the name. Apply all of the other usual rules of nomenclature.


Sample Problem 19.1

Give the IUPAC name of each compound.



Solution


- a. [1] Find and name the longest chain containing COOH:



hexane → **hexanoic acid**
(6 C's)

The COOH contributes one C to the longest chain.


- [2] Number and name the substituents:



two methyl substituents on C4 and C5


Answer: 4,5-dimethylhexanoic acid

- b. [1] Find and name the ring bonded to COOH.



cyclohexane + carboxylic acid
(6 C's)

- [2] Number and name the substituents:




[Number to put COOH at C1 and give the second substituent (CH₃) the lower number (C2).]

Answer: 2,5,5-trimethylcyclohexanecarboxylic acid

Problem 19.1

Give the IUPAC name for each compound.

- a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{COOH}$
- b. $\text{CH}_3\text{CH}(\text{Cl})\text{CH}_2\text{CH}_2\text{COOH}$
- c. $(\text{CH}_3\text{CH}_2)_2\text{CHCH}_2\text{CH}(\text{CH}_2\text{CH}_3)\text{COOH}$



Problem 19.2

Give the structure corresponding to each IUPAC name.

- a. 2-bromobutanoic acid
- b. 2,3-dimethylpentanoic acid
- c. 3,3,4-trimethylheptanoic acid
- d. 2-sec-butyl-4,4-diethylnonanoic acid
- e. 3,4-diethylcyclohexanecarboxylic acid
- f. 1-isopropylcyclobutanecarboxylic acid

19.2B Common Names

Most simple carboxylic acids have common names that are more widely used than their IUPAC names.

- A common name is formed by using a common parent name followed by the suffix *-ic acid*.

Table 19.1 lists common parent names for some simple carboxylic acids. These parent names are used in the nomenclature of many other compounds with carbonyl groups (Chapters 21 and 22).

Greek letters are used to designate the location of substituents in common names.

- The carbon adjacent to the COOH is called the α carbon.
- The carbon bonded to the α carbon is the β carbon, followed by the γ (gamma) carbon, the δ (delta) carbon, and so forth down the chain. The last carbon in the chain is sometimes called the Ω (omega) carbon.

The α carbon in the common system is numbered C2 in the IUPAC system.




Table 19.1 Common Names for Some Simple Carboxylic Acids

Number of C atoms	Structure	Parent name	Common name
1		form-	formic acid
2		acet-	acetic acid
3		propion-	propionic acid
4		butyr-	butyric acid
5		valer-	valeric acid
6		capro-	caproic acid
		benzo-	benzoic acid

Figure 19.3

Two molecules of acetic acid (CH_3COOH) held together by two hydrogen bonds





Table 19.2 Physical Properties of Carboxylic Acids

Property	Observation												
Boiling point and melting point	<ul style="list-style-type: none"> Carboxylic acids have higher boiling points and melting points than other compounds of comparable molecular weight. <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ VDW</th> <th>$\text{CH}_3\text{CH}_2\text{CHO}$ VDW, DD</th> <th>$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB</th> <th>CH_3COOH VDW, DD, two HB</th> </tr> </thead> <tbody> <tr> <td>MW = 58</td> <td>MW = 58</td> <td>MW = 60</td> <td>MW = 60</td> </tr> <tr> <td>bp 0 °C</td> <td>bp 48 °C</td> <td>bp 97 °C</td> <td>bp 118 °C</td> </tr> </tbody> </table> <p style="text-align: center;"> Increasing strength of intermolecular forces Increasing boiling point </p>	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ VDW	$\text{CH}_3\text{CH}_2\text{CHO}$ VDW, DD	$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB	CH_3COOH VDW, DD, two HB	MW = 58	MW = 58	MW = 60	MW = 60	bp 0 °C	bp 48 °C	bp 97 °C	bp 118 °C
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$ VDW	$\text{CH}_3\text{CH}_2\text{CHO}$ VDW, DD	$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB	CH_3COOH VDW, DD, two HB										
MW = 58	MW = 58	MW = 60	MW = 60										
bp 0 °C	bp 48 °C	bp 97 °C	bp 118 °C										
Solubility	<ul style="list-style-type: none"> Carboxylic acids are soluble in organic solvents regardless of size. Carboxylic acids having ≤ 5 C's are water soluble because they can hydrogen bond with H_2O (Section 3.4C). Carboxylic acids having > 5 C's are water insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent. These "fatty" acids dissolve in a nonpolar fat-like environment but do not dissolve in water. 												

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding, MW = molecular weight

Problem 19.7 Rank the following compounds in order of increasing boiling point. Which compound is the most water soluble? Which compound is the least water soluble?




19.4 Spectroscopic Properties

Carboxylic acids have very characteristic IR and NMR absorptions. In the IR, carboxylic acids show two strong absorptions.

- The **C=O group absorbs at about 1710 cm^{-1}** , in the usual region for a carbonyl.
- The **O–H absorption** occurs from **$2500\text{--}3500\text{ cm}^{-1}$** . This very broad absorption sometimes obscures the C–H peak at 3000 cm^{-1} .

Figure 19.4

The IR spectrum of butanoic acid, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$



- A strong C=O absorption occurs at 1712 cm^{-1} .
- The broad O–H absorption ($2500\text{--}3500\text{ cm}^{-1}$) nearly obscures the C–H peak at $\sim 3000\text{ cm}^{-1}$.


The IR spectrum of butanoic acid in Figure 19.4 illustrates these characteristic peaks.

Carboxylic acids have two noteworthy ^1H NMR absorptions and one noteworthy ^{13}C NMR absorption.

- **The highly deshielded OH proton absorbs in the ^1H NMR spectrum somewhere between 10 and 12 ppm**, farther *downfield* than all other absorptions of common organic compounds. Like the OH signal of an alcohol, the exact location depends on the degree of hydrogen bonding and the concentration of the sample.
- The protons on the α carbon to the carboxy group are somewhat deshielded, absorbing at 2–2.5 ppm.
- In the ^{13}C NMR spectrum, the carbonyl absorption is highly deshielded, appearing at 170–210 ppm.

Figure 19.5 illustrates the ^1H and ^{13}C NMR spectra of propanoic acid.

Problem 19.8 Explain how you could use IR spectroscopy to distinguish among the following three compounds.



Problem 19.9 Identify the structure of a compound of molecular formula $\text{C}_4\text{H}_8\text{O}_2$ that gives the following ^1H NMR data: 0.95 (triplet, 3 H), 1.65 (multiplet, 2 H), 2.30 (triplet, 2 H), and 11.8 (singlet, 1 H) ppm.


Problem 19.10 How could ^1H NMR spectroscopy be used to distinguish between formic acid (HCO_2H) and malonic acid [$\text{CH}_2(\text{CO}_2\text{H})_2$]?

19.5 Interesting Carboxylic Acids

Several simple carboxylic acids have characteristic odors and flavors.



HCOOH




- **Formic acid**, a carboxylic acid with an acrid odor and a biting taste, is responsible for the sting of some types of ants. The name is derived from the Latin word *formica*, meaning “ant.”

Figure 19.5

The ^1H and ^{13}C NMR spectra of propanoic acid


Sometimes the OH absorption of a carboxylic acid is very broad, so that it is almost buried in the baseline of the ^1H NMR spectrum, making it difficult to see.



- **^1H NMR spectrum:** There are three signals due to three different kinds of H atoms. The H_a and H_b signals are split into a triplet and quartet, respectively, but the H_c signal is a singlet.
- **^{13}C NMR spectrum:** There are three signals due to three different kinds of carbon atoms.


Pure acetic acid is often called *glacial* acetic acid, because it freezes just below room temperature (mp = 17 °C), forming white crystals reminiscent of the ice in a glacier.

CH₃COOH



- **Acetic acid** is the sour-tasting component of vinegar. The name comes from the Latin *acetum*, meaning “vinegar.” The air oxidation of ethanol to acetic acid is the process that makes “bad” wine taste sour. Acetic acid is an industrial starting material for polymers used in paints and adhesives.

CH₃CH₂CH₂COOH




- **Butanoic acid** is an oxidation product that contributes to the disagreeable smell of body odor. Its common name, butyric acid, is derived from the Latin word *butyrum*, meaning “butter,” because butyric acid gives rancid butter its peculiar odor and taste.




Although oxalic acid is toxic, you would have to eat about nine pounds of spinach at one time to ingest a fatal dose.

Oxalic acid and **lactic acid** are simple carboxylic acids quite prevalent in nature. Oxalic acid occurs naturally in spinach and rhubarb. Lactic acid gives sour milk its distinctive taste.




oxalic acid




lactic acid

4-Hydroxybutanoic acid, known by its common name γ -hydroxybutyric acid (GHB), is an illegal recreational drug that depresses the central nervous system and results in intoxication. GHB is highly addictive and widely abused, and because its taste can be easily masked in an alcoholic beverage, it has been used as a “date rape” drug.


4-hydroxybutanoic acid
GHB

Salts of carboxylic acids are commonly used as preservatives. **Sodium benzoate**, a fungal growth inhibitor, is a preservative used in soft drinks, and potassium sorbate is an additive that prolongs the shelf-life of baked goods and other foods.

Soaps, the sodium salts of fatty acids, were discussed in Section 3.6.



sodium benzoate







potassium sorbate

19.6 Aspirin, Arachidonic Acid, and Prostaglandins



The word *aspirin* is derived from the prefix *a-* for *acetyl* + *spir* from the Latin name *spirea* for the meadowsweet plant.


Recall from Chapter 2 that **aspirin (acetylsalicylic acid)** is a synthetic carboxylic acid, similar in structure to **salicin**, a naturally occurring compound isolated from willow bark, and **salicylic acid**, found in meadowsweet.

aspirin
(acetylsalicylic acid)salicin
(isolated from
willow bark)salicylic acid
(isolated from
meadowsweet)sodium salicylate
(sweet carboxylate salt)


Both salicylic acid and sodium salicylate (its sodium salt) were widely used analgesics in the nineteenth century, but both had undesirable side effects. Salicylic acid irritated the mucous membranes of the mouth and stomach, and sodium salicylate was too sweet for most patients. Aspirin, a synthetic compound, was first sold in 1899 after Felix Hoffman, a German chemist at Bayer Company, developed a feasible commercial synthesis. Hoffman’s work was motivated by personal reasons; his father suffered from rheumatoid arthritis and was unable to tolerate the sweet taste of sodium salicylate.

Aspirin is the most widely used pain reliever and anti-inflammatory agent in the world, yet its mechanism of action remained unknown until the 1970s. John Vane, Bengt Samuelsson, and Sune Bergstrom shared the 1982 Nobel Prize in Physiology or Medicine for unraveling the details of its mechanism.


How does aspirin relieve pain and reduce inflammation? Aspirin blocks the synthesis of **prostaglandins**, 20-carbon fatty acids with a five-membered ring that are responsible for pain, inflammation, and a wide variety of other biological functions. $\text{PGF}_{2\alpha}$ contains the typical carbon skeleton of a prostaglandin.



Prostaglandins are not stored in cells. Rather they are synthesized from arachidonic acid, a polyunsaturated fatty acid having four cis double bonds. Unlike hormones, which are transported in the bloodstream to their sites of action, prostaglandins act where they are synthesized. Aspirin acts by blocking the synthesis of prostaglandins from arachidonic acid. Aspirin inactivates cyclooxygenase, an enzyme that converts arachidonic acid to PGG_2 , an unstable precursor of $\text{PGF}_{2\alpha}$ and other prostaglandins. **Aspirin lessens pain and decreases inflammation because it prevents the synthesis of prostaglandins, the compounds responsible for both of these physiological responses.**



Although prostaglandins have a wide range of biological activity, their inherent instability often limits their usefulness as drugs. Consequently, more stable analogues with useful medicinal properties have been synthesized. For example, latanoprost (trade name Xalatan) and bimatoprost (trade name Lumigan) are prostaglandin analogues used to reduce eye pressure in individuals with glaucoma.



Problem 19.11 How many tetrahedral stereogenic centers does $\text{PGF}_{2\alpha}$ contain? Draw its enantiomer. How many of its double bonds can exhibit cis-trans isomerism? Considering both its double bonds and its tetrahedral stereogenic centers, how many stereoisomers are possible for $\text{PGF}_{2\alpha}$?

19.7 Preparation of Carboxylic Acids


Our discussion of the reactions involving carboxylic acids begins with a brief list of reactions that synthesize them. This list serves as a reminder of where you have seen this functional group before. In these reactions, the carboxy group is formed in the *product*, and many different functional groups serve as starting materials. Reactions that produce a particular functional group are called **preparations**.

In the remainder of Chapter 19 (and Chapters 20 and 22) we discuss reactions in which a carboxylic acid is a *starting material* that may be converted to a variety of different products. Keep in mind that **reactions of a particular functional group follow a common theme**. For example, alkenes undergo addition reactions. As a result, these reactions are easier to learn than the list of preparations, in which vastly different functional groups undergo a wide variety of reactions to form the same kind of product.

Where have we encountered carboxylic acids as reaction products before? The carbonyl carbon is highly oxidized, because it has three C–O bonds, so **carboxylic acids are typically prepared by oxidation reactions**. Three oxidation methods are summarized below. Two other useful methods to prepare carboxylic acids are presented in Chapter 20.


[1] By oxidation of 1° alcohols (Section 12.12B)

1° Alcohols are converted to carboxylic acids with $\text{Na}_2\text{Cr}_2\text{O}_7$, $\text{K}_2\text{Cr}_2\text{O}_7$, or CrO_3 in the presence of H_2O and H_2SO_4 .



[2] By oxidation of alkyl benzenes (Section 18.14A)


Alkyl benzenes having at least one benzylic C–H bond are oxidized with KMnO_4 to benzoic acid.



Benzoic acid is *always* the product regardless of the alkyl benzene used as starting material.


[3] By oxidative cleavage of alkynes (Section 12.11)

Both internal and terminal alkynes are oxidatively cleaved with ozone to give carboxylic acids.







With internal alkynes two carboxylic acids are formed as products. With terminal alkynes, the *sp* hybridized C–H bond is converted to CO_2 .

Problem 19.12 What alcohol can be oxidized to each carboxylic acid?




Problem 19.13 Identify **A–D** in the following reactions.



19.8 Reactions of Carboxylic Acids—General Features


The polar C–O and O–H bonds, nonbonded electron pairs on oxygen, and the π bond give a carboxylic acid many reactive sites, complicating its chemistry somewhat. By far, **the most important reactive feature of a carboxylic acid is its polar O–H bond, which is readily cleaved with base.**

- Carboxylic acids react as Brønsted–Lowry acids—that is, as proton donors.




Much of the rest of Chapter 19 is devoted to the acidity of carboxylic acids, as well as some related acid–base reactions. Two other structural features are less important in the reactions of carboxylic acids, but they play a role in the reactions of Chapters 20 and 22.

The nonbonded electron pairs on oxygen create electron-rich sites that can be protonated by strong acids (H–A). Protonation occurs at the carbonyl oxygen because the resulting conjugate acid is resonance stabilized (Possibility [1]). The product of protonation of the OH group (Possibility [2]) cannot be resonance stabilized. As a result, **carboxylic acids are weakly basic—they react with strong acids by protonation of the carbonyl oxygen.** This reaction plays an important role in several mechanisms in Chapter 22.





Possibility [2]
Protonation of the OH group



Finally, the polar C–O bonds make the carboxy carbon electrophilic, so carboxylic acids react with nucleophiles. Nucleophilic attack occurs at an sp^2 hybridized carbon atom, so it results in the cleavage of the π bond, as well. This reaction is also discussed in Chapter 22.


Nucleophilic attack at the carboxy carbon






How resonance affects acidity was first discussed in Section 2.5C.

Experimental data support this resonance description of acetate. **The acetate anion has two C–O bonds of equal length** (127 pm) and intermediate between the length of a C–O single bond (136 pm) and C=O (121 pm).



Resonance stabilization accounts for why carboxylic acids are more acidic than other compounds with O–H bonds—namely, alcohols and phenols. For example, the pK_a values of ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) and phenol ($\text{C}_6\text{H}_5\text{OH}$) are 16 and 10, respectively, both higher than the pK_a of acetic acid (4.8).



To understand the relative acidity of ethanol, phenol, and acetic acid, we must compare the stability of their conjugate bases and use the following rule:


- Anything that stabilizes a conjugate base A^- makes the starting acid H–A more acidic.

Ethoxide, the conjugate base of ethanol, bears a negative charge on an oxygen atom, but there are no additional factors to further stabilize the anion. Because ethoxide is less stable than acetate, **ethanol is a weaker acid than acetic acid.**



The resonance hybrid of phenoxide illustrates that its negative charge is dispersed over four atoms—three C atoms and one O atom.

Like acetate, **phenoxide** ($\text{C}_6\text{H}_5\text{O}^-$, the conjugate base of phenol) is also resonance stabilized. In the case of phenoxide, however, there are five resonance structures that disperse the negative charge over a total of four different atoms (three different carbons and the oxygen).




Five resonance structures delocalize the negative charge over four atoms.

Phenoxide is more stable than ethoxide, but less stable than acetate, because acetate has two electronegative oxygen atoms upon which to delocalize the negative charge, whereas phenoxide has only one. Additionally, phenoxide resonance structures 2–4 have the negative charge on a carbon, a less electronegative element than oxygen. As a result, structures 2–4 are less stable than structures 1 and 5, which have the negative charge on oxygen.

Figure 19.6

The relative energies of the five resonance structures for phenoxide and its hybrid



Moreover, resonance structures **1** and **5** have intact aromatic rings, whereas structures **2–4** do not. This, too, makes structures **2–4** less stable than **1** and **5**. Figure 19.6 summarizes this information about phenoxide by displaying the approximate relative energies of its five resonance structures and its hybrid.

As a result, resonance stabilization of the conjugate base is important in determining acidity, but **the absolute number of resonance structures alone is not what's important**. We must evaluate their relative contributions to predict the relative stability of the conjugate bases.




- Because of their O–H bond, RCOOH, ROH, and C₆H₅OH are more acidic than most organic hydrocarbons.
- A carboxylic acid is a stronger acid than an alcohol or phenol because its conjugate base is most effectively resonance stabilized.

Keep in mind that although carboxylic acids are strong organic acids, they are still much weaker than strong inorganic acids like HCl and H₂SO₄, which have pK_a values < 0.

The relationship between acidity and stability of the conjugate base is summarized for acetic acid, phenol, and ethanol in Figure 19.7.


Because alcohols and phenols are weaker acids than carboxylic acids, stronger bases are needed to deprotonate them. To deprotonate C₆H₅OH (pK_a = 10), a base whose conjugate acid has a pK_a > 10 is needed. Thus, of the bases listed in Table 19.3, NaOCH₃, NaOH, NaOCH₂CH₃, and NaH are strong enough. To deprotonate CH₃CH₂OH (pK_a = 16), only NaH is strong enough.

Problem 19.14 Draw the products of each acid–base reaction.



Problem 19.15 Given the pK_a values in Appendix A, which of the following bases are strong enough to deprotonate CH₃COOH: (a) F[−]; (b) (CH₃)₃CO[−]; (c) CH₃[−]; (d) [−]NH₂; (e) Cl[−]?


Problem 19.16 Rank the labeled protons (H_a–H_c) in mandelic acid, a naturally occurring carboxylic acid in plums and peaches, in order of increasing acidity. Explain in detail why you chose this order.



mandelic acid

Figure 19.7

Summary: The relationship between acidity and conjugate base stability for acetic acid, phenol, and ethanol



- **Acetate is the most stable conjugate base** because it has two equivalent resonance structures, both of which place a negative charge on an O atom.
- **Phenoxide** has only one O atom to accept the negative charge. The two resonance structures that contain an intact aromatic ring and place a negative charge on an O atom are major contributors to the hybrid. Resonance stabilizes phenoxide but not as much as resonance stabilizes acetate.
- **Ethoxide is the least stable conjugate base** because it has no additional resonance stabilization.


19.10 Inductive Effects in Aliphatic Carboxylic Acids

The $\text{p}K_{\text{a}}$ of a carboxylic acid is affected by nearby groups that inductively donate or withdraw electron density.

- **Electron-withdrawing groups stabilize a conjugate base, making a carboxylic acid more acidic.**
- **Electron-donating groups destabilize the conjugate base, making a carboxylic acid less acidic.**

The relative acidity of CH_3COOH , ClCH_2COOH , and $(\text{CH}_3)_3\text{CCOOH}$ illustrates these principles in the following equations.


We first learned about inductive effects and acidity in Section 2.5B.




- ClCH_2COOH is more acidic ($pK_a = 2.8$) than CH_3COOH ($pK_a = 4.8$) because its conjugate base is stabilized by the electron-withdrawing inductive effect of the electronegative Cl.
- $(\text{CH}_3)_3\text{CCOOH}$ is less acidic ($pK_a = 5.1$) than CH_3COOH because the three polarizable CH_3 groups donate electron density and destabilize the conjugate base.

The number, electronegativity, and location of substituents also affect acidity.


- The larger the number of electronegative substituents, the stronger the acid.



- The more electronegative the substituent, the stronger the acid.



- The closer the electron-withdrawing group to the COOH, the stronger the acid.



Problem 19.17

Match each of the following pK_a values (3.2, 4.9, and 0.2) to the appropriate carboxylic acid: (a) $\text{CH}_3\text{CH}_2\text{COOH}$; (b) CF_3COOH ; (c) ICH_2COOH .

Problem 19.18 Explain why HCOOH (formic acid) has a lower pK_a than acetic acid (3.8 versus 4.8).

Problem 19.19 Rank the compounds in each group in order of increasing acidity.

a. CH_3COOH , HSCH_2COOH , HOCH_2COOH b. ICH_2COOH , I_2CHCOOH , $\text{ICH}_2\text{CH}_2\text{COOH}$

19.11 Substituted Benzoic Acids

Recall from Chapter 18 that substituents on a benzene ring either donate or withdraw electron density, depending on the balance of their inductive and resonance effects. These same effects also determine the acidity of substituted benzoic acids. There are two rules to keep in mind.

Rule [1] Electron-donor groups destabilize a conjugate base, making an acid less acidic.

An **electron-donor group destabilizes a conjugate base** by donating electron density onto a negatively charged carboxylate anion. A benzoic acid substituted by an electron-donor group has a higher pK_a than benzoic acid ($pK_a = 4.2$).

D = Electron-donor group



This acid is less acidic than benzoic acid.

$$pK_a > 4.2$$

D destabilizes the carboxylate anion.

Rule [2] Electron-withdrawing groups stabilize a conjugate base, making an acid more acidic.

An **electron-withdrawing group stabilizes a conjugate base** by removing electron density from the negatively charged carboxylate anion. A benzoic acid substituted by an electron-withdrawing group has a lower pK_a than benzoic acid ($pK_a = 4.2$).

W = Electron-withdrawing group



This acid is more acidic than benzoic acid.

$$pK_a < 4.2$$

W stabilizes the carboxylate anion.

How do we know which groups are electron donating or electron withdrawing on a benzene ring? We already learned the characteristics of electron-donating and electron-withdrawing groups in Chapter 18, and how they affect the rate of electrophilic aromatic substitution. These principles can now be extended to substituted benzoic acids.




Figure 19.8 illustrates how common electron-donating and electron-withdrawing groups affect both the rate of reaction of a benzene ring towards electrophiles and the acidity of substituted benzoic acids.

Figure 19.8


How common substituents affect the reactivity of a benzene ring towards electrophiles and the acidity of substituted benzoic acids

	Substituent	Effect in electrophilic substitution	Effect on acidity of substituted benzoic acids
electron-donating groups	$-\dot{\text{N}}\text{H}_2$ [$\dot{\text{N}}\text{HR}$, $\dot{\text{N}}\text{R}_2$]	activating groups	These groups make a benzoic acid less acidic .
	$-\ddot{\text{O}}\text{H}$		
	$-\ddot{\text{O}}\text{R}$		
	$-\dot{\text{N}}\text{HCOR}$		
	$-\text{R}$		
electron-withdrawing groups	$-\ddot{\text{X}}:$ [$\text{X} = \text{F}, \text{Cl}, \text{Br}, \text{I}$]	deactivating groups	These groups make a benzoic acid more acidic .
	$-\text{CHO}$		
	$-\text{COR}$		
	$-\text{COOR}$		
	$-\text{COOH}$		
	$-\text{CN}$		
	$-\text{SO}_3\text{H}$		
	$-\text{NO}_2$		
	$-\text{NR}_3^+$		


- **Groups that donate electron density activate a benzene ring towards electrophilic attack and make a benzoic acid *less* acidic.** Common electron-donating groups are R groups, or groups that have an N or O atom (with a lone pair) bonded to the benzene ring.
- **Groups that withdraw electron density deactivate a benzene ring towards electrophilic attack, and make a benzoic acid *more* acidic.** Common electron-withdrawing groups are the halogens, or groups with an atom Y (with a full or partial positive charge) bonded to the benzene ring.

Sample Problem 19.2


Rank the following three carboxylic acids in order of increasing acidity.

**Solution**

***p*-Methoxybenzoic acid (B):** The CH_3O group is an electron-donor group because its electron-donating resonance effect is stronger than its electron-withdrawing inductive effect (Section 18.6). This destabilizes the conjugate base by donating electron density to the negatively charged carboxylate anion, making **B** less acidic than benzoic acid **A**.




***p*-Nitrobenzoic acid (C):** The NO₂ group is an electron-withdrawing group because of both inductive effects and resonance (Section 18.6). This stabilizes the conjugate base by removing electron density from the negatively charged carboxylate anion, making **C** more acidic than benzoic acid **A**.




By this analysis, the order of acidity is **B** < **A** < **C**.

Problem 19.20 Rank the compounds in each group in order of increasing acidity.




Problem 19.21



Poison ivy contains the irritant urushiol.

Substituted phenols show substituent effects similar to substituted benzoic acids. Should the pK_a of phenol **A**, one of the naturally occurring phenols called urushiols isolated from poison ivy, be higher or lower than the pK_a of phenol (C₆H₅OH, $pK_a = 10$)? Explain.



19.12 Extraction

An organic chemist in the laboratory must separate and purify mixtures of compounds. One particularly useful technique is **extraction**, which uses solubility differences and acid–base principles to separate and purify compounds.

Two solvents are used in extraction: water or an aqueous solution such as 10% NaHCO₃ or 10% NaOH; and an organic solvent such as dichloromethane (CH₂Cl₂), diethyl ether, or hexane. **Compounds are separated by their solubility differences in an aqueous and organic solvent.**

An item of glassware called a **separatory funnel**, depicted in Figure 19.9, is used for the extraction. When two insoluble liquids are added to the separatory funnel, two layers form, with the less dense liquid on top and the more dense liquid on the bottom.


Suppose a mixture of benzoic acid (C₆H₅COOH) and NaCl is added to a separatory funnel containing H₂O and CH₂Cl₂. The benzoic acid would dissolve in the organic layer and the NaCl would dissolve in the water layer. Separating the organic and aqueous layers and placing them in different flasks separates the benzoic acid and NaCl from each other.

How could we separate a mixture of benzoic acid and cyclohexanol? Both compounds are organic, and as a result, both are soluble in an organic solvent such as CH₂Cl₂ and insoluble in water. If a mixture of benzoic acid and cyclohexanol were added to a separatory funnel with CH₂Cl₂ and water, both would dissolve in the CH₂Cl₂ layer, and the two compounds would *not* be separated from each other. Is it possible to use extraction to separate two compounds of this sort that have similar solubility properties?

Extraction has long been and remains the first step in isolating a natural product from its source.


Figure 19.9

Using a separatory funnel for extraction




- When two insoluble liquids are added to a separatory funnel, two layers are visible, and the less dense liquid forms the upper layer.
- To separate the layers, the lower layer can be drained from the bottom of the separatory funnel by opening the stopcock. The top layer can then be poured out the top neck of the funnel.

Recall from Tables 9.1 and 19.2 that alcohols and carboxylic acids having more than five carbons are water insoluble.




If a carboxylic acid is one of the compounds, the answer is *yes*, because we can use acid–base chemistry to change its solubility properties.

When benzoic acid (a strong organic acid) is treated with aqueous NaOH, benzoic acid is deprotonated, forming sodium benzoate. Because sodium benzoate is ionic, it is soluble in water, but insoluble in organic solvents.




The solubility properties of the conjugate base are *different* from those of the starting acid.

A similar acid–base reaction does *not* occur when cyclohexanol is treated with NaOH because organic alcohols are much weaker organic acids, so they can be deprotonated only by a *very strong base* such as NaH. NaOH is not strong enough to form significant amounts of the sodium alkoxide.



This difference in acid–base chemistry can be used to separate benzoic acid and cyclohexanol by the stepwise extraction procedure illustrated in Figure 19.10. This extraction scheme relies on two basic principles:

Figure 19.10
Separation of benzoic acid and cyclohexanol by an extraction procedure



- Both compounds dissolve in the organic solvent CH_2Cl_2 .
- Adding 10% aqueous NaOH solution forms two layers. When the two layers are mixed, the NaOH deprotonates $\text{C}_6\text{H}_5\text{COOH}$ to form $\text{C}_6\text{H}_5\text{COO}^-\text{Na}^+$, which dissolves in the aqueous layer.
- The cyclohexanol remains in the CH_2Cl_2 layer.
- Draining the lower layer out the bottom stopcock separates the two layers, and the separation process is complete.
- Cyclohexanol (dissolved in CH_2Cl_2) is in one flask. The sodium salt of benzoic acid, $\text{C}_6\text{H}_5\text{COO}^-\text{Na}^+$ (dissolved in water) is in another flask.

• Extraction can separate only compounds having different solubility properties. One compound must dissolve in the aqueous layer and one must dissolve in the organic layer.

• A carboxylic acid can be separated from other organic compounds by converting it to a water-soluble carboxylate anion by an acid–base reaction.

Thus, the water-soluble salt, $\text{C}_6\text{H}_5\text{COO}^-\text{Na}^+$ (derived from $\text{C}_6\text{H}_5\text{COOH}$ by an acid–base reaction) can be separated from water-insoluble cyclohexanol by an extraction procedure.

Problem 19.22 Which of the following pairs of compounds can be separated from each other by an extraction procedure?

- $\text{CH}_3(\text{CH}_2)_6\text{COOH}$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$
- $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$ and $(\text{CH}_3\text{CH}_2\text{CH}_2)_2\text{O}$
- $\text{CH}_3(\text{CH}_2)_6\text{COOH}$ and NaCl
- NaCl and KCl


Problem 19.23 Explain why two low molecular weight organic compounds such as $\text{CH}_3\text{CH}_2\text{COOH}$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ can't be separated by an aqueous extraction technique.

19.13 Sulfonic Acids


Although much less common than carboxylic acids, **sulfonic acids** constitute a useful group of organic acids. Sulfonic acids have the general structure RSO_3H . The most widely used sulfonic acid, *p*-toluenesulfonic acid, was first discussed in Section 2.6.

Recall from Section 9.13 that $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_2^-$ is called a **tosyl group**, abbreviated by the letters **Ts**. For this reason, *p*-toluenesulfonic acid (also called tosic acid) is abbreviated as **TsOH**.


General structure



Example



Sulfonic acids are very strong acids ($\text{p}K_{\text{a}}$ values ≈ -7) because their conjugate bases are resonance stabilized, and all the resonance structures delocalize a negative charge on oxygen. The conjugate base of a sulfonic acid is called a **sulfonate anion**.



Because sulfonate anions are such weak bases, they make **good leaving groups** in nucleophilic substitution reactions, as we learned in Section 9.13.

Problem 19.24

Two other commonly used sulfonic acids are methanesulfonic acid ($\text{CH}_3\text{SO}_3\text{H}$) and trifluoromethanesulfonic acid ($\text{CF}_3\text{SO}_3\text{H}$). Which has the weaker conjugate base? Which conjugate base is the better leaving group? Which of these acids has the higher $\text{p}K_{\text{a}}$?


19.14 Amino Acids

Amino acids, one of four kinds of small biomolecules that have important biological functions in the cell (Section 3.9), also undergo proton transfer reactions.



19.14A Introduction

Chapter 28 discusses the synthesis of amino acids and their conversion to proteins.

Amino acids contain two functional groups—an amino group (NH_2) and a carboxy group (COOH). In most naturally occurring amino acids, the amino group is bonded to the α carbon, and so they are called **α -amino acids**. Amino acids are the building blocks of proteins, biomolecules that comprise muscle, hair, fingernails, and many other biological tissues.



The 20 amino acids that occur naturally in proteins differ in the identity of the R group bonded to the α carbon. **The simplest amino acid, called glycine, has $\text{R} = \text{H}$.** When the R group is any other substituent, **the α carbon is a stereogenic center**, and there are two possible enantiomers.

Simplest amino acid, $\text{R} = \text{H}$ Two possible enantiomers when $\text{R} \neq \text{H}$ 

Only this isomer occurs in proteins.



Humans can synthesize only 10 of the 20 amino acids needed for protein synthesis. The remaining 10, called **essential amino acids**, must be obtained from the diet and consumed on a regular, almost daily basis. Because no one plant source has sufficient amounts of all the essential amino acids, vegetarian diets must be carefully balanced. Grains—wheat, rice, and corn—are low in lysine, and legumes—beans, peas, and peanuts—are low in methionine, but a combination of these foods provides all the needed amino acids. Thus, a diet of corn tortillas and beans, or rice and tofu, provides all essential amino acids. A peanut butter sandwich on wheat bread does, too.

Table 19.4 Representative Amino Acids



R group	Name	Three-letter abbreviation	One-letter abbreviation
H	glycine	Gly	G
CH ₃	alanine	Ala	A
CH ₂ C ₆ H ₅	phenylalanine	Phe	F
CH ₂ OH	serine	Ser	S
CH ₂ SH	cysteine	Cys	C
CH ₂ CH ₂ SCH ₃	methionine	Met	M
CH ₂ CH ₂ COOH	glutamic acid	Glu	E
(CH ₂) ₄ NH ₂	lysine	Lys	K

Amino acids exist in nature as only one of these enantiomers. Except when the R group is CH₂SH, the stereogenic center on the α carbon has the *S* configuration. An older system of nomenclature names the **naturally occurring enantiomer of an amino acid as the L isomer, and its unnatural enantiomer the D isomer.**

The R group of an amino acid can be H, alkyl, aryl, or an alkyl chain containing an N, O, or S atom. Representative examples are listed in Table 19.4. All amino acids have common names, which are abbreviated by a three-letter or one-letter designation. For example, glycine is often written as the three-letter abbreviation **Gly**, or the one-letter abbreviation **G**. These abbreviations are also given in Table 19.4. A complete list of the 20 naturally occurring amino acids is found in Figure 28.2.

Problem 19.25

Draw both enantiomers of each amino acid and label them as *R* or *S*: (a) phenylalanine; (b) methionine.


19.14B Acid–Base Properties

An amino acid is both an acid and a base.

- The NH₂ group has a nonbonded electron pair, making it a base.
- The COOH group has an acidic proton, making it an acid.


Amino acids are never uncharged neutral compounds. They exist as salts, so they have very high melting points and are very soluble in water.

- Proton transfer from the acidic carboxy group to the basic amino group forms a salt called a **zwitterion**, which contains both a positive and a negative charge.




In actuality, an amino acid can exist in three different forms, depending on the pH of the aqueous solution in which it is dissolved.


When the pH of a solution is ~6, alanine (R = CH₃) exists in its zwitterionic form (**A**), having no net charge. In this form the carboxy group bears a negative charge—it is a **carboxylate anion**—and the amino group bears a net positive charge (an **ammonium cation**).



When strong acid is added to lower the pH (≤ 2), the carboxylate anion is protonated and the **amino acid has a net positive charge** (form **B**).



When strong base is added to **A** to raise the pH (≥ 10), the ammonium cation is deprotonated and the **amino acid has a net negative charge** (form **C**).



Thus, **alanine exists in one of three different forms depending on the pH of the solution in which it is dissolved**. If the pH of a solution is gradually increased from 2 to 10, the following process occurs.

- At low pH alanine has a net (+) charge (form B).
- As the pH is increased to ~6, the carboxy group is deprotonated, and the amino acid exists as a zwitterion with no overall charge (form A).
- At high pH, the ammonium cation is deprotonated, and the amino acid has a net (-) charge (form C).


These reactions are summarized in Figure 19.11.

Problem 19.26

Explain why amino acids, unlike most other organic compounds, are insoluble in organic solvents like diethyl ether.

Figure 19.11

Summary of the acid–base reactions of alanine



Problem 19.27 Draw the positively charged, neutral, and negatively charged forms for the amino acid glycine. Which species predominates at pH 11? Which species predominates at pH 1?

19.14C Isoelectric Point

Because a protonated amino acid has at least two different protons that can be removed, a pK_a value is reported for each of these protons. For example, the pK_a of the carboxy proton of alanine is 2.35 and the pK_a of the ammonium proton is 9.87. Table 28.1 lists these values for all 20 amino acids.

- The pH at which the amino acid exists primarily in its neutral form is called its *isoelectric point*, abbreviated as *pI*.

Generally, the isoelectric point is the average of both pK_a values of an amino acid:

$$\text{Isoelectric point} = pI = \frac{pK_a(\text{COOH}) + pK_a(\text{NH}_3^+)}{2}$$

$$\text{For alanine: } pI = \frac{2.35 + 9.87}{2} = 6.12 \text{ (pI of alanine)}$$

More information on the isoelectric point can be found in Section 28.1 and Problem 28.35.

Problem 19.28 The pK_a values for the carboxy and ammonium protons of phenylalanine are 2.58 and 9.24, respectively. What is the isoelectric point of phenylalanine? Draw the structure of phenylalanine at its isoelectric point.

Problem 19.29 Explain why the pK_a of the COOH group of glycine is much lower than the pK_a of the COOH of acetic acid (2.35 compared to 4.8).

KEY CONCEPTS

Carboxylic Acids and the Acidity of the O–H Bond


General Facts

- Carboxylic acids contain a carboxy group (COOH). The central carbon is sp^2 hybridized and trigonal planar (19.1).
- Carboxylic acids are identified by the suffixes *-oic acid*, *carboxylic acid*, or *-ic acid* (19.2).
- Carboxylic acids are polar compounds that exhibit hydrogen bonding interactions (19.3).

Summary of Spectroscopic Absorptions (19.4)

IR absorptions	C=O	$\sim 1710 \text{ cm}^{-1}$
	O–H	$3500\text{--}2500 \text{ cm}^{-1}$ (very broad and strong)
^1H NMR absorptions	O–H	10–12 ppm (highly deshielded proton)
	C–H α to COOH	2–2.5 ppm (somewhat deshielded $C_{sp^3}\text{--H}$)
^{13}C NMR absorption	C=O	170–210 ppm (highly deshielded carbon)

General Acid–Base Reaction of Carboxylic Acids (19.9)




- Carboxylic acids are especially acidic because carboxylate anions are resonance stabilized.
- For equilibrium to favor the products, the base must have a conjugate acid with a $pK_a > 5$. Common bases are listed in Table 19.3.

Factors That Affect Acidity

Resonance effects

- A carboxylic acid is more acidic than an alcohol or phenol because its conjugate base is more effectively stabilized by resonance (19.9).




Inductive effects

- Acidity increases with the presence of electron-withdrawing groups (like the electronegative halogens) and decreases with the presence of electron-donating groups (like polarizable alkyl groups) (19.10).

Substituted benzoic acids

- Electron-donor groups (**D**) make a substituted benzoic acid less acidic than benzoic acid.
- Electron-withdrawing groups (**W**) make a substituted benzoic acid more acidic than benzoic acid.












Other Facts

- Extraction is a useful technique for separating compounds having different solubility properties. Carboxylic acids can be separated from other organic compounds by extraction, because aqueous base converts a carboxylic acid into a water-soluble carboxylate anion (19.12).
- A sulfonic acid (RSO_3H) is a strong acid because it forms a weak, resonance-stabilized conjugate base on deprotonation (19.13).
- Amino acids have an amino group on the α carbon to the carboxy group [$\text{RCH}(\text{NH}_2)\text{COOH}$]. Amino acids exist as zwitterions at $\text{pH} \approx 6$. Adding acid forms a species with a net (+1) charge [$\text{RCH}(\text{NH}_3^+)\text{COOH}$]. Adding base forms a species with a net (-1) charge [$\text{RCH}(\text{NH}_2)\text{COO}^-$] (19.14).



PROBLEMS

Nomenclature

19.30 Give the IUPAC name for each compound.



19.31 Draw the structure corresponding to each name.







- 19.32 Draw the structures and give the IUPAC names for the carboxylic acids having molecular formula $C_5H_{10}O_2$. Then give the IUPAC names for the sodium salts that result from treatment of each carboxylic acid with NaOH.

Physical Properties



- 19.33 Rank the compounds in each group in order of increasing boiling point.
- $CH_3CH_2CH_2CH_2COOH$, $(CH_3CH_2CH_2)_2O$, $CH_3(CH_2)_5OH$
 - $CH_3COCH_2CH(CH_3)_2$, $(CH_3)_2CHCH_2COOH$, $(CH_3)_2CHCH_2CH(OH)CH_3$

Preparation of Carboxylic Acids

- 19.34 Draw the organic products formed in each reaction.






- 19.35 Identify the lettered compounds in each reaction sequence.








Acid–Base Reactions; General Questions on Acidity




- 19.36 Using the pK_a table in Appendix A, determine whether each of the following bases is strong enough to deprotonate the three compounds listed below. Bases: [1] ^-OH ; [2] $CH_3CH_2^-$; [3] $^-NH_2$; [4] NH_3 ; [5] $HC\equiv C^-$.




- 19.37 Draw the products of each acid–base reaction, and using the pK_a table in Appendix A, determine if equilibrium favors the reactants or products.




- 19.38 Which compound in each pair has the lower pK_a ? Which compound in each pair has the stronger conjugate base?



19.39 Rank the compounds in each group in order of increasing acidity.




19.40 Rank the compounds in each group in order of increasing basicity.



19.41 Match the pK_a values to the appropriate structure. pK_a values: 0.28, 1.24, 2.66, 2.86, and 3.12. Compounds: (a) FCH_2COOH ; (b) CF_3COOH ; (c) F_2CHCOOH ; (d) ICH_2COOH ; (e) BrCH_2COOH .

19.42 Although codeine occurs in low concentration in the opium poppy, most of the codeine used in medicine is prepared from morphine (the principal component of opium) by the following reaction. Explain why selective methylation occurs at only one OH in morphine to give codeine. Codeine is a less potent and less addictive analgesic than morphine.




19.43 Explain each statement.


- a. The pK_a of *p*-nitrophenol is lower than the pK_a of phenol (7.2 vs. 10).
 b. The pK_a of *p*-nitrophenol is lower than the pK_a of *m*-nitrophenol (7.2 vs. 8.3).

19.44 Explain the following statement. Although 2-methoxyacetic acid ($\text{CH}_3\text{OCH}_2\text{COOH}$) is a stronger acid than acetic acid (CH_3COOH), *p*-methoxybenzoic acid ($\text{CH}_3\text{OC}_6\text{H}_4\text{COOH}$) is a weaker acid than benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$).


19.45 Explain why the pK_a of compound **A** is lower than the pK_a 's of both compounds **B** and **C**.




19.46 Rank the following compounds in order of increasing acidity and explain in detail your choice of order.




19.47 Explain the following result. Acetic acid (CH_3COOH), labeled at its OH oxygen with the uncommon ^{18}O isotope, was treated with aqueous base, and then the solution was acidified. Two products having the ^{18}O label at different locations were formed.



- 19.48** Draw all resonance structures of the conjugate bases formed by removal of the labeled protons (H_a , H_b , and H_c) in 1,3-cyclohexanedione and acetanilide. For each compound, rank these protons in order of increasing acidity and explain the order you chose.



1,3-cyclohexanedione




acetanilide

- 19.49** As we will see in Chapter 23, C–H bonds are sometimes more acidic than O–H bonds. Explain why the pK_a of $CH_2(CHO)_2$ is lower than the pK_a of $HO(CH_2)_3OH$ (9 vs. 16).

- 19.50** Explain why sulfonic acids (RSO_3H) are stronger organic acids than carboxylic acids ($RCOOH$).



- 19.51** Identify **X** in the following equation, and explain how hexanoic acid, the chapter-opening molecule, is formed by this stepwise reaction sequence.



- 19.52** The pK_a of acetamide (CH_3CONH_2) is 16. Draw the structure for its conjugate base and explain why acetamide is less acidic than CH_3COOH .

Extraction

- 19.53** Write out the steps needed to separate hydrocarbon **A** and carboxylic acid **B** by using an extraction procedure.

**A****B**

- 19.54** Because phenol (C_6H_5OH) is less acidic than a carboxylic acid, it can be deprotonated by $NaOH$ but not by the weaker base $NaHCO_3$. Using this information, write out an extraction sequence that can be used to separate C_6H_5OH from cyclohexanol. Show what compound is present in each layer at each stage of the process, and if it is present in its neutral or ionic form.

- 19.55** Can octane and 1-octanol be separated using an aqueous extraction procedure? Explain why or why not.



Spectroscopy

- 19.56** Identify each compound from its spectral data.



- Molecular formula: $C_3H_5ClO_2$
IR: $3500\text{--}2500\text{ cm}^{-1}$, 1714 cm^{-1}
 1H NMR data: 2.87 (triplet, 2 H), 3.76 (triplet, 2 H), and 11.8 (singlet, 1 H) ppm
- Molecular formula: $C_8H_8O_3$
IR: $3500\text{--}2500\text{ cm}^{-1}$, 1688 cm^{-1}
 1H NMR data: 3.8 (singlet, 3 H), 7.0 (doublet, 2 H), 7.9 (doublet, 2 H), and 12.7 (singlet, 1 H) ppm
- Molecular formula: $C_8H_8O_3$
IR: $3500\text{--}2500\text{ cm}^{-1}$, 1710 cm^{-1}
 1H NMR data: 4.7 (singlet, 2 H), 6.9–7.3 (multiplet, 5 H), and 11.3 (singlet, 1 H) ppm

19.57 Use the ^1H NMR and IR spectra given below to identify the structures of two isomers (**A** and **B**) having molecular formula $\text{C}_4\text{H}_8\text{O}_2$.


Compound **A**:



Compound **B**:




19.58 An unknown compound **C** (molecular formula $\text{C}_4\text{H}_8\text{O}_3$) exhibits IR absorptions at 3600–2500 and 1734 cm^{-1} , as well as the following ^1H NMR spectrum. What is the structure of **C**?



19.59 Propose a structure for **D** (molecular formula $C_9H_9ClO_2$) consistent with the given spectroscopic data.

^{13}C NMR signals at 30, 36, 128, 130, 133, 139, and 179 ppm



19.60 What is the structure of a carboxylic acid (molecular formula $C_6H_{12}O_2$) that gives three singlets in its 1H NMR spectrum at 1.1, 2.2, and 11.9 ppm?




19.61 A monomer needed to synthesize polyethylene terephthalate (PET), a polymer used to make plastic sheeting and soft drink bottles (Section 22.16), shows a strong absorption in its IR spectrum at 1692 cm^{-1} and two singlets in its 1H NMR spectrum at 8.2 and 10.0 ppm. What is the structure of this monomer (molecular formula $C_8H_6O_4$)?

19.62 Match the ^{13}C NMR data to the appropriate structure.

Spectrum [1]: signals at 14, 22, 27, 34, 181 ppm

Spectrum [2]: signals at 27, 39, 186 ppm


Spectrum [3]: signals at 22, 26, 43, 180 ppm



19.63 γ -Butyrolactone ($C_4H_6O_2$, GBL) is a biologically inactive compound that is converted to the biologically active recreational drug GHB (Section 19.5) by a lactonase enzyme in the body. Since γ -butyrolactone is more fat soluble than GHB, it is more readily absorbed by tissues and thus produces a faster onset of physiological symptoms. γ -Butyrolactone shows an absorption in its IR spectrum at 1770 cm^{-1} and the following 1H NMR spectral data: 2.28 (multiplet, 2 H), 2.48 (triplet, 2 H), and 4.35 (triplet, 2 H) ppm. What is the structure of γ -butyrolactone?

Amino Acids


19.64 Threonine is a naturally occurring amino acid that has two stereogenic centers.



a. Draw the four possible stereoisomers using wedges and dashes.

b. The naturally occurring amino acid has the $2S,3R$ configuration at its two stereogenic centers. Which structure does this correspond to?

19.65 Proline is an unusual amino acid because its N atom on the α carbon is part of a five-membered ring.



a. Draw both enantiomers of proline.

b. Draw proline in its zwitterionic form.

19.66 For each amino acid $[RCH(NH_2)COOH]$, draw its neutral, positively charged, and negatively charged forms. Which form predominates at pH = 1, 6, and 11? What is the structure of each amino acid at its isoelectric point?

a. methionine ($R = CH_2CH_2SCH_3$)

b. serine ($R = CH_2OH$)

19.67 Calculate the isoelectric point for each amino acid.

a. cysteine: $pK_a(\text{COOH}) = 2.05$; $pK_a(\alpha\text{-NH}_3^+) = 10.25$

b. methionine: $pK_a(\text{COOH}) = 2.28$; $pK_a(\alpha\text{-NH}_3^+) = 9.21$

Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

20

- 20.1 Introduction
- 20.2 General reactions of carbonyl compounds
- 20.3 A preview of oxidation and reduction
- 20.4 Reduction of aldehydes and ketones
- 20.5 The stereochemistry of carbonyl reduction
- 20.6 Enantioselective carbonyl reductions
- 20.7 Reduction of carboxylic acids and their derivatives
- 20.8 Oxidation of aldehydes
- 20.9 Organometallic reagents
- 20.10 Reaction of organometallic reagents with aldehydes and ketones
- 20.11 Retrosynthetic analysis of Grignard products
- 20.12 Protecting groups
- 20.13 Reaction of organometallic reagents with carboxylic acid derivatives
- 20.14 Reaction of organometallic reagents with other compounds
- 20.15 α,β -Unsaturated carbonyl compounds
- 20.16 Summary—The reactions of organometallic reagents
- 20.17 Synthesis



Juvenile hormones are a group of structurally related molecules that regulate the complex life cycle of an insect. In particular, they maintain the juvenile stage until an insect is ready for adulthood. This property has been exploited to control mosquitoes and other pests infecting both livestock and crops. Application of synthetic juvenile hormones to the egg or larva of an insect prevents maturation. With no sexually mature adults to propagate the next generation, the insect population is reduced. Juvenile hormones are synthesized by reactions that form new carbon-carbon bonds, thus allowing complex organic molecules to be prepared from simple starting materials. In Chapter 20 we learn about these useful organic reactions.

Chapters 20 through 24 of this text discuss carbonyl compounds—aldehydes, ketones, acid halides, esters, amides, and carboxylic acids. **The carbonyl group is perhaps the most important functional group in organic chemistry**, because its electron-deficient carbon and easily broken π bond make it susceptible to a wide variety of useful reactions.


We begin by examining the similarities and differences between two broad classes of carbonyl compounds. We will then spend the remainder of Chapter 20 on reactions that are especially important in organic synthesis. Chapters 21 and 22 present specific reactions that occur at the carbonyl carbon, and Chapters 23 and 24 concentrate on reactions occurring at the α carbon to the carbonyl group.

Although Chapter 20 is “jam-packed” with reactions, most of them follow one of two general pathways, so they can be classified in a well-organized fashion, provided you remember a few basic principles. Keep in mind the following fundamental themes about reactions:


- Nucleophiles attack electrophiles.
- π Bonds are easily broken.
- Bonds to good leaving groups are easily cleaved heterolytically.

20.1 Introduction

Two broad classes of compounds contain a **carbonyl group**:




[1] Compounds that have only carbon and hydrogen atoms bonded to the carbonyl group



- An **aldehyde** has at least one H atom bonded to the carbonyl group.
- A **ketone** has two alkyl or aryl groups bonded to the carbonyl group.

[2] Compounds that contain an electronegative atom bonded to the carbonyl group




These include **carboxylic acids**, **acid chlorides**, **esters**, and **amides**, as well as other similar compounds discussed in Chapter 22. Each of these compounds contains an electronegative atom (Cl, O, or N) capable of acting as a **leaving group**. Acid chlorides, esters, and amides are often called **carboxylic acid derivatives**, because they can be synthesized from carboxylic acids (Chapter 22). Since each compound contains an acyl group ($\text{RCO}-$), they are also called **acyl derivatives**.

- The presence or absence of a leaving group on the carbonyl carbon determines the type of reactions these compounds undergo (Section 20.2).

The carbonyl carbon atom is sp^2 hybridized and trigonal planar, and all bond angles are $\sim 120^\circ$. The double bond of a carbonyl group consists of one σ bond and one π bond. The π bond is

Figure 20.1

Electrostatic potential map of formaldehyde, $\text{CH}_2=\text{O}$




- An electrostatic potential map shows the electron-deficient carbon and the electron-rich oxygen atom of the carbonyl group.


formed by the overlap of two p orbitals, and extends above and below the plane. In these features the carbonyl group resembles the trigonal planar, sp^2 hybridized carbons of a $\text{C}-\text{C}$ double bond.




The aldehyde α -sinensal (Problem 20.1) is the major compound responsible for the orange-like odor of mandarin oil, obtained from the mandarin tree in southern China.



In one important way, though, a $\text{C}=\text{O}$ and $\text{C}=\text{C}$ are very different. **The electronegative oxygen atom in the carbonyl group means that the bond is polarized, making the carbonyl carbon electron deficient.** Using a resonance description, the carbonyl group is represented by two resonance structures, with a charge-separated resonance structure a minor contributor to the hybrid. An electrostatic potential plot for formaldehyde, the simplest aldehyde, is shown in Figure 20.1. It clearly indicates the polarized carbonyl group.




Problem 20.1



- What orbitals are used to form the indicated bonds in α -sinensal?
- In what type of orbitals do the lone pairs on O reside?


20.2 General Reactions of Carbonyl Compounds

With what types of reagents should a carbonyl group react? The electronegative oxygen makes the carbonyl carbon electrophilic, and because it is trigonal planar, a carbonyl carbon is uncrowded. Moreover, a carbonyl group has an easily broken π bond.




As a result, **carbonyl compounds react with nucleophiles.** The outcome of nucleophilic attack, however, depends on the identity of the carbonyl starting material.

- Aldehydes and ketones undergo nucleophilic addition.



- Carbonyl compounds that contain leaving groups undergo nucleophilic substitution.




Let's examine each of these general reactions individually.

20.2A Nucleophilic Addition to Aldehydes and Ketones

Aldehydes and ketones react with nucleophiles to form addition products by the two-step process shown in Mechanism 20.1: **nucleophilic attack** followed by **protonation**.



Mechanism 20.1 Nucleophilic Addition—A Two-Step Process




- In Step [1], the nucleophile (**:Nu⁻**) attacks the **electrophilic carbonyl**. As the new bond to the nucleophile forms, the π bond is broken, moving an electron pair out on the oxygen atom. This forms an sp^3 hybridized intermediate.
- In Step [2], **protonation of the negatively charged oxygen atom by H₂O** (or another proton source) forms the addition product.

More examples of nucleophilic addition to aldehydes and ketones are discussed in Chapter 21.

The net result is that the π bond is broken, two new σ bonds are formed, and the elements of H and Nu are added across the π bond. Nucleophilic addition with two different nucleophiles—**hydride (H⁻)** and **carbanions (R⁻)**—is discussed in Chapter 20.

Aldehydes are more reactive than ketones towards nucleophilic attack for both steric and electronic reasons.




- The two R groups bonded to the ketone carbonyl group make it more crowded, so nucleophilic attack is more difficult.
- The two electron-donor R groups stabilize the partial charge on the carbonyl carbon of a ketone, making it more stable and less reactive.

20.2B Nucleophilic Substitution of RCOZ (Z = Leaving Group)

Carbonyl compounds with leaving groups react with nucleophiles to form substitution products by the two-step process shown in Mechanism 20.2: **nucleophilic attack**, followed by **loss of the leaving group**.



Mechanism 20.2 Nucleophilic Substitution—A Two-Step Process



- In Step [1], the nucleophile (:Nu⁻) attacks the electrophilic carbonyl, forming an sp^3 hybridized intermediate. This step is identical to nucleophilic addition.
- Step [2] is different. Because the intermediate contains an electronegative atom Z, **Z can act as a leaving group**. To do so, an electron pair on O re-forms the π bond, and Z leaves with the electron pair in the C–Z bond.


The net result is that Nu replaces Z—a nucleophilic substitution reaction. This reaction is often called **nucleophilic acyl substitution** to distinguish it from the nucleophilic substitution reactions at sp^3 hybridized carbons discussed in Chapter 7. Nucleophilic substitution with two different nucleophiles—**hydride (H⁻)** and **carbanions (R⁻)**—is discussed in Chapter 20. Other nucleophiles are examined in Chapter 22.

Carboxylic acid derivatives differ greatly in their reactivity towards nucleophiles. The order in which they react parallels the leaving group ability of the group Z bonded to the carbonyl carbon.

Recall from Section 7.7 that the weaker the base, the better the leaving group.

- The better the leaving group Z, the more reactive RCOZ is in nucleophilic acyl substitution.


Thus, the following trends result:



- Acid chlorides (RCOCl), which have the best leaving group (Cl^-), are the most reactive carboxylic acid derivatives, and amides (RCONH₂), which have the worst leaving group ($^-NH_2$), are the least reactive.
- Carboxylic acids (RCOOH) and esters (RCOOR'), which have leaving groups of similar basicity (^-OH and $^-OR'$), fall in the middle.


Nucleophilic addition and nucleophilic acyl substitution involve the *same* first step—**nucleophilic attack on the electrophilic carbonyl group** to form a tetrahedral intermediate. The difference between them is what then happens to this intermediate. Aldehydes and ketones cannot undergo substitution because they have no leaving group bonded to the newly formed sp^3

hybridized carbon. Nucleophilic substitution with an aldehyde, for example, would form H^- , an extremely strong base and therefore a very poor (and highly unlikely) leaving group.



Problem 20.2 Which compounds undergo nucleophilic addition and which undergo substitution?
 a. $(\text{CH}_3)_2\text{C}=\text{O}$ b. $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCl}$ c. $\text{CH}_3\text{COOCH}_3$ d. $\text{C}_6\text{H}_5\text{CHO}$

Problem 20.3 Which carbonyl groups in the anticancer drug taxol (Section 5.5) will undergo nucleophilic addition and which will undergo nucleophilic substitution?



Problem 20.4 Which compound in each pair is more reactive towards nucleophilic attack?
 a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ or $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCH}_3$ c. $\text{CH}_3\text{CH}_2\text{COCl}$ or $\text{CH}_3\text{COOCH}_3$
 b. $\text{CH}_3\text{CH}_2\text{COCH}_3$ or $\text{CH}_3\text{CH}(\text{CH}_3)\text{COCH}_2\text{CH}_3$ d. $\text{CH}_3\text{COOCH}_3$ or $\text{CH}_3\text{CONHCH}_3$


To show how these general principles of nucleophilic substitution and addition apply to carbonyl compounds, we are going to discuss oxidation and reduction reactions, and reactions with organometallic reagents—compounds that contain carbon–metal bonds. We begin with reduction to build on what you learned previously in Chapter 12.

20.3 A Preview of Oxidation and Reduction

Recall the definitions of oxidation and reduction presented in Section 12.1:


- Oxidation results in an increase in the number of C–Z bonds (usually C–O bonds) or a decrease in the number of C–H bonds.
- Reduction results in a decrease in the number of C–Z bonds (usually C–O bonds) or an increase in the number of C–H bonds.

Carbonyl compounds are either reactants or products in many of these reactions, as illustrated in the accompanying diagram. For example, because aldehydes fall in the middle of this scheme, they can be both oxidized and reduced. Carboxylic acids and their derivatives (RCOZ), on the other hand, are already highly oxidized, so their only useful reaction is reduction.




The three most useful oxidation and reduction reactions of carbonyl starting materials can be summarized as follows:

[1] Reduction of aldehydes and ketones to alcohols (Sections 20.4–20.6)




Aldehydes and ketones are reduced to 1° and 2° alcohols, respectively.

[2] Reduction of carboxylic acid derivatives (Section 20.7)



The reduction of carboxylic acids and their derivatives gives a variety of products, depending on the identity of Z and the nature of the reducing agent. The usual products are aldehydes or 1° alcohols.

[3] Oxidation of aldehydes to carboxylic acids (Section 20.8)




The most useful oxidation reaction of carbonyl compounds is the oxidation of aldehydes to carboxylic acids.

We begin with reduction, because the mechanisms of reduction reactions follow directly from the general mechanisms for nucleophilic addition and substitution.

20.4 Reduction of Aldehydes and Ketones


LiAlH₄ and NaBH₄ serve as a source of H⁻, but there are no free H⁻ ions present in reactions with these reagents.

The most useful reagents for reducing aldehydes and ketones are the metal hydride reagents (Section 12.2). The two most common metal hydride reagents are **sodium borohydride (NaBH₄)** and **lithium aluminum hydride (LiAlH₄)**. These reagents contain a polar metal–hydrogen bond that serves as a source of the nucleophile hydride, **H⁻**. LiAlH₄ is a stronger reducing agent than NaBH₄, because the Al–H bond is more polar than the B–H bond.




20.4A Reduction with Metal Hydride Reagents

Treating an aldehyde or a ketone with NaBH_4 or LiAlH_4 , followed by water or some other proton source, affords an **alcohol**. This is an addition reaction because the elements of H_2 are added across the π bond, but it is also a reduction because the product alcohol has fewer C–O bonds than the starting carbonyl compound.




LiAlH_4 reductions must be carried out under anhydrous conditions, because water reacts violently with the reagent. Water is added to the reaction mixture (to serve as a proton source) *after* the reduction with LiAlH_4 is complete.

The product of this reduction reaction is a **1° alcohol** when the starting carbonyl compound is an aldehyde, and a **2° alcohol** when it is a ketone.




NaBH_4 selectively reduces aldehydes and ketones in the presence of most other functional groups. Reductions with NaBH_4 are typically carried out in CH_3OH as solvent. LiAlH_4 reduces aldehydes and ketones and many other functional groups as well (Sections 12.6 and 20.7).

Problem 20.5 What alcohol is formed when each compound is treated with NaBH_4 in CH_3OH ?




Problem 20.6 What aldehyde or ketone is needed to prepare each alcohol by metal hydride reduction?



Problem 20.7 Why can't 1-methylcyclohexanol be prepared from a carbonyl compound by reduction?

20.4B The Mechanism of Hydride Reduction

Hydride reduction of aldehydes and ketones occurs via the general mechanism of nucleophilic addition—that is, **nucleophilic attack** followed by **protonation**. Mechanism 20.3 is shown using LiAlH_4 , but an analogous mechanism can be written for NaBH_4 .


Mechanism 20.3 LiAlH_4 Reduction of RCHO and $\text{R}_2\text{C}=\text{O}$ 

- In Step [1], the nucleophile (AlH_4^-) donates H^- to the carbonyl group, cleaving the π bond, and moving an electron pair onto oxygen. This forms a new C–H bond.
- In Step [2], the alkoxide is protonated by H_2O (or CH_3OH) to form the alcohol reduction product. This acid–base reaction forms a new O–H bond.

- The net result of adding H^- (from NaBH_4 or LiAlH_4) and H^+ (from H_2O) is the addition of the elements of H_2 to the carbonyl π bond.

20.4C Catalytic Hydrogenation of Aldehydes and Ketones


Catalytic hydrogenation also reduces aldehydes and ketones to 1° and 2° alcohols, respectively, using H_2 and Pd-C (or another metal catalyst). H_2 adds to the $\text{C}=\text{O}$ in much the same way that it adds to the $\text{C}=\text{C}$ of an alkene (Section 12.3). The metal catalyst (Pd-C) provides a surface that binds the carbonyl starting material and H_2 , and two H atoms are sequentially transferred with cleavage of the π bond.



When a compound contains both a carbonyl group and a carbon–carbon double bond, selective reduction of one functional group can be achieved by proper choice of reagent.

- A $\text{C}=\text{C}$ is reduced faster than a $\text{C}=\text{O}$ with H_2 (Pd-C).
- A $\text{C}=\text{O}$ is readily reduced with NaBH_4 and LiAlH_4 , but a $\text{C}=\text{C}$ is inert.

Thus, 2-cyclohexenone, a compound that contains both a carbon–carbon double bond and a carbonyl group, can be reduced to three different compounds—an allylic alcohol, a carbonyl compound, or an alcohol—depending on the reagent.



Problem 20.8

Draw the products formed when $\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}=\text{CH}_2$ is treated with each reagent: (a) LiAlH_4 , then H_2O ; (b) NaBH_4 in CH_3OH ; (c) H_2 (1 equiv), Pd-C; (d) H_2 (excess), Pd-C; (e) NaBH_4 (excess) in CH_3OH ; (f) NaBD_4 in CH_3OH .

The reduction of aldehydes and ketones is a common reaction used in the synthesis of many useful natural products. Two examples are shown in Figure 20.2.


20.5 The Stereochemistry of Carbonyl Reduction

Recall from Section 9.15 that an achiral starting material gives a racemic mixture when a new stereogenic center is formed.


The stereochemistry of carbonyl reduction follows the same principles we have previously learned. Reduction converts a **planar sp^2 hybridized carbonyl carbon to a tetrahedral sp^3 hybridized carbon**. What happens when a new stereogenic center is formed in this process? With an achiral reagent like NaBH_4 or LiAlH_4 , a racemic product is obtained. For example, NaBH_4 in CH_3OH solution reduces 2-butanone, an achiral ketone, to 2-butanol, an alcohol that contains a new stereogenic center. Both enantiomers of 2-butanol are formed in equal amounts.

Figure 20.2


NaBH_4 reductions used in organic synthesis




The male musk deer, a small antlerless deer found in the mountain regions of China and Tibet, has long been hunted for its musk, a strongly scented liquid used in early medicine and later in perfumery.



- **Muscone** is the major compound in musk, one of the oldest known ingredients in perfumes. Musk was originally isolated from the male musk deer, but it can now be prepared synthetically in the laboratory in a variety of ways.






Why is a racemic mixture formed? Because the carbonyl carbon is sp^2 hybridized and planar, hydride can approach the double bond with equal probability from both sides of the plane, forming two alkoxides, which are **enantiomers** of each other. Protonation of the alkoxides gives an equal amount of two alcohols, which are also **enantiomers**.



- **Conclusion:** Hydride reduction of an achiral ketone with LiAlH_4 or NaBH_4 gives a racemic mixture of two alcohols when a new stereogenic center is formed.

Problem 20.9 Draw the products formed (including stereoisomers) when each compound is reduced with NaBH_4 in CH_3OH .




20.6 Enantioselective Carbonyl Reductions

20.6A CBS Reagents

One enantiomer can be formed selectively from the reduction of a carbonyl group, provided a **chiral reducing agent** is used. This strategy is identical to that employed in the Sharpless asymmetric epoxidation reaction (Section 12.15). A reduction that forms one enantiomer predominantly or exclusively is an **enantioselective** or **asymmetric reduction**.


Many different chiral reducing agents have now been prepared for this purpose. One such reagent, formed by reacting borane (BH_3) with a heterocycle called an **oxazaborolidine**, has one stereogenic center (and thus two enantiomers).

Two enantiomers of the chiral CBS reducing agent



(*S*)-2-methyl-CBS-oxazaborolidine

(*S*)-CBS reagent



(*R*)-2-methyl-CBS-oxazaborolidine

(*R*)-CBS reagent

These reagents are called the (*S*)-CBS reagent and the (*R*)-CBS reagent, named for Corey, Bakshi, and Shibata, the chemists who developed these versatile reagents. One B–H bond of BH_3 serves as the source of hydride in this reduction. The stereochemistry of the new stereogenic center in the product is often predictable. For ketones having the general structure $\text{C}_6\text{H}_5\text{COR}$, draw the starting material with the aryl group on the left side of the carbonyl, as shown with acetophenone. Then, to draw the product, keep in mind:

- The (*S*)-CBS reagent delivers hydride (H^-) from the front side of the $\text{C}=\text{O}$. This generally affords the *R* alcohol as the major product.
- The (*R*)-CBS reagent delivers hydride (H^-) from the back side of the $\text{C}=\text{O}$. This generally affords the *S* alcohol as the major product.







Figure 20.3 Enantioselective reduction—A key step in the synthesis of salmeterol

- (*R*)-Salmeterol is a long-acting bronchodilator used for the treatment of asthma.
- In this example, the (*R*)-CBS reagent adds the new H atom from behind, the same result observed with acetophenone and propiophenone. In this case, however, alcohol **A** has the *R* configuration using the rules for assigning priority in Chapter 5.

These reagents are highly enantioselective. Treatment of propiophenone with the (*S*)-CBS reagent forms the *R* alcohol in 97% enantiomeric excess (*ee*). Enantioselective reductions are key steps in the synthesis of several widely used drugs, including salmeterol, a long-acting bronchodilator shown in Figure 20.3. This new technology provides access to single enantiomers of biologically active compounds, often previously available only as a racemic mixture.




Problem 20.10 What reagent is needed to reduce **A** to **B**, an intermediate in the synthesis of the antidepressant (*R*)-fluoxetine (trade name: Prozac)?




20.6B Enantioselective Biological Reduction

Although laboratory reduction reactions often do not proceed with 100% enantioselectivity, biological reductions that occur in cells *always* proceed with complete selectivity, forming a single enantiomer. NaBH_4 or chiral boranes are not the reducing agents for these processes. In cells, the reducing agent is **NADH**.




NADH is a **coenzyme**, an organic molecule that can function only in the presence of an enzyme. The active site of the enzyme binds both the carbonyl substrate and NADH, keeping them in close proximity. **NADH then donates H^-** in much the same way as a metal hydride reagent; that is, reduction consists of nucleophilic attack followed by protonation.

- In Step [1], **NADH donates H^- to the carbonyl group** to form an alkoxide. In the process, NADH is converted to NAD^+ .
- In Step [2], the alkoxide is protonated by the aqueous medium.




Pyruvic acid is formed during the metabolism of glucose. During periods of strenuous exercise, when there is insufficient oxygen to metabolize pyruvic acid to CO_2 , pyruvic acid is reduced to lactic acid. The tired feeling of sore muscles is a result of lactic acid accumulation.

This reaction is completely enantioselective. For example, reduction of pyruvic acid with NADH catalyzed by lactate dehydrogenase affords a single enantiomer of lactic acid with the *S* configuration. NADH reduces a variety of different carbonyl compounds in biological systems. The configuration of the product (*R* or *S*) depends on the enzyme used to catalyze the process.



Niacin can be obtained from foods such as soybeans, which contain it naturally, and from breakfast cereals, which are fortified with it.

NAD^+ , the oxidized form of NADH, is a **biological oxidizing agent** capable of oxidizing alcohols to carbonyl compounds (it forms NADH in the process). NAD^+ is synthesized from the vitamin niacin, which can be obtained from soybeans among other dietary sources. Breakfast cereals are fortified with niacin to help people consume their recommended daily allowance of this B vitamin.




20.7 Reduction of Carboxylic Acids and Their Derivatives

The reduction of carboxylic acids and their derivatives (RCOZ) is complicated because the products obtained depend on the identity of both the leaving group (Z) and the reducing agent. Metal hydride reagents are the most useful reducing reagents. **Lithium aluminum hydride is a strong reducing agent that reacts with all carboxylic acid derivatives.** Two other related but milder reducing agents are also used.


- [1] **Diisobutylaluminum hydride**, $[(\text{CH}_3)_2\text{CHCH}_2]_2\text{AlH}$, abbreviated as **DIBAL-H**, has two bulky isobutyl groups, which make this reagent less reactive than LiAlH_4 . The single H atom is donated as H^- in hydride reductions.
- [2] **Lithium tri-*tert*-butoxyaluminum hydride**, $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]_3$, has three electronegative oxygen atoms bonded to aluminum, which make this reagent less nucleophilic than LiAlH_4 .

LiAlH_4 is a strong, nonselective reducing agent. DIBAL-H and $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]_3$ are milder, more selective reducing agents.




20.7A Reduction of Acid Chlorides and Esters


Acid chlorides and esters can be reduced to either aldehydes or 1° alcohols, depending on the reagent.



- LiAlH_4 converts RCOCl and RCOOR' to 1° alcohols.
- A milder reducing agent (**DIBAL-H** or $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]_3$) converts RCOCl or RCOOR' to RCHO at low temperatures.



In the reduction of an acid chloride, Cl^- comes off as the leaving group.




In the reduction of the ester, CH_3O^- comes off as the leaving group, which is then protonated by H_2O to form CH_3OH .

Mechanism 20.4 illustrates why two different products are possible. It can be conceptually divided into two parts: **nucleophilic substitution** to form an aldehyde, followed by **nucleophilic addition** to the aldehyde to form a 1° alcohol. A general mechanism is drawn using LiAlH_4 as reducing agent.




Mechanism 20.4 Reduction of RCOCl and RCOOR' with a Metal Hydride Reagent

Part [1] Nucleophilic substitution forms an aldehyde.



- **Nucleophilic attack of H^-** (from LiAlH_4) in Step [1] forms a tetrahedral intermediate with a leaving group Z.
- In Step [2], the π bond is re-formed and **Z^- comes off**. The overall result of the addition of H^- and elimination of Z^- is the substitution of H for Z.


Part [2] Nucleophilic addition forms a 1° alcohol.




- **Nucleophilic attack of H^-** (from LiAlH_4) in Step [3] forms an alkoxide.
- **Protonation of the alkoxide** by H_2O in Step [4] forms the alcohol reduction product. The overall result of Steps [3] and [4] is addition of H_2 .

With milder reducing agents such as DIBAL-H and $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]_3$, the process stops after reaction with one equivalent of H^- and the aldehyde is formed as product. With a stronger reducing agent like LiAlH_4 , two equivalents of H^- are added and a 1° alcohol is formed.

Problem 20.11 Draw a stepwise mechanism for the following reaction.



Problem 20.12 Draw the structure of both an acid chloride and an ester that can be used to prepare each compound by reduction.



Selective reductions are routinely used in the synthesis of highly complex natural products such as **ciguatoxin CTX3C**, a potent neurotoxin found in more than 400 species of warm-water fish. Interest in providing a practical supply of ciguatoxin CTX3C for biological studies led to its laboratory synthesis in 2001. One reaction in the synthesis involved the reduction of an ester to an aldehyde using DIBAL-H, as shown in Figure 20.4.

20.7B Reduction of Carboxylic Acids and Amides


Carboxylic acids are reduced to 1° alcohols with LiAlH_4 . LiAlH_4 is too strong a reducing agent to stop the reaction at the aldehyde stage, but milder reagents are not strong enough to initiate the reaction in the first place, so this is the only useful reduction reaction of carboxylic acids.

Figure 20.4

The DIBAL-H reduction of an ester to an aldehyde in the synthesis of the marine neurotoxin ciguatoxin CTX3C




Thousands of people contract ciguatera seafood poisoning each year from ingesting tropical reef fish containing ciguatoxin. Even very low concentrations of ciguatoxin CTX3C cause gastrointestinal and neurological problems, leading to paralysis and sometimes death.




- One step in a lengthy synthesis of ciguatoxin CTX3C involved selective reduction of an ester to an aldehyde using DIBAL-H.

Reduction of a carboxylic acid




Example



Two C—O bonds are replaced by C—H bonds.

Unlike the LiAlH_4 reduction of all other carboxylic acid derivatives, which affords 1° alcohols, the LiAlH_4 reduction of amides forms amines.


Reduction of an amide



Both C—O bonds are replaced by C—H bonds.

Both C–O bonds are reduced to C–H bonds by LiAlH_4 , and any H atom or R group bonded to the amide nitrogen atom remains bonded to it in the product. Because NH_2^- (or NHR^- or NR_2^-) is a poorer leaving group than Cl^- or OR^- , NH_2^- is never lost during reduction, and therefore it forms an amine in the final product.

Examples




Imines and related compounds are discussed in Chapter 21.

The mechanism, illustrated in Mechanism 20.5 with RCONH_2 as starting material, is somewhat different than the previous reductions of carboxylic acid derivatives. Amide reduction proceeds with formation of an intermediate *imine*, a compound containing a C–N double bond, which is then further reduced to an amine.


Mechanism 20.5 Reduction of an Amide to an Amine with LiAlH_4

Part [1] Reduction of an amide to an imine




- In Part [1], the amide is converted to an **imine** by proton transfer, nucleophilic attack of H^- , and loss of $(\text{AlH}_3\text{O})^{2-}$.
- The polarized $\text{C}=\text{N}$ of an imine makes it susceptible to nucleophilic attack.

Part [2] Reduction of an imine to an amine




- **Nucleophilic attack of H^-** (from LiAlH_4) in Step [5], followed by protonation forms the **amine**. The overall result of Steps [5] and [6] is addition of H_2 to the intermediate imine.

Problem 20.13 Draw the products formed from LiAlH_4 reduction of each compound.



Problem 20.14 What amide(s) will form each of the following amines on treatment with LiAlH_4 ?




20.7C A Summary of the Reagents for Reduction

The many available metal hydride reagents reduce a wide variety of functional groups. Keep in mind that LiAlH_4 is such a strong reducing agent that it *nonselectively* reduces most polar functional groups. All other metal hydride reagents are milder, and each has its particular reactions that best utilize its reduced reactivity. The reagents and their uses are summarized in Table 20.1.

Table 20.1 A Summary of Metal Hydride Reducing Agents

	Reagent	Starting material	→	Product
strong reagent	LiAlH ₄	RCHO	→	RCH ₂ OH
		R ₂ CO	→	R ₂ CHOH
		RCOOH	→	RCH ₂ OH
		RCOOR'	→	RCH ₂ OH
		RCOCl	→	RCH ₂ OH
		RCONH ₂	→	RCH ₂ NH ₂
milder reagents	NaBH ₄	RCHO	→	RCH ₂ OH
		R ₂ CO	→	R ₂ CHOH
	LiAlH[OC(CH ₃) ₃] ₃	RCOCl	→	RCHO
		DIBAL-H	RCOOR'	→

Problem 20.15 What product is formed when each compound is treated with either LiAlH₄ (followed by H₂O), or NaBH₄ in CH₃OH?



Problem 20.16 What product is formed when (CH₃)₂C=CHCH₂COCH₂CH₂CO₂CH₂CH₃ is treated with each reagent: (a) H₂ (1 equiv), Pd-C; (b) H₂ (2 equiv), Pd-C; (c) LiAlH₄, followed by H₂O; (d) NaBH₄, CH₃OH?


20.8 Oxidation of Aldehydes



Aldehydes give a positive Tollens test; that is, they react with Ag⁺ to form RCOOH and Ag. When the reaction is carried out in a glass flask, a silver mirror is formed on its walls. Other functional groups give a negative Tollens test, because no silver mirror forms.


The most common oxidation reaction of carbonyl compounds is the oxidation of **aldehydes to carboxylic acids**. A variety of oxidizing agents can be used, including CrO₃, Na₂Cr₂O₇, K₂Cr₂O₇, and KMnO₄. Cr⁶⁺ reagents are also used to oxidize 1° and 2° alcohols, as discussed in Section 12.12. Because ketones have no H on the carbonyl carbon, they do not undergo this oxidation reaction.

Aldehydes are oxidized selectively in the presence of other functional groups using **silver(I) oxide in aqueous ammonium hydroxide (Ag₂O in NH₄OH)**. This is called **Tollens reagent**. Oxidation with Tollens reagent provides a distinct color change, because the Ag⁺ reagent is reduced to silver metal (Ag), which precipitates out of solution.



Problem 20.17 What product is formed when each compound is treated with either Ag₂O, NH₄OH or Na₂Cr₂O₇, H₂SO₄, H₂O: (a) C₆H₅CH₂OH; (b) CH₃CH(OH)CH₂CH₂CH₂CHO?

Problem 20.18 Review the oxidation reactions using Cr^{6+} reagents in Section 12.12. Then, draw the product formed when compound **B** is treated with each reagent.




- a. NaBH_4 , CH_3OH d. Ag_2O , NH_4OH
 b. $[1] \text{LiAlH}_4$; $[2] \text{H}_2\text{O}$ e. CrO_3 , H_2SO_4 , H_2O
 c. PCC


20.9 Organometallic Reagents

We will now discuss the reactions of carbonyl compounds with organometallic reagents, another class of nucleophiles.

- **Organometallic reagents contain a carbon atom bonded to a metal.**



Lithium, magnesium, and copper are the most commonly used metals in organometallic reagents, but others (such as Sn, Si, Tl, Al, Ti, and Hg) are known. General structures of the three common organometallic reagents are shown. R can be alkyl, aryl, allyl, benzyl, sp^2 hybridized, and with M = Li or Mg, sp hybridized. Because metals are *more electropositive* (less electronegative) than carbon, they donate electron density towards carbon, so that **carbon bears a partial negative charge**.




- **The more polar the carbon–metal bond, the more reactive the organometallic reagent.**


Because both Li and Mg are very electropositive metals, **organolithium (RLi)** and **organomagnesium reagents (RMgX)** contain very polar carbon–metal bonds and are therefore very reactive reagents. Organomagnesium reagents are called **Grignard reagents**, after Victor Grignard, who received the Nobel Prize in Chemistry in 1912 for his work with them.

Organocopper reagents (R_2CuLi), also called **organocuprates**, have a less polar carbon–metal bond and are therefore less reactive. Although organocuprates contain two alkyl groups bonded to copper, only one R group is utilized in a reaction.

Regardless of the metal, organometallic reagents are useful synthetically because they react as if they were free carbanions; that is, carbon bears a partial *negative* charge, so the **reagents react as bases and nucleophiles**.



Electronegativity values for carbon and the common metals in R–M reagents are C (2.5), Li (1.0), Mg (1.3), and Cu (1.8).




Problem 20.20 1-Octyne ($\text{HC}\equiv\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$) reacts rapidly with NaH , forming a gas that bubbles out of the reaction mixture, as one product. 1-Octyne also reacts rapidly with CH_3MgBr , and a different gas is produced. Write balanced equations for both reactions and identify the gases formed.


20.9C Reaction as a Base

- Organometallic reagents are strong bases that readily abstract a proton from water to form hydrocarbons.


The electron pair in the carbon–metal bond is used to form a new bond to the proton. Equilibrium favors the products of this acid–base reaction because H_2O is a much stronger acid than the alkane product.







Similar reactions occur for the same reason with the O–H proton in alcohols and carboxylic acids, and the N–H protons of amines.



Because organolithium and Grignard reagents are themselves prepared from alkyl halides, a two-step method converts an alkyl halide into an alkane (or another hydrocarbon).




Problem 20.21 Draw the product formed when each organometallic reagent is treated with H_2O .



20.9D Reaction as a Nucleophile


Organometallic reagents are also strong nucleophiles that react with electrophilic carbon atoms to form new carbon–carbon bonds. These reactions are very valuable in forming the carbon skeletons of complex organic molecules. The following reactions of organometallic reagents are examined in Sections 20.10, 20.13, and 20.14:

[1] Reaction of R–M with aldehydes and ketones to afford alcohols (Section 20.10)




Aldehydes and ketones are converted to 1°, 2°, or 3° alcohols with R''Li or R''MgX.

[2] Reaction of R–M with carboxylic acid derivatives (Section 20.13)



Acid chlorides and esters can be converted to ketones or 3° alcohols with organometallic reagents. The identity of the product depends on the identity of R''–M and the leaving group Z.


[3] Reaction of R–M with other electrophilic functional groups (Section 20.14)



Organometallic reagents also react with CO₂ to form carboxylic acids and with epoxides to form alcohols.

20.10 Reaction of Organometallic Reagents with Aldehydes and Ketones

Treatment of an aldehyde or ketone with either an organolithium or Grignard reagent followed by water forms an alcohol with a new carbon–carbon bond. This reaction is an **addition reaction** because the elements of R'' and H are added across the π bond.




20.10A General Features

This reaction follows the general mechanism for nucleophilic addition (Section 20.2A)—that is, **nucleophilic attack** by a carbanion followed by **protonation**. Mechanism 20.6 is shown using R''MgX, but the same steps occur with organolithium reagents and acetylide anions.




Mechanism 20.6 Nucleophilic Addition of $R''MgX$ to $RCHO$ and $R_2C=O$



- In Step [1], the **nucleophile (R''^-) attacks the carbonyl carbon** and the π bond cleaves, forming an alkoxide. **This step forms a new carbon-carbon bond.**
- In Step [2], **protonation of the alkoxide by H_2O** forms the alcohol addition product. This acid-base reaction forms a new O-H bond.
- The overall result is addition of (R''^-) (from $R''MgX$) and H^+ (from H_2O) to the carbonyl group.

This reaction is used to prepare **1°, 2°, and 3° alcohols**, depending on the number of alkyl groups bonded to the carbonyl carbon of the aldehyde or ketone.




[1] Addition of $R''MgX$ to formaldehyde ($CH_2=O$) forms a 1° alcohol.

[2] Addition of $R''MgX$ to all other aldehydes forms a 2° alcohol.


[3] Addition of $R''MgX$ to ketones forms a 3° alcohol.

Each reaction results in addition of one new alkyl group to the carbonyl carbon, and forms one new carbon-carbon bond. The reaction is general for all organolithium and Grignard reagents, and works for acetylide anions as well, as illustrated in Equations [1]–[3].




Because organometallic reagents are strong bases that rapidly react with H₂O (Section 20.9C), the addition of the new alkyl group must be carried out under anhydrous conditions to prevent traces of water from reacting with the reagent, thus reducing the yield of the desired alcohol. Water is added after the addition to protonate the alkoxide.

Problem 20.22 Draw the product formed when each carbonyl compound is treated with C₆H₅MgBr, followed by protonation with H₂O.




Problem 20.23 Draw the product of each reaction.



20.10B Stereochemistry


Like reduction, addition of organometallic reagents converts an *sp*² hybridized carbonyl carbon to a tetrahedral *sp*³ hybridized carbon. Addition of R–M always occurs from both sides of the trigonal planar carbonyl group. **When a new stereogenic center is formed from an achiral starting material, an equal mixture of enantiomers results**, as shown in Sample Problem 20.1.

Sample Problem 20.1 Draw all stereoisomers formed in the following reaction.



Solution

The Grignard reagent adds from both sides of the trigonal planar carbonyl group, forming two alkoxides, each containing a new stereogenic center. Protonation with water yields **an equal amount of two enantiomers—a racemic mixture**.



Problem 20.24 Draw the products (including stereochemistry) of the following reactions.





Figure. 20.5
The synthesis of ethynylestradiol



20.10C Applications in Synthesis

Many syntheses of useful compounds utilize the nucleophilic addition of a Grignard or organolithium reagent to form carbon–carbon bonds. For example, a key step in the synthesis of ethynylestradiol (Section 11.4), an oral contraceptive component, is the addition of lithium acetylide to a ketone, as shown in Figure 20.5.

The synthesis of C_{18} juvenile hormone, the molecule that opened Chapter 20, is another example. The last steps of the synthesis are outlined in Figure 20.6.

Although juvenile hormone itself is too unstable in light and too expensive to synthesize for use in controlling insect populations, related compounds, called **juvenile hormone mimics**, have been used effectively. The best known example is called **methoprene**, sold under such trade names as Altocid, Precor, and Diacon. Methoprene is used in cattle salt blocks to control hornflies, in stored tobacco to control pests, and on dogs and cats to control fleas.





Figure 20.6
 C_{18} juvenile hormone




- Addition of CH_3MgCl to ketone **A** gives an alkoxide, **B**, which is protonated with H_2O to form 3° alcohol **C**. Although the ester group ($-COOCH_3$) can also react with the Grignard reagent (Section 20.13), it is less reactive than the ketone carbonyl. Thus, with control of reaction conditions, nucleophilic addition occurs selectively at the ketone.
- Treatment of halohydrin **C** with K_2CO_3 forms the C_{18} juvenile hormone in one step. Conversion of a halohydrin to an epoxide was discussed in Section 9.6.

20.11 Retrosynthetic Analysis of Grignard Products

To use the Grignard addition in synthesis, you must be able to determine what carbonyl and Grignard components are needed to prepare a given compound—that is, **you must work backwards, in the retrosynthetic direction**. This involves a two-step process:


Step [1] Find the carbon bonded to the OH group in the product.

Step [2] Break the molecule into two components: One alkyl group bonded to the carbon with the OH group comes from the organometallic reagent. The rest of the molecule comes from the carbonyl component.




For example, to synthesize 3-pentanol [(CH₃CH₂)₂CHOH] by a Grignard reaction, locate the carbon bonded to the OH group, and then break the molecule into two components at this carbon. Thus, retrosynthetic analysis shows that one of the ethyl groups on this carbon comes from a Grignard reagent (CH₃CH₂MgX), and the rest of the molecule comes from the carbonyl component, a three-carbon aldehyde.

Retrosynthetic analysis for preparing 3-pentanol



Then, writing the reaction in the synthetic direction—that is, from starting material to product—shows whether the analysis is correct. In this example, a three-carbon aldehyde reacts with CH₃CH₂MgBr to form an alkoxide, which can then be protonated by H₂O to form 3-pentanol, the desired alcohol.


In the synthetic direction:



There is often more than one way to synthesize a 2° alcohol by Grignard addition, as shown in Sample Problem 20.2.

Sample Problem 20.2

Show two different methods to synthesize 2-butanol using a Grignard reaction.




Linalool and lavandulol (Problem 20.26) are two of the over 300 compounds that determine the characteristic fragrance of lavender.

Solution

Because 2-butanol has two different alkyl groups bonded to the carbon bearing the OH group, there are two different ways to form a new carbon-carbon bond by Grignard addition.


Possibility [1] Use CH_3MgX and a three-carbon aldehyde.

Form this new C-C bond.




Possibility [2] Use $\text{CH}_3\text{CH}_2\text{MgX}$ and a two-carbon aldehyde.

Form this new C-C bond.




Both methods give the desired product, 2-butanol, as can be seen by writing the reactions from starting material to product.




Problem 20.25

What Grignard reagent and carbonyl compound are needed to prepare each alcohol? As shown in part (d), 3° alcohols with three different R groups on the carbon bonded to the OH group can be prepared by three different Grignard reactions.




Problem 20.26

Linalool and lavandulol are two of the major components of lavender oil. (a) What organolithium reagent and carbonyl compound can be used to make each alcohol? (b) How might lavandulol be formed by reduction of a carbonyl compound? (c) Why can't linalool be prepared by a similar pathway?



Problem 20.27 What Grignard reagent and carbonyl compound can be used to prepare the antidepressant venlafaxine (trade name Effexor)?



20.12 Protecting Groups


Although the addition of organometallic reagents to carbonyls is a very versatile reaction, it cannot be used with molecules that contain both a carbonyl group and N–H or O–H bonds.

Rapid acid–base reactions occur between organometallic reagents and all of the following functional groups: ROH, RCOOH, RNH₂, R₂NH, RCONH₂, RCONHR, and RSH.


- Carbonyl compounds that also contain N–H or O–H bonds undergo an acid–base reaction with organometallic reagents, not nucleophilic addition.

Suppose, for example, that you wanted to add methylmagnesium chloride (CH₃MgCl) to the carbonyl group of 5-hydroxy-2-pentanone to form a diol. Nucleophilic addition will *not* occur with this substrate. Instead, **because Grignard reagents are strong bases and proton transfer reactions are fast, CH₃MgCl removes the O–H proton before nucleophilic addition takes place.** The stronger acid and base react to form the weaker conjugate acid and conjugate base, as we learned in Section 20.9C.

CH₃MgCl does *not* add to the carbonyl group.



CH₃MgCl acts like a base, *not* a nucleophile.



Solving this problem requires a three-step strategy:


Step [1] Convert the OH group into another functional group that does not interfere with the desired reaction. This new blocking group is called a protecting group, and the reaction that creates it is called **protection**.

Step [2] Carry out the desired reaction.

Step [3] Remove the protecting group. This reaction is called **deprotection**.


Application of the general strategy to the Grignard addition of CH₃MgCl to 5-hydroxy-2-pentanone is illustrated in Figure 20.7.

Figure 20.7
General strategy for using a protecting group




- In Step [1], the OH proton in 5-hydroxy-2-pentanone is replaced with a protecting group, written as **PG**. Because the product of Step [1] no longer has an OH proton, it can now undergo nucleophilic addition.
- In Step [2], CH_3MgCl adds to the carbonyl group to yield a 3° alcohol after protonation with water.
- Removal of the protecting group in Step [3] forms the desired product, 4-methyl-1,4-pentanediol.


A common OH protecting group is a **silyl ether**. A silyl ether has a new O–Si bond in place of the O–H bond of the alcohol. The most widely used silyl ether protecting group is the ***tert*-butyldimethylsilyl ether**.




tert-Butyldimethylsilyl ethers are prepared from alcohols by reaction with *tert*-butyldimethylsilyl chloride and an amine base, usually imidazole.



The silyl ether is typically removed with a fluoride salt, usually **tetrabutylammonium fluoride** ($(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_4\text{N}^+ \text{F}^-$).




The use of a *tert*-butyldimethylsilyl ether as a protecting group makes possible the synthesis of 4-methyl-1,4-pentanediol by a three-step sequence.



- **Step [1] Protect the OH group** as a *tert*-butyldimethylsilyl ether by reaction with *tert*-butyldimethylsilyl chloride and imidazole.
- **Step [2] Carry out nucleophilic addition** by using CH_3MgCl , followed by protonation.
- **Step [3] Remove the protecting group** with tetrabutylammonium fluoride to form the desired addition product.


Protecting groups block interfering functional groups, and in this way, a wider variety of reactions can take place with a particular substrate. For more on protecting groups, see the discussion of acetals in Section 21.15.

Problem 20.28 Using protecting groups, show how estrone can be converted to ethynylestradiol, a widely used oral contraceptive.



20.13 Reaction of Organometallic Reagents with Carboxylic Acid Derivatives



Organometallic reagents react with carboxylic acid derivatives (RCOZ) to form two different products, depending on the identity of both the leaving group Z and the reagent $\text{R}-\text{M}$. The most useful reactions are carried out with esters and acid chlorides, forming either **ketones** or **3° alcohols**.




- Keep in mind that RLi and RMgX are very reactive reagents, whereas R_2CuLi is much less reactive. This reactivity difference makes selective reactions possible.

20.13A Reaction of RLi and RMgX with Esters and Acid Chlorides

Both esters and acid chlorides form 3° alcohols when treated with two equivalents of either Grignard or organolithium reagents. Two new carbon–carbon bonds are formed in the product.



Problem 20.29 Draw the product formed when each compound is treated with two equivalents of $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{MgBr}$ followed by H_2O .




The mechanism for this addition reaction resembles the mechanism for the metal hydride reduction of acid chlorides and esters discussed in Section 20.7A. The mechanism is conceptually divided into two parts: **nucleophilic substitution** to form a ketone, followed by **nucleophilic addition** to form a 3° alcohol, as shown in Mechanism 20.7.




Mechanism 20.7 Reaction of $\text{R}''\text{MgX}$ or $\text{R}''\text{Li}$ with RCOCl and RCOOR'

Part [1] Nucleophilic substitution forms a ketone.



- **Nucleophilic attack of $(\text{R}'')^-$** (from $\text{R}''\text{MgX}$) in Step [1] forms a tetrahedral intermediate with a leaving group Z.
- In Step [2], the π bond is re-formed and **Z^- comes off**. The overall result of the addition of $(\text{R}'')^-$ and elimination of Z^- is the substitution of R'' for Z.
- Because the product of Part [1] is a ketone, it can react with a second equivalent of $\text{R}''\text{MgX}$ to form an alcohol by nucleophilic addition in Part [2].

Part [2] Nucleophilic addition forms a 3° alcohol.




- **Nucleophilic attack of $(\text{R}'')^-$** (from $\text{R}''\text{MgX}$) in Step [3] forms an alkoxide.
- **Protonation of the alkoxide** by H_2O in Step [4] forms a 3° alcohol.

Organolithium and Grignard reagents always afford 3° alcohols when they react with esters and acid chlorides. As soon as the ketone forms by addition of one equivalent of reagent to RCOZ (Part [1] of the mechanism), it reacts with a second equivalent of reagent to form the 3° alcohol.


This reaction is more limited than the Grignard addition to aldehydes and ketones, because only 3° alcohols having two identical alkyl groups can be prepared. Nonetheless, it is still a valuable reaction because it forms two new carbon–carbon bonds.

Sample Problem 20.3 What ester and Grignard reagent are needed to prepare the following alcohol?




Solution

A 3° alcohol formed from an ester and Grignard reagent must have **two identical R groups**, and these R groups come from RMgX. The remainder of the molecule comes from the ester.






All other C's come from the ester.
R' = any alkyl group

Checking in the synthetic direction:




Problem 20.30 What ester and Grignard reagent are needed to prepare each alcohol?




20.13B Reaction of R_2CuLi with Acid Chlorides

To form a ketone from a carboxylic acid derivative, a less reactive organometallic reagent—namely an **organocuprate**—is needed. **Acid chlorides, which have the best leaving group (Cl^-) of the carboxylic acid derivatives, react with $\text{R}'_2\text{CuLi}$, to give a ketone as product.** Esters, which contain a poorer leaving group (^-OR), do not react with $\text{R}'_2\text{CuLi}$.

General reaction




Example




This reaction results in nucleophilic substitution of an alkyl group R' for the leaving group Cl, forming one new carbon–carbon bond.

Problem 20.31 What organocuprate reagent is needed to convert $\text{CH}_3\text{CH}_2\text{COCl}$ to each ketone?




Problem 20.32 What reagent is needed to convert $(\text{CH}_3)_2\text{CHCH}_2\text{COCl}$ into each compound?



A ketone with two different R groups bonded to the carbonyl carbon can be made by two different methods, as illustrated in Sample Problem 20.4.

Sample Problem 20.4 Show two different ways to prepare 2-pentanone from an acid chloride and an organocuprate reagent.




Solution

In each case, one alkyl group comes from the organocuprate and one comes from the acid chloride.


Possibility [1] Use $(\text{CH}_3)_2\text{CuLi}$ and a four-carbon acid chloride.

Form this new C–C bond.



Possibility [2] Use $(\text{CH}_3\text{CH}_2\text{CH}_2)_2\text{CuLi}$ and a two-carbon acid chloride.

Form this new C–C bond.




Problem 20.33 Draw two different ways to prepare each ketone from an acid chloride and an organocuprate reagent: (a) $\text{C}_6\text{H}_5\text{COCH}_3$; (b) $(\text{CH}_3)_2\text{CHCOC}(\text{CH}_3)_3$.

20.14 Reaction of Organometallic Reagents with Other Compounds


Because organometallic reagents are strong nucleophiles, they react with many other electrophiles in addition to carbonyl groups. Because these reactions always lead to the formation of new carbon–carbon bonds, they are also valuable in organic synthesis. In Section 20.14, we examine the reactions of organometallic reagents with **carbon dioxide** and **epoxides**.

20.14A Reaction of Grignard Reagents with Carbon Dioxide

Grignard reagents react with CO_2 to give carboxylic acids after protonation with aqueous acid. This reaction, called **carboxylation**, forms a carboxylic acid with one more carbon atom than the Grignard reagent from which it is prepared.




Because Grignard reagents are made from alkyl halides, an alkyl halide can be converted to a carboxylic acid having one more carbon atom by a two-step reaction sequence: **formation of a Grignard reagent**, followed by **reaction with CO₂**.



The mechanism resembles earlier reactions of nucleophilic Grignard reagents with carbonyl groups, as shown in Mechanism 20.8.




Mechanism 20.8 Carboxylation—Reaction of RMgX with CO₂



- In Step [1], the nucleophilic Grignard reagent attacks the electrophilic carbon atom of CO₂, cleaving a π bond and forming a new carbon–carbon bond.
- The carboxylate anion is protonated with aqueous acid in Step [2] to form the carboxylic acid.


Problem 20.34 What carboxylic acid is formed from each alkyl halide on treatment with [1] Mg; [2] CO₂; [3] H₃O⁺?




20.14B Reaction of Organometallic Reagents with Epoxides

Like other strong nucleophiles, **organometallic reagents**—RLi, RMgX, and R₂CuLi—**open epoxide rings to form alcohols**.

General reaction




Example




The opening of epoxide rings with negatively charged nucleophiles was discussed in Section 9.15A.

The reaction follows the same two-step process as the opening of epoxide rings with other negatively charged nucleophiles—that is, **nucleophilic attack from the back side of the epoxide ring**, followed by **protonation of the resulting alkoxide**. In unsymmetrical epoxides, nucleophilic attack occurs at the less substituted carbon atom.




Problem 20.35 What epoxide is needed to convert $\text{CH}_3\text{CH}_2\text{MgBr}$ to each of the following alcohols, after quenching with water?



20.15 α,β -Unsaturated Carbonyl Compounds


α,β -Unsaturated carbonyl compounds are conjugated molecules containing a carbonyl group and a carbon-carbon double bond, separated by a single σ bond.




α,β -unsaturated carbonyl compound

Both functional groups of α,β -unsaturated carbonyl compounds have π bonds, but individually, they react with very different kinds of reagents. Carbon-carbon double bonds react with electrophiles (Chapter 10) and carbonyl groups react with nucleophiles (Section 20.2). What happens, then, when these two functional groups having opposite reactivity are in close proximity?


Because the two π bonds are conjugated, the electron density in an α,β -unsaturated carbonyl compound is delocalized over four atoms. Three resonance structures show that the carbonyl carbon and the β carbon bear a partial positive charge. This means that **α,β -unsaturated carbonyl compounds can react with nucleophiles at two different sites.**



- Addition of a nucleophile to the carbonyl carbon, called 1,2-addition, adds the elements of H and Nu across the $\text{C}=\text{O}$, forming an allylic alcohol.



- Addition of a nucleophile to the β carbon, called 1,4-addition or conjugate addition, forms a carbonyl compound.

**1,4-Addition
(conjugate addition)**


The nucleophile attacks at the β carbon.

a carbonyl compound with a new substituent on the β carbon


Both 1,2- and 1,4-addition result in nucleophilic addition of the elements of H and Nu.

20.15A The Mechanisms for 1,2-Addition and 1,4-Addition

The steps for the mechanism of 1,2-addition are exactly the same as those for the nucleophilic addition to an aldehyde or ketone—that is, **nucleophilic attack**, followed by **protonation** (Section 20.2A), as shown in Mechanism 20.9.



Mechanism 20.9 1,2-Addition to an α,β -Unsaturated Carbonyl Compound




The mechanism for 1,4-addition also begins with nucleophilic attack, and then protonation and tautomerization add the elements of H and Nu to the α and β carbons of the carbonyl compound, as shown in Mechanism 20.10.




Mechanism 20.10 1,4-Addition to an α,β -Unsaturated Carbonyl Compound

Part [1] Nucleophilic attack at the β carbon



- In Part [1], nucleophilic attack at the β carbon forms a resonance-stabilized anion called an **enolate**. Either resonance structure can be used to continue the mechanism in Part [2].

Part [2] Protonation and tautomerization




- Protonation on the carbon end of the enolate forms the 1,4-addition product directly.
- Protonation of the oxygen end of the enolate forms an **enol**. Recall from Section 11.9 that enols are unstable and tautomerize (by a two-step process) to carbonyl compounds. Tautomerization forms the same 1,4-addition product that results from protonation on carbon.

20.15B Reaction of α,β -Unsaturated Carbonyl Compounds with Organometallic Reagents



The identity of the metal in an organometallic reagent determines whether it reacts with an α,β -unsaturated aldehyde or ketone by 1,2-addition or 1,4-addition.

Why is conjugate addition also called 1,4-addition? If the atoms of the enol are numbered beginning with the O atom, then the elements of H and Nu are bonded to atoms "1" and "4," respectively.



The enol has H and Nu added to atoms 1 and 4.



- Organolithium and Grignard reagents form 1,2-addition products.





- Organocuprate reagents form 1,4-addition products.



Sample Problem 20.5


Draw the products of each reaction.




Solution

The characteristic reaction of α,β -unsaturated carbonyl compounds is nucleophilic addition. The reagent determines the mode of addition (1,2- or 1,4-).


- a. **Grignard reagents undergo 1,2-addition.** CH_3MgBr adds a new CH_3 group at the carbonyl carbon.




- b. **Organocuprate reagents undergo 1,4-addition.** The cuprate reagent adds a new vinyl group ($\text{CH}_2=\text{CH}$) at the β carbon.



Problem 20.36 Draw the product when each compound is treated with either $(\text{CH}_3)_2\text{CuLi}$, followed by H_2O , or $\text{HC}\equiv\text{CLi}$, followed by H_2O .




Problem 20.37 How can $(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$ (mesityl oxide) be converted to each compound?



20.16 Summary—The Reactions of Organometallic Reagents


We have now seen many different reactions of organometallic reagents with a variety of functional groups, and you may have some difficulty keeping them all straight. Rather than memorizing them all, keep in mind the following three concepts:

[1] Organometallic reagents ($\text{R}-\text{M}$) attack electrophilic carbon atoms, especially the carbonyl carbon.



[2] After an organometallic reagent adds to a carbonyl group, the fate of the intermediate depends on the presence or absence of a leaving group.


- Without a leaving group, the characteristic reaction is *nucleophilic addition*.
- With a leaving group, it is *nucleophilic substitution*.



[3] The polarity of the $\text{R}-\text{M}$ bond determines the reactivity of the reagents.

- RLi and RMgX are very reactive reagents.
- R_2CuLi is much less reactive.

Figure 20.8
Conversion of 2-hexanol into
other compounds




20.17 Synthesis

The reactions learned in Chapter 20 have proven extremely useful in organic synthesis. Oxidation and reduction reactions interconvert two functional groups that differ in oxidation state. Organometallic reagents form new carbon–carbon bonds.


Synthesis is perhaps the most difficult aspect of organic chemistry. It requires you to remember both the new reactions you've just learned, and the ones you've encountered in previous chapters. In a successful synthesis, you must also put these reactions in a logical order. Don't be discouraged. Learn the basic reactions and then practice them over and over again with synthesis problems.

In Sample Problems 20.6–20.8 that follow, keep in mind that the products formed by the reactions of Chapter 20 can themselves be transformed into many other functional groups. For example, 2-hexanol, the product of Grignard addition of butylmagnesium chloride to acetaldehyde, can be transformed into a variety of other compounds, as shown in Figure 20.8.




Before proceeding with Sample Problems 20.6–20.8, you should review the stepwise strategy for designing a synthesis found in Section 11.12.

Sample Problem 20.6 Synthesize 1-methylcyclohexene from cyclohexanone and any organic alcohol.



Retrosynthetic Analysis




Thinking backwards:

- [1] Form the double bond by dehydration of an alcohol.
- [2] Make the 3° alcohol by Grignard addition of CH_3MgX .
- [3] Prepare the Grignard reagent stepwise from an alcohol.


Synthesis

Four steps are needed:




- Conversion of CH_3OH to the Grignard reagent CH_3MgBr requires two steps: formation of an alkyl halide (Step [1]), followed by reaction with Mg (Step [2]).
- Addition of CH_3MgBr to cyclohexanone followed by protonation forms an alcohol in Step [3].
- Acid-catalyzed elimination of water in Step [4] forms a mixture of alkenes, with the desired trisubstituted alkene as the major product.

Sample Problem 20.7 Synthesize 2,4-dimethyl-3-hexanone from four-carbon alcohols.



Retrosynthetic Analysis




Thinking backwards:


- [1] Form the ketone by oxidation of a 2° alcohol.
- [2] Make the 2° alcohol by Grignard addition to an aldehyde. Both of these compounds have 4 C's, and each must be synthesized from an alcohol.

Synthesis

First, make both components needed for the Grignard reaction.




Then, complete the synthesis with Grignard addition, followed by oxidation of the alcohol to the ketone.




Sample Problem 20.8

Synthesize isopropylcyclopentane from alcohols having ≤ 5 C's.



Retrosynthetic Analysis




Thinking backwards:


- [1] Form the alkane by hydrogenation of an alkene.
- [2] Introduce the double bond by dehydration of an alcohol.
- [3] Form the 3° alcohol by Grignard addition to a ketone. Both components of the Grignard reaction must then be synthesized.

Synthesis

First make both components needed for the Grignard reaction.




Complete the synthesis with Grignard addition, dehydration, and hydrogenation.



Problem 20.38

Synthesize each compound from cyclohexanol, ethanol, and any other needed reagents.





KEY CONCEPTS


Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

Reduction Reactions


[1] Reduction of aldehydes and ketones to 1° and 2° alcohols (20.4)

[2] Reduction of α,β -unsaturated aldehydes and ketones (20.4C)


[3] Enantioselective ketone reduction (20.6)



[4] Reduction of acid chlorides (20.7A)




[5] Reduction of esters (20.7A)




• LiAlH₄, a strong reducing agent, reduces an ester to a 1° alcohol.

• With DIBAL-H, a milder reducing agent, reduction stops at the aldehyde stage.

[6] Reduction of carboxylic acids to 1° alcohols (20.7B)




[7] Reduction of amides to amines (20.7B)



Oxidation Reactions



Oxidation of aldehydes to carboxylic acids (20.8)



• All Cr⁶⁺ reagents except PCC oxidize RCHO to RCOOH.

• Tollens reagent (Ag₂O + NH₄OH) oxidizes RCHO only. Primary (1°) and 2° alcohols do not react with Tollens reagent.

Preparation of Organometallic Reagents (20.9)




Reactions with Organometallic Reagents

[1] Reaction as a base (20.9C)




- RM = RLi, RMgX, R₂CuLi
- This acid–base reaction occurs with H₂O, ROH, RNH₂, R₂NH, RSH, RCOOH, RCONH₂, and RCONHR.


[2] Reaction with aldehydes and ketones to form 1°, 2°, and 3° alcohols (20.10)



[3] Reaction with esters to form 3° alcohols (20.13A)




[4] Reaction with acid chlorides (20.13B)




- More reactive organometallic reagents—R''Li and R''MgX—add two equivalents of R'' to an acid chloride to form a 3° alcohol with two identical R'' groups.

- Less reactive organometallic reagents—R'₂CuLi—add only one equivalent of R' to an acid chloride to form a ketone.


[5] Reaction with carbon dioxide—Carboxylation (20.14A)



[6] Reaction with epoxides (20.14B)



[7] Reaction with α,β -unsaturated aldehydes and ketones (20.15B)




• More reactive organometallic reagents— $R'Li$ and $R'MgX$ —react with α,β -unsaturated carbonyls by 1,2-addition.


• Less reactive organometallic reagents— R'_2CuLi —react with α,β -unsaturated carbonyls by 1,4-addition.

Protecting Groups (20.12)

[1] Protecting an alcohol as a *tert*-butyldimethylsilyl ether



[2] Deprotecting a *tert*-butyldimethylsilyl ether to re-form an alcohol



PROBLEMS

Reactions and Reagents

20.39 Draw the product formed when pentanal ($CH_3CH_2CH_2CH_2CHO$) is treated with each reagent. With some reagents, no reaction occurs.









- | | |
|---------------------------------------|---|
| a. $NaBH_4$, CH_3OH | g. [1] CH_3MgBr ; [2] H_2O |
| b. [1] $LiAlH_4$; [2] H_2O | h. [1] C_6H_5Li ; [2] H_2O |
| c. H_2 , Pd-C | i. [1] $(CH_3)_2CuLi$; [2] H_2O |
| d. PCC | j. [1] $HC\equiv CNa$; [2] H_2O |
| e. $Na_2Cr_2O_7$, H_2SO_4 , H_2O | k. [1] $CH_3C\equiv Cl$; [2] H_2O |
| f. Ag_2O , NH_4OH | l. The product in (a), then TBDMS-Cl, imidazole |

20.40 Repeat Problem 20.39 using 2-pentanone ($CH_3COCH_2CH_2CH_3$) as the starting material.





20.41 Draw the product formed when 1-bromobutane is treated with each reagent.

- | | |
|---------------------------------------|---|
| a. Li (2 equiv) | d. The answer in (a), then H_2O |
| b. Mg in $(CH_3CH_2)_2O$ solvent | e. The answer in (b), then D_2O |
| c. Li (2 equiv), then CuI (0.5 equiv) | f. The answer in (a), then $CH_3C\equiv CH$ |

20.42 Draw the product formed when $\text{CH}_3\text{CH}_2\text{CH}_2\text{MgBr}$ is treated with each compound.










20.43 Draw the product formed when $(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{CuLi}$ is treated with each compound. In some cases, no reaction occurs.






20.44 The stereochemistry of the products of reduction depends on the reagent used, as you learned in Sections 20.5 and 20.6. With this in mind, how would you convert 3,3-dimethyl-2-butanone [$\text{CH}_3\text{COC}(\text{CH}_3)_2$] to: (a) racemic 3,3-dimethyl-2-butanol [$\text{CH}_3\text{CH}(\text{OH})\text{C}(\text{CH}_3)_2$]; (b) only (2*R*)-3,3-dimethyl-2-butanol; (c) only (2*S*)-3,3-dimethyl-2-butanol?




20.45 Draw the product formed when the α,β -unsaturated ketone **A** is treated with each reagent.



20.46 What reagent is needed to convert $(\text{CH}_3)_2\text{CHCH}_2\text{CH}_2\text{COCl}$ to each compound?







20.47 What reagent is needed to convert $\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_2\text{CH}_3$ to each compound?




20.48 As discussed in Sections 12.12 and 20.8, some oxidizing agents selectively oxidize a particular functional group, whereas others oxidize many different functional groups. Draw the product formed when $\text{HOCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$ is treated with each reagent: (a) CrO_3 , H_2SO_4 , H_2O ; (b) PCC; (c) Ag_2O , NH_4OH ; (d) $\text{Na}_2\text{Cr}_2\text{O}_7$, H_2SO_4 , H_2O .


20.49 Draw the products of each reduction reaction.




20.50 Draw the products of the following reactions with organometallic reagents.




20.51 Draw all stereoisomers formed in each reaction.




20.52 Explain why metal hydride reduction gives an endo alcohol as the major product in one reaction and an exo alcohol as the major product in the other reaction.




20.53 A student tried to carry out the following reaction sequence, but none of diol **A** was formed. Explain what was wrong with this plan, and design a successful stepwise synthesis of **A**.



20.57 Draw a stepwise mechanism for the following reaction. Account for the formation of all of the products.




20.58 Tertiary alcohols can be formed by the reaction of dimethyl carbonate $[(\text{CH}_3\text{O})_2\text{C}=\text{O}]$ with excess Grignard reagent. Draw a stepwise mechanism for the following transformation.




Synthesis


20.59 What Grignard reagent and aldehyde (or ketone) are needed to prepare each alcohol? Show all possible routes.




20.60 What ester and Grignard reagent are needed to synthesize each alcohol?



20.61 What organolithium reagent and carbonyl compound can be used to prepare each of the following compounds? You may use aldehydes, ketones, or esters as carbonyl starting materials.



20.62 What epoxide and organometallic reagent are needed to synthesize each alcohol?







20.63 Propose at least three methods to convert $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{Br}$ to $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_3$.





20.64 Devise a synthesis of each alcohol from organic alcohols having one or two carbons and any required reagents: (a) $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$; (b) $(\text{CH}_3)_3\text{COH}$; (c) $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$; (d) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$.

20.65 Propose two different methods to synthesize 1-octen-3-ol $[\text{CH}_3(\text{CH}_2)_4\text{CH}(\text{OH})\text{CH}=\text{CH}_2]$ using a Grignard reagent and a carbonyl compound. 1-Octen-3-ol is commonly called matsutake alcohol because it was first isolated from the Japanese matsutake mushroom.

20.66 Synthesize each compound from cyclohexanol using any other organic or inorganic compounds.






(Each cyclohexane ring must come from cyclohexanol.)




(Each cyclohexane ring must come from cyclohexanol.)

20.67 Convert 2-propanol [(CH₃)₂CHOH] into each compound. You may use any other organic or inorganic compounds.

- (CH₃)₂CHCl
- (CH₃)₂C=O
- (CH₃)₂CHCH(OH)CH₃
- (CH₃)₂CHCH₂CH₂OH
- (CH₃)₂CHCOOH
- (CH₃)₂CHCHO
- (CH₃)₂CHD



20.68 Devise a synthesis of mestranol, a synthetic estrogen used in oral contraceptives, from the female sex hormone estradiol. You may use any needed organic compounds or inorganic reagents.








20.69 Devise three different methods to prepare each compound from benzene: (a) C₆H₅CH₂CH₂OH (2-phenylethanol); (b) C₆H₅COCH₃ (acetophenone). You may also use organic compounds that have one or two carbons, and any required inorganic reagents. At least two of the three methods must use a reaction of the organometallic reagents described in this chapter.






20.70 Carry out each of the following syntheses from the given starting material. You may use any other needed organic compounds or inorganic reagents.

- (CH₃)₂CHCH₂CO₂H \implies (CH₃)₂C=CH₂
- (CH₃)₂CHCH₂CH₂CH₂OH \implies (CH₃)₂C=CH₂
- (CH₃)₃CCH₂OH \implies (CH₃)₃CH






20.71 Convert benzene into each compound. You may also use any inorganic reagents and organic alcohols having three carbons or fewer. One step of the synthesis must use a Grignard reagent.



20.72 Design a synthesis of each compound from alcohols having four carbons or fewer as the only organic starting materials. You may use any other inorganic reagents you choose.





20.73 Synthesize each compound from the given starting material. You may use any other required inorganic reagents.



Spectroscopy

20.74 An unknown compound **A** (molecular formula $C_7H_{14}O$) was treated with $NaBH_4$ in CH_3OH to form compound **B** (molecular formula $C_7H_{16}O$). Compound **A** has a strong absorption in its IR spectrum at 1716 cm^{-1} . Compound **B** has a strong absorption in its IR spectrum at $3600\text{--}3200\text{ cm}^{-1}$. The 1H NMR spectra of **A** and **B** are given. What are the structures of **A** and **B**?



20.75 Treatment of compound **C** (molecular formula C_4H_8O) with C_6H_5MgBr , followed by H_2O , affords compound **D** (molecular formula $C_{10}H_{14}O$). Compound **D** has a strong peak in its IR spectrum at $3600\text{--}3200\text{ cm}^{-1}$. The 1H NMR spectral data of **C** and **D** are given. What are the structures of **C** and **D**?

Compound **C** signals at 1.3 (singlet, 6 H) and 2.4 (singlet, 2 H) ppm


Compound **D** signals at 1.2 (singlet, 6 H), 1.6 (singlet, 1 H), 2.7 (singlet, 2 H), and 7.2 (multiplet, 5 H) ppm

20.76 Treatment of compound **E** (molecular formula $C_4H_8O_2$) with excess CH_3CH_2MgBr yields compound **F** (molecular formula $C_6H_{14}O$) after protonation with H_2O . **E** shows a strong absorption in its IR spectrum at 1743 cm^{-1} . **F** shows a strong IR absorption at $3600\text{--}3200\text{ cm}^{-1}$. The 1H NMR spectral data of **E** and **F** are given. What are the structures of **E** and **F**?


Compound **E** signals at 1.2 (triplet, 3 H), 2.0 (singlet, 3 H), and 4.1 (quartet, 2 H) ppm


Compound **F** signals at 0.9 (triplet, 6 H), 1.1 (singlet, 3 H), 1.5 (quartet, 4 H), and 1.55 (singlet, 1 H) ppm

- 20.77** Reaction of butanenitrile ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CN}$) with methylmagnesium bromide (CH_3MgBr), followed by treatment with aqueous acid, forms compound **G**. **G** has a molecular ion in its mass spectrum at $m/z = 86$ and a base peak at $m/z = 43$. **G** exhibits a strong absorption in its IR spectrum at 1721 cm^{-1} and has the $^1\text{H NMR}$ spectrum given below. What is the structure of **G**? We will learn about the details of this reaction in Chapter 22.




- 20.78** Treatment of isobutene [$(\text{CH}_3)_2\text{C}=\text{CH}_2$] with $(\text{CH}_3)_3\text{CCl}$ forms a carbanion that reacts with $\text{CH}_2=\text{O}$ to form **H** after water is added to the reaction mixture. **H** has a molecular ion in its mass spectrum at $m/z = 86$, and shows fragments at 71 and 68. **H** exhibits absorptions in its IR spectrum at $3600\text{--}3200$ and 1651 cm^{-1} , and has the $^1\text{H NMR}$ spectrum given below. What is the structure of **H**?




- 20.79**  oxetane Reaction of oxetane with CH_3MgBr followed by H_2O forms a compound that shows a strong absorption in its IR spectrum at $3600\text{--}3200\text{ cm}^{-1}$ and the following $^1\text{H NMR}$ data: 0.94 (triplet, 3 H), 1.39 (multiplet, 2 H), 1.53 (multiplet, 2 H), 2.24 (singlet, 1 H), and 3.63 (triplet, 2 H) ppm. Draw the structure of the product and write a mechanism that illustrates its formation.

Challenge Problems


- 20.80** Design a synthesis of (*R*)-salmeterol (Figure 20.3) from the following starting materials.




- 20.81** Lithium tri-*sec*-butylborohydride, also known as L-selectride, is a metal hydride reagent that contains three *sec*-butyl groups bonded to boron. When this reagent is used to reduce cyclic ketones, one stereoisomer often predominates as product. Explain why the reduction of 4-*tert*-butylcyclohexanone with L-selectride forms the *cis* alcohol as the major product.



- 20.82** Explain why the β carbon of an α,β -unsaturated carbonyl compound absorbs farther downfield in the ^{13}C NMR spectrum than the α carbon, even though the α carbon is closer to the electron-withdrawing carbonyl group. For example, the β carbon of mesityl oxide absorbs at 150.5 ppm, while the α carbon absorbs at 122.5 ppm.




- 20.83** Identify **X** and **Y**, two of the intermediates in a synthesis of the antidepressant venlafaxine (trade name Effexor), in the following reaction scheme. Write a mechanism for the formation of **X** from **W**.



21

Aldehydes and Ketones—
Nucleophilic Addition

- 21.1 Introduction
- 21.2 Nomenclature
- 21.3 Physical properties
- 21.4 Spectroscopic properties
- 21.5 Interesting aldehydes and ketones
- 21.6 Preparation of aldehydes and ketones
- 21.7 Reactions of aldehydes and ketones—General considerations
- 21.8 Nucleophilic addition of H^- and R^- —A review
- 21.9 Nucleophilic addition of ^-CN
- 21.10 The Wittig reaction
- 21.11 Addition of 1° amines
- 21.12 Addition of 2° amines
- 21.13 Addition of H_2O —Hydration
- 21.14 Addition of alcohols—Acetal formation
- 21.15 Acetals as protecting groups
- 21.16 Cyclic hemiacetals
- 21.17 An introduction to carbohydrates



The natural product **digoxin** has been prescribed since the 1960s for patients with congestive heart failure, a condition that results when fluid builds up in the body because the heart's pumping action is weak. Unlike many commercial medications that are synthesized from simple precursors, digoxin is still obtained by extraction of the leaves of the woolly foxglove plant, which is grown in the Netherlands and shipped to the United States for processing. One thousand kilograms of dried leaves yield one kilogram of digoxin, sold under the trade name of Lanoxin. Digoxin contains three acetal units, which are formed by addition reactions to carbonyl groups. In Chapter 21, we learn about nucleophilic addition, the characteristic reaction of aldehydes and ketones.

In Chapter 21 we continue the study of carbonyl compounds with a detailed look at **aldehydes** and **ketones**. We will first learn about the nomenclature, physical properties, and spectroscopic absorptions that characterize aldehydes and ketones. The remainder of Chapter 21 is devoted to **nucleophilic addition** reactions. Although we have already learned two examples of this reaction in Chapter 20, nucleophilic addition to aldehydes and ketones is a general reaction that occurs with many nucleophiles, forming a wide variety of products.


Every new reaction in Chapter 21 involves nucleophilic addition, so the challenge lies in learning the specific reagents and mechanisms that characterize each reaction.

21.1 Introduction


An aldehyde is often written as **RCHO**. Remember that the **H atom is bonded to the carbon atom**, not the oxygen.

Likewise, a ketone is written as **RCOR**, or if both alkyl groups are the same, **R₂CO**. Each structure must contain a C=O for every atom to have an octet.

As we learned in Chapter 20, **aldehydes and ketones contain a carbonyl group**. An aldehyde contains at least one H atom bonded to the carbonyl carbon, whereas a ketone has two alkyl or aryl groups bonded to it.




Two structural features determine the chemistry and properties of aldehydes and ketones.




- The carbonyl group is sp^2 hybridized and trigonal planar, making it relatively uncrowded.
- The electronegative oxygen atom polarizes the carbonyl group, making the carbonyl carbon electrophilic.

As a result, **aldehydes and ketones react with nucleophiles**. The relative reactivity of the carbonyl group is determined by the number of R groups bonded to it. **As the number of R groups around the carbonyl carbon increases, the reactivity of the carbonyl compound decreases**, resulting in the following order of reactivity:

Increasing the number of alkyl groups on the carbonyl carbon decreases reactivity for both steric and electronic reasons, as discussed in Section 20.2B.



Problem 21.1 Rank the compounds in each group in order of increasing reactivity towards nucleophilic attack.



Problem 21.2 Explain why benzaldehyde is less reactive than cyclohexanecarbaldehyde towards nucleophilic attack.



21.2 Nomenclature

Both IUPAC and common names are used for aldehydes and ketones.

21.2A Naming Aldehydes in the IUPAC System

In IUPAC nomenclature, aldehydes are identified by a suffix added to the parent name of the longest chain. Two different suffixes are used, depending on whether the CHO group is bonded to a chain or a ring.

To name an aldehyde using the IUPAC system:

- [1] If the CHO is bonded to a chain of carbons, find the longest chain containing the CHO group, and change the *-e* ending of the parent alkane to the suffix *-al*. If the CHO group is bonded to a ring, name the ring and add the suffix *-carbaldehyde*.
- [2] Number the chain or ring to put the CHO group at C1, but omit this number from the name. Apply all of the other usual rules of nomenclature.


Sample Problem 21.1

Give the IUPAC name for each compound.




Solution


- a. [1] Find and name the longest chain containing the CHO:




- [2] Number and name substituents:



- b. [1] Find and name the ring bonded to the CHO group:




- [2] Number and name substituents:



Problem 21.3

Give the IUPAC name for each aldehyde.



Problem 21.4

Give the structure corresponding to each IUPAC name.


- a. 2-isobutyl-3-isopropylhexanal
- b. *trans*-3-methylcyclopentanecarbaldehyde
- c. 1-methylcyclopropanecarbaldehyde
- d. 3,6-diethylnonanal

21.2B Common Names for Aldehydes

Like carboxylic acids, many simple aldehydes have common names that are widely used.

- A common name for an aldehyde is formed by taking the common parent name and adding the suffix *-aldehyde*.

The common parent names are similar to those used for carboxylic acids, listed in Table 19.1. The common names **formaldehyde**, **acetaldehyde**, and **benzaldehyde** are virtually always used instead of their IUPAC names.



Greek letters are used to designate the location of substituents in common names. **The carbon adjacent to the CHO group is the α carbon**, and so forth down the chain.




Figure 21.1 gives the common and IUPAC names for three aldehydes.

21.2C Naming Ketones in the IUPAC System


- In the IUPAC system all ketones are identified by the suffix *-one*.

To name an acyclic ketone using IUPAC rules:

- [1] Find the longest chain containing the carbonyl group, and change the *-e* ending of the parent alkane to the suffix *-one*.
- [2] Number the carbon chain to give the carbonyl carbon the lower number. Apply all of the other usual rules of nomenclature.


With cyclic ketones, numbering always begins at the carbonyl carbon, but the “1” is usually omitted from the name. The ring is then numbered clockwise or counterclockwise to give the first substituent the lower number.

Figure 21.1
Three examples of aldehyde nomenclature




(Common names are in parentheses.)

Sample Problem 21.2 Give the IUPAC name for each ketone.




Solution


a. [1] Find and name the longest chain containing the carbonyl group:




[2] Number and name substituents:




b. [1] Name the ring:



[2] Number and name substituents:




Problem 21.5 Give the IUPAC name for each ketone.



21.2D Common Names for Ketones

Most common names for ketones are formed by **naming both alkyl groups** on the carbonyl carbon, **arranging them alphabetically**, and adding the word *ketone*. Using this method, the common name for 2-butanone becomes ethyl methyl ketone.



Three widely used common names for some simple ketones do not follow this convention:






Figure 21.2 gives acceptable names for two ketones.

Figure 21.2
Two examples of
ketone nomenclature




IUPAC name: 2-methyl-3-pentanone
Common name: ethyl isopropyl ketone



m-bromoacetophenone
or
3-bromoacetophenone

21.2E Additional Nomenclature Facts


Do not confuse a **benzyl** group with a **benzoyl** group.




benzyl group



formyl group




acetyl group




benzoyl group

Sometimes acyl groups (RCO–) must be named as substituents. To name an acyl group, take either the IUPAC or common parent name and add the suffix **-yl** or **-oyl**. The three most common acyl groups are drawn below.

Compounds containing both a C–C double bond and an aldehyde are named as **enals**, and compounds that contain both a C–C double bond and a ketone are named as **enones**. The chain is numbered to give the carbonyl group the lower number.





2,2-dimethyl-3-butenal



4-methyl-3-penten-2-one

Problem 21.6 Give the structure corresponding to each name: (a) *sec*-butyl ethyl ketone; (b) methyl vinyl ketone; (c) *p*-ethylacetophenone; (d) 3-benzoyl-2-benzylcyclopentanone; (e) 6,6-dimethyl-2-cyclohexenone; (f) 3-ethyl-5-hexenal.

Problem 21.7 Give the IUPAC name (including any *E,Z* designation) for each unsaturated aldehyde. Neral is obtained from lemon grass, and the unmistakable odor of a freshly cut cucumber is due largely to cucumber aldehyde.




21.3 Physical Properties

Aldehydes and ketones exhibit dipole–dipole interactions because of their polar carbonyl group. Because they have no O–H bond, two molecules of RCHO or RCOR are incapable of intermolecular hydrogen bonding, making them less polar than alcohols and carboxylic acids. How these intermolecular forces affect the physical properties of aldehydes and ketones is summarized in Table 21.1.

Problem 21.8 The boiling point of 2-butanone (80 °C) is significantly higher than the boiling point of diethyl ether (35 °C), even though both compounds exhibit dipole–dipole interactions and have comparable molecular weights. Offer an explanation.

Table 21.1 Physical Properties of Aldehydes and Ketones

Property	Observation
Boiling point and melting point	<ul style="list-style-type: none"> For compounds of comparable molecular weight, bp's and mp's follow the usual trend: The stronger the intermolecular forces, the higher the bp or mp. <div style="display: flex; justify-content: space-around; align-items: center; margin-top: 10px;"> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ VDW MW = 72 bp 36 °C </div> <div style="border: 1px solid black; padding: 5px; text-align: center;"> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$ VDW, DD MW = 72 bp 76 °C $\text{CH}_3\text{CH}_2\text{COCH}_3$ VDW, DD MW = 72 bp 80 °C </div> <div style="text-align: center;"> $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ VDW, DD, HB MW = 74 bp 118 °C </div> </div> <div style="text-align: center; margin-top: 10px;">  <p>Increasing strength of intermolecular forces Increasing boiling point</p> </div>
Solubility	<ul style="list-style-type: none"> RCHO and RCOR are soluble in organic solvents regardless of size. RCHO and RCOR having ≤ 5 C's are H₂O soluble because they can hydrogen bond with H₂O (Section 3.4C). RCHO and RCOR having > 5 C's are H₂O insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H₂O solvent.

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding, MW = molecular weight

21.4 Spectroscopic Properties

The presence of the carbonyl group in aldehydes and ketones gives them characteristic absorptions in their IR and NMR spectra.

21.4A IR Spectra

Aldehydes and ketones exhibit the following characteristic IR absorptions:

- Like all carbonyl compounds, **aldehydes and ketones give a strong peak at $\sim 1700\text{ cm}^{-1}$ due to the C=O.**
- The sp^2 hybridized C–H bond of an aldehyde shows one or two peaks at $\sim 2700\text{--}2830\text{ cm}^{-1}$.**

The IR spectrum of propanal in Figure 21.3 illustrates these characteristic peaks.

The exact position of the carbonyl absorption often provides additional information about a compound. For example, most aldehydes have a C=O peak around 1730 cm^{-1} , whereas for ketones, it is typically around 1715 cm^{-1} . Two other structural features—**ring size** (for cyclic ketones) and **conjugation**—affect the location of the carbonyl absorption in a predictable manner.

[1] The carbonyl absorption of cyclic ketones shifts to higher wavenumber as the size of the ring decreases and the ring strain increases.





Figure 21.3
The IR spectrum of propanal, $\text{CH}_3\text{CH}_2\text{CHO}$



- A strong $\text{C}=\text{O}$ occurs at 1739 cm^{-1} .
- The $\text{sp}^2\text{ C-H}$ of the CHO appears as two peaks at 2813 and 2716 cm^{-1} .

[2] Conjugation of the carbonyl group with a $\text{C}=\text{C}$ or a benzene ring shifts the absorption to lower wavenumber by $\sim 30\text{ cm}^{-1}$.

The effect of conjugation on the frequency of the $\text{C}=\text{O}$ absorption is explained by **resonance**. An α,β -unsaturated carbonyl compound can be written as three resonance structures, two of which place a single bond between the carbon and oxygen atoms of the carbonyl group. Thus, the π bond of the carbonyl group is delocalized, giving the conjugated carbonyl group some single bond character, and making it somewhat **weaker** than an unconjugated $\text{C}=\text{O}$. **Weaker bonds absorb at lower frequency (lower wavenumber) in an IR spectrum.**





Figure 21.4 illustrates the effects of conjugation on the location of the carbonyl absorption in some representative compounds.

Problem 21.9 Which carbonyl group in each pair absorbs at a higher frequency?



Problem 21.10 Why does the carbonyl group of a ketone absorb at lower wavenumber in IR spectroscopy than the carbonyl group of an aldehyde? For example, the carbonyl absorptions for hexanal ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$) and 3-hexanone ($\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_2\text{CH}_3$) are 1730 and 1715 cm^{-1} , respectively.

Figure 21.4
The effect of conjugation on the carbonyl absorption in an IR spectrum



21.4B NMR Spectra

Aldehydes and ketones exhibit the following characteristic ^1H and ^{13}C NMR absorptions:


- The sp^2 hybridized C–H proton of an aldehyde is highly deshielded and absorbs far downfield at 9–10 ppm. Splitting occurs with protons on the α carbon, but the coupling constant is often very small ($J = 1\text{--}3$ Hz).
- Protons on the α carbon to the carbonyl group absorb at 2–2.5 ppm. Methyl ketones, for example, give a characteristic singlet at ~ 2.1 ppm.
- In a ^{13}C NMR spectrum, the carbonyl carbon is highly deshielded, appearing in the 190–215 ppm region.

The ^1H and ^{13}C NMR spectra of propanal are illustrated in Figure 21.5.

Problem 21.11

Draw the structure of all constitutional isomers that contain a ketone and have molecular formula $\text{C}_5\text{H}_{10}\text{O}$. Give the IUPAC name for each isomer and state how ^{13}C NMR spectroscopy could be used to distinguish these isomers.

Figure 21.5
The ^1H and ^{13}C NMR spectra of propanal, $\text{CH}_3\text{CH}_2\text{CHO}$




- **^1H NMR:** There are three signals due to the three different kinds of hydrogens, labeled H_a , H_b , and H_c . The deshielded CHO proton occurs downfield at 9.8 ppm. The H_c signal is split into a triplet by the adjacent CH_2 group, but the coupling constant is small.
- **^{13}C NMR:** There are three signals due to the three different kinds of carbons, labeled C_a , C_b , and C_c . The deshielded carbonyl carbon absorbs downfield at 203 ppm.

21.5 Interesting Aldehydes and Ketones



formaldehyde
 $\text{CH}_2=\text{O}$



acetone
 $(\text{CH}_3)_2\text{C}=\text{O}$

Because it is a starting material for the synthesis of many resins and plastics, billions of pounds of **formaldehyde** are produced annually in the United States by the oxidation of methanol (CH_3OH). Formaldehyde is also sold as a 37% aqueous solution called **formalin**, which has been used as a disinfectant, antiseptic, and preservative for biological specimens. Formaldehyde, a product of the incomplete combustion of coal and other fossil fuels, is partly responsible for the irritation caused by smoggy air.

Acetone is an industrial solvent and a starting material in the synthesis of some organic polymers. Acetone is produced *in vivo* during the breakdown of fatty acids. In diabetes, a common endocrine disease in which normal metabolic processes are altered because of the inadequate secretion of insulin, individuals often have unusually high levels of acetone in their bloodstreams. The characteristic odor of acetone can be detected on the breath of diabetic patients when their disease is poorly controlled.

Many aldehydes with characteristic odors occur in nature, as shown in Figure 21.6.

Many steroid hormones contain a carbonyl along with other functional groups. **Cortisone** and **prednisone** are two anti-inflammatory steroids with closely related structures. Cortisone is secreted by the body's adrenal gland, whereas prednisone is a synthetic analogue used in the treatment of inflammatory diseases such as arthritis and asthma.





Figure 21.6 Some naturally occurring aldehydes with strong odors




21.6 Preparation of Aldehydes and Ketones

Aldehydes and ketones can be prepared by a variety of methods. Because these reactions are needed for many multistep syntheses, Section 21.6 briefly summarizes earlier reactions that synthesize an aldehyde or ketone.


21.6A Common Methods to Synthesize Aldehydes

Aldehydes are prepared from 1° alcohols, esters, acid chlorides, and alkynes.




21.6B Common Methods to Synthesize Ketones

Ketones are prepared from 2° alcohols, acid chlorides, and alkynes.




Aldehydes and ketones are also both obtained as products of the oxidative cleavage of alkenes (Section 12.10).



Problem 21.12 What reagents are needed to convert each compound into butanal ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$):
 (a) $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOCH}_3$; (b) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$; (c) $\text{HC}\equiv\text{CCH}_2\text{CH}_3$;
 (d) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}=\text{CHCH}_2\text{CH}_2\text{CH}_3$?

Problem 21.13 What reagents are needed to convert each compound into acetophenone ($\text{C}_6\text{H}_5\text{COCH}_3$):
 (a) benzene; (b) $\text{C}_6\text{H}_5\text{COCl}$; (c) $\text{C}_6\text{H}_5\text{C}\equiv\text{CH}$?

Problem 21.14 What alkene would yield 2,2-dimethoxy-1,3-cyclopentanedicarbaldehyde on treatment with O_3 followed by $(\text{CH}_3)_2\text{S}$?




2,2-dimethoxy-1,3-cyclopentanedicarbaldehyde

21.7 Reactions of Aldehydes and Ketones—General Considerations

Let's begin our discussion of carbonyl reactions by looking at the two general kinds of reactions that aldehydes and ketones undergo.

[1] Reaction at the carbonyl carbon

Recall from Chapter 20 that the uncrowded, electrophilic carbonyl carbon makes aldehydes and ketones susceptible to **nucleophilic addition** reactions.




The elements of H and Nu are added to the carbonyl group. In Chapter 20 you learned about this reaction with hydride (H^-) and carbanions (R^-) as nucleophiles. In Chapter 21, we will discuss similar reactions with other nucleophiles.

[2] Reaction at the α carbon

A second general reaction of aldehydes and ketones involves reaction at the **α carbon**. A C–H bond on the α carbon to a carbonyl group is more acidic than many other C–H bonds, because reaction with base forms a resonance-stabilized enolate anion.

- Enolates are nucleophiles, and so they react with electrophiles to form new bonds on the α carbon.



Chapters 23 and 24 are devoted to reactions at the α carbon to a carbonyl group.


- Aldehydes and ketones react with nucleophiles at the carbonyl carbon.
- Aldehydes and ketones form enolates that react with electrophiles at the α carbon.

21.7A The General Mechanism of Nucleophilic Addition

Two general mechanisms are usually drawn for nucleophilic addition, depending on the nucleophile (negatively charged versus neutral) and the presence or absence of an acid catalyst. With negatively charged nucleophiles, nucleophilic addition follows the two-step process first discussed in Chapter 20—**nucleophilic attack** followed by **protonation**, as shown in Mechanism 21.1.



Mechanism 21.1 General Mechanism—Nucleophilic Addition



- In Step [1], the **nucleophile attacks the carbonyl group**, cleaving the π bond and moving an electron pair onto oxygen. This forms an sp^3 hybridized intermediate with a new C–Nu bond.
- In Step [2], **protonation of the negatively charged O atom by H_2O** forms the addition product.


In this mechanism **nucleophilic attack precedes protonation**. This process occurs with strong neutral or negatively charged nucleophiles.

With some neutral nucleophiles, however, nucleophilic addition does not occur unless an acid catalyst is added. The general mechanism for this reaction consists of three steps (not two), but the same product results because H and Nu add across the carbonyl π bond. In this mechanism, **protonation precedes nucleophilic attack**. Mechanism 21.2 is shown with the neutral nucleophile H–Nu: and a general acid H–A.




Mechanism 21.2 General Mechanism—Acid-Catalyzed Nucleophilic Addition


Step [1] Protonation of the carbonyl group



Steps [2]–[3] Nucleophilic attack and deprotonation



The effect of protonation is to convert a neutral carbonyl group to one having a net positive charge. **This protonated carbonyl group is much more electrophilic**, and much more susceptible to attack by a nucleophile. This step is unnecessary with strong nucleophiles like hydride (H^-) that were used in Chapter 20. With weaker nucleophiles, however, nucleophilic attack does not occur unless the carbonyl group is first protonated.




This step is a specific example of a general phenomenon.

- Any reaction involving a carbonyl group and a strong acid begins with the same first step—protonation of the carbonyl oxygen.

21.7B The Nucleophile

What nucleophiles add to carbonyl groups? This cannot be predicted solely on the trends in nucleophilicity learned in Chapter 7. Only *some* of the nucleophiles that react well in nucleophilic substitution at sp^3 hybridized carbons give reasonable yields of nucleophilic addition products.

Cl^- , Br^- , and I^- are good nucleophiles in substitution reactions at sp^3 hybridized carbons, but they are ineffective nucleophiles in addition. Addition of Cl^- to a carbonyl group, for example, would cleave the C–O π bond, forming an alkoxide. Because Cl^- is a much weaker base than the alkoxide formed, equilibrium favors the starting materials (the weaker base, Cl^-), *not* the addition product.



The situation is further complicated because some of the initial nucleophilic addition adducts are unstable and undergo elimination to form a stable product. For example, amines (RNH_2) add to carbonyl groups in the presence of mild acid to form unstable **carbinolamines**, which readily lose water to form **imines**. This addition–elimination sequence replaces a $\text{C}=\text{O}$ by a $\text{C}=\text{N}$. The details of this process are discussed in Section 21.11.





Figure 21.7 lists nucleophiles that add to a carbonyl group, as well as the products obtained from nucleophilic addition using cyclohexanone as a representative ketone. These reactions are discussed in the remaining sections of Chapter 21. In cases in which the initial addition adduct is unstable, it is enclosed within brackets, followed by the final product.

Problem 21.15 Why does equilibrium favor the product when H^- adds to a carbonyl group?


Figure 21.7
Specific examples of nucleophilic addition




21.8 Nucleophilic Addition of H^- and R^- —A Review

We begin our study of nucleophilic additions to aldehydes and ketones by briefly reviewing nucleophilic addition of hydride and carbanions, two reactions examined in Sections 20.4 and 20.10, respectively.


Treatment of an aldehyde or ketone with either NaBH_4 or LiAlH_4 followed by protonation forms a 1° or 2° alcohol. NaBH_4 and LiAlH_4 serve as a source of **hydride, H^-** —the **nucleophile**—and the reaction results in addition of the elements of H_2 across the $\text{C}=\text{O}$ π bond. Addition of H_2 reduces the carbonyl group to an alcohol.



Hydride reduction of aldehydes and ketones occurs via the two-step mechanism of nucleophilic addition—that is, **nucleophilic attack of H^-** followed by **protonation**—shown previously in Section 20.4B.




Treatment of an aldehyde or ketone with either an organolithium ($\text{R}''\text{Li}$) or Grignard reagent ($\text{R}''\text{MgX}$) followed by water forms a 1°, 2°, or 3° alcohol containing a new carbon-carbon bond. $\text{R}''\text{Li}$ and $\text{R}''\text{MgX}$ serve as a source of a **carbanion (R''^-)**—the **nucleophile**—and the reaction results in addition of the elements of R'' and H across the $\text{C}=\text{O}$ π bond.




The stereochemistry of hydride reduction and Grignard addition was discussed previously in Sections 20.5 and 20.10B, respectively.

The nucleophilic addition of carbanions to aldehydes and ketones occurs via the two-step mechanism of nucleophilic addition—that is, **nucleophilic attack of (R''^-)** followed by **protonation**—shown previously in Section 20.10.




In both reactions, the nucleophile—either hydride or a carbanion—attacks the trigonal planar sp^2 hybridized carbonyl from both sides, so that when a new stereogenic center is formed, a mixture of stereoisomers results, as shown in Sample Problem 21.3.

Sample Problem 21.3 Draw the products (including the stereochemistry) formed in the following reaction.




Solution

The Grignard reagent adds CH_3^- from both sides of the trigonal planar carbonyl group, yielding a mixture of 3° alcohols after protonation with water. In this example, the starting ketone and both alcohol products are chiral. The two products, which contain two stereogenic centers, are stereoisomers but not mirror images—that is, they are **diastereomers**.




Problem 21.16 Draw the products of each reaction. Include all stereoisomers formed.




21.9 Nucleophilic Addition of $^- \text{CN}$

Treatment of an aldehyde or ketone with NaCN and a strong acid such as HCl adds the elements of HCN across the carbon–oxygen π bond, forming a **cyanohydrin**.



This reaction adds one carbon to the aldehyde or ketone, forming a new carbon–carbon bond.




21.9A The Mechanism

The mechanism of cyanohydrin formation involves the usual two steps of nucleophilic addition: **nucleophilic attack followed by protonation** as shown in Mechanism 21.3.




Mechanism 21.3 Nucleophilic Addition of CN^- —Cyanohydrin Formation



- In Step [1], **nucleophilic attack of CN^-** forms a new carbon–carbon bond with cleavage of the C–O π bond.
- In Step [2], **protonation of the negatively charged O atom by HCN** forms the addition product. The hydrogen cyanide (HCN) used in this step is formed by the acid–base reaction of cyanide (CN^-) with the strong acid, HCl.


This reaction does not occur with HCN alone. The **cyanide anion** makes addition possible because it is a strong nucleophile that attacks the carbonyl group.

Cyanohydrins can be reconverted to carbonyl compounds by treatment with base. This process is just the reverse of the addition of HCN: **deprotonation followed by elimination of CN^-** .




Note the difference between two similar terms. **Hydration** results in *adding* water to a compound. **Hydrolysis** results in *cleaving bonds* with water.

The cyano group (CN) of a cyanohydrin is readily hydrolyzed to a carboxy group (COOH) by heating with aqueous acid or base. Hydrolysis replaces the three C–N bonds by three C–O bonds.



Problem 21.17 Draw the products of each reaction.




21.9B Application: Naturally Occurring Cyanohydrin Derivatives



Peach and apricot pits are a natural source of the cyanohydrin derivative amygdalin.

Although the cyanohydrin is an uncommon functional group, **linamarin** and **amygdalin** are two naturally occurring cyanohydrin derivatives. Both contain a carbon atom bonded to both an oxygen atom and a cyano group, analogous to a cyanohydrin.



Linamarin is isolated from cassava, a woody shrub grown as a root crop in the humid tropical regions of South America and Africa. **Amygdalin** is present in the seeds and pits of apricots,




Cassava is a widely grown root crop, first introduced to Africa by Portuguese traders from Brazil in the sixteenth century. The peeled root is eaten after boiling or roasting. If the root is eaten without processing, illness and even death can result from high levels of HCN.

peaches, and wild cherries. Amygdalin, sometimes called **laetrile**, was once touted as an anti-cancer drug, and is still available in some countries for this purpose, although its effectiveness is unproven.

Both **linamarin** and **amygdalin** are toxic compounds because they are metabolized to cyanohydrins, which are hydrolyzed to carbonyl compounds and toxic HCN gas, a cellular poison with a characteristic almond odor. This second step is merely the reconversion of a cyanohydrin to a carbonyl compound, a process that occurs with base in reactions run in the laboratory (Section 21.9A). If cassava root is processed with care, linamarin is enzymatically metabolized by this reaction sequence and the toxic HCN is released before the root is ingested, making it safe to eat.

The breakdown of linamarin to HCN




Problem 21.18 What cyanohydrin and carbonyl compound are formed when amygdalin is metabolized in a similar manner?


21.10 The Wittig Reaction

The additions of H^- , R^- , and CN^- all involve the same two steps—nucleophilic attack followed by protonation. Other examples of nucleophilic addition in Chapter 21 are somewhat different. Although they still involve attack of a nucleophile, the initial addition adduct is converted to another product by one or more reactions.

The first reaction in this category is the **Wittig reaction**, named for German chemist Georg Wittig, who was awarded the Nobel Prize in Chemistry in 1979 for its discovery. The Wittig reaction uses a carbon nucleophile, the **Wittig reagent**, to form **alkenes**. When a carbonyl compound is treated with a Wittig reagent, the carbonyl oxygen atom is replaced by the negatively charged alkyl group bonded to the phosphorus—that is, **the $\text{C}=\text{O}$ is converted to a $\text{C}=\text{C}$** .




A Wittig reaction forms two new carbon–carbon bonds—one new σ bond and one new π bond—as well as a phosphorus by-product, $\text{Ph}_3\text{P}=\text{O}$ (triphenylphosphine oxide).



21.10A The Wittig Reagent

A **Wittig reagent** is an **organophosphorus reagent**—a reagent that contains a carbon–phosphorus bond. A typical Wittig reagent has a phosphorus atom bonded to three phenyl groups, plus another alkyl group that bears a negative charge.




Phosphorus ylides are also called **phosphoranes**.


A Wittig reagent is an **ylide**, a species that contains two oppositely charged atoms bonded to each other, and both atoms have octets. In a Wittig reagent, a negatively charged carbon atom is bonded to a positively charged phosphorus atom.

Because phosphorus is a third-row element, it can be surrounded by more than eight electrons. As a result, a second resonance structure can be drawn that places a double bond between carbon and phosphorus. Regardless of which resonance structure is drawn, a **Wittig reagent has no net charge**. In one resonance structure, though, the carbon atom bears a net negative charge, so it is nucleophilic.

Two resonance structures for the Wittig reagent




Wittig reagents are synthesized by a two-step procedure.

Step [1] $\text{S}_{\text{N}}2$ reaction of triphenylphosphine with an alkyl halide forms a phosphonium salt.

Because phosphorus is located below nitrogen in the periodic table, a neutral phosphorus atom with three bonds also has a lone pair of electrons.

Triphenylphosphine ($\text{Ph}_3\text{P}:$), which contains a lone pair of electrons on P, is the nucleophile. Because the reaction follows an $\text{S}_{\text{N}}2$ mechanism, it works best with unhindered CH_3X and 1° alkyl halides (RCH_2X). Secondary alkyl halides (R_2CHX) can also be used, although yields are often lower.

Step [2] Deprotonation of the phosphonium salt with a strong base (:B) forms the ylide.




Section 20.9C discussed the reaction of organometallic reagents as strong bases.

Because removal of a proton from a carbon bonded to phosphorus generates a resonance-stabilized carbanion (the ylide), this proton is somewhat more acidic than other protons on an


alkyl group in the phosphonium salt. Very strong bases are still needed, though, to favor the products of this acid–base reaction. Common bases used for this reaction are the organolithium reagents such as butyllithium, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Li}$, abbreviated as BuLi.

To synthesize the Wittig reagent, $\text{Ph}_3\text{P}=\text{CH}_2$, use these two steps:




- **Step [1]** Form the phosphonium salt by $\text{S}_{\text{N}}2$ reaction of $\text{Ph}_3\text{P:}$ and CH_3Br .
- **Step [2]** Form the ylide by removal of a proton using BuLi as a strong base.

Problem 21.19 Draw the products of the following Wittig reactions.



Problem 21.20 Outline a synthesis of each Wittig reagent from Ph_3P and an alkyl halide.




21.10B Mechanism of the Wittig Reaction

The currently accepted mechanism of the Wittig reaction involves two steps. Like other nucleophiles, the Wittig reagent attacks an electrophilic carbonyl carbon, but then the initial addition adduct undergoes elimination to form an alkene. Mechanism 21.4 is drawn using $\text{Ph}_3\text{P}=\text{CH}_2$.




Mechanism 21.4 The Wittig Reaction

Step [1] Nucleophilic addition forms a four-membered ring.



- Step [1] forms two bonds and generates a four-membered ring. The negatively charged carbon atom of the ylide attacks the carbonyl carbon to form a new carbon–carbon σ bond, while the carbonyl O atom attacks the positively charged P atom.
- This process generates an **oxaphosphetane**, a four-membered ring containing a strong P–O bond.

Step [2] Elimination of $\text{Ph}_3\text{P}=\text{O}$ forms the alkene.



- In Step [2], **$\text{Ph}_3\text{P}=\text{O}$ (triphenylphosphine oxide) is eliminated**, forming two new π bonds. The formation of the very strong P–O double bond provides the driving force for the Wittig reaction.

One limitation of the Wittig reaction is that a mixture of alkene stereoisomers sometimes forms. For example, reaction of propanal ($\text{CH}_3\text{CH}_2\text{CHO}$) with a Wittig reagent forms the mixture of *E* and *Z* isomers shown.





Figure 21.8
A Wittig reaction used to synthesize β -carotene



- The more stable *E* alkene is the major product in this Wittig reaction.

Because the Wittig reaction forms two carbon–carbon bonds in a single reaction, it has been used to synthesize many natural products, including β -carotene, shown in Figure 21.8.


Problem 21.21 Draw the products (including stereoisomers) formed when benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$) is treated with each Wittig reagent: (a) $\text{Ph}_3\text{P}=\text{CHCH}_2\text{CH}_3$; (b) $\text{Ph}_3\text{P}=\text{CHC}_6\text{H}_5$; (c) $\text{Ph}_3\text{P}=\text{CHCOOCH}_3$.

21.10C Retrosynthetic Analysis


To use the Wittig reaction in synthesis, you must be able to determine what carbonyl compound and Wittig reagent are needed to prepare a given compound—that is, **you must work backwards, in the retrosynthetic direction**. There can be two different Wittig routes to a given alkene, but one is often preferred on steric grounds.

HOW TO Determine the Starting Materials for a Wittig Reaction Using Retrosynthetic Analysis

Example What starting materials are needed to synthesize alkene A by a Wittig reaction?



Step [1] Cleave the carbon–carbon double bond into two components.




- Part of the molecule becomes the carbonyl component and the other part becomes the Wittig reagent.

—Continued


HOW TO, continued . . .

There are usually two routes to a given alkene using a Wittig reaction:

Possibility [1]




Possibility [2]



Step [2] Compare the Wittig reagents. The preferred pathway uses a Wittig reagent derived from an unhindered alkyl halide— CH_3X or RCH_2X .

Determine what alkyl halide is needed to prepare each Wittig reagent:




Because the synthesis of the Wittig reagent begins with an $\text{S}_{\text{N}}2$ reaction, **the preferred pathway begins with an unhindered methyl halide or 1° alkyl halide.** In this example, retrosynthetic analysis of both Wittig reagents indicates that only one of them ($\text{Ph}_3\text{P}=\text{CHCH}_3$) can be synthesized from a 1° alkyl halide, making Possibility [1] the preferred pathway.

Problem 21.22 What starting materials are needed to prepare each alkene by a Wittig reaction? When there are two possible routes, indicate which route, if any, is preferred: (a) $(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{CH}_3$; (b) $\text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$; (c) $\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_3$.

21.10D Comparing Methods of Alkene Synthesis


An advantage in using the Wittig reaction over other elimination methods to synthesize alkenes is that **you always know the location of the double bond.** Whereas other methods of alkene synthesis often give a mixture of constitutional isomers, the Wittig reaction always gives a single constitutional isomer.

For example, two methods can be used to convert cyclohexanone into alkene **B** (methylenecyclohexane): a two-step method consisting of Grignard addition followed by dehydration, or a one-step Wittig reaction.




In a two-step method, treatment of cyclohexanone with CH_3MgBr forms a 3° alcohol after protonation. Dehydration of the alcohol with H_2SO_4 forms a mixture of alkenes, in which the desired disubstituted alkene is the minor product.

Recall from Section 9.8 that the major product formed in acid-catalyzed dehydration of an alcohol is the more substituted alkene.




By contrast, reaction of cyclohexanone with $\text{Ph}_3\text{P}=\text{CH}_2$ affords the desired alkene as the only product. The newly formed double bond always joins the carbonyl carbon with the negatively charged carbon of the Wittig reagent. In other words, **the position of the double bond is always unambiguous in the Wittig reaction.** This makes the Wittig reaction an especially attractive method for preparing many alkenes.




Problem 21.23

Show two methods to synthesize each alkene: a one-step method using a Wittig reagent, and a two-step method that forms a carbon-carbon bond with an organometallic reagent in one of the steps.



21.11 Addition of 1° Amines


We now move on to the reaction of aldehydes and ketones with nitrogen and oxygen heteroatoms. **Amines, for example, are organic nitrogen compounds that contain a nonbonded electron pair on the N atom.** Amines are classified as 1°, 2°, or 3° by the number of alkyl groups bonded to the *nitrogen* atom.




Both 1° and 2° amines react with aldehydes and ketones. We begin by examining the reaction of aldehydes and ketones with 1° amines.

21.11A Formation of Imines

Treatment of an aldehyde or ketone with a 1° amine affords an **imine** (also called a **Schiff base**). Nucleophilic attack of the 1° amine on the carbonyl group forms an unstable **carbinolamine**, which loses water to form an imine. The overall reaction results in **replacement of C=O by C=NR**.



Because the N atom of an imine is surrounded by three groups (two atoms and a lone pair), it is sp^2 hybridized, making the C-N-R'' bond angle $\sim 120^\circ$ (not 180°). Imine formation is fastest when the reaction medium is weakly acidic.




The mechanism of imine formation (Mechanism 21.5) can be divided into two distinct parts: **nucleophilic addition of the 1° amine, followed by elimination of H₂O**. Each step involves a reversible equilibrium, so that the reaction is driven to completion by removing H₂O.




Mechanism 21.5 Imine Formation from an Aldehyde or Ketone

Part [1] Nucleophilic addition forms a carbinolamine.



- **Nucleophilic attack** of the amine followed by proton transfer forms the unstable carbinolamine (Steps [1]–[2]). These steps result in the addition of H and NHR'' to the carbonyl group.


Part [2] Elimination of H₂O forms an imine.



- Elimination of H₂O forms the imine in three steps. Protonation of the OH group in Step [3] forms a good leaving group, leading to **loss of water** in Step [4], giving a resonance-stabilized **iminium ion**. Loss of a proton forms the imine in Step [5].
- Except for Steps [1] (nucleophilic addition) and [4] (H₂O elimination), all other steps in the mechanism are acid–base reactions—that is, moving a proton from one atom to another.




Imine formation is most rapid at pH 4–5. Mild acid is needed for protonation of the hydroxy group in Step [3] to form a **good leaving group**. Under strongly acidic conditions, the reaction rate decreases because the amine nucleophile is protonated. With no free electron pair, it is no longer a nucleophile, and so nucleophilic addition cannot occur.

Protonation makes a good leaving group...but at low pH, the basic amine is protonated.





Problem 21.24

Draw the product formed when CH₃CH₂CH₂CH₂NH₂ reacts with each carbonyl compound in the presence of mild acid.




Problem 21.25 What 1° amine and carbonyl compound are needed to prepare each imine?



21.11B Application: Retinal, Rhodopsin, and the Chemistry of Vision


Many imines play vital roles in biological systems. A key molecule in the chemistry of vision is the highly conjugated imine **rhodopsin**, which is synthesized in the rod cells of the eye from **11-*cis*-retinal** and a 1° amine in the protein **opsin**.



The central role of rhodopsin in the visual process was delineated by Nobel Laureate George Wald of Harvard University.

The complex process of vision centers around this imine derived from retinal (Figure 21.9). The 11-*cis* double bond in rhodopsin creates crowding in the rather rigid side chain. When light strikes the rod cells of the retina, it is absorbed by the conjugated double bonds of rhodopsin, and the 11-*cis* double bond is isomerized to the 11-*trans* arrangement. This isomerization is accompanied by a drastic change in shape in the protein, altering the concentration of Ca^{2+} ions moving across the cell membrane, and sending a nerve impulse to the brain, which is then processed into a visual image.

Figure 21.9
The key reaction in the chemistry of vision




- Rhodopsin is a light-sensitive compound located in the membrane of the rod cells in the retina of the eye. Rhodopsin contains the protein opsin bonded to 11-*cis*-retinal via an imine linkage. When light strikes this molecule, the crowded 11-*cis* double bond isomerizes to the 11-*trans* isomer, and a nerve impulse is transmitted to the brain by the optic nerve.

Problem 21.26 Draw a stepwise mechanism for the formation of rhodopsin from 11-*cis*-retinal and an NH₂ group in opsin.


21.12 Addition of 2° Amines

21.12A Formation of Enamines

A 2° amine reacts with an aldehyde or ketone to give an **enamine**. *Enamines have a nitrogen atom bonded to a double bond (alkene + amine = enamine)*.



Like imines, enamines are also formed by the addition of a nitrogen nucleophile to a carbonyl group followed by elimination of water. In this case, however, elimination occurs across two adjacent *carbon* atoms to form a new carbon–carbon π bond.




The mechanism for enamine formation (Mechanism 21.6) is identical to the mechanism for imine formation except for the last step, involving formation of the π bond. The mechanism can be divided into two distinct parts: **nucleophilic addition of the 2° amine, followed by elimination of H₂O**. Each step involves a reversible equilibrium once again, so that the reaction is driven to completion by removing H₂O.




Mechanism 21.6 Enamine Formation from an Aldehyde or Ketone

Part [1] Nucleophilic addition forms a carbinolamine.




- **Nucleophilic attack** of the amine followed by proton transfer forms the unstable carbinolamine (Steps [1]–[2]).

Part [2] Elimination of H₂O forms an enamine.



- Protonation of the OH group in Step [3] forms a good leaving group, leading to **loss of water** in Step [4], giving a resonance-stabilized **iminium ion**.
- Removal of a proton from the adjacent C–H bond forms the enamine in Step [5].

Figure 21.10
The formation of imines and enamines compared




- With a **1° amine**, the intermediate iminium ion still has a proton on the N atom that may be removed to form a C=N .
- With a **2° amine**, the intermediate iminium ion has no proton on the N atom. A proton must be removed from an adjacent C-H bond, and this forms a C=C .

The mechanisms illustrate why **the reaction of 1° amines with carbonyl compounds forms imines, but the reaction with 2° amines forms enamines**. In Figure 21.10, the last step of both mechanisms is compared using cyclohexanone as starting material. The position of the double bond depends on which proton is removed in the last step. Removal of an N-H proton forms a C=N , whereas removal of a C-H proton forms a C=C .

21.12B Imine and Enamine Hydrolysis

Because imines and enamines are formed by a set of reversible reactions, both can be converted back to carbonyl compounds by hydrolysis with mild acid. The mechanism of these reactions is exactly the *reverse* of the mechanism written for the formation of imines and enamines. In the hydrolysis of enamines, the carbonyl carbon in the product comes from the sp^2 hybridized carbon bonded to the N atom in the starting material.


- Hydrolysis of imines and enamines forms aldehydes and ketones.



Problem 21.27 What two enamines are formed when 2-methylcyclohexanone is treated with $(\text{CH}_3)_2\text{NH}$?


Problem 21.28 Explain why benzaldehyde cannot form an enamine with a 2° amine.

Problem 21.29 What carbonyl compound and amine are formed by the hydrolysis of each compound?




21.13 Addition of H₂O—Hydration

Treatment of a carbonyl compound with H₂O in the presence of an acid or base catalyst **adds the elements of H and OH across the carbon–oxygen π bond**, forming a *gem*-diol or hydrate.



Hydration of a carbonyl group gives a good yield of *gem*-diol only with an unhindered aldehyde like formaldehyde, and with aldehydes containing nearby electron-withdrawing groups.




21.13A The Thermodynamics of Hydrate Formation

Whether addition of H₂O to a carbonyl group affords a good yield of the *gem*-diol depends on the relative energies of the starting material and the product. With less stable carbonyl starting materials, equilibrium favors the hydrate product, whereas with more stable carbonyl starting materials, equilibrium favors the carbonyl starting material. Because **alkyl groups stabilize a carbonyl group** (Section 20.2B):

- Increasing the number of alkyl groups on the carbonyl carbon decreases the amount of hydrate at equilibrium.

This can be illustrated by comparing the amount of hydrate formed from formaldehyde, acetaldehyde, and acetone.




Formaldehyde, the least stable carbonyl compound, forms the largest percentage of hydrate. On the other hand, acetone and other ketones, which have two electron-donor R groups, form < 1% of the hydrate at equilibrium. Other electronic factors come into play as well.

- Electron-donating groups near the carbonyl carbon stabilize the carbonyl group, decreasing the amount of the hydrate at equilibrium.
- Electron-withdrawing groups near the carbonyl carbon destabilize the carbonyl group, increasing the amount of hydrate at equilibrium.

Chloral hydrate, a sedative sometimes administered to calm a patient prior to a surgical procedure, has also been used for less reputable purposes. Adding it to an alcoholic beverage makes a so-called knock-out drink, causing an individual who drinks it to pass out. Because it is addictive and care must be taken in its administration, it is a controlled substance.

This explains why chloral (trichloroacetaldehyde) forms a large amount of hydrate at equilibrium. Three electron-withdrawing Cl atoms place a partial positive charge on the α carbon to the carbonyl, destabilizing the carbonyl group, and therefore increasing the amount of hydrate at equilibrium.



Having two similar charges (δ^+) on adjacent atoms destabilizes the carbonyl group.

A less stable carbonyl compound means more hydrate at equilibrium.

Problem 21.30


Which compound in each pair forms the higher percentage of *gem*-diol at equilibrium: (a) CH₃CH₂CH₂CHO or CH₃CH₂COCH₃; (b) CH₃CF₂CHO or CH₃CH₂CHO?

21.13B The Kinetics of Hydrate Formation

Although H₂O itself adds slowly to a carbonyl group, both acid and base catalyze the addition. In base, the nucleophile is ⁻OH, and the mechanism follows the usual two steps for nucleophilic addition: **nucleophilic attack followed by protonation**, as shown in Mechanism 21.7.



Mechanism 21.7 Base-Catalyzed Addition of H₂O to a Carbonyl Group




- In Step [1], the nucleophile (⁻OH) attacks the carbonyl group, cleaving the π bond, and moving an electron pair onto oxygen.
- In Step [2], protonation of the negatively charged O atom by H₂O forms the *gem*-diol.

The acid-catalyzed addition follows the general mechanism presented in Section 21.7A. For a poorer nucleophile like H₂O to attack a carbonyl group, the **carbonyl must be protonated by acid first; thus, protonation precedes nucleophilic attack**. The overall mechanism has three steps, as shown in Mechanism 21.8.




Mechanism 21.8 Acid-Catalyzed Addition of H₂O to a Carbonyl Group

Step [1] Protonation of the carbonyl group



- Protonation of the carbonyl oxygen forms a resonance-stabilized cation that bears a full positive charge.

Steps [2]–[3] Nucleophilic attack and deprotonation




- In Step [2], the nucleophile (H₂O) attacks, and then deprotonation forms the neutral addition product in Step [3].
- The overall result is the addition of H and OH to the carbonyl group and regeneration of the acid catalyst.

Acid and base increase the rate of reaction for different reasons.

- **Base converts H_2O into ^-OH , a stronger nucleophile.**
- **Acid protonates the carbonyl group, making it more electrophilic towards nucleophilic attack.**

These catalysts increase the rate of the reaction, but they do not affect the equilibrium constant. Starting materials that give a low yield of *gem*-diol do so whether or not a catalyst is present. Because these reactions are reversible, the conversion of *gem*-diols to aldehydes and ketones is also catalyzed by acid and base, and the steps of the mechanism are reversed.


Problem 21.31 Draw a stepwise mechanism for the following reaction.



21.14 Addition of Alcohols—Acetal Formation


The term *acetal* refers to any compound derived from an aldehyde or ketone, having two OR groups bonded to a single carbon. The term *ketal* is sometimes used when the starting carbonyl compound is a ketone; that is, the carbon bonded to the alkoxy groups is *not* bonded to a H atom and the general structure is $\text{R}_2\text{C}(\text{OR}')_2$. Since ketals are considered a subclass of acetals in the IUPAC system, we will use the single general term *acetal* for any compound having two OR groups on a carbon atom.

Aldehydes and ketones react with two equivalents of alcohol to form acetals. In an acetal, the carbonyl carbon from the aldehyde or ketone is now singly bonded to two OR'' (alkoxy) groups.




This reaction differs from other additions we have seen thus far, because **two equivalents of alcohol are added to the carbonyl group**, and two new C–O σ bonds are formed. Acetal formation is catalyzed by acids, commonly *p*-toluenesulfonic acid (TsOH).


Example



When a diol such as ethylene glycol is used in place of two equivalents of ROH, a cyclic acetal is formed. Both oxygen atoms in the cyclic acetal come from the diol.



Acetals are not ethers, even though both functional groups contain a C–O σ bond. Having two C–O σ bonds on the *same* carbon atom makes an acetal very different from an ether.



Like *gem*-diol formation, the synthesis of acetals is reversible, and often the equilibrium favors reactants, not products. In acetal synthesis, however, water is formed as a by-product, so the equilibrium can be driven to the right by removing the water as it is formed. This can be done in a variety of ways in the laboratory. A drying agent can be added that reacts with the water, or more commonly, the water can be distilled from the reaction mixture as it is formed by using a Dean–Stark trap, as pictured in Figure 21.11. Driving an equilibrium to the right by removing one of the products is an application of Le Châtelier's principle (see Section 9.8).

Problem 21.32 Draw the products of each reaction.





Figure 21.11
A Dean–Stark trap for removing water




- A **Dean–Stark trap** is an apparatus used for removing water from a reaction mixture. To use a Dean–Stark trap to convert a carbonyl compound to an acetal:

The carbonyl compound, an alcohol, and an acid are dissolved in benzene. As the mixture is heated, the carbonyl compound is converted to the acetal with water as a by-product. Benzene and water co-distill from the reaction mixture. When the hot vapors reach the cold condenser, they condense, forming a liquid that then collects in the glass tube below. Water, the more dense liquid, forms the lower layer, so that as it collects, it can be drained through the stopcock into a flask. In this way, water can be removed from a reaction mixture, driving the equilibrium.

21.14A The Mechanism

The mechanism for acetal formation can be divided into two parts: **the addition of one equivalent of alcohol** to form a **hemiacetal**, followed by the **conversion of the hemiacetal to the acetal**. A **hemiacetal** has a carbon atom bonded to one OH group and one OR group.




Like *gem*-diols, hemiacetals are often higher in energy than their carbonyl starting materials, making the direction of equilibrium unfavorable for hemiacetal formation. The elimination of H₂O, which can be removed from the reaction mixture to drive the equilibrium to favor product, occurs during the conversion of the hemiacetal to the acetal. This explains why two equivalents of ROH react with a carbonyl compound, forming the acetal as product.

The mechanism is written in two parts (Mechanisms 21.9 and 21.10) with a general acid HA.




Mechanism 21.9 Acetal Formation—Part [1] Formation of a Hemiacetal

Step [1] Protonation of the carbonyl group



- **Protonation** of the carbonyl oxygen forms a resonance-stabilized cation that bears a full positive charge.

Steps [2]–[3] Nucleophilic attack and deprotonation




- In Step [2], the **nucleophile (ROH) attacks**, and then deprotonation forms the neutral addition product in Step [3].
- **The overall result is the addition of H and OR to the carbonyl group.**

Formation of the acetal requires a second nucleophilic attack of ROH. For this to occur, however, **H₂O must first come off as a leaving group**, as shown in Mechanism 21.10.




Mechanism 21.10 Acetal Formation—Part [2] Formation of the Acetal

Steps [4]–[5] Elimination of H₂O



- Protonation of the OH group in the hemiacetal in Step [4] forms a **good leaving group (H₂O)**. Loss of H₂O in Step [5] forms a resonance-stabilized cation.

Steps [6]–[7] Nucleophilic attack and deprotonation







- Nucleophilic attack on the cation in Step [6] followed by loss of a proton forms the acetal.
- **The overall result of Steps [4]–[7] is the addition of a second OR group to the carbonyl group.**


Although this mechanism is lengthy—there are seven steps altogether—there are only three different kinds of reactions: **addition of a nucleophile, elimination of a leaving group, and proton transfer**. Steps [2] and [6] involve nucleophilic attack and Step [5] eliminates H₂O. The other four steps in the mechanism shuffle protons from one oxygen atom to another, to make a better leaving group or a more electrophilic carbonyl group.

Problem 21.33

Label each compound as an acetal, a hemiacetal, or an ether.





Problem 21.34 Draw a stepwise mechanism for the following reaction.




21.14B Hydrolysis of Acetals

Conversion of an aldehyde or ketone to an acetal is a **reversible reaction**, so **an acetal can be hydrolyzed to an aldehyde or ketone by treatment with aqueous acid**. Because this reaction is also an equilibrium process, it is driven to the right by using a large excess of water for hydrolysis.



The mechanism for this reaction is the reverse of acetal synthesis, as illustrated in Sample Problem 21.4.

Sample Problem 21.4 Draw a stepwise mechanism for the following reaction.




Solution

The mechanism is the reverse of acetal formation and involves two parts—conversion of the acetal to a hemiacetal, followed by conversion of the hemiacetal to the carbonyl compound.


Part [1] Conversion of the acetal to a hemiacetal

To convert this acetal to a hemiacetal, one molecule of CH₃OH must be eliminated and one molecule of H₂O must be added.




Part [2] Conversion of the hemiacetal to the carbonyl compound

To convert the hemiacetal to a carbonyl compound, one molecule of CH_3OH must be eliminated and the $\text{C}=\text{O}$ π bond must be formed.




Steps [2] and [6] involve loss of the leaving group (CH_3OH), and Step [3] involves nucleophilic attack of H_2O . The other four steps in the mechanism shuffle protons from one oxygen atom to another. Steps [2] and [6] form resonance-stabilized carbocations, but only one resonance structure is drawn.

Acetal hydrolysis requires a strong acid to make a good leaving group (ROH). In Sample Problem 21.4, H_2SO_4 converts CH_3O^- into CH_3OH , a weak base and neutral leaving group. Acetal hydrolysis does not occur in base.

Problem 21.35 Draw the products of each reaction.**Problem 21.36**


Sassafras, source of safrole

Safrole is a naturally occurring acetal isolated from sassafras plants. Once used as a common food additive in root beer and other beverages, it is now banned because it is carcinogenic. What compounds are formed when safrole is hydrolyzed with aqueous acid?

**21.15 Acetals as Protecting Groups**

Just as the *tert*-butyldimethylsilyl ethers are used as protecting groups for alcohols (Section 20.12), **acetals are valuable protecting groups for aldehydes and ketones.**


Suppose a starting material **A** contains both a ketone and an ester, and it is necessary to selectively reduce the ester to an alcohol (6-hydroxy-2-hexanone), leaving the ketone untouched. Such a selective reduction is *not* possible in one step. Because ketones are more readily reduced, methyl 5-hydroxyhexanoate is formed instead.



To solve this problem we can use a protecting group to block the more reactive ketone carbonyl group. The overall process requires three steps.

- [1] **Protect the interfering functional group—the ketone carbonyl.**
- [2] **Carry out the desired reaction—reduction.**
- [3] **Remove the protecting group.**


The following three-step sequence using a cyclic acetal leads to the desired product.



- **Step [1]** The ketone carbonyl is protected as a cyclic acetal by reaction of the starting material with $\text{HOCH}_2\text{CH}_2\text{OH}$ and TsOH .
- **Step [2]** Reduction of the ester is then carried out with LiAlH_4 , followed by treatment with H_2O .
- **Step [3]** The acetal is then converted back to a ketone carbonyl group with aqueous acid.

Acetals are widely used protecting groups for aldehydes and ketones because they are easy to add and easy to remove, and they are stable to a wide variety of reaction conditions. Acetals do not react with base, oxidizing agents, reducing agents, or nucleophiles. Good protecting groups must survive a variety of reaction conditions that take place at other sites in a molecule, but they must also be selectively removed under mild conditions when needed.


Problem 21.37 How would you use a protecting group to carry out the following transformation?



21.16 Cyclic Hemiacetals


Cyclic hemiacetals are also called **lactols**.

Although acyclic hemiacetals are generally unstable and therefore not present in appreciable amounts at equilibrium, **cyclic hemiacetals containing five- and six-membered rings are stable compounds** that are readily isolated.




21.16A Forming Cyclic Hemiacetals

All hemiacetals are formed by nucleophilic addition of a hydroxy group to a carbonyl group. In the same way, cyclic hemiacetals are formed by **intramolecular cyclization of hydroxy aldehydes**.



Such intramolecular reactions to form five- and six-membered rings are faster than the corresponding intermolecular reactions. The two reacting functional groups, in this case OH and C=O, are held in close proximity, increasing the probability of reaction.


Problem 21.38 What lactol (cyclic hemiacetal) is formed from intramolecular cyclization of each hydroxy aldehyde?



Hemiacetal formation is catalyzed by both acid and base. The acid-catalyzed mechanism is identical to Mechanism 21.9, except that the reaction occurs in an **intramolecular** fashion, as shown for the acid-catalyzed cyclization of 5-hydroxypentanal to form a six-membered cyclic hemiacetal in Mechanism 21.11.




Mechanism 21.11 Acid-Catalyzed Cyclic Hemiacetal Formation




- Protonation of the carbonyl oxygen in Step [1] followed by **intramolecular nucleophilic attack** in Step [2] forms the six-membered ring.
- Deprotonation in Step [3] forms the neutral cyclic hemiacetal.

Intramolecular cyclization of a hydroxy aldehyde forms a **hemiacetal with a new stereogenic center**, so that an equal amount of two enantiomers results.




Re-drawing the starting material and products in a three-dimensional representation results in the following:



21.16B The Conversion of Hemiacetals to Acetals

Cyclic hemiacetals can be converted to acetals by treatment with an alcohol and acid. This reaction converts the OH group that is part of the hemiacetal to an OR group.




Mechanism 21.12 for this reaction is identical to Mechanism 21.10, which illustrates the conversion of an acyclic hemiacetal to an acetal.




Mechanism 21.12 A Cyclic Acetal from a Cyclic Hemiacetal

Steps [1]–[2] Protonation and loss of the leaving group




Steps [3]–[4] Nucleophilic attack and deprotonation




The overall result of this reaction is the **replacement of the hemiacetal OH group by an OCH₃ group**. This substitution reaction readily occurs because the carbocation formed in Step [2] is stabilized by resonance. This fact makes the OH group of a hemiacetal different from the hydroxy group in other alcohols.

Thus, when a compound that contains both an alcohol OH group and a hemiacetal OH group is treated with an alcohol and acid, only the hemiacetal OH group reacts to form an acetal. The alcohol OH group does *not* react.


The conversion of cyclic hemiacetals to acetals is an important reaction in carbohydrate chemistry, as discussed in Chapter 27.



Problem 21.39 Draw the products of each reaction.



Problem 21.40 Two naturally occurring compounds that contain stable cyclic hemiacetals and acetals are monensin and digoxin, the chapter-opening molecule. Monensin, a polyether antibiotic produced by *Streptomyces cinamonensis*, is used as an additive in cattle feed. Digoxin is a widely prescribed cardiac drug used to increase the force of heart contractions. Label each acetal, hemiacetal, and ether in both compounds.




21.17 An Introduction to Carbohydrates


Carbohydrates, commonly referred to as sugars and starches, are polyhydroxy aldehydes and ketones, or compounds that can be hydrolyzed to them. Along with proteins, fatty acids, and nucleotides, they form one of the four main groups of biomolecules responsible for the structure and function of all living cells.

Many carbohydrates contain cyclic acetals or hemiacetals. Examples include **glucose**, the most common simple sugar, and **lactose**, the principal carbohydrate in milk.

Glucose is the carbohydrate that is transported in the blood to individual cells. The hormone insulin regulates the level of glucose in the blood. Diabetes is a common disease that results from a deficiency of insulin, resulting in increased glucose levels in the blood and other metabolic abnormalities. Insulin injections control glucose levels.



Hemiacetals in sugars are formed in the same way that other hemiacetals are formed—that is, by **cyclization of hydroxy aldehydes**. Thus, the hemiacetal of glucose is formed by cyclization of an acyclic *polyhydroxy aldehyde* **A**, as shown in the accompanying equation. This process illustrates two important features.




- When the OH group on C5 is the nucleophile, **cyclization yields a six-membered ring**, and this ring size is preferred.
- **Cyclization forms a new stereogenic center**, exactly analogous to the cyclization of the simpler hydroxy aldehyde (5-hydroxypentanal) in Section 21.16A. **The new OH group of the hemiacetal can occupy either the equatorial or axial position.**

For glucose, this results in two cyclic forms, called **β -D-glucose** (having an equatorial OH group) and **α -D-glucose** (having an axial OH group). Because β -D-glucose has the new OH group in the more roomy equatorial position, this cyclic form of glucose is the major product. At equilibrium, only a trace of the acyclic hydroxy aldehyde **A** is present.

Many more details on this process and other aspects of carbohydrate chemistry are presented in Chapter 27.

Problem 21.41



- How many stereogenic centers are present in α -D-galactose?
- Label the hemiacetal carbon in α -D-galactose.
- Draw the structure of β -D-galactose.
- Draw the structure of the polyhydroxy aldehyde that cyclizes to α - and β -D-galactose.
- From what you learned in Section 21.16B, what product(s) is (are) formed when α -D-galactose is treated with CH_3OH and an acid catalyst?

KEY CONCEPTS

Aldehydes and Ketones—Nucleophilic Addition


General Facts

- Aldehydes and ketones contain a carbonyl group bonded to only H atoms or R groups. The carbonyl carbon is sp^2 hybridized and trigonal planar (21.1).
- Aldehydes are identified by the suffix *-al*, whereas ketones are identified by the suffix *-one* (21.2).
- Aldehydes and ketones are polar compounds that exhibit dipole–dipole interactions (21.3).


Summary of Spectroscopic Absorptions of RCHO and R_2CO (21.4)

IR absorptions	C=O	$\sim 1715\text{ cm}^{-1}$ for ketones (increasing frequency with decreasing ring size) $\sim 1730\text{ cm}^{-1}$ for aldehydes
		• For both RCHO and R_2CO , the frequency decreases with conjugation.
^1H NMR absorptions	$\text{C}_{sp^2}\text{-H}$ of CHO	$\sim 2700\text{--}2830\text{ cm}^{-1}$ (one or two peaks)
	CHO	9–10 ppm (highly deshielded proton)
	C–H α to C=O	2–2.5 ppm (somewhat deshielded $\text{C}_{sp^3}\text{-H}$)
^{13}C NMR absorption	C=O	190–215 ppm


Nucleophilic Addition Reactions

[1] Addition of hydride (H^-) (21.8)

- The mechanism has two steps.
- H^- adds to the planar $\text{C}=\text{O}$ from both sides.


[2] Addition of organometallic reagents (R^-) (21.8)

- The mechanism has two steps.
- $(\text{R}'')^-$ adds to the planar $\text{C}=\text{O}$ from both sides.


[3] Addition of cyanide (CN^-) (21.9)

- The mechanism has two steps.
- CN^- adds to the planar $\text{C}=\text{O}$ from both sides.


[4] Wittig reaction (21.10)




- The reaction forms a new $\text{C}-\text{C}$ σ bond and a new $\text{C}-\text{C}$ π bond.
- $\text{Ph}_3\text{P}=\text{O}$ is formed as by-product.

[5] Addition of 1° amines (21.11)

- The reaction is fastest at pH 4–5.
- The intermediate carbinolamine is unstable, and loses H_2O to form the $\text{C}=\text{N}$.


[6] Addition of 2° amines (21.12)

- The reaction is fastest at pH 4–5.
- The intermediate carbinolamine is unstable, and loses H_2O to form the $\text{C}=\text{C}$.

[7] Addition of H_2O —Hydration (21.13)

- The reaction is reversible. Equilibrium favors the product only with less stable carbonyl compounds (e.g., H_2CO and Cl_3CCHO).
- The reaction is catalyzed by either H^+ or ^-OH .


[8] Addition of alcohols (21.14)



- The reaction is reversible.
- The reaction is catalyzed by acid.
- Removal of H_2O drives the equilibrium to favor the products.


Other Reactions

[1] Synthesis of Wittig reagents (21.10A)




- Step [1] is best with CH_3X and RCH_2X because the reaction follows an $\text{S}_{\text{N}}2$ mechanism.
- A strong base is needed for proton removal in Step [2].

[2] Conversion of cyanohydrins to aldehydes and ketones (21.9)




- This reaction is the reverse of cyanohydrin formation.


[3] Hydrolysis of nitriles (21.9)



[4] Hydrolysis of imines and enamines (21.12)



[5] Hydrolysis of acetals (21.14)














- The reaction is acid catalyzed and is the reverse of acetal synthesis.
- A large excess of H_2O drives the equilibrium to favor the products.

PROBLEMS

Nomenclature

21.42 Give the IUPAC name for each compound.




21.43 Give the structure corresponding to each name.

- | | | |
|--|---------------------------------------|---|
| a. 2-methyl-3-phenylbutanal | e. 3-benzoylcyclopentanone | i. 2-sec-butyl-3-cyclopentenone |
| b. dipropyl ketone | f. 2-formylcyclopentanone | j. 5,6-dimethyl-1-cyclohexanecarbaldehyde |
| c. 3,3-dimethylcyclohexanecarbaldehyde | g. (3 <i>R</i>)-3-methyl-2-heptanone | |
| d. α -methoxypropionaldehyde | h. <i>m</i> -acetylbenzaldehyde | |

21.44 Including stereoisomers, draw the 11 aldehydes with molecular formula $C_6H_{12}O$, and give the IUPAC name (with any needed *R,S* designation) for each compound.


Reactions

21.45 Draw the product formed when phenylacetaldehyde ($C_6H_5CH_2CHO$) is treated with each reagent. Phenylacetaldehyde is partly responsible for the fragrance of the flowers of the plumeria tree, which is native to the tropical and subtropical Americas.


- | | | |
|--------------------------------|---------------------------------|--|
| a. $NaBH_4, CH_3OH$ | e. $Ph_3P=CHCH_3$ | i.  NH, mild acid |
| b. [1] $LiAlH_4$; [2] H_2O | f. $(CH_3)_2CHNH_2$, mild acid | j. $HOCH_2CH_2OH, H^+$ |
| c. [1] CH_3MgBr ; [2] H_2O | g. $(CH_3CH_2)_2NH$, mild acid | |
| d. $NaCN, HCl$ | h. CH_3CH_2OH (excess), H^+ | |

21.46 Answer Problem 21.45 using 2-heptanone ($CH_3COCH_2CH_2CH_2CH_2CH_3$) as starting material. 2-Heptanone is partly responsible for the odor of bleu cheese.

21.47 Draw the products formed in each Wittig reaction. Draw all stereoisomers formed when a mixture of products results.




21.48 Draw the products formed in each reaction sequence.




21.49 What alkyl halide is needed to prepare each Wittig reagent?

- a. $Ph_3P=CHCH_2CH_2CH_3$ b. $Ph_3P=C(CH_2CH_2CH_3)_2$ c. $Ph_3P=CHCH=CH_2$


21.50 Fill in the lettered reagents (A–G) in the following reaction scheme.




21.51 Draw the products of each reaction.




21.52 What carbonyl compound and alcohol are formed by hydrolysis of each acetal?



21.53 Identify the lettered intermediates in the following reaction sequence. When a mixture of ortho and para products results in electrophilic aromatic substitution, consider the para product only. The ^1H NMR spectrum of **G** shows two singlets at 2.6 and 8.18 ppm.




21.54 Draw all stereoisomers formed in each reaction.



21.55 Hydroxy aldehydes **A** and **B** readily cyclize to form hemiacetals. Draw the stereoisomers formed in this reaction from both **A** and **B**. Explain why this process gives an optically inactive product mixture from **A**, and an optically active product mixture from **B**.




21.56 Etoposide, sold under the trade name of Etopophos, is used for the treatment of lung cancer, testicular cancer, and lymphomas. (a) Locate the acetals in etoposide. (b) What products are formed when all of the acetals are hydrolyzed with aqueous acid?



Properties of Aldehydes and Ketones

21.57 Rank the compounds in each group in order of increasing reactivity in nucleophilic addition.




21.58 Explain why a *gem*-diol is the major species present at equilibrium when cyclopropanone is dissolved in H_2O .


- 21.59** Consider the para-substituted aromatic ketones, $\text{NO}_2\text{C}_6\text{H}_4\text{COCH}_3$ (*p*-nitroacetophenone) and $\text{CH}_3\text{OC}_6\text{H}_4\text{COCH}_3$ (*p*-methoxyacetophenone).
- Which carbonyl compound is more stable?
 - Which compound forms the higher percentage of hydrate at equilibrium?
 - Which compound exhibits a carbonyl absorption at higher wavenumber in its IR spectrum? Explain your reasoning in each part.

Synthesis

- 21.60** What Wittig reagent and carbonyl compound are needed to prepare each alkene? When two routes are possible, indicate which route, if any, is preferred.




- 21.61** What carbonyl compound and amine or alcohol are needed to prepare each product?




- 21.62** What reagents are needed to convert each compound to benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$)? More than one step may be required.

- $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$
- $\text{C}_6\text{H}_5\text{COCl}$
- $\text{C}_6\text{H}_5\text{COOCH}_3$
- $\text{C}_6\text{H}_5\text{COOH}$
- $\text{C}_6\text{H}_5\text{CH}_3$
- $\text{C}_6\text{H}_5\text{CH}=\text{CH}_2$
- $\text{C}_6\text{H}_5\text{CH}=\text{NCH}_2\text{CH}_2\text{CH}_3$
- $\text{C}_6\text{H}_5\text{CH}(\text{OCH}_2\text{CH}_3)_2$


- 21.63** What reagents are needed to convert each compound to 2-butanone ($\text{CH}_3\text{COCH}_2\text{CH}_3$)?




- 21.64** Show two different methods to carry out each transformation: a one-step method using a Wittig reagent, and a two-step method using a Grignard reagent. Which route, if any, is preferred for each compound?




- 21.65** Devise a synthesis of each alkene using a Wittig reaction to form the double bond. You may use benzene and organic alcohols having four or fewer carbons as starting materials and any required reagents.




- 21.66** Devise a synthesis of each compound from cyclohexene and organic alcohols. You may use any other required organic or inorganic reagents.




- 21.67** Devise a synthesis of each compound from the given starting materials. You may also use organic alcohols having four or fewer carbons, and any organic or inorganic reagents.




21.68 Devise a synthesis of each compound from 5-hydroxy-2-pentanone ($\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_2\text{OH}$) as starting material. You may also use alcohols having three or fewer carbons and any required organic or inorganic reagents.



21.69 Devise a synthesis of each compound from ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) as the only source of carbon atoms. You may use any other organic or inorganic reagents you choose.




21.70 Albuterol is a bronchodilator used to treat the symptoms of asthma. Devise a synthesis of albuterol from **X** using any required organic compounds or inorganic reagents.




Protecting Groups

21.71 Design a stepwise synthesis to convert cyclopentanone and 4-bromobutanal to hydroxy aldehyde **A**.




21.72 Besides the *tert*-butyldimethylsilyl ethers introduced in Chapter 20, there are many other widely used alcohol protecting groups. For example, an alcohol such as cyclohexanol can be converted to a methoxy methyl ether (a MOM protecting group) by treatment with base and chloromethyl methyl ether, $\text{ClCH}_2\text{OCH}_3$. The protecting group can be removed by treatment with aqueous acid.




- Write a stepwise mechanism for the formation of a MOM ether from cyclohexanol.
- What functional group comprises a MOM ether?
- Besides cyclohexanol, what other products are formed by aqueous hydrolysis of the MOM ether? Draw a stepwise mechanism that accounts for formation of each product.

Mechanism


21.73 Draw a stepwise mechanism for each reaction.




- 21.74** Treatment of $(\text{HOCH}_2\text{CH}_2\text{CH}_2\text{CH}_2)_2\text{CO}$ with acid forms a product of molecular formula $\text{C}_9\text{H}_{16}\text{O}_2$ and a molecule of water. Draw the structure of the product and explain how it is formed.
- 21.75** When acetone is dissolved in aqueous acid containing isotopically labeled H_2O (H_2^{18}O), the carbonyl group becomes labeled with ^{18}O . Draw a mechanism that explains this observation.




- 21.76** Draw a stepwise mechanism for each reaction.




- 21.77** Another way to synthesize cyclic acetals uses enol ethers (not carbonyl compounds) as starting materials. Draw a stepwise mechanism for the following synthesis of an acetal from an enol ether and ethylene glycol.




- 21.78** Salsolinol is a naturally occurring compound found in bananas, chocolate, and several foods derived from plant sources. Salsolinol is also formed in the body when acetaldehyde, an oxidation product of the ethanol ingested in an alcoholic beverage, reacts with dopamine, a neurotransmitter. Draw a stepwise mechanism for the formation of salsolinol in the following reaction.




- 21.79** Sulfur ylides, like Wittig reagents, are useful intermediates in organic synthesis. Sulfur ylides are formed by the treatment of sulfonium salts with butyllithium. They react with carbonyl compounds to form epoxides. Draw the mechanism for formation of epoxide **X** from cyclohexanone using a sulfur ylide.



- 21.80** (a) Explain how NaBH_4 in CH_3OH can reduce hemiacetal **A** to 1,4-butanediol ($\text{HOCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$). (b) What product is formed when **A** is treated with $\text{Ph}_3\text{P}=\text{CHCH}_2\text{CH}(\text{CH}_3)_2$?



- 21.81** Reaction of 5,5-dimethoxy-2-pentanone with methylmagnesium iodide followed by treatment with aqueous acid forms cyclic hemiacetal **Y**. Draw a stepwise mechanism that illustrates how **Y** is formed.




Spectroscopy

21.82 Although the carbonyl absorption of cyclic ketones generally shifts to higher wavenumber with decreasing ring size, the C=O of cyclopropanone absorbs at lower wavenumber in its IR spectrum than the C=O of cyclohexanone. Explain this observation by using the principles of aromaticity learned in Chapter 17.



cyclopropanone
(1640 cm^{-1})




2-cyclohexenone
(1685 cm^{-1})

21.83 How would the compounds in each pair differ in their IR spectra: (a) $\text{CH}_3(\text{CH}_2)_4\text{CHO}$ and $\text{CH}_3(\text{CH}_2)_3\text{COCH}_3$; (b) $\text{C}_6\text{H}_5\text{CH}_2\text{COCH}_3$ and $\text{C}_6\text{H}_5\text{COCH}_2\text{CH}_3$; (c) cyclohexanone and 2-methylcyclopentanone?





21.84 Use the ^1H NMR and IR data to determine the structure of each compound.

Compound A	Molecular formula:	$\text{C}_5\text{H}_{10}\text{O}$
	IR absorptions at	1728, 2791, and 2700 cm^{-1}
	^1H NMR data:	1.08 (singlet, 9 H) and 9.48 (singlet, 1 H) ppm
Compound B	Molecular formula:	$\text{C}_5\text{H}_{10}\text{O}$
	IR absorption at	1718 cm^{-1}
	^1H NMR data:	1.10 (doublet, 6 H), 2.14 (singlet, 3 H), and 2.58 (septet, 1 H) ppm
Compound C	Molecular formula:	$\text{C}_{10}\text{H}_{12}\text{O}$
	IR absorption at	1686 cm^{-1}
	^1H NMR data:	1.21 (triplet, 3 H), 2.39 (singlet, 3 H), 2.95 (quartet, 2 H), 7.24 (doublet, 2 H), and 7.85 (doublet, 2 H) ppm
Compound D	Molecular formula:	$\text{C}_{10}\text{H}_{12}\text{O}$
	IR absorption at	1719 cm^{-1}
	^1H NMR data:	1.02 (triplet, 3 H), 2.45 (quartet, 2 H), 3.67 (singlet, 2 H), and 7.06–7.48 (multiplet, 5 H) ppm


21.85 A solution of acetone [$(\text{CH}_3)_2\text{C}=\text{O}$] in ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) in the presence of a trace of acid was allowed to stand for several days, and a new compound of molecular formula $\text{C}_7\text{H}_{16}\text{O}_2$ was formed. The IR spectrum showed only one major peak in the functional group region around 3000 cm^{-1} , and the ^1H NMR spectrum is given here. What is the structure of the product?




21.86 Compounds **A** and **B** have molecular formula $C_9H_{10}O$. Identify their structures from the 1H NMR and IR spectra given.



21.87 An unknown compound **C** of molecular formula $C_6H_{12}O_3$ exhibits a strong absorption in its IR spectrum at 1718 cm^{-1} and the given 1H NMR spectrum. What is the structure of **C**?




- 21.88** An unknown compound **D** exhibits a strong absorption in its IR spectrum at 1692 cm^{-1} . The mass spectrum of **D** shows a molecular ion at $m/z = 150$ and a base peak at 121. The $^1\text{H NMR}$ spectrum of **D** is shown below. What is the structure of **D**?




Carbohydrates

- 21.89** Draw the structure of the acyclic polyhydroxy aldehyde that cyclizes to each hemiacetal.




- 21.90** β -D-Glucose, a hemiacetal, can be converted to a mixture of acetals on treatment with CH_3OH in the presence of acid. Draw a stepwise mechanism for this reaction. Explain why two acetals are formed from a single starting material.




Challenge Problems

- 21.91** Draw a stepwise mechanism for the following reaction.




- 21.92** Brevicommin, the aggregation pheromone of the western pine bark beetle, contains a bicyclic bridged ring system and is prepared by the acid-catalyzed cyclization of 6,7-dihydroxy-2-nonanone.

- a. Suggest a structure for brevicommin.
 b. Devise a synthesis of 6,7-dihydroxy-2-nonanone from 6-bromo-2-hexanone. You may also use three-carbon alcohols and any required organic or inorganic reagents.



21.93 Maltose is a carbohydrate present in malt, the liquid obtained from barley and other grains. Although maltose has numerous functional groups, its reactions are explained by the same principles we have already encountered.



- Label the acetal and hemiacetal carbons.
- What products are formed when maltose is treated with each of the following reagents: [1] H_3O^+ ; [2] CH_3OH and HCl ; [3] excess NaH , then excess CH_3I ?
- Draw the products formed when the compound formed in Reaction [3] of part (b) is treated with aqueous acid.

The reactions in parts (b) and (c) are used to determine structural features of carbohydrates like maltose. We will learn much more about maltose and similar carbohydrates in Chapter 27.

Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

22

- 22.1 Introduction
- 22.2 Structure and bonding
- 22.3 Nomenclature
- 22.4 Physical properties
- 22.5 Spectroscopic properties
- 22.6 Interesting esters and amides
- 22.7 Introduction to nucleophilic acyl substitution
- 22.8 Reactions of acid chlorides
- 22.9 Reactions of anhydrides
- 22.10 Reactions of carboxylic acids
- 22.11 Reactions of esters
- 22.12 Application: Lipid hydrolysis
- 22.13 Reactions of amides
- 22.14 Application: The mechanism of action of β -lactam antibiotics
- 22.15 Summary of nucleophilic acyl substitution reactions
- 22.16 Natural and synthetic fibers
- 22.17 Biological acylation reactions
- 22.18 Nitriles





Nylon 6,6, a polymer composed of many amide bonds, was the first synthetic fiber and one of the most lucrative products ever invented at DuPont. Because nylon is strong and durable, and can be spun into a fiber that resembles the silk produced by silkworms, nylon found immediate use in making parachutes, clothing, stockings, and rope. Nylon 6,6 is a polyamide that can be prepared by joining a diacid chloride and a diamine together. In Chapter 22, we learn about amides and other acyl derivatives of carboxylic acids.


Chapter 22 continues the study of carbonyl compounds with a detailed look at **nucleophilic acyl substitution**, a key reaction of carboxylic acids and their derivatives. Substitution at sp^2 hybridized carbon atoms was introduced in Chapter 20 with reactions involving carbon and hydrogen nucleophiles. In Chapter 22, we learn that nucleophilic acyl substitution is a general reaction that occurs with a variety of heteroatomic nucleophiles. This reaction allows the conversion of one carboxylic acid derivative into another. **Every reaction in Chapter 22 that begins with a carbonyl compound involves nucleophilic substitution.** Chapter 22 also discusses the properties and chemical reactions of **nitriles**, compounds that contain a carbon–nitrogen triple bond. Nitriles are in the same carbon oxidation state as carboxylic acids, and they undergo reactions that form related products.

22.1 Introduction


Chapter 22 focuses on carbonyl compounds that contain an **acyl group bonded to an electronegative atom**. These include the **carboxylic acids**, as well as carboxylic acid derivatives that can be prepared from them: **acid chlorides, anhydrides, esters, and amides**.




Anhydrides contain two carbonyl groups joined by a single oxygen atom. **Symmetrical anhydrides** have two identical alkyl groups bonded to the carbonyl carbons, and **mixed anhydrides** have two different alkyl groups. **Cyclic anhydrides** are also known.



Amides are classified as 1° , 2° , or 3° depending on the number of carbon atoms bonded directly to the *nitrogen* atom.




Cyclic esters and amides are called **lactones** and **lactams**, respectively. The ring size of the heterocycle is indicated by a Greek letter. An amide in a four-membered ring is called a **β -lactam**, because the β carbon to the carbonyl is bonded to the heteroatom. An ester in a five-membered ring is called a **γ -lactone**.




Nucleophilic acyl substitution was first discussed in Chapter 20 with R^- and H^- as the nucleophiles. This substitution reaction is general for a variety of nucleophiles, making it possible to form many different substitution products, as discussed in Sections 22.8–22.13.

All of these compounds contain an acyl group bonded to an electronegative atom Z that can serve as a **leaving group**. As a result, these compounds undergo **nucleophilic acyl substitution**. Recall from Chapters 20 and 21 that aldehydes and ketones do *not* undergo nucleophilic substitution because they have no leaving group on the carbonyl carbon.



Nitriles are compounds that contain a **cyano group**, $C\equiv N$, bonded to an **alkyl group**. Nitriles have no carbonyl group, so they are structurally distinct from carboxylic acids and their derivatives. The carbon atom of the cyano group, however, has the same oxidation state as the carbonyl carbon of carboxylic acid derivatives, so there are certain parallels in their chemistry.



22.3 Nomenclature


The names of carboxylic acid derivatives are formed from the names of the parent carboxylic acids discussed in Section 19.2. Keep in mind that the common names **formic acid**, **acetic acid**, and **benzoic acid** are virtually always used for the parent acid, so these common parent names are used for their derivatives as well.

22.3A Naming an Acid Chloride—RCOCl


Acid chlorides are named by naming the acyl group and adding the word **chloride**. Two different methods are used.

- [1] For acyclic acid chlorides: Change the suffix **-ic acid** of the parent carboxylic acid to the suffix **-yl chloride**; or
- [2] When the -COCl group is bonded to a ring: Change the suffix **-carboxylic acid** to **-carbonyl chloride**.


Naming acid chlorides



acetyl chloride




cyclohexanecarbonyl chloride



2-methylbutanoyl chloride


22.3B Naming an Anhydride

The word *anhydride* means “without water.” Removing one molecule of water from two molecules of carboxylic acid forms an anhydride.




Symmetrical anhydrides are named by changing the **acid** ending of the parent carboxylic acid to the word **anhydride**. **Mixed anhydrides**, which are derived from two different carboxylic acids, are named by alphabetizing the names for both acids and replacing the word **acid** by the word **anhydride**.

Symmetrical anhydride



Mixed anhydride



22.3C Naming an Ester—RCOOR'


Esters are often written as RCOOR' , where the alkyl group (R') is written *last*. When an ester is named, however, the R' group appears *first* in the name.

An ester has two parts to its structure, each of which must be named: an **acyl group** (RCO-) and an **alkyl group** (designated as R') bonded to an oxygen atom.

- In the IUPAC system, esters are identified by the suffix **-ate**.


HOW TO Name an Ester (RCO₂R') Using the IUPAC System

Example Give a systematic name for each ester:




Step [1] Name the R' group bonded to the oxygen atom as an alkyl group.

- The name of the alkyl group, ending in the suffix *-yl*, becomes the *first* part of the ester name.



Step [2] Name the acyl group (RCO-) by changing the *-ic acid* ending of the parent carboxylic acid to the suffix *-ate*.

- The name of the acyl group becomes the *second* part of the name.




Answer: ethyl acetate

Answer: *tert*-butyl cyclohexanecarboxylate

22.3D Naming an Amide

All 1° amides are named by replacing the *-ic acid*, *-oic acid*, or *-ylic acid* ending with the suffix **amide**.


Naming 1° amides



A 2° or 3° amide has two parts to its structure: an **acyl group** that contains the carbonyl group (RCO-) and one or two **alkyl groups** bonded to the nitrogen atom.

HOW TO Name a 2° or 3° Amide

Example Give a systematic name for each amide:




—Continued


HOW TO, continued . . .

Step [1] Name the alkyl group (or groups) bonded to the N atom of the amide. Use the prefix “*N*-” preceding the name of each alkyl group.

- The names of the alkyl groups form the first part of each amide name.
- For 3° amides, use the prefix **di-** if the two alkyl groups on N are the same. If the two alkyl groups are different, **alphabetize** their names. One “*N*-” is needed for each alkyl group, even if both R groups are identical.




- The compound is a 2° amide with one ethyl group → *N*-ethyl.




- The compound is a 3° amide with two methyl groups.
- Use the prefix **di-** and two “*N*-” to begin the name → *N,N*-dimethyl.

Step [2] Name the acyl group (RCO–) with the suffix *-amide*.



derived from
formic acid -----> formamide

- Change the *-ic acid* or *-oic acid* suffix of the parent carboxylic acid to the suffix *-amide*.
- Put the two parts of the name together.
- **Answer: *N*-ethylformamide**



derived from
benzoic acid -----> benzamide

- Change *benzoic acid* to *benzamide*.
- Put the two parts of the name together.
- **Answer: *N,N*-dimethylbenzamide**

22.3E Naming a Nitrile

In contrast to the carboxylic acid derivatives, **nitriles are named as alkane derivatives**. To name a nitrile using IUPAC rules:

- Find the longest chain that contains the CN and add the word **nitrile** to the name of the parent alkane. Number the chain to put CN at C1, but omit this number from the name.

In naming a nitrile, the CN carbon is one carbon atom of the longest chain. **CH₃CH₂CN is propanenitrile, not ethanenitrile.**

Common names for nitriles are derived from the names of the carboxylic acid having the same number of carbon atoms by replacing the *-ic acid* ending of the carboxylic acid by the suffix ***-onitrile***.


When CN is named as a substituent, it is called a ***cyano*** group. Figure 22.1 illustrates features of nitrile nomenclature.

Table 22.2 summarizes the most important points about the nomenclature of carboxylic acid derivatives.


Figure 22.1

Summary of nitrile nomenclature

a. IUPAC name for a nitrile



b. Common name for a nitrile



c. CN as a substituent









Table 22.2 Summary: Nomenclature of Carboxylic Acid Derivatives and Nitriles


Compound	Name ending	Example	Name
acid chloride	-yl chloride or -carbonyl chloride		benzoyl chloride
anhydride	anhydride		benzoic anhydride
ester	-ate		ethyl benzoate
amide	-amide		N-methylbenzamide
nitrile	-nitrile or -onitrile	$C_6H_5-C\equiv N$	benzonitrile

Sample Problem 22.1


Give the IUPAC name for each compound.

**Solution**a. The functional group is an acid chloride bonded to a chain of atoms, so the name ends in **-yl chloride**.


[1] Find and name the longest chain containing the COCl:

**hexanoic acid** → **hexanoyl chloride**
(6 C's)


[2] Number and name the substituents:

**Answer:**
2,4-dimethylhexanoyl chlorideb. The functional group is an ester, so the name ends in **-ate**.


[1] Find and name the longest chain containing the carbonyl group:

**pentanoic acid** → **pentanoate**
(5 C's)

[2] Number and name the substituents:

**Answer: isopropyl 2-methylpentanoate**The name of the alkyl group on the O atom goes **first** in the name.**Problem 22.4**

Give an IUPAC or common name for each compound.



Problem 22.5 Draw the structure corresponding to each name.

- | | |
|---|--|
| a. 5-methylheptanoyl chloride | e. 3-methylpentanenitrile |
| b. isopropyl propanoate | f. <i>o</i> -cyanobenzoic acid |
| c. acetic formic anhydride | g. <i>sec</i> -butyl 2-methylhexanoate |
| d. <i>N</i> -isobutyl- <i>N</i> -methylbutanamide | h. <i>N</i> -ethylhexanamide |

22.4 Physical Properties

Because all carbonyl compounds have a polar carbonyl group, they exhibit **dipole–dipole interactions**. Nitriles also have dipole–dipole interactions because they have a polar $C\equiv N$ group. Because they contain one or two $N-H$ bonds, 1° and 2° amides are capable of intermolecular hydrogen bonding. The $N-H$ bond of one amide intermolecularly hydrogen bonds to the $C=O$ of another amide, as shown using two acetamide molecules (CH_3CONH_2) in Figure 22.2.

How these factors affect the physical properties of carboxylic acid derivatives is summarized in Table 22.3.

Problem 22.6 Explain why the boiling point of CH_3CONH_2 ($221^\circ C$) is significantly higher than the boiling point of $CH_3CON(CH_3)_2$ ($166^\circ C$), even though the latter compound has a higher molecular weight and more surface area.


Table 22.3 Physical Properties of Carboxylic Acid Derivatives

Property	Observation
Boiling point and melting point	<ul style="list-style-type: none"> Primary (1°) and 2° amides have <i>higher</i> boiling points and melting points than compounds of comparable molecular weight. The boiling points and melting points of other carboxylic acid derivatives are similar to those of other polar compounds of comparable size and shape. <div style="text-align: center; margin-top: 10px;"> <p style="text-align: center;"> CH_3COCl CH_3COOCH_3 $CH_3COCH_2CH_3$ $CH_3CH_2CONH_2$ MW = 78.5 MW = 74 MW = 72 MW = 73 bp 52 °C bp 58 °C bp 80 °C bp 213 °C ~ ~ < </p> <p style="text-align: center;"> similar boiling points higher boiling point 1° amide </p> </div>
Solubility	<ul style="list-style-type: none"> Carboxylic acid derivatives are soluble in organic solvents regardless of size. Most carboxylic acid derivatives having ≤ 5 C's are H_2O soluble because they can hydrogen bond with H_2O (Section 3.4C). Carboxylic acid derivatives having > 5 C's are H_2O insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent.

Key: MW = molecular weight

Figure 22.2

Intermolecular hydrogen bonding between two CH_3CONH_2 molecules



22.5 Spectroscopic Properties

22.5A IR Spectra

The most prominent IR absorptions for carboxylic acid derivatives and nitriles are as follows:

- [1] Like all carbonyl compounds, carboxylic acid derivatives have a **strong C=O absorption between 1600 and 1850 cm⁻¹**.
- [2] **Primary (1°) and 2° amides** have two additional absorptions due to the N–H bonds:
 - one or two N–H stretching peaks at **3200–3400 cm⁻¹**.
 - an N–H bending absorption at **~1640 cm⁻¹**.
- [3] **Nitriles** have an absorption at **2250 cm⁻¹ for the C≡N**.

The exact location of the carbonyl absorption varies with the identity of Z in the carbonyl compound RCOZ. As detailed in Section 22.2, as the basicity of Z increases, resonance stabilization of RCOZ increases, resulting in the following trend:

- As the carbonyl π bond becomes more delocalized, the C=O absorption shifts to lower frequency.

Thus, the carbonyl group of an acid chloride and anhydride, which are least stabilized by resonance, absorb at higher frequency than the carbonyl group of an amide, which is more stabilized by resonance. Table 22.4 lists specific values for the carbonyl absorptions of the carboxylic acid derivatives.

Conjugation and ring size affect the location of these carbonyl absorptions.

- Conjugation shifts a carbonyl absorption to lower frequencies.
- For cyclic carboxylic acid derivatives, decreasing ring size shifts a carbonyl absorption to higher frequencies.

The effects of conjugation and ring size on the location of a carbonyl absorption were first discussed in Section 21.4A.

Problem 22.7

How would the compounds in each pair differ in their IR spectra?








Table 22.4 IR Absorptions for the Carbonyl Group of Carboxylic Acid Derivatives

Compound type	Structure (RCOZ)	Carbonyl absorption ($\tilde{\nu}$)
acid chloride		~1800
anhydride		1820 and 1760 (2 peaks)
ester		1735–1745
amide	 R' = H or alkyl	1630–1680

Increasing basicity of Z
Increasing $\tilde{\nu}$ of absorption

22.5B NMR Spectra

Carboxylic acid derivatives have two characteristic ^1H NMR absorptions.

- [1] Protons on the α carbon to the carbonyl absorb at 2–2.5 ppm.
- [2] The N–H protons of 1° and 2° amides absorb at 7.5–8.5 ppm.

In their ^{13}C NMR spectra, carboxylic acid derivatives give a highly deshielded peak at 160–180 ppm due to the carbonyl carbon. This is somewhat upfield from the carbonyl absorption of aldehydes and ketones, which occurs at 190–215 ppm.

Nitriles give a peak at 115–120 ppm in their ^{13}C NMR spectrum due to the sp hybridized carbon. This is farther downfield than the signal due to the sp hybridized carbon of an alkyne, which occurs at 65–100 ppm.

Problem 22.8

Deduce the structures of compounds **A** and **B**, two of the major components of jasmine oil, from the given data.

Compound **A**: $\text{C}_9\text{H}_{10}\text{O}_2$; IR absorptions at 3091–2895 and 1743 cm^{-1} ; ^1H NMR signals at 2.06 (singlet, 3 H), 5.08 (singlet, 2 H), and 7.33 (broad singlet, 5 H) ppm.

Compound **B**: $\text{C}_{14}\text{H}_{12}\text{O}_2$; IR absorptions at 3091–2953 and 1718 cm^{-1} ; ^1H NMR signals at 5.35 (singlet, 2 H) and 7.26–8.15 (multiplets, 10 H) ppm.



22.6 Interesting Esters and Amides

22.6A Esters





The characteristic odor of many fruits is due to low molecular weight esters.



Many low molecular weight esters have pleasant and very characteristic odors.



isoamyl acetate
odor of banana




ethyl butanoate
odor of mango




methyl 2-methylbutanoate
odor of pineapple


Several esters have important biological activities.




- **Vitamin C** (or **ascorbic acid**) is a water-soluble vitamin containing a five-membered lactone that we first discussed in Section 3.5B. Although vitamin C is synthesized in plants, humans do not have the necessary enzymes to make it, and so they must obtain it from their diet.



The coca plant, *Erythroxylon coca*, is the source of the addictive drug cocaine.



- **Cocaine** is an addictive stimulant obtained from the leaves of the coca plant. Chewing coca leaves for pleasure has been practiced by the indigenous peoples of South America for over a thousand years, and coca leaves were a very minor ingredient in Coca-Cola for the first 20 years of its production. Cocaine is a widely abused recreational drug, and the possession and use of cocaine is currently illegal in most countries.




- **FK506** is an immunosuppressant known by the trade name Prograf. One functional group in its complex structure is a cyclic ester (shown in red) in a 21-membered ring. A cyclic ester contained in such a large ring is called a **macrocyclic lactone** or a **macrolide**. FK506 is used to suppress rejection after organ transplants.

22.6B Amides

An important group of naturally occurring amides consists of **proteins, polymers of amino acids joined together by amide linkages**. Proteins differ in the length of the polymer chain, as well as in the identity of the R groups bonded to it. The word *protein* is usually reserved for high molecular weight polymers composed of 40 or more amino acid units, while the designation *peptide* is given to polymers of lower molecular weight.

Portion of a protein molecule




[Amide bonds are shown in red.]


The R groups come from amino acids joined together to form the protein.

Peptides and proteins are discussed in detail in Chapter 28.

Proteins and peptides have diverse functions in the cell. They form the structural components of muscle, connective tissue, hair, and nails. They catalyze reactions and transport ions and molecules across cell membranes. **Met-enkephalin**, for example, a peptide with four amide bonds found predominately in nerve tissue cells, relieves pain and acts as an opiate by producing morphine-like effects.




Several useful drugs are amides. For example, **Gleevec** (generic name: imatinib mesylate), an amide sold as a salt with methanesulfonic acid ($\text{CH}_3\text{SO}_3\text{H}$), is an anticancer drug used for the treatment of chronic myeloid leukemia as well as certain gastrointestinal tumors. Gleevec represents a new approach to cancer chemotherapy, which targets a single molecule to disable the molecular mechanism responsible for a specific type of cancer.



Trade name: Gleevec
Generic name: imatinib mesylate


Penicillins are a group of structurally related antibiotics, known since the pioneering work of Sir Alexander Fleming led to the discovery of penicillin G in the 1920s. All penicillins contain a strained β -lactam fused to a five-membered ring, as well as a second amide located α to the β -lactam carbonyl group. Particular penicillins differ in the identity of the R group in the amide side chain.

General structure—Penicillin



Cephalosporins represent a second group of β -lactam antibiotics that contain a four-membered ring fused to a six-membered ring. Cephalosporins are generally active against a broader range of bacteria than penicillins.


General structure—Cephalosporin



Problem 22.9 For both amoxicillin and cephalixin: (a) How many stereogenic centers does each compound contain? (b) What is the maximum number of stereoisomers possible? (c) Draw the enantiomer of each compound.

22.7 Introduction to Nucleophilic Acyl Substitution

The characteristic reaction of carboxylic acid derivatives is *nucleophilic acyl substitution*. This is a general reaction that occurs with both negatively charged nucleophiles (Nu^-) and neutral nucleophiles (HNu):



- Carboxylic acid derivatives (RCOZ) react with nucleophiles because they contain an electrophilic, unhindered carbonyl carbon.
- Substitution occurs, *not* addition, because carboxylic acid derivatives (RCOZ) have a leaving group Z on the carbonyl carbon.


22.7A The Mechanism

The general mechanism for nucleophilic acyl substitution is a two-step process: **nucleophilic attack** followed by **loss of the leaving group**, as shown in Mechanism 22.1.

The mechanism for nucleophilic acyl substitution was first presented in Section 20.2.




Mechanism 22.1 General Mechanism—Nucleophilic Acyl Substitution




[Z = Cl, OCOR, OH, OR', NR'₂]

- In Step [1], **the nucleophile attacks the carbonyl group**, cleaving the π bond, and forming a tetrahedral intermediate with a new C–Nu bond.
- In Step [2], **elimination of the leaving group** forms the substitution product.

The overall result of addition of a nucleophile and elimination of a leaving group is substitution of the nucleophile for the leaving group. Recall from Chapter 20 that nucleophilic substitution occurs with carbanions (R^-) and hydride (H^-) as nucleophiles. A variety of oxygen and nitrogen nucleophiles also participate in this reaction.




Nucleophilic acyl substitution using heteroatomic nucleophiles results in the conversion of one carboxylic acid derivative into another, as shown in two examples.



Each reaction results in the replacement of the leaving group by the nucleophile, regardless of the identity of or charge on the nucleophile. To draw any nucleophilic acyl substitution product:

- **Find the sp^2 hybridized carbon with the leaving group.**
- **Identify the nucleophile.**
- **Substitute the nucleophile for the leaving group.** With a neutral nucleophile a proton must be lost to obtain a neutral substitution product.

Problem 22.10 Draw the products of each reaction.



22.7B Relative Reactivity of Carboxylic Acids and Their Derivatives


As discussed in Section 20.2B, carboxylic acids and their derivatives differ greatly in reactivity toward nucleophiles. The order of reactivity parallels the leaving group ability of the group Z.

- **The better the leaving group, the more reactive RCOZ is in nucleophilic acyl substitution.**

Recall that the **best leaving group is the weakest base**.


The relative basicity of the common leaving groups, Z, is given in Table 22.1.

Thus, the following trends result:



Based on this order of reactivity, **more reactive acyl compounds (acid chlorides and anhydrides) can be converted to less reactive ones (carboxylic acids, esters, and amides). The reverse is not usually true.**


To see why this is so, recall that nucleophilic addition to a carbonyl group forms a tetrahedral intermediate with two possible leaving groups, Z^- or :Nu^- . The group that is subsequently eliminated is the *better* of the two leaving groups. For a reaction to form a substitution product, therefore, Z^- must be the better leaving group, making the starting material RCOZ a more reactive acyl compound.



To evaluate whether a nucleophilic substitution reaction will occur, **compare the leaving group ability of the incoming nucleophile and the departing leaving group**, as shown in Sample Problem 22.2.

Sample Problem 22.2

Determine whether each nucleophilic acyl substitution is likely to occur.



Solution

- a. Conversion of CH_3COCl to $\text{CH}_3\text{COOCH}_2\text{CH}_3$ requires the substitution of Cl^- by $\text{OCH}_2\text{CH}_3^-$. Because Cl^- is a weaker base and therefore a better leaving group than $\text{OCH}_2\text{CH}_3^-$, **this reaction occurs.**
- b. Conversion of $\text{C}_6\text{H}_5\text{CONH}_2$ to $(\text{C}_6\text{H}_5\text{CO})_2\text{O}$ requires the substitution of NH_2^- by $\text{OCOC}_6\text{H}_5^-$. Because NH_2^- is a stronger base and therefore a poorer leaving group than $\text{OCOC}_6\text{H}_5^-$, **this reaction does not occur.**

Problem 22.11

Without reading ahead in Chapter 22, state whether it should be possible to carry out each of the following nucleophilic substitution reactions.

- a. $\text{CH}_3\text{COCl} \longrightarrow \text{CH}_3\text{COOH}$ c. $\text{CH}_3\text{COOCH}_3 \longrightarrow \text{CH}_3\text{COCl}$
 b. $\text{CH}_3\text{CONHCH}_3 \longrightarrow \text{CH}_3\text{COOCH}_3$ d. $(\text{CH}_3\text{CO})_2\text{O} \longrightarrow \text{CH}_3\text{CONH}_2$

Learn the order of reactivity of carboxylic acid derivatives.

Keeping this in mind allows you to organize a very large number of reactions.

To summarize:

- Nucleophilic substitution occurs when the leaving group Z^- is a weaker base and therefore better leaving group than the attacking nucleophile :Nu^- .

- More reactive acyl compounds can be converted to less reactive acyl compounds by nucleophilic substitution.

Problem 22.12 Rank the compounds in each group in order of increasing reactivity in nucleophilic acyl substitution.

a. $\text{C}_6\text{H}_5\text{COOCH}_3$, $\text{C}_6\text{H}_5\text{COCl}$, $\text{C}_6\text{H}_5\text{CONH}_2$
 b. $\text{CH}_3\text{CH}_2\text{COOH}$, $(\text{CH}_3\text{CH}_2\text{CO})_2\text{O}$, $\text{CH}_3\text{CH}_2\text{CONHCH}_3$


Problem 22.13 Explain why trichloroacetic anhydride $[(\text{Cl}_3\text{CCO})_2\text{O}]$ is more reactive than acetic anhydride $[(\text{CH}_3\text{CO})_2\text{O}]$ in nucleophilic acyl substitution reactions.

22.7C A Preview of Specific Reactions

Sections 22.8–22.14 are devoted to specific examples of nucleophilic acyl substitution using heteroatoms as nucleophiles. There are a great many reactions, and it is easy to confuse them unless you learn the general order of reactivity of carboxylic acid derivatives. **Keep in mind that every reaction that begins with an acyl starting material involves nucleophilic substitution.**

In this text, all of the nucleophilic substitution reactions are grouped according to the carboxylic acid derivative used as a starting material. We begin with the reactions of acid chlorides, the most reactive acyl compounds, then proceed to less and less reactive carboxylic acid derivatives, ending with amides. Acid chlorides undergo many reactions, because they have the best leaving group of all acyl compounds, whereas amides undergo only one reaction, which must be carried out under harsh reaction conditions, because amides have a poor leaving group.



In general, we will examine nucleophilic acyl substitution with four different nucleophiles, as shown in the following equations.



These reactions are used to make anhydrides, carboxylic acids, esters, and amides, but not acid chlorides, from other acyl compounds. Acid chlorides are the most reactive acyl compounds (they have the best leaving group), so they are not easily formed as a product of nucleophilic substitution reactions. They can only be prepared from carboxylic acids using special reagents, as discussed in Section 22.10A.


22.8 Reactions of Acid Chlorides

Acid chlorides readily react with nucleophiles to form nucleophilic substitution products, with HCl usually formed as a reaction by-product. A weak base like pyridine is added to the reaction mixture to remove this strong acid, forming an ammonium salt.




The reaction of acid chlorides with water is rapid. Exposure of an acid chloride to moist air on a humid day leads to some hydrolysis, giving the acid chloride a very acrid odor, due to the HCl formed as a by-product.

Acid chlorides react with oxygen nucleophiles to form anhydrides, carboxylic acids, and esters.




Insect repellents containing DEET have become particularly popular because of the recent spread of many insect-borne diseases such as West Nile virus and Lyme disease. DEET does not kill insects—it repels them. It is thought that DEET somehow confuses insects so that they can no longer sense the warm moist air that surrounds a human body.

Acid chlorides also react with ammonia and 1° and 2° amines to form 1°, 2°, and 3° amides, respectively. Two equivalents of NH₃ or amine are used. One equivalent acts as a nucleophile to replace Cl and form the substitution product, while the second equivalent reacts as a base with the HCl by-product to form an ammonium salt.



As an example, reaction of an acid chloride with diethylamine forms the 3° amide *N,N*-diethyl-*m*-toluamide, popularly known as **DEET**. DEET, the active ingredient in the most widely used insect repellents, is effective against mosquitoes, fleas, and ticks.




Problem 22.14 Draw the products formed when benzoyl chloride ($\text{C}_6\text{H}_5\text{COCl}$) is treated with each nucleophile: (a) H_2O , pyridine; (b) CH_3COO^- ; (c) NH_3 (excess); (d) $(\text{CH}_3)_2\text{NH}$ (excess).

With a carboxylate nucleophile the mechanism follows the general, two-step mechanism discussed in Section 22.7: **nucleophilic attack followed by loss of the leaving group**, as shown in Mechanism 22.2.




Mechanism 22.2 Conversion of Acid Chlorides to Anhydrides



Nucleophilic substitution with the neutral nucleophiles (H_2O , $\text{R}'\text{OH}$, NH_3 , and so forth) requires an additional step for proton transfer. For example, the reaction of an acid chloride with H_2O as nucleophile converts an acid chloride to a carboxylic acid in three steps (Mechanism 22.3).




Mechanism 22.3 Conversion of Acid Chlorides to Carboxylic Acids



The exact same three-step process can be written for any neutral nucleophile that reacts with acid chlorides.

Problem 22.15 Draw a stepwise mechanism for the formation of **A** from an alcohol and acid chloride. **A** was converted in one step to blattellaquinone, the sex pheromone of the female German cockroach *Blattella germanica*.




22.9 Reactions of Anhydrides




A short laboratory synthesis of blattellaquinone (Problem 22.15), the sex pheromone of the female German cockroach, opens new possibilities for cockroach population control using pheromone-baited traps.

Nucleophilic substitution occurs only when the leaving group is a weaker base and therefore a better leaving group than the attacking nucleophile.

Although somewhat less reactive than acid chlorides, anhydrides nonetheless readily react with most nucleophiles to form substitution products. Nucleophilic substitution reactions of anhydrides are no different than the reactions of other carboxylic acid derivatives, even though anhydrides contain two carbonyl groups. **Nucleophilic attack occurs at one carbonyl group, while the second carbonyl becomes part of the leaving group.**



Anhydrides can't be used to make acid chlorides, because RCOO^- is a stronger base and therefore a poorer leaving group than Cl^- . Anhydrides can be used to make all other acyl derivatives, however. Reaction with water and alcohols yields **carboxylic acids** and **esters**, respectively. Reaction with two equivalents of NH_3 or amines forms **1°, 2°, and 3° amides**. A molecule of carboxylic acid (or a carboxylate salt) is always formed as a by-product.




Problem 22.16 Draw the products formed when benzoic anhydride $[(\text{C}_6\text{H}_5\text{CO})_2\text{O}]$ is treated with each nucleophile: (a) H_2O ; (b) CH_3OH ; (c) NH_3 (excess); (d) $(\text{CH}_3)_2\text{NH}$ (excess).

The conversion of an anhydride to an amide illustrates the mechanism of nucleophilic acyl substitution with an anhydride as starting material (Mechanism 22.4). Besides the usual steps of **nucleophilic addition** and **elimination of the leaving group**, an additional proton transfer is needed.



Mechanism 22.4 Conversion of an Anhydride to an Amide




- In Step [1], **nucleophilic attack by NH_3** forms a tetrahedral intermediate.
- Removal of a proton followed by **elimination of the leaving group, RCOO^-** (Steps [2]–[3]), forms the substitution product, a **1° amide**.

Acetaminophen reduces pain and fever, but it is not anti-inflammatory, so it is ineffective in treating conditions like arthritis, which have a significant inflammatory component. In large doses, acetaminophen causes liver damage, so dosage recommendations must be carefully followed.




Opium has been widely used as a recreational drug and pain-killing remedy for centuries. The analgesic and narcotic effects of opium are largely due to morphine. Poppy seed tea, which contains morphine, was used as a folk remedy in parts of England until World War II.

Anhydrides react with alcohols and amines with ease, so they are often used in the laboratory to prepare esters and amides. For example, acetic anhydride is used to prepare two analgesics, **acetylsalicylic acid** (aspirin) and **acetaminophen** (the active ingredient in Tylenol).



These are called **acetylation** reactions because they result in the transfer of an acetyl group, $\text{CH}_3\text{CO}-$, from one heteroatom to another.

Heroin is prepared by the acetylation of morphine, an analgesic compound isolated from the opium poppy. Both OH groups of morphine are readily acetylated with acetic anhydride to form the diester present in heroin.




Problem 22.17

Why does acetylation of *p*-aminophenol ($\text{HOC}_6\text{H}_4\text{NH}_2$) occur on the NH_2 group rather than the OH group in the synthesis of acetaminophen (Section 22.9)?

22.10 Reactions of Carboxylic Acids

Carboxylic acids are strong organic acids. Because acid–base reactions proceed rapidly, any nucleophile that is also a strong base will react with a carboxylic acid by removing a proton *first*, before any nucleophilic substitution reaction can take place.




An acid–base reaction (Reaction [1]) occurs with OH^- , NH_3 , and amines, all common nucleophiles used in nucleophilic acyl substitution reactions. Nonetheless, carboxylic acids can be converted to a variety of other acyl derivatives using special reagents, with acid catalysis, or sometimes, by using rather forcing reaction conditions. These reactions are summarized in Figure 22.3 and detailed in Sections 22.10A–22.10D.

22.10A Conversion of RCOOH to RCOCl


Carboxylic acids can't be converted to acid chlorides by using Cl^- as a nucleophile, because the attacking nucleophile Cl^- is a weaker base than the departing leaving group, OH^- . But

Figure 22.3

Nucleophilic acyl substitution reactions of carboxylic acids



carboxylic acids *can* be converted to acid chlorides using thionyl chloride, SOCl_2 , a reagent that was introduced in Section 9.12 to convert alcohols to alkyl chlorides.




This reaction converts a less reactive acyl derivative (a carboxylic acid) into a more reactive one (an acid chloride). This is possible because thionyl chloride converts the OH group of the acid into a better leaving group, and because it provides the nucleophile (Cl^-) to displace the leaving group. The steps in the process are illustrated in Mechanism 22.5.




Mechanism 22.5 Conversion of Carboxylic Acids to Acid Chlorides

Steps [1] and [2] Conversion of the OH group into a good leaving group




- Reaction of the OH group with SOCl_2 forms an intermediate that loses a proton in Step [2]. This two-step process converts the OH group into OSOCl , a **good leaving group**.

Steps [3] and [4] Substitution of the leaving group by Cl^-




- **Nucleophilic attack by Cl^- and loss of the leaving group ($\text{SO}_2 + \text{Cl}^-$)** forms the acid chloride.

Problem 22.18 Draw the products of each reaction.




22.10B Conversion of RCOOH to (RCO)₂O

Carboxylic acids cannot be readily converted to anhydrides, but dicarboxylic acids can be converted to cyclic anhydrides by heating to high temperatures. This is a **dehydration** reaction because a water molecule is lost from the diacid.




22.10C Conversion of RCOOH to RCOOR'

Treatment of a carboxylic acid with an alcohol in the presence of an acid catalyst forms an ester. This reaction is called a **Fischer esterification**.



This reaction is an equilibrium. According to Le Châtelier's principle, it is driven to the right by using excess alcohol or by removing the water as it is formed.

Ethyl acetate is a common organic solvent with a characteristic odor. It is used in nail polish remover and model airplane glue.




The mechanism for the Fischer esterification involves the usual two steps of nucleophilic acyl substitution—that is, **addition of a nucleophile followed by elimination of a leaving group**. Because the reaction is acid catalyzed, however, there are additional protonation and deprotonation steps. As always, though, the first step of any mechanism with an oxygen-containing starting material and an acid is to **protonate an oxygen atom** as shown with a general acid HA in Mechanism 22.6.




Mechanism 22.6 Fischer Esterification—Acid-Catalyzed Conversion of Carboxylic Acids to Esters

Part [1] Addition of the nucleophile R'OH




- **Protonation** in Step [1] makes the carbonyl group more electrophilic.
- **Nucleophilic addition of R'OH** forms a tetrahedral intermediate, and loss of a proton forms the neutral addition product (Steps [2]–[3]).

Part [2] Elimination of the leaving group H₂O




- Protonation of an OH group in Step [4] forms a good leaving group that is **eliminated in Step [5]**.
- Loss of a proton in Step [6] forms the ester.
- Only one resonance structure is drawn for the resonance-stabilized cations formed in Steps [1] and [5].





Esterification of a carboxylic acid occurs in the presence of acid but not in the presence of base. Base removes a proton from the carboxylic acid, forming a carboxylate anion, which does not react with an electron-rich nucleophile.



Intramolecular esterification of γ - and δ -hydroxy carboxylic acids forms five- and six-membered lactones.




Problem 22.19 Draw the products of each reaction.




Problem 22.20 Draw the products formed when benzoic acid (C₆H₅COOH) is treated with CH₃OH having its O atom labeled with ¹⁸O (CH₃¹⁸OH). Indicate where the labeled oxygen atom resides in the products.

Problem 22.21 Draw a stepwise mechanism for the following reaction.



22.10D Conversion of RCOOH to RCONR'_2

The direct conversion of a carboxylic acid to an amide with NH_3 or an amine is very difficult, even though a more reactive acyl compound is being transformed into a less reactive one. The problem is that carboxylic acids are strong organic acids and NH_3 and amines are bases, so they undergo an **acid–base reaction to form an ammonium salt** before any nucleophilic substitution occurs.




Heating at high temperature ($>100\text{ }^\circ\text{C}$) dehydrates the resulting ammonium salt of the carboxylate anion to form an amide, though the yield can be low.


Therefore, the overall conversion of RCOOH to RCONH_2 requires two steps:

- [1] **Acid–base reaction of RCOOH with NH_3 to form an ammonium salt**
- [2] **Dehydration at high temperature ($>100\text{ }^\circ\text{C}$)**


Amides are much more easily prepared from acid chlorides and anhydrides, as discussed in Sections 22.8 and 22.9.



A carboxylic acid and an amine readily react to form an amide in the presence of an additional reagent, **dicyclohexylcarbodiimide (DCC)**, which is converted to the by-product dicyclohexylurea in the course of the reaction.



DCC is a dehydrating agent. The dicyclohexylurea by-product is formed by adding the elements of H_2O to DCC. DCC promotes amide formation by converting the carboxy OH group into a better leaving group.




The mechanism consists of two parts: [1] conversion of the OH group into a better leaving group, followed by [2] **addition of the nucleophile and loss of the leaving group** to form the product of nucleophilic acyl substitution (Mechanism 22.7).




Mechanism 22.7 Conversion of Carboxylic Acids to Amides with DCC

Part [1] Conversion of OH into a better leaving group



- The first part of the mechanism consists of two steps that convert the carboxy OH group into a **better leaving group**.
- This process *activates* the carboxy group towards nucleophilic attack in Part [2].

Part [2] Addition of the nucleophile and loss of the leaving group



- In Steps [3] and [4], **nucleophilic attack of the amine** on the activated carboxy group, followed by **elimination of dicyclohexylurea** as the leaving group, forms the amide.

The reaction of an acid and an amine with DCC is often used in the laboratory to form the amide bond in peptides, as is discussed in Chapter 28.


Problem 22.22 What product is formed when acetic acid is treated with each reagent: (a) CH_3NH_2 ; (b) CH_3NH_2 , then heat; (c) $\text{CH}_3\text{NH}_2 + \text{DCC}$?

22.11 Reactions of Esters

Esters can be converted into carboxylic acids and amides.


- Esters are hydrolyzed with water in the presence of either acid or base to form carboxylic acids or carboxylate anions.

Ester hydrolysis



- Esters react with NH_3 and amines to form 1°, 2°, or 3° amides.


Reaction with nitrogen nucleophiles



22.11A Ester Hydrolysis in Aqueous Acid

The first step in acid-catalyzed ester hydrolysis is **protonation on oxygen**, the same first step of any mechanism involving an oxygen-containing starting material and an acid.

The hydrolysis of esters in aqueous acid is a reversible equilibrium reaction that is driven to the right by using a large excess of water.




The mechanism of ester hydrolysis in acid (shown in Mechanism 22.8) is the reverse of the mechanism of ester synthesis from carboxylic acids (Mechanism 22.6). Thus, the mechanism consists of the addition of the nucleophile and the elimination of the leaving group, the two steps common to all nucleophilic acyl substitutions, as well as several proton transfers, because the reaction is acid-catalyzed.




Mechanism 22.8 Acid-Catalyzed Hydrolysis of an Ester to a Carboxylic Acid

Part [1] Addition of the nucleophile H_2O



- **Protonation** in Step [1] makes the carbonyl group more electrophilic.
- **Nucleophilic addition of H_2O** forms a tetrahedral intermediate, and loss of a proton forms the neutral addition product (Steps [2]–[3]).

Part [2] Elimination of the leaving group $\text{R}'\text{OH}$



- Protonation of the OR' group in Step [4] forms a **good leaving group ($\text{R}'\text{OH}$) that is eliminated** in Step [5].
- Loss of a proton in Step [6] forms the carboxylic acid.


Problem 22.23

In Mechanism 22.8, only one Lewis structure is drawn for each intermediate. Draw all other resonance structures for the resonance-stabilized intermediates.

22.11B Ester Hydrolysis in Aqueous Base

The word **saponification** comes from the Latin *sapo* meaning **soap**. Soap is prepared by hydrolyzing esters in fats with aqueous base, as explained in Section 22.12B.


Esters are hydrolyzed in aqueous base to form carboxylate anions. Basic hydrolysis of an ester is called **saponification**.



The mechanism for this reaction has the usual two steps of the general mechanism for nucleophilic acyl substitution presented in Section 22.7—**addition of the nucleophile** followed by **loss of a leaving group**—plus an additional step involving proton transfer (Mechanism 22.9).




Mechanism 22.9 Base-Promoted Hydrolysis of an Ester to a Carboxylic Acid



- Steps [1] and [2] result in **addition of the nucleophile**, OH^- , followed by **elimination of the leaving group**, OR'^- . These two steps, which form the carboxylic acid, are reversible, because the stability of the reactants and products is comparable.
- Next, the carboxylic acid is a strong organic acid and the leaving group (OR'^-) is a strong base, so an **acid–base reaction** occurs in Step [3] to form the carboxylate anion.

Hydrolysis is base promoted, not base catalyzed, because the base (OH^-) is the nucleophile that adds to the ester and forms part of the product. It participates in the reaction and is not regenerated later.



The carboxylate anion is resonance stabilized, and this drives the equilibrium in its favor. Once the reaction is complete and the carboxylate anion is formed, it can be protonated with strong acid to form the neutral carboxylic acid.



Where do the oxygen atoms in the product come from? The $\text{C}-\text{OR}'$ bond in the ester is cleaved, so the OR' group becomes the alcohol by-product ($\text{R}'\text{OH}$) and one of the oxygens in the carboxylate anion product comes from OH^- (the nucleophile).



Problem 22.24

Sunscreens that contain esters can slowly degrade over time because the ester can be hydrolyzed. Draw the products formed when each commercial sunscreen undergoes hydrolysis with [1] H_3O^+ or [2] H_2O , OH^- .



Problem 22.25

When each isotopically labeled starting material is hydrolyzed with aqueous base, where does the label end up in the products?



22.12 Application: Lipid Hydrolysis

22.12A Olestra—A Synthetic Fat

The most prevalent naturally occurring esters are the **triacylglycerols**, which were first discussed in Section 10.6. **Triacylglycerols are the lipids that comprise animal fats and vegetable oils.**

- Each triacylglycerol is a triester, containing three long hydrocarbon side chains.
- Unsaturated triacylglycerols have one or more double bonds in their long hydrocarbon chains, whereas saturated triacylglycerols have none.





Figure 22.4 contains a ball-and-stick model of a saturated fat.

Animals store energy in the form of triacylglycerols, kept in a layer of fat cells below the surface of the skin. This fat serves to insulate the organism, as well as provide energy for its metabolic needs for long periods. The first step in the metabolism of a triacylglycerol is hydrolysis of the ester bonds to form glycerol and three fatty acids. **This reaction is simply ester hydrolysis.** In cells, this reaction is carried out with enzymes called **lipases**.




[The three bonds drawn in red are cleaved in hydrolysis.]

The fatty acids produced on hydrolysis are then oxidized in a stepwise fashion, ultimately yielding CO_2 and H_2O , as well as a great deal of energy. Oxidation of fatty acids yields twice as much energy per gram as oxidation of an equivalent weight of carbohydrate.

Figure 22.4

The three-dimensional structure of a saturated triacylglycerol





- This triacylglycerol has no double bonds in the three R groups (each with 11 C's) bonded to the ester carbonyls, making it a saturated fat.

Diets high in fat content lead to a large amount of stored fat, ultimately causing an individual to be overweight. One recent attempt to reduce calories in common snack foods has been to substitute “fake fats” such as **olestra** (trade name: **Olean**) for triacylglycerols.



Some snack foods contain the “fake fat” olestra, giving them fewer calories for the calorie-conscious consumer.



Olestra is a polyester formed from long-chain fatty acids and sucrose, the sweet-tasting carbohydrate in table sugar. Naturally occurring triacylglycerols are also polyesters formed from long-chain fatty acids, but olestra has so many ester units clustered together in close proximity that they are too hindered to be hydrolyzed. As a result, olestra is not metabolized. Instead, it passes through the body unchanged, providing no calories to the consumer.

Thus, olestra’s many C–C and C–H bonds make it similar in solubility to naturally occurring triacylglycerols, but its three-dimensional structure makes it inert to hydrolysis because of steric hindrance.

Problem 22.26 How would you synthesize olestra from sucrose?


22.12B The Synthesis of Soap

Soap has been previously discussed in Section 3.6.


Soap is prepared by the basic hydrolysis or saponification of a triacylglycerol. Heating an animal fat or vegetable oil with aqueous base hydrolyzes the three esters to form glycerol and sodium salts of three fatty acids. These carboxylate salts are **soaps**, which clean away dirt because of their two structurally different regions. The nonpolar tail dissolves grease and oil and the polar head makes it soluble in water (Figure 3.6). Most triacylglycerols have two or three different R groups in their hydrocarbon chains, so soaps are usually mixtures of two or three different carboxylate salts.



All soaps are salts of fatty acids. The main difference between soaps is the addition of other ingredients that do not alter their cleaning properties: dyes for color, scents for a pleasing odor, and oils for lubrication. Soaps that float are aerated so that they are less dense than water.




For example:



Soaps are typically made from lard (from hogs), tallow (from cattle or sheep), coconut oil, or palm oil. All soaps work in the same way, but have somewhat different properties depending on the lipid source. The length of the carbon chain in the fatty acids and the number of degrees of unsaturation affect the properties of the soap to some extent.

Problem 22.27


What is the composition of the soap prepared by hydrolysis of this triacylglycerol?



22.13 Reactions of Amides


Because amides have the poorest leaving group of all the carboxylic acid derivatives, they are the least reactive. Under strenuous reaction conditions, **amides are hydrolyzed in acid or base to form carboxylic acids or carboxylate anions.**

Amide hydrolysis



In acid, the amine by-product is protonated as an ammonium ion, whereas in base, a neutral amine is formed.

Examples



The relative lack of reactivity of the amide bond is notable in proteins, which are polymers of amino acids connected by amide linkages (Section 22.6B). Proteins are stable in aqueous


solution in the absence of acid or base, so they can perform their various functions in the aqueous cellular environment without breaking down. The hydrolysis of the amide bonds in proteins requires a variety of specific enzymes.

The mechanism of amide hydrolysis in acid is exactly the same as the mechanism of ester hydrolysis in acid (Section 22.11A) except that the leaving group is different.

The mechanism of amide hydrolysis in base has the usual two steps of the general mechanism for nucleophilic acyl substitution—**addition of the nucleophile** followed by **loss of a leaving group**—plus an additional proton transfer. The initially formed carboxylic acid reacts further under basic conditions to form the resonance-stabilized carboxylate anion, and this drives the reaction to completion. Mechanism 22.10 is written for a 1° amide.




Mechanism 22.10 Amide Hydrolysis in Base




- Steps [1] and [2] result in **addition of the nucleophile**, OH^- , followed by **elimination of the leaving group**, NH_2^- .
- Because the carboxylic acid is a strong organic acid and the leaving group (NH_2^-) is a strong base, an **acid-base reaction** occurs in Step [3] to form the carboxylate anion.

Step [2] of Mechanism 22.10 deserves additional comment. For amide hydrolysis to occur, the tetrahedral intermediate must lose NH_2^- , a stronger base and therefore poorer leaving group than OH^- (Step [2]). This means that loss of NH_2^- does not often happen. Instead, OH^- is lost as the leaving group most of the time, and the starting material is regenerated. But, when NH_2^- is occasionally eliminated, the carboxylic acid product is converted to a lower energy carboxylate anion in Step [3], and this drives the equilibrium to favor its formation.


Problem 22.28 Draw a stepwise mechanism for the following reaction.



Problem 22.29 With reference to the structures of acetylsalicylic acid (aspirin, Chapter 2 opening molecule) and acetaminophen (the active ingredient in Tylenol), explain each statement: (a) Acetaminophen tablets can be stored in the medicine cabinet for years, but aspirin slowly decomposes over time; (b) Children's Tylenol can be sold as a liquid (acetaminophen dissolved in water), but aspirin cannot.




acetylsalicylic acid




acetaminophen

22.14 Application: The Mechanism of Action of β -Lactam Antibiotics

Penicillin and related β -lactams kill bacteria by a nucleophilic acyl substitution reaction. All penicillins have an unreactive amide side chain and a very reactive amide that is part of a β -lactam. The β -lactam is more reactive than other amides because it is part of a strained, four-membered ring that is readily opened with nucleophiles.




The antibiotic properties of penicillin were discovered when bacteriologist Sir Alexander Fleming noticed that a mold of the genus *Penicillium* inhibited the growth of certain bacteria.



Unlike mammalian cells, bacterial cells are surrounded by a fairly rigid cell wall, which allows the bacterium to live in many different environments. This protective cell wall is composed of carbohydrates linked together by peptide chains containing amide linkages, formed using the enzyme **glycopeptide transpeptidase**.


Penicillin interferes with the synthesis of the bacterial cell wall. A nucleophilic OH group of the glycopeptide transpeptidase enzyme cleaves the β -lactam ring of penicillin by a nucleophilic acyl substitution reaction. The opened ring of the penicillin molecule remains covalently bonded to the enzyme, thus deactivating the enzyme, halting cell wall construction, and killing the bacterium. Penicillin has no effect on mammalian cells because they are surrounded by a flexible membrane composed of a lipid bilayer (Chapter 3) and not a cell wall.



Thus, penicillin and other β -lactam antibiotics are biologically active precisely because they undergo a nucleophilic acyl substitution reaction with an important bacterial enzyme.

Problem 22.30

Some penicillins cannot be administered orally because their β -lactam is rapidly hydrolyzed by the acidic environment of the stomach. What product is formed in the following hydrolysis reaction?



22.15 Summary of Nucleophilic Acyl Substitution Reactions

To help you organize and remember all of the nucleophilic acyl substitution reactions that can occur at a carbonyl carbon, keep in mind the following two principles:

- The better the leaving group, the more reactive the carboxylic acid derivative.
- More reactive acyl compounds can always be converted to less reactive ones. The reverse is not usually true.

This results in the following order of reactivity:




Table 22.5 summarizes the specific nucleophilic acyl substitution reactions. Use it as a quick reference to remind you which products can be formed from a given starting material.

Table 22.5 Summary of the Nucleophilic Substitution Reactions of Carboxylic Acids and Their Derivatives

Starting material	Product				
	RCOCl	(RCO) ₂ O	RCOOH	RCOOR'	RCONR' ₂
[1] RCOCl →	-	✓	✓	✓	✓
[2] (RCO) ₂ O →	✗	-	✓	✓	✓
[3] RCOOH →	✓	✓	-	✓	✓
[4] RCOOR' →	✗	✗	✓	-	✓
[5] RCONR' ₂ →	✗	✗	✓	✗	-


Table key: ✓ = A reaction occurs.
✗ = No reaction occurs.

22.16 Natural and Synthetic Fibers

All natural and synthetic fibers are high molecular weight polymers. Natural fibers are obtained from either plant or animal sources, and this determines the fundamental nature of their chemical structure. Fibers like **wool and silk obtained from animals are proteins**, and so they are formed from amino acids joined together by many amide linkages. **Cotton and linen**, on the other hand, are derived from plants and so they are **carbohydrates having the general structure of cellulose**, formed from glucose monomers. General structures for these polymers are shown in Figure 22.5.

An important practical application of organic chemistry has been the synthesis of synthetic fibers, many of which have properties that are different from and sometimes superior to their naturally occurring counterparts. The two most common classes of synthetic polymers are based on polyamides and polyesters.

Figure 22.5
The general structure of the common natural fibers




22.16A Nylon—A Polyamide




DuPont built the first commercial nylon plant in 1938. Although it was initially used by the military to make parachutes, nylon quickly replaced silk in many common clothing articles after World War II.


The search for a synthetic fiber with properties similar to silk led to the discovery of **nylon** (the chapter-opening molecule), a **polyamide**. There are several different kinds of nylon, but the most well known is called nylon 6,6.



Nylon 6,6 can be synthesized from two six-carbon monomers (hence its name)—adipoyl chloride ($\text{ClOCCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{COCl}$) and hexamethylenediamine ($\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$). This diacid chloride and diamine react together to form new amide bonds, yielding the polymer. Nylon is called a **condensation polymer** because a small molecule, in this case HCl , is eliminated during its synthesis.




Problem 22.31 What two monomers are needed to prepare nylon 6,10?




22.16B Polyesters

Polyesters constitute a second major class of condensation polymers. The most common polyester is polyethylene terephthalate (**PET**), which is sold under a variety of trade names (Dacron, Terylene, and Mylar) depending on its use.



Ester bonds (in red) join the carbon skeleton together.


One method of synthesizing a polyester is by acid-catalyzed esterification of a diacid with a diol (Fischer esterification).



Three new ester bonds are shown.

Because these polymers are easily and cheaply prepared and form strong and chemically stable materials, they have been used in clothing, films, tires, and many other products.


Problem 22.32 Draw the structure of Kodel, a polyester formed from 1,4-dihydroxymethylcyclohexane and terephthalic acid. Explain why fabrics made from Kodel are stiff and crease resistant.



Problem 22.33 Poly(lactic acid) (PLA) has received much recent attention because the lactic acid monomer [CH₃CH(OH)COOH] from which it is made can be obtained from carbohydrates rather than petroleum. This makes PLA a more “environmentally friendly” polyester. (A more in-depth discussion of green polymer synthesis is presented in Section 30.8.) Draw the structure of PLA.


22.17 Biological Acylation Reactions

Nucleophilic acyl substitution is a common reaction in biological systems. These acylation reactions are called **acyl transfer reactions** because they result in the transfer of an acyl group from one atom to another (from Z to Nu in this case).




The acyl group is transferred from Z to Nu.


In cells, such acylations occur with the sulfur analogue of an ester, called a **thioester**, having the general structure **RCOSR'**. The most common thioester is called **acetyl coenzyme A**, often referred to merely as **acetyl CoA**.



- A thioester (RCOSR') has a good leaving group ('SR'), so, like other acyl compounds, it undergoes substitution reactions with other nucleophiles.




For example, acetyl CoA undergoes enzyme-catalyzed nucleophilic acyl substitution with choline, forming acetylcholine, a charged compound that transmits nerve impulses between nerve cells.



Many other acyl transfer reactions are important cellular processes. Thioesters of fatty acids react with cholesterol, forming **cholesteryl esters** in an enzyme-catalyzed reaction (Figure 22.6). These esters are the principal form in which cholesterol is stored and transported in the body. Because cholesterol is a lipid, insoluble in the aqueous environment of the blood, it travels

Figure 22.6
Cholesteryl esters and lipoprotein particles




through the bloodstream in particles that also contain proteins and phospholipids. These particles are classified by their density.

- **LDL particles** (low density lipoproteins) transport cholesterol from the liver to the tissues.
- **HDL particles** (high density lipoproteins) transport cholesterol from the tissues back to the liver, where it is metabolized or converted to other steroids.

Atherosclerosis is a disease that results from the buildup of fatty deposits on the walls of arteries, forming deposits called **plaque**. Plaque is composed largely of the cholesterol (esterified as an ester) of LDL particles. LDL is often referred to as “bad cholesterol” for this reason. In contrast, HDL particles are called “good cholesterol” because they reduce the amount of cholesterol in the bloodstream by transporting it back to the liver.

Problem 22.34 Glucosamine is a dietary supplement available in many over-the-counter treatments for osteoarthritis. Reaction of acetyl CoA with glucosamine forms NAG, *N*-acetylglucosamine, the monomer used to form chitin, the carbohydrate that forms the rigid shells of lobsters and crabs. What is the structure of NAG?




22.18 Nitriles


We end Chapter 22 with the chemistry of **nitriles** ($\text{RC}\equiv\text{N}$). Nitriles have a carbon atom in the same oxidation state as in the acyl compounds that are the principal focus of Chapter 22. Moreover, the chemical reactions of nitriles illustrate some of the concepts first discussed earlier in Chapter 22 and in Chapters 20 and 21.

In addition to the cyanohydrins discussed in Section 21.9, two useful biologically active nitriles are **letrozole** and **anastrozole**, new drugs that reduce the recurrence of breast cancer in women whose tumors are estrogen positive.

Letrozole and anastrozole are called **aromatase inhibitors** because they block the activity of the aromatase enzyme, which is responsible for estrogen synthesis. This inhibits tumor growth in those forms of breast cancer that are stimulated by estrogen.




Generic name: letrozole
Trade name: Femara




Generic name: anastrozole
Trade name: Arimidex

Nitriles are readily prepared by $\text{S}_{\text{N}}2$ substitution reactions of unhindered methyl and 1° alkyl halides with CN^- . This reaction adds one carbon to the alkyl halide and **forms a new carbon-carbon bond**.




Because nitriles have no leaving group, they do not undergo nucleophilic substitution reactions like carboxylic acid derivatives. Because the cyano group contains an electrophilic carbon atom

that is part of a multiple bond, a nitrile reacts with nucleophiles by a **nucleophilic addition reaction**. The nature of the nucleophile determines the structure of the product.




The reactions of nitriles with water, hydride, and organometallic reagents as nucleophiles are as follows:




22.18A Hydrolysis of Nitriles

Nitriles are hydrolyzed with water in the presence of acid or base to yield **carboxylic acids** or **carboxylate anions**. In this reaction, the three C–N bonds are replaced by three C–O bonds.



The mechanism of this reaction involves the formation of an amide tautomer. Two tautomers can be drawn for any carbonyl compound, and those for a 1° amide are as follows:




Recall from Chapter 11 that tautomers are constitutional isomers that differ in the location of a double bond and a proton.

- The amide form is the more stable tautomer, having a C=O and an N–H bond.
- The imidic acid tautomer is the less stable form, having a C=N and an O–H bond.

The imidic acid and amide tautomers are interconverted by treating with acid or base, analogous to the keto–enol tautomers of other carbonyl compounds. In fact, the two amide tautomers are exactly the same as keto–enol tautomers except that a nitrogen atom replaces a carbon atom bonded to the carbonyl group.

Recall from Chapter 11 that the keto and enol tautomers of a carbonyl compound are in equilibrium, but the keto form is lower in energy, so it is highly favored in most cases.




The mechanism of nitrile hydrolysis in both acid and base consists of three parts: [1] **nucleophilic addition** of H_2O or OH^- to form the imidic acid tautomer; [2] **tautomerization** to form the amide, and [3] **hydrolysis of the amide** to form RCOOH or RCOO^- . The mechanism is shown for the basic hydrolysis of RCN to RCOO^- (Mechanism 22.11).




Mechanism 22.11 Hydrolysis of a Nitrile in Base

Part [1] Addition of the nucleophile (OH^-) to form an imidic acid




- **Nucleophilic attack of OH^-** followed by protonation forms an imidic acid.

Part [2] Tautomerization of the imidic acid to an amide




- Tautomerization occurs by a two-step sequence—**deprotonation followed by protonation**.

Part [3] Hydrolysis of the 1° amide to a carboxylate anion




- Conversion of the amide to the carboxylate anion occurs by the multistep sequence detailed in Section 22.13.

Problem 22.35 Draw the products of each reaction.




Problem 22.36 Draw a tautomer of each compound.




22.18B Reduction of Nitriles

Nitriles are reduced with metal hydride reagents to form either 1° amines or aldehydes, depending on the reducing agent.

- Treatment of a nitrile with LiAlH_4 followed by H_2O adds two equivalents of H_2 across the triple bond, forming a 1° amine.




- Treatment of a nitrile with a milder reducing agent such as DIBAL-H followed by H_2O forms an aldehyde.




The mechanism of both reactions involves **nucleophilic addition of hydride (H^-) to the polarized C–N triple bond**. Mechanism 22.12 illustrates that reduction of a nitrile to an amine requires addition of two equivalents of H^- from LiAlH_4 . It is likely that intermediate nitrogen anions complex with AlH_3 (formed in situ) to facilitate the addition. Protonation of the dianion in Step [4] forms the amine.

Mechanism 22.12 Reduction of a Nitrile with LiAlH_4

Part [1] Addition of one equivalent of hydride




Part [2] Addition of a second equivalent of hydride




With **DIBAL-H**, nucleophilic addition of one equivalent of hydride forms an anion (Step [1]), which is protonated with water to generate an **imine**, as shown in Mechanism 22.13. As described in Section 21.12, imines are hydrolyzed in water to form aldehydes. Mechanism 22.13 is written without complexation of aluminum with the anion formed in Step [1], to emphasize the identity of intermediates formed during reduction.



Mechanism 22.13 Reduction of a Nitrile with DIBAL-H




Problem 22.37 Draw the product of each reaction.



22.18C Addition of Grignard and Organolithium Reagents to Nitriles


Both Grignard and organolithium reagents react with nitriles to form ketones with a new carbon-carbon bond.




The reaction occurs by nucleophilic addition of the organometallic reagent to the polarized C–N triple bond to form an anion (Step [1]), which is protonated with water to form an **imine**. Water then hydrolyzes the imine, replacing the C=N by C=O as described in Section 21.12. The final product is a ketone with a new carbon-carbon bond (Mechanism 22.14).



Mechanism 22.14 Addition of Grignard and Organolithium Reagents (R–M) to Nitriles



Problem 22.38 Draw the products of each reaction.



Problem 22.39 What reagents are needed to convert phenylacetonitrile ($\text{C}_6\text{H}_5\text{CH}_2\text{CN}$) to each compound: (a) $\text{C}_6\text{H}_5\text{CH}_2\text{COCH}_3$; (b) $\text{C}_6\text{H}_5\text{CH}_2\text{COC}(\text{CH}_3)_3$; (c) $\text{C}_6\text{H}_5\text{CH}_2\text{CHO}$; (d) $\text{C}_6\text{H}_5\text{CH}_2\text{COOH}$?

Problem 22.40 Outline two different ways that 2-butanone can be prepared from a nitrile and a Grignard reagent.

KEY CONCEPTS

Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

Summary of Spectroscopic Absorptions of RCOZ (22.5)

IR absorptions

- All RCOZ compounds have a C=O absorption in the region 1600–1850 cm^{-1} .
 - RCOCl: 1800 cm^{-1}
 - (RCO)₂O: 1820 and 1760 cm^{-1} (two peaks)
 - RCOOR': 1735–1745 cm^{-1}
 - RCONR'₂: 1630–1680 cm^{-1}
- Additional amide absorptions occur at 3200–3400 cm^{-1} (N–H stretch) and 1640 cm^{-1} (N–H bending).
- Decreasing the ring size of a cyclic lactone, lactam, or anhydride increases the frequency of the C=O absorption.
- Conjugation shifts the C=O to lower wavenumber.

¹H NMR absorptions

- C–H α to the C=O absorbs at 2–2.5 ppm.
- N–H of an amide absorbs at 7.5–8.5 ppm.

¹³C NMR absorption

- C=O absorbs at 160–180 ppm.

Summary of Spectroscopic Absorptions of RCN (22.5)


IR absorption

- C \equiv N absorbs at ~2250 cm^{-1} .

¹³C NMR absorption

- C \equiv N absorbs at 115–120 ppm.

Summary: The Relationship between the Basicity of Z⁻ and the Properties of RCOZ




General Features of Nucleophilic Acyl Substitution

- The characteristic reaction of compounds having the general structure RCOZ is nucleophilic acyl substitution (22.1).
- The mechanism consists of two steps (22.7A):
 - [1] Addition of a nucleophile to form a tetrahedral intermediate
 - [2] Elimination of a leaving group
- More reactive acyl compounds can be used to prepare less reactive acyl compounds. The reverse is not necessarily true (22.7B).

Nucleophilic Acyl Substitution Reactions


[1] Reactions that produce acid chlorides (RCOCl)

[2] Reactions that produce anhydrides [(RCO)₂O]

a. From RCOCl (22.8):





b. From dicarboxylic acids (22.10B):





[3] Reactions that produce carboxylic acids (RCOOH)

a. From RCOCl (22.8):



b. From (RCO)₂O (22.9):

c. From RCOOR' (22.11):


d. From RCONR'₂ (R' = H or alkyl, 22.13):

[4] Reactions that produce esters (RCOOR')





a. From RCOCl (22.8):

b. From (RCO)₂O (22.9):

c. From RCOOH (22.10C):




[5] Reactions that produce amides (RCONH_2) [The reactions are written with NH_3 as nucleophile to form RCONH_2 . Similar reactions occur with $\text{R}'\text{NH}_2$ to form RCONHR' , and with $\text{R}'_2\text{NH}$ to form RCONR'_2 .]




Nitrile Synthesis (22.18)

Nitriles are prepared by $\text{S}_{\text{N}}2$ substitution using unhindered alkyl halides as starting materials.




Reactions of Nitriles


[1] Hydrolysis (22.18A)



[2] Reduction (22.18B)













[3] Reaction with organometallic reagents (22.18C)



PROBLEMS

Nomenclature

22.41 Give the IUPAC or common name for each compound.



22.42 Give the structure corresponding to each name.

a. propanoic anhydride

b. α -chlorobutyryl chloride

c. cyclohexyl propanoate

d. cyclohexanecarboxamide

e. isopropyl formate

f. *N*-cyclopentylpentanamide

g. 4-methylheptanenitrile

h. vinyl acetate

i. benzoic propanoic anhydride

j. 3-methylhexanoyl chloride

k. octyl butanoate

l. *N,N*-dibenzylformamide

Properties of Carboxylic Acid Derivatives

22.43 Rank the compounds in each group in order of increasing reactivity in nucleophilic acyl substitution.


a. $\text{CH}_3\text{CH}_2\text{CH}_2\text{CONH}_2$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCl}$, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOCH}_2\text{CH}_2\text{CH}_3$

b. $(\text{CH}_3\text{CH}_2\text{CO})_2\text{O}$, $(\text{CF}_3\text{CO})_2\text{O}$, $\text{CH}_3\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$


c. CH_3COOH , CH_3COSH , CH_3COCl

22.44 Explain each statement.

a. Ester **A** is more reactive than ester **B** in nucleophilic acyl substitution.



b. Imidazolides are much more reactive than other amides in nucleophilic acyl substitution.



22.45 Explain why CH_3CONH_2 is a stronger acid and a weaker base than $\text{CH}_3\text{CH}_2\text{NH}_2$.

Reactions

22.46 Draw the product formed when pentanoyl chloride ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{COCl}$) is treated with each reagent.

a. H_2O , pyridine

c. CH_3COO^-

e. $(\text{CH}_3\text{CH}_2)_2\text{NH}$ (excess)

b. $\text{CH}_3\text{CH}_2\text{OH}$, pyridine

d. NH_3 (excess)

f. $\text{C}_6\text{H}_5\text{NH}_2$ (excess)

22.47 Draw the product formed when pentanoic anhydride [$(\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CO})_2\text{O}$] is treated with each reagent. With some reagents, no reaction occurs.

a. SOCl_2

c. CH_3OH

e. $(\text{CH}_3\text{CH}_2)_2\text{NH}$ (excess)

b. H_2O

d. NaCl

f. $\text{CH}_3\text{CH}_2\text{NH}_2$ (excess)

22.48 Draw the product formed when phenylacetic acid ($C_6H_5CH_2COOH$) is treated with each reagent. With some reagents, no reaction occurs.

- | | | |
|--------------|----------------------|---|
| a. $NaHCO_3$ | e. NH_3 (1 equiv) | i. [1] $NaOH$; [2] CH_3COCl |
| b. $NaOH$ | f. NH_3, Δ | j. CH_3NH_2, DCC |
| c. $SOCl_2$ | g. CH_3OH, H_2SO_4 | k. [1] $SOCl_2$; [2] $CH_3CH_2CH_2NH_2$ (excess) |
| d. $NaCl$ | h. $CH_3OH, ^-OH$ | l. [1] $SOCl_2$; [2] $(CH_3)_2CHOH$ |

22.49 Draw the product formed when ethyl butanoate ($CH_3CH_2CH_2COOCH_2CH_3$) is treated with each reagent. With some reagents, no reaction occurs.

- a. $SOCl_2$ b. H_3O^+ c. $H_2O, ^-OH$ d. NH_3 e. $CH_3CH_2NH_2$











22.50 Draw the product formed when phenylacetamide ($C_6H_5CH_2CONH_2$) is treated with each reagent.


- a. H_3O^+ b. $H_2O, ^-OH$

22.51 Draw the product formed when phenylacetonitrile ($C_6H_5CH_2CN$) is treated with each reagent.

- | | | |
|-----------------|----------------------------------|-------------------------------|
| a. H_3O^+ | c. [1] CH_3MgBr ; [2] H_2O | e. [1] DIBAL-H; [2] H_2O |
| b. $H_2O, ^-OH$ | d. [1] CH_3CH_2Li ; [2] H_2O | f. [1] $LiAlH_4$; [2] H_2O |


22.52 Draw the organic products formed in each reaction.




22.53  What products are formed when **X**, which contains both a lactone and an acetal, is treated with each reagent: (a) H_3O^+ ; (b) $NaOH, H_2O$?

X


22.54 Identify compounds **A–M** in the following reaction sequence.



22.55 Draw the products of each reaction and indicate the stereochemistry at any stereogenic centers.




22.56 What products are formed when all of the amide and ester bonds are hydrolyzed in each of the following compounds? **Tamiflu** [part (a)] is the trade name of the antiviral agent oseltamivir, thought to be the most effective agent in treating influenza. Governments are stockpiling the drug in the event of an influenza pandemic. **Aspartame** [part (b)] is the artificial sweetener used in Equal and many diet beverages. One of the products of this hydrolysis reaction is the amino acid phenylalanine. Infants afflicted with phenylketonuria cannot metabolize this amino acid, so it accumulates, causing mental retardation. When the affliction is identified early, a diet limiting the consumption of phenylalanine (and compounds like aspartame that are converted to it) can make a normal life possible.




Mechanism


22.57 Draw a stepwise mechanism for each reaction.




22.58 When acetic acid (CH_3COOH) is treated with a trace of acid in water labeled with ^{18}O , the label gradually appears in both oxygen atoms of the carboxylic acid. Draw a mechanism that explains this phenomenon.




22.59 Although γ -butyrolactone (Problem 19.63) is a biologically inactive compound, it is converted in the body to 4-hydroxybutanoic acid (GHB), an addictive and intoxicating recreational drug (Section 19.5). Draw a stepwise mechanism for this conversion in the presence of acid.




22.75 Devise a synthesis of each compound from benzene and organic alcohols containing four or fewer carbons. You may also use any required organic or inorganic reagents.



22.76 Devise a synthesis of each ester from benzene, organic alcohols, and any other needed inorganic reagents.



22.77 (a) Both monomers needed for the synthesis of nylon 6,6 can be prepared from 1,4-dichlorobutane. Write out the steps illustrating these syntheses. (b) Devise a synthesis of adipic acid from cyclohexene.




22.78 Devise a synthesis of each labeled compound using H_2^{18}O and $\text{CH}_3^{13}\text{CH}_2\text{OH}$ as the only sources of labeled starting materials. You may use any other unlabeled organic compounds and inorganic reagents.




Polymers

22.79 What polyester or polyamide can be prepared from each pair of monomers?





22.80 What two monomers are needed to prepare each polymer?



General Problem


22.81 Taxotere is the trade name for the synthetic anticancer drug docetaxel, whose structure closely resembles the naturally occurring compound taxol, which is isolated from the Pacific yew tree (Section 5.5).




- Taxol's limited water solubility led to an extensive search for related compounds with increased water solubility. What structural features give docetaxel a better water solubility profile?
- Docetaxel contains a carbamate (labeled in red), a functional group with a carbonyl group bonded to a nitrogen and oxygen atom. Draw three more resonance structures for a carbamate (in addition to the Lewis structure with all neutral atoms given). Rank all four resonance structures in order of increasing stability.
- A carbamate with a *tert*-butoxy group [(CH₃)₃CO-] is hydrolyzed according to the given equation. Draw a stepwise mechanism for the hydrolysis of a carbamate to the three products shown.
- Assuming that all ester and carbamate bonds are cleaved, draw all products formed when docetaxel is hydrolyzed with aqueous acid.

Spectroscopy

22.82 How can IR spectroscopy be used to distinguish between each pair of isomers?



22.83 Rank the compounds in each group in order of increasing frequency of the C=O absorption in their IR spectra.




22.84 Identify the structures of each compound from the given data.

- Molecular formula: C₆H₁₂O₂
 IR absorption: 1738 cm⁻¹
¹H NMR: 1.12 (triplet, 3 H), 1.23 (doublet, 6 H), 2.28 (quartet, 2 H), and 5.00 (septet, 1 H) ppm
- Molecular formula: C₄H₇N
 IR absorption: 2250 cm⁻¹
¹H NMR: 1.08 (triplet, 3 H), 1.70 (multiplet, 2 H), and 2.34 (triplet, 2 H) ppm
- Molecular formula: C₈H₉NO
 IR absorptions: 3328 and 1639 cm⁻¹
¹H NMR: 2.95 (singlet, 3 H), 6.95 (singlet, 1 H), and 7.3–7.7 (multiplet, 5 H) ppm
- Molecular formula: C₄H₇ClO
 IR absorption: 1802 cm⁻¹
¹H NMR: 0.95 (triplet, 3 H), 1.07 (multiplet, 2 H), and 2.90 (triplet, 2 H) ppm
- Molecular formula: C₅H₁₀O₂
 IR absorption: 1750 cm⁻¹
¹H NMR: 1.20 (doublet, 6 H), 2.00 (singlet, 3 H), and 4.95 (septet, 1 H) ppm


- f. Molecular formula $C_{10}H_{12}O_2$
 IR absorption: 1740 cm^{-1}
 $^1\text{H NMR}$: 1.2 (triplet, 3 H), 2.4 (quartet, 2 H), 5.1 (singlet, 2 H), and 7.1–7.5 (multiplet, 5 H) ppm
- g. Molecular formula $C_8H_{14}O_3$
 IR absorptions: 1810 and 1770 cm^{-1}
 $^1\text{H NMR}$: 1.25 (doublet, 12 H) and 2.65 (septet, 2 H) ppm

22.85 Identify the structures of **A** and **B**, isomers of molecular formula $C_{10}H_{12}O_2$, from their IR data and $^1\text{H NMR}$ spectra.



a. IR absorption for **A** at 1718 cm^{-1}




b. IR absorption for **B** at 1740 cm^{-1}



22.86 Phenacetin is an analgesic compound having molecular formula $C_{10}H_{13}NO_2$. Once a common component in over-the-counter pain relievers such as APC (aspirin, phenacetin, caffeine), phenacetin is no longer used because of its liver toxicity. Deduce the structure of phenacetin from its $^1\text{H NMR}$ and IR spectra.




22.87 Identify the structure of compound **C** (molecular formula $C_{11}H_{15}NO_2$), which has an IR absorption at 1699 cm^{-1} and the $^1\text{H NMR}$ spectrum shown below.




22.88 Identify the structures of **D** and **E**, isomers of molecular formula $C_6H_{12}O_2$, from their IR and $^1\text{H NMR}$ data. Signals at 1.35 and 1.60 ppm in the $^1\text{H NMR}$ spectrum of **D** and 1.90 ppm in the $^1\text{H NMR}$ spectrum of **E** are multiplets.

a. IR absorption for **D** at 1743 cm^{-1}




b. IR absorption for **E** at 1746 cm^{-1}




Challenge Problems

- 22.89** The ^1H NMR spectrum of 2-chloroacetamide ($\text{ClCH}_2\text{CONH}_2$) shows three signals at 4.02, 7.35, and 7.60 ppm. What protons give rise to each signal? Explain why three signals are observed.
- 22.90** Compelling evidence for the existence of a tetrahedral intermediate in nucleophilic acyl substitution was obtained in a series of elegant experiments carried out by Myron Bender in 1951. The key experiment was the reaction of aqueous OH^- with ethyl benzoate ($\text{C}_6\text{H}_5\text{COOCH}_2\text{CH}_3$) labeled at the carbonyl oxygen with ^{18}O . Bender did not allow the hydrolysis to go to completion, and then examined the presence of a label in the *recovered starting material*. He found that some of the recovered ethyl benzoate no longer contained a label at the carbonyl oxygen. With reference to the accepted mechanism of nucleophilic acyl substitution, explain how this provides evidence for a tetrahedral intermediate.

The starting material contains a label on the carbonyl oxygen.



- 22.91** Draw a stepwise mechanism for the following reactions, two steps in R. B. Woodward's classic synthesis of reserpine in 1958. Reserpine, which is isolated from the extracts of the Indian snakeroot *Rauwolfia serpentina Benth*, has been used to manage mild hypertension associated with anxiety.



23

Substitution Reactions of Carbonyl Compounds at the α Carbon

- 23.1 Introduction
- 23.2 Enols
- 23.3 Enolates
- 23.4 Enolates of unsymmetrical carbonyl compounds
- 23.5 Racemization at the α carbon
- 23.6 A preview of reactions at the α carbon
- 23.7 Halogenation at the α carbon
- 23.8 Direct enolate alkylation
- 23.9 Malonic ester synthesis
- 23.10 Acetoacetic ester synthesis



Tamoxifen is a potent anticancer drug used widely in the treatment of breast cancer. Tamoxifen binds to estrogen receptors, and in this way inhibits the growth of breast cancers that are estrogen dependent. One method to synthesize tamoxifen forms a new carbon-carbon bond on the α carbon to a carbonyl group using an intermediate enolate. In Chapter 23 we learn about these and other carbon-carbon bond-forming reactions that occur at the α carbon.


Chapters 23 and 24 focus on reactions that occur at the α carbon to a carbonyl group. These reactions are different from the reactions of Chapters 20–22, all of which involved nucleophilic attack at the electrophilic carbonyl carbon. In reactions at the α carbon, the carbonyl compound serves as a *nucleophile* that reacts with a carbon or halogen electrophile to form a new bond to the α carbon.

Chapter 23 concentrates on **substitution reactions at the α carbon**, whereas Chapter 24 concentrates on reactions between two carbonyl compounds, one of which serves as the nucleophile and one of which is the electrophile. Many of the reactions in Chapter 23 form new carbon–carbon bonds, thus adding to your repertoire of reactions that can be used to synthesize more complex organic molecules from simple precursors. As you will see, the reactions introduced in Chapter 23 have been used to prepare a wide variety of interesting and useful compounds.


23.1 Introduction

Up to now, the discussion of carbonyl compounds has centered on their reactions with nucleophiles at the electrophilic carbonyl carbon. **Two general reactions are observed**, depending on the structure of the carbonyl starting material.

- **Nucleophilic addition** occurs when there is no electronegative atom Z on the carbonyl carbon (as with aldehydes and ketones).




- **Nucleophilic acyl substitution** occurs when there is an electronegative atom Z on the carbonyl carbon (as with carboxylic acids and their derivatives).




Reactions can also occur at the α carbon to the carbonyl group. These reactions proceed by way of **enols or enolates**, two electron-rich intermediates that react with electrophiles, forming a new bond on the α carbon. This reaction results in the **substitution of the electrophile E for hydrogen**.

Hydrogen atoms on the α carbon are called **α hydrogens**.




23.2 Enols

Recall from Chapter 11 that **enol and keto forms are tautomers of the carbonyl group that differ in the position of a double bond and a proton**. These constitutional isomers are in equilibrium with each other.




- A keto tautomer has a $C=O$ and an additional $C-H$ bond.
- An enol tautomer has an $O-H$ group bonded to a $C=C$.

Equilibrium favors the keto form for most carbonyl compounds largely because a $C=O$ is much stronger than a $C=C$. For simple carbonyl compounds, $< 1\%$ of the enol is present at equilibrium. With unsymmetrical ketones, moreover, two different enols are possible, yet they still total $< 1\%$.




With compounds containing two carbonyl groups separated by a single carbon (called β -dicarbonyl compounds or 1,3-dicarbonyl compounds), however, the concentration of the enol form sometimes exceeds the concentration of the keto form.



Two factors stabilize the enol of β -dicarbonyl compounds: **conjugation** and **intramolecular hydrogen bonding**. The $C=C$ of the enol is conjugated with the carbonyl group, allowing delocalization of the electron density in the π bonds. Moreover, the OH of the enol can hydrogen bond to the oxygen of the nearby carbonyl group. Such intramolecular hydrogen bonds are especially stabilizing when they form a six-membered ring, as in this case.


Sample Problem 23.1

Convert each compound to its enol or keto tautomer.




Solution






- a. To convert a carbonyl compound to its enol tautomer, draw a double bond between the carbonyl carbon and the α carbon, and change the $C=O$ to $C-OH$. In this case, both α carbons are identical, so only one enol is possible.



- b. To convert an enol to its keto tautomer, change the $C-OH$ to $C=O$ and add a proton to the other end of the $C=C$.



Problem 23.1 Draw the enol or keto tautomer(s) of each compound.




Problem 23.2 Ignoring stereoisomers, draw the two possible enols for 2-butanone ($\text{CH}_3\text{COCH}_2\text{CH}_3$), and predict which one is more stable.

23.2A The Mechanism of Tautomerization

Tautomerization, the process of converting one tautomer into another, is catalyzed by both acid and base. Tautomerization always requires two steps (**protonation** and **deprotonation**), but the order of these steps depends on whether the reaction takes place in acid or base. In Mechanisms 23.1 and 23.2 for tautomerization, the keto form is converted to the enol form. All of the steps are reversible, though, so they equally apply to the conversion of the enol form to the keto form.




Mechanism 23.1 Tautomerization in Acid



- With acid, **protonation precedes deprotonation**.
- **Protonation** of the carbonyl oxygen forms a resonance-stabilized cation in Step [1], and deprotonation in Step [2] forms the enol. The net result of these two steps is the movement of a double bond and a proton.




Mechanism 23.2 Tautomerization in Base



- With base, **deprotonation precedes protonation**.
- **Removal of a proton from the α carbon** forms a resonance-stabilized enolate in Step [1].
- **Protonation** of the enolate with H_2O forms the enol in Step [2].


Problem 23.3 Draw a stepwise mechanism for the following reaction.




23.2B How Enols React

Like other compounds with carbon-carbon double bonds, **enols are electron rich, so they react as nucleophiles**. Enols are even more electron rich than alkenes, though, because the OH group has a powerful electron-donating resonance effect. A second resonance structure can be drawn

for the enol that places a negative charge on one of the carbon atoms. As a result, this carbon atom is especially nucleophilic, and it can react with an electrophile E^+ to form a new bond to carbon. Loss of a proton then forms a neutral product.



- Reaction of an enol with an electrophile E^+ forms a new C–E bond on the α carbon. The net result is substitution of H by E on the α carbon.




Problem 23.4 When phenylacetaldehyde ($C_6H_5CH_2CHO$) is dissolved in D_2O with added DCl, the hydrogen atoms α to the carbonyl are gradually replaced by deuterium atoms. Write a mechanism for this process that involves enols as intermediates.


23.3 Enolates

Enolates are formed when a base removes a proton on the α carbon to a carbonyl group. A C–H bond on the α carbon is more acidic than many other sp^3 hybridized C–H bonds, because the resulting enolate is resonance stabilized. Moreover, one of the resonance structures is especially stable because it places a negative charge on an electronegative oxygen atom.

Forming enolates from carbonyl compounds was first discussed in Section 21.7.






Enolates are always formed by removal of a proton on the α carbon.



The pK_a of the α hydrogen in an aldehyde or ketone is ~ 20 . As shown in Table 23.1, this makes it considerably more acidic than the C–H bonds in CH_3CH_3 and $CH_3CH=CH_2$. Although C–H bonds α to a carbonyl are more acidic than many other C–H bonds, they are still less acidic than

Table 23.1 A Comparison of pK_a Values

	Compound	pK_a	Conjugate base	Structural features of the conjugate base
Increasing acidity  Increasing stability of the conjugate base	CH_3CH_3	50	$\text{CH}_3\text{C}^-\text{H}_2$	• The conjugate base has a (-) charge on C, but is not resonance stabilized.
	$\text{CH}_2=\text{CHCH}_3$	43	$\text{CH}_2=\text{CH}-\text{C}^-\text{H}_2 \leftrightarrow \text{C}^-\text{H}_2-\text{CH}=\text{CH}_2$	• The conjugate base has a (-) charge on C, and is resonance stabilized.
	$(\text{CH}_3)_2\text{C}=\text{O}$	19.2		• The conjugate base has two resonance structures, one of which has a (-) charge on O.
	$\text{CH}_3\text{CH}_2\text{OH}$	16	$\text{CH}_3\text{CH}_2\text{O}^-$	• The conjugate base has a (-) charge on O, but is not resonance stabilized.
	$\text{CH}_3\text{CO}_2\text{H}$	4.8		• The conjugate base has two resonance structures, both of which have a (-) charge on O.

- **Resonance stabilization of the conjugate base increases acidity.**
 - $\text{CH}_2=\text{CHCH}_3$ is more acidic than $\text{CH}_3\text{CH}_2\text{CH}_3$.
 - CH_3COOH is more acidic than $\text{CH}_3\text{CH}_2\text{OH}$.
- **Placing a negative charge on O in the conjugate base increases acidity.**
 - $\text{CH}_3\text{CH}_2\text{OH}$ is more acidic than $\text{CH}_3\text{CH}_2\text{CH}_3$.
 - CH_3COCH_3 is more acidic than $\text{CH}_2=\text{CHCH}_3$.
 - CH_3COOH (with two O atoms) is more acidic than CH_3COCH_3 .


O–H bonds that always place the negative charge of the conjugate base on an electronegative oxygen atom (c.f. $\text{CH}_3\text{CH}_2\text{OH}$ and CH_3COOH in Table 23.1).

The electrostatic potential plots in Figure 23.1 compare the electron density of the acetone enolate, which is resonance stabilized and delocalized, with that of $(\text{CH}_3)_2\text{CHO}^-$, an alkoxide that is not resonance stabilized.


23.3A Examples of Enolates and Related Anions

In addition to enolates from aldehydes and ketones, **enolates from esters and 3° amides can be formed as well**, although the α hydrogen is somewhat less acidic. **Nitriles** also have acidic protons on the carbon atom adjacent to the cyano group, because the negative charge of the conjugate base is stabilized by delocalization onto an electronegative nitrogen atom.


Figure 23.1
Electron density in an enolate and an alkoxide




- The acetone enolate is resonance stabilized. The negative charge is delocalized on the oxygen atom (pale red) and the carbon atom (pale green).
- The alkoxide anion is not resonance stabilized. The negative charge is concentrated on the oxygen atom only (deep red).



The protons on the carbon between the two carbonyl groups of a β -dicarbonyl compound are especially acidic because resonance delocalizes the negative charge on two different oxygen atoms. Table 23.2 lists $\text{p}K_a$ values for β -dicarbonyl compounds as well as other carbonyl compounds and nitriles.



Problem 23.5 Draw additional resonance structures for each anion.



Problem 23.6 Which C–H bonds in the following molecules are acidic because the resulting conjugate base is resonance stabilized?





Table 23.2 $\text{p}K_a$ Values for Some Carbonyl Compounds and Nitriles

Compound type	Example	$\text{p}K_a$	Compound type	Example	$\text{p}K_a$
[1] Amide		30	[6] 1,3-Diester		13.3
[2] Nitrile	$\text{CH}_3\text{C}\equiv\text{N}$	25	[7] 1,3-Dinitrile	$\text{N}\equiv\text{C}-\text{CH}_2-\text{C}\equiv\text{N}$	11
[3] Ester		25	[8] β -Keto ester		10.7
[4] Ketone		19.2	[9] β -Diketone		9
[5] Aldehyde		17			


Problem 23.7

Rank the protons in the indicated CH₂ groups in order of increasing acidity, and explain why you chose this order.

**23.3B The Base**

The formation of an enolate is an acid–base equilibrium, so the **stronger the base, the more enolate that forms**.

**Enolate formation—
An acid–base equilibrium**



Stronger bases drive the equilibrium to the right.

We have now used the term **amide** in two different ways—first as a functional group (e.g., the carboxylic acid derivative RCONH₂) and now as a base (e.g., ⁻NH₂, which can be purchased as a sodium or lithium salt, NaNH₂ or LiNH₂, respectively). In Chapter 23 we will use dialkylamides, ⁻NR₂, in which the two H atoms of ⁻NH₂ have been replaced by R groups.



Enolate formation with LDA is typically carried out at $-78\text{ }^{\circ}\text{C}$, a convenient temperature to maintain in the laboratory because it is the temperature at which dry ice (solid CO₂) sublimates. A low-temperature cooling bath can be made by adding dry ice to acetone until the acetone cools to $-78\text{ }^{\circ}\text{C}$. Immersing a reaction flask in this cooling bath keeps its contents at a constant low temperature.

We can predict the extent of an acid–base reaction by comparing the pK_a of the starting acid (the carbonyl compound in this case) with the pK_a of the conjugate acid formed. **The equilibrium favors the side with the weaker acid (the acid with the higher pK_a value)**. The pK_a of many carbonyl compounds is ~ 20 , so a significant amount of enolate will form only if the pK_a of the conjugate acid is > 20 .


The common bases used to form enolates are hydroxide (⁻OH), various alkoxides (⁻OR), hydride (⁻H), and dialkylamides (⁻NR₂). How much enolate is formed using each of these bases is indicated in Table 23.3.

When the pK_a of the conjugate acid is < 20 , as it is for ⁻OH and all ⁻OR (entries 1–3), only a small amount of enolate is formed at equilibrium. These bases are more useful in forming enolates when more acidic 1,3-dicarbonyl compounds are used as starting materials. They are also used when both the enolate and the carbonyl starting material are involved in the reaction, as is the case for reactions described in Chapter 24.

To form an enolate in essentially 100% yield, a much stronger base such as lithium diisopropylamide, Li⁺ ⁻N[CH(CH₃)₂]₂, abbreviated as **LDA**, is used (entry 5). **LDA is a strong nonnucleophilic base**. Like the other nonnucleophilic bases (Sections 7.8B and 8.1), its bulky isopropyl groups make the nitrogen atom too hindered to serve as a nucleophile. It is still able, though, to remove a proton in an acid–base reaction.


Table 23.3 Enolate Formation with Various Bases:
RCOCH₃ ($pK_a \approx 20$) + B: → RCOCH₂⁻ + HB⁺

	Base (B:)	Conjugate acid (HB ⁺)	pK_a of HB ⁺	% Enolate
[1]	Na ⁺ ⁻ OH	H ₂ O	15.7	< 1%
[2]	Na ⁺ ⁻ OCH ₂ CH ₃	CH ₃ CH ₂ OH	16	< 1%
[3]	K ⁺ ⁻ OC(CH ₃) ₃	(CH ₃) ₃ COH	18	1–10% (depending on the carbonyl compound)
[4]	Na ⁺ ⁻ H	H ₂	35	100%
[5]	Li ⁺ ⁻ N[CH(CH ₃) ₂] ₂	HN[CH(CH ₃) ₂] ₂	40	100%



lithium diisopropylamide

LDA



The N atom is too crowded to be a nucleophile.


THF

tetrahydrofuran

a polar aprotic

solvent


LDA quickly deprotonates essentially all of the carbonyl starting material, even at -78°C , to form the enolate product. THF is the typical solvent for these reactions.



Equilibrium greatly favors the products.
Essentially all of the ketone is converted to enolate.





LDA can be prepared by deprotonating diisopropylamine with an organolithium reagent such as butyllithium, and then used immediately in a reaction.

Preparation of LDA



LDA


Problem 23.8 Draw the product formed when each starting material is treated with LDA in THF solution at -78°C .



Problem 23.9 When ethyl acetoacetate ($\text{CH}_3\text{COCH}_2\text{CO}_2\text{CH}_2\text{CH}_3$) is treated with one equivalent of CH_3MgBr , a gas is evolved from the reaction mixture, and after adding aqueous acid, ethyl acetoacetate is recovered in high yield. Identify the gas formed and explain why the starting material was recovered in this reaction.

23.3C General Reactions of Enolates

Enolates are nucleophiles, and as such they react with many electrophiles. Because an enolate is resonance stabilized, however, it has two reactive sites—the carbon and oxygen atoms that bear the negative charge. **A nucleophile with two reactive sites is called an ambident nucleophile.** In theory, each of these atoms could react with an electrophile to form two different products, one with a new bond to carbon, and one with a new bond to oxygen.




Because enolates usually react at carbon instead of oxygen, the resonance structure that places the negative charge on oxygen will often be omitted in multistep mechanisms.

An enolate usually reacts at the carbon end, however, because this site is more nucleophilic. Thus, enolates generally react with electrophiles on the α carbon so that many reactions in Chapter 23 follow a two-step path:

- [1] Reaction of a carbonyl compound with base forms an enolate.
- [2] Reaction of the enolate with an electrophile forms a new bond on the α carbon.

23.4 Enolates of Unsymmetrical Carbonyl Compounds

What happens when an unsymmetrical carbonyl compound like 2-methylcyclohexanone is treated with base? **Two enolates are possible**, one formed by removal of a 2° hydrogen, and one formed by removal of a 3° hydrogen.



Path [1] occurs *faster* than Path [2] because it results in removal of the less hindered 2° hydrogen, forming an enolate on the less substituted α carbon. Path [2] results in removal of a 3° hydrogen, forming the *more stable* enolate with the more substituted double bond. This enolate predominates at equilibrium.


- The kinetic enolate is formed faster because it is the less substituted enolate.
- The thermodynamic enolate is lower in energy because it is the more substituted enolate.

It is possible to regioselectively form one or the other enolate by the proper use of reaction conditions, because the base, solvent, and reaction temperature all affect the identity of the enolate formed.

Kinetic Enolates

The kinetic enolate forms faster, so mild reaction conditions favor it over slower processes with higher energies of activation. It is the less stable enolate, so it must not be allowed to equilibrate to the more stable thermodynamic enolate. **The kinetic enolate is favored by:**

- [1] **A strong nonnucleophilic base.** A strong base assures that the enolate is formed rapidly. A bulky base like LDA removes the more accessible proton on the less substituted carbon much faster than a more hindered proton.
- [2] **Polar aprotic solvent.** The solvent must be polar to dissolve the polar starting materials and intermediates. It must be aprotic so that it does not protonate any enolate that is formed. THF is both polar and aprotic.
- [3] **Low temperature.** The temperature must be low (-78°C) to prevent the kinetic enolate from equilibrating to the thermodynamic enolate.




- A kinetic enolate is formed with a strong, nonnucleophilic base (LDA) in a polar aprotic solvent (THF) at low temperature ($-78\text{ }^\circ\text{C}$).

Thermodynamic Enolates

A **thermodynamic enolate is favored by equilibrating conditions**. This is often achieved using a **strong base in a protic solvent**. A strong base yields both enolates, but in a protic solvent, enolates can also be protonated to re-form the carbonyl starting material. At equilibrium, the lower energy intermediate always wins out, so that **the more stable, more substituted enolate is present in higher concentration**. Thus, the **thermodynamic enolate is favored by:**



- [1] **A strong base.** $\text{Na}^+\text{OCH}_2\text{CH}_3$, $\text{K}^+\text{OC}(\text{CH}_3)_3$, or other alkoxides are common.
- [2] **Protic solvent.** $\text{CH}_3\text{CH}_2\text{OH}$ or other alcohols.
- [3] **Room temperature ($25\text{ }^\circ\text{C}$).**



- A thermodynamic enolate is formed with a strong base (RO^-) in a polar protic solvent (ROH) at room temperature.


Sample Problem 23.2

What is the major enolate formed in each reaction?




Solution


- a. LDA is a strong, nonnucleophilic base that removes a proton on the less substituted α carbon to form the kinetic enolate.



- b. $\text{NaOCH}_2\text{CH}_3$ (a strong base) and $\text{CH}_3\text{CH}_2\text{OH}$ (a protic solvent) favor removal of a proton from the more substituted α carbon to form the thermodynamic enolate.



Problem 23.10 What enolate is formed when each ketone is treated with LDA in THF solution? What enolate is formed when these same ketones are treated with NaOCH_3 in CH_3OH solution?



23.5 Racemization at the α Carbon


Recall from Section 16.5 that an enolate can be stabilized by the delocalization of electron density only if it possesses the proper geometry and hybridization.

- The electron pair on the carbon adjacent to the $\text{C}=\text{O}$ must occupy a p orbital that overlaps with the two other p orbitals of the $\text{C}=\text{O}$, making an enolate conjugated.

- Thus, all three atoms of the enolate are sp^2 hybridized and trigonal planar.

These bonding features are shown in the acetone enolate in Figure 23.2.


When the α carbon to the carbonyl is a stereogenic center, treatment with aqueous base leads to **racemization** by a two-step process: **deprotonation to form an enolate and protonation to re-form the carbonyl compound**. For example, chiral ketone **A** reacts with aqueous OH^- to form an achiral enolate having an sp^2 hybridized α carbon. Because the enolate is planar, it can be protonated with H_2O with equal probability from both directions, yielding a racemic mixture of two ketones.



Problem 23.11 Explain each observation: (a) When (2*R*)-2-methylcyclohexanone is treated with NaOH in H_2O , the optically active solution gradually loses optical activity. (b) When (3*R*)-3-methylcyclohexanone is treated with NaOH in H_2O , the solution remains optically active.

Figure 23.2

The hybridization and geometry of the acetone enolate ($\text{CH}_3\text{COCH}_2^-$)




- The O atom and both C's of the enolate are sp^2 hybridized and lie in a plane.
- Each atom has a p orbital extending above and below the plane; these orbitals overlap to delocalize electron density.

23.6 A Preview of Reactions at the α Carbon

Having learned about the synthesis and properties of enolates, we can now turn our attention to their reactions. Like enols, **enolates are nucleophiles**, but because they are negatively charged, enolates are much more nucleophilic than neutral enols. Consequently, they undergo a wider variety of reactions.


Two general types of reactions of enolates—**substitutions** and **reactions with other carbonyl compounds**—will be discussed in the remainder of Chapter 23 and in Chapter 24. Both reactions form new bonds to the carbon α to the carbonyl.

- Enolates react with electrophiles to afford substitution products.



Two different kinds of substitution reactions are examined: **halogenation** with X_2 and **alkylation** with alkyl halides RX . These reactions are detailed in Sections 23.7–23.10.


- Enolates react with other carbonyl groups at the electrophilic carbonyl carbon.



These reactions are more complicated because the initial addition adduct goes on to form different products depending on the structure of the carbonyl group. These reactions form the subject of Chapter 24.

23.7 Halogenation at the α Carbon

The first substitution reaction we examine is **halogenation**. Treatment of a ketone or aldehyde with halogen and either acid or base results in **substitution of X for H on the α carbon**, forming an **α -halo aldehyde or ketone**. Halogenation readily occurs with Cl_2 , Br_2 , and I_2 .




The mechanisms of halogenation in acid and base are somewhat different.

- Reactions done in acid generally involve *enol* intermediates.
- Reactions done in base generally involve *enolate* intermediates.

23.7A Halogenation in Acid


Halogenation is often carried out by treating a carbonyl compound with a halogen in acetic acid. In this way, acetic acid is both the solvent and the acid catalyst for the reaction.



The mechanism of acid-catalyzed halogenation consists of two parts: **tautomerization** of the carbonyl compound to the enol form, and **reaction of the enol with halogen**. Mechanism 23.3 illustrates the reaction of $(\text{CH}_3)_2\text{C}=\text{O}$ with Br_2 in CH_3COOH .


Mechanism 23.3 Acid-Catalyzed Halogenation at the α Carbon

Part [1] Tautomerization to the enol




- In Part [1], the ketone is converted to its enol tautomer by the usual two-step process: **protonation** of the carbonyl oxygen, followed by **deprotonation** of the α carbon atom.

Part [2] Reaction of the enol with halogen




- In Part [2], addition of the halogen to the enol followed by deprotonation forms the neutral substitution product (Steps [3]–[4]). **The overall process results in substitution of H by Br on the α carbon.**

Problem 23.12 Draw the products of each reaction.



23.7B Halogenation in Base

Halogenation in base is much less useful, because it is often difficult to stop the reaction after addition of just one halogen atom to the α carbon. For example, treatment of propiophenone with Br_2 and aqueous OH^- yields a dibromo ketone.




Reactions of carbonyl compounds with base invariably involve enolates because the α hydrogens of the carbonyl compound are easily removed.

The mechanism for introduction of each Br atom involves the same two steps: **deprotonation with base followed by reaction with Br_2** to form a new C–Br bond, as shown in Mechanism 23.4.




Mechanism 23.4 Halogenation at the α Carbon in Base




- Treatment of the ketone with base forms a nucleophilic enolate in Step [1], which reacts with Br_2 (the electrophile) to form the monosubstitution product—that is, one α H is replaced by one Br on the α carbon.

Only a small amount of the enolate forms at equilibrium using OH^- as base, but the enolate is such a strong nucleophile that it readily reacts with Br_2 , thus driving the equilibrium to the right. Then, the same two steps introduce the second Br atom on the α carbon: **deprotonation** followed by **nucleophilic attack**.




It is difficult to stop this reaction after the addition of one Br atom because the electron-withdrawing inductive effect of Br stabilizes the second enolate. As a result, the α H of α -bromopropiophenone is more acidic than the α H atoms of propiophenone, making it easier to remove with base.

Halogenation of a methyl ketone with excess halogen, called the **haloform reaction**, results in cleavage of a carbon-carbon σ bond and formation of two products, a carboxylate anion and CHX_3 (commonly called **haloform**).



Although all ketones with α hydrogens react with base and I_2 , only **methyl** ketones form CHI_3 (iodoform), a pale yellow solid that precipitates from the reaction mixture. This reaction is the basis of the **iodoform test**, once a common chemical method to detect methyl ketones. Methyl ketones give a positive iodoform test (appearance of a yellow solid), whereas other ketones give a negative iodoform test (no change in the reaction mixture).




In the haloform reaction, the three H atoms of the CH_3 group are successively replaced by X to form an intermediate that is oxidatively cleaved with base. Mechanism 23.5 is written with I_2 as halogen, forming CHI_3 (iodoform) as product.




Mechanism 23.5 The Haloform Reaction

Part [1] The conversion of CH_3 to CI_3



- In Part [1], each of the three hydrogen atoms of the CH_3 group is replaced with iodine via the two-step mechanism previously discussed—that is, deprotonation followed by halogenation to form the substitution product.
- Steps [1] and [2] are then repeated twice more to form the triiodo substitution product.

Part [2] Oxidative cleavage with OH^-



- In Step [3], OH^- adds to the carbonyl group in a typical **nucleophilic addition** reaction of a ketone, but the three I atoms give this ketone a good leaving group, CI_3^- .
- Elimination of CI_3^- in Step [4] results in cleavage of a carbon–carbon bond**, and then in the last step, proton transfer forms the carboxylate anion and iodoform.

Steps [3] and [4] result in a **nucleophilic substitution** reaction of a *ketone*. Because ketones normally undergo **nucleophilic addition**, this two-step sequence makes the haloform reaction unique. Substitution occurs because the three electronegative halogen atoms make CX_3 (CI_3 in the example) a good leaving group.

Figure 23.3 summarizes the three possible outcomes of halogenation at the α carbon, depending on the substrate and chosen reaction conditions.

Problem 23.13

Draw the products of each reaction. Assume excess halogen is present.





Figure 23.3


Summary: Halogenation reactions at the α carbon to a carbonyl group

[1] Halogenation in acid




monosubstitution on the α carbon

[2] General halogenation in base



polysubstitution on the α carbon

[3] Halogenation of *methyl* ketones with excess X_2 and base




oxidative cleavage

23.7C Reactions of α -Halo Carbonyl Compounds

α -Halo carbonyl compounds undergo two useful reactions—**elimination** with base and **substitution** with nucleophiles.

For example, treatment of 2-bromocyclohexanone with the base Li_2CO_3 in the presence of LiBr in the polar aprotic solvent DMF [$\text{HCON}(\text{CH}_3)_2$] affords 2-cyclohexenone by elimination of the elements of Br and H from the α and β carbons, respectively. Thus, a two-step method can convert a carbonyl compound such as cyclohexanone into an α,β -unsaturated carbonyl compound such as 2-cyclohexenone.




α,β -Unsaturated carbonyl compounds undergo a variety of 1,2- and 1,4-addition reactions as discussed in Section 20.15.

[1] **Bromination at the α carbon is accomplished with Br_2 in CH_3COOH .**

[2] **Elimination of Br and H occurs with Li_2CO_3 and LiBr in DMF.**

α -Halo carbonyl compounds also react with nucleophiles by $\text{S}_{\text{N}}2$ reactions. For example, reaction of 2-bromocyclohexanone with CH_3NH_2 affords the substitution product **A**. A related intramolecular nucleophilic substitution of an α -halo ketone was a key step in the synthesis of the antimalarial drug quinine, as shown in Figure 23.4.



Problem 23.14 Draw the organic products formed when 2-bromo-3-pentanone ($\text{CH}_3\text{CH}_2\text{COCHBrCH}_3$) is treated with each reagent: (a) Li_2CO_3 , LiBr , DMF; (b) $\text{CH}_3\text{CH}_2\text{NH}_2$; (c) CH_3SH .

Problem 23.15 Identify the product **M** of the following two-step reaction sequence. **M** was converted to the hallucinogen LSD (Chapter 18 opening molecule) in several steps.





Figure 23.4


Intramolecular nucleophilic substitution in the synthesis of quinine



- Intramolecular $\text{S}_{\text{N}}2$ reaction of a nitrogen nucleophile with an α -halo ketone affords a compound that can be converted to quinine in a single step. The new $\text{C}-\text{N}$ bond on the α carbon is labeled in red.


23.8 Direct Enolate Alkylation

Treatment of an aldehyde or ketone with base and an alkyl halide (RX) results in **alkylation**—the **substitution of R for H on the α carbon atom**. Alkylation forms a new carbon–carbon bond on the α carbon.



23.8A General Features

We will begin with the most direct method of alkylation, and then (in Sections 23.9 and 23.10) examine two older, multistep methods that are still used today. Direct alkylation is carried out by a two-step process:




[1] **Deprotonation:** Base removes a proton from the α carbon to generate an enolate. The reaction works best with a strong nonnucleophilic base like LDA in THF solution at low temperature ($-78\text{ }^\circ\text{C}$).


[2] **Nucleophilic attack:** The nucleophilic enolate attacks the alkyl halide, displacing the halide (a good leaving group) and forming the alkylation product by an $\text{S}_{\text{N}}2$ reaction.

Because Step [2] is an $\text{S}_{\text{N}}2$ reaction, it works best with unhindered methyl and 1° alkyl halides. Hindered alkyl halides and those with halogens bonded to sp^2 hybridized carbons do not undergo substitution.


R_3CX , $\text{CH}_2=\text{CHX}$, and $\text{C}_6\text{H}_5\text{X}$ do not undergo alkylation reactions with enolates, because they are unreactive in $\text{S}_{\text{N}}2$ reactions.




Ester enolates and carbanions derived from nitriles are also alkylated under these conditions.




Problem 23.16 What product is formed when each compound is treated first with LDA in THF solution at low temperature, followed by $\text{CH}_3\text{CH}_2\text{I}$?




The stereochemistry of enolate alkylation follows the general rule governing the stereochemistry of reactions: **an achiral starting material yields an achiral or racemic product.** For example, when cyclohexanone (an achiral starting material) is converted to 2-ethylcyclohexanone by treatment with base and $\text{CH}_3\text{CH}_2\text{I}$, a new stereogenic center is introduced, and both enantiomers of the product are formed in equal amounts—that is, a **racemic mixture**.



Problem 23.17 Draw the products obtained (including stereochemistry) when each compound is treated with LDA, followed by CH_3I .



Problem 23.18 The analgesic naproxen can be prepared by a stepwise reaction sequence from ester **A**. Using enolate alkylation in one step, what reagents are needed to convert **A** to naproxen? Write the structure of each intermediate. Explain why a racemic product is formed.




23.8B Alkylation of Unsymmetrical Ketones


An unsymmetrical ketone can be regioselectively alkylated to yield one major product. The strategy depends on the use of the appropriate base, solvent, and temperature to form the kinetic or thermodynamic enolate (Section 23.4), which is then treated with an alkyl halide to form the alkylation product.

For example, 2-methylcyclohexanone can be converted to either 2,6-dimethylcyclohexanone (**A**) or 2,2-dimethylcyclohexanone (**B**) by proper choice of reaction conditions.


- **Treatment of 2-methylcyclohexanone with LDA in THF solution at -78°C gives the less substituted kinetic enolate, which then reacts with CH_3I to form **A**.**




- Treatment of 2-methylcyclohexanone with $\text{NaOCH}_2\text{CH}_3$ in $\text{CH}_3\text{CH}_2\text{OH}$ solution at room temperature forms the more substituted thermodynamic enolate, which then reacts with CH_3I to form **B**.



Problem 23.19 How can 2-pentanone be converted into each compound?




23.8C Application of Enolate Alkylation: Tamoxifen Synthesis




Tamoxifen has been commercially available since the 1970s, sold under the brand name of Nolvadex.

Tamoxifen, the chapter-opening molecule, is a potent anticancer drug that has been used to treat certain forms of breast cancer for many years. One step in the synthesis of tamoxifen involves the treatment of ketone **A** with NaH as base to form an enolate. Alkylation of this enolate with $\text{CH}_3\text{CH}_2\text{I}$ forms **B** in high yield. **B** is converted to tamoxifen in several steps, some of which are reactions you have already learned.



Problem 23.20 Identify **A**, **B**, and **C**, intermediates in the synthesis of the five-membered ring called an α -methylene- γ -butyrolactone. This heterocyclic ring system is present in some antitumor agents.




23.9 Malonic Ester Synthesis

Besides the direct method of enolate alkylation discussed in Section 23.8, a new alkyl group can also be introduced on the α carbon using the malonic ester synthesis and the acetoacetic ester synthesis.

- The malonic ester synthesis prepares carboxylic acids having two general structures:




- The acetoacetic ester synthesis prepares methyl ketones having two general structures:




23.9A Background for the Malonic Ester Synthesis


The malonic ester synthesis is a stepwise method for converting diethyl malonate into a carboxylic acid having one or two alkyl groups on the α carbon. To simplify the structures, the CH_3CH_2 groups of the esters are abbreviated as Et.




Before writing out the steps in the malonic ester synthesis, recall from Section 22.11 that esters are hydrolyzed by aqueous acid. Thus, heating diethyl malonate with acid and water hydrolyzes both esters to carboxy groups, forming a β -diacid (1,3-diacid).




β -Diacids are unstable to heat. They **decarboxylate** (lose CO_2), resulting in cleavage of a carbon-carbon bond and formation of a carboxylic acid. Decarboxylation is not a general reaction of all carboxylic acids. It occurs with β -diacids, however, because CO_2 can be eliminated through a cyclic, six-atom transition state. This forms an enol of a carboxylic acid, which in turn tautomerizes to the more stable keto form.




The net result of decarboxylation is cleavage of a carbon-carbon bond on the α carbon, with loss of CO_2 .



Decarboxylation occurs readily whenever a carboxy group (COOH) is bonded to the α carbon of another carbonyl group. For example, β -keto acids also readily lose CO_2 on heating to form ketones.




Problem 23.21 Which of the following compounds will readily lose CO_2 when heated?




23.9B Steps in the Malonic Ester Synthesis

The malonic ester synthesis converts diethyl malonate to a carboxylic acid in three steps.




[1] **Deprotonation.** Treatment of diethyl malonate with ^-OEt removes the acidic α proton between the two carbonyl groups. Recall from Section 23.3A that these protons are more acidic than other α protons because three resonance structures can be drawn for the enolate, instead of the usual two. Thus, ^-OEt , rather than the stronger base LDA, can be used for this reaction.




[2] **Alkylation.** The nucleophilic enolate reacts with an alkyl halide in an $\text{S}_{\text{N}}2$ reaction to form a substitution product. Because the mechanism is $\text{S}_{\text{N}}2$, the yields are higher when R is CH_3 or a 1° alkyl group.

[3] **Hydrolysis and decarboxylation.** Heating the diester with aqueous acid hydrolyzes the diester to a β -diacid, which loses CO_2 to form a carboxylic acid.


The synthesis of butanoic acid ($\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$) from diethyl malonate illustrates the basic process:





If the first two steps of the reaction sequence are repeated *prior* to hydrolysis and decarboxylation, then a carboxylic acid having *two new alkyl groups* on the α carbon can be synthesized. This is illustrated in the synthesis of 2-benzylbutanoic acid [$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_2\text{C}_6\text{H}_5)\text{COOH}$] from diethyl malonate:



An intramolecular malonic ester synthesis can be used to form rings having three to six atoms, provided the appropriate dihalide is used as starting material. For example, cyclopentanecarboxylic acid can be prepared from diethyl malonate and 1,4-dibromobutane ($\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$) by the following sequence of reactions:



Problem 23.22 Draw the products of each reaction.




Problem 23.23 What cyclic product is formed from each dihalide using the malonic ester synthesis: (a) $\text{ClCH}_2\text{CH}_2\text{CH}_2\text{Cl}$; (b) $(\text{BrCH}_2\text{CH}_2)_2\text{O}$?

23.9C Retrosynthetic Analysis

To use the malonic ester synthesis you must be able to determine what starting materials are needed to prepare a given compound—that is, you must **work backwards in the retrosynthetic direction**. This involves a two-step process:


- [1] Locate the α carbon to the COOH group, and identify all alkyl groups bonded to the α carbon.
- [2] Break the molecule into two (or three) components: Each alkyl group bonded to the α carbon comes from an alkyl halide. The remainder of the molecule comes from $\text{CH}_2(\text{COOEt})_2$.

**Sample Problem 23.3**


What starting materials are needed to prepare 2-methylhexanoic acid [$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)\text{COOH}$] using a malonic ester synthesis?

Solution


The target molecule has two different alkyl groups bonded to the α carbon, so three components are needed for the synthesis:



Writing the synthesis in the synthetic direction:

**Problem 23.24**


What alkyl halides are needed to prepare each carboxylic acid by the malonic ester synthesis?

**Problem 23.25**


Explain why each of the following carboxylic acids cannot be prepared by a malonic ester synthesis: (a) $(\text{CH}_3)_3\text{CCH}_2\text{COOH}$; (b) $\text{C}_6\text{H}_5\text{CH}_2\text{COOH}$; (c) $(\text{CH}_3)_3\text{CCOOH}$.

23.10 Acetoacetic Ester Synthesis


The acetoacetic ester synthesis is a stepwise method for converting ethyl acetoacetate into a ketone having one or two alkyl groups on the α carbon.



23.10A Steps in the Acetoacetic Ester Synthesis




The steps in the acetoacetic ester synthesis are exactly the same as those in the malonic ester synthesis. Because the starting material, $\text{CH}_3\text{COCH}_2\text{COOEt}$, is a β -keto ester, the final product is a **ketone**, not a carboxylic acid.




- [1] **Deprotonation.** Treatment of ethyl acetoacetate with OEt^- removes the acidic proton between the two carbonyl groups.
- [2] **Alkylation.** The nucleophilic enolate reacts with an alkyl halide (RX) in an $\text{S}_{\text{N}}2$ reaction to form a substitution product. Because the mechanism is $\text{S}_{\text{N}}2$, the yields are higher when R is CH_3 or a 1° alkyl group.
- [3] **Hydrolysis and decarboxylation.** Heating the β -keto ester with aqueous acid hydrolyzes the ester to a β -keto acid, which loses CO_2 to form a ketone.

If the first two steps of the reaction sequence are repeated *prior* to hydrolysis and decarboxylation, then a ketone having *two new alkyl groups* on the α carbon can be synthesized.




Problem 23.26 What ketones are prepared by the following reactions?




To determine what starting materials are needed to prepare a given ketone using the acetoacetic ester synthesis, you must again work in the **retrosynthetic** direction. This involves a two-step process:

- [1] Identify the alkyl groups bonded to the α carbon to the carbonyl group.
- [2] Break the molecule into two (or three) components: Each alkyl group bonded to the α carbon comes from an alkyl halide. The remainder of the molecule comes from $\text{CH}_3\text{COCH}_2\text{COOEt}$.




Sample Problem 23.4

What starting materials are needed to synthesize 2-heptanone using the acetoacetic ester synthesis?




Solution

2-Heptanone has only one alkyl group bonded to the α carbon, so only one alkyl halide is needed in the acetoacetic ester synthesis.






Writing the acetoacetic ester synthesis in the synthetic direction:



Problem 23.27

What alkyl halides are needed to prepare each ketone using the acetoacetic ester synthesis?




Problem 23.28




The jack-o'-lantern, source of the antitumor agent illudin-S


Treatment of ethyl acetoacetate with NaOEt (2 equiv) and $\text{BrCH}_2\text{CH}_2\text{Br}$ forms compound **X**. This reaction is the first step in the synthesis of illudin-S, an antitumor substance isolated from the jack-o'-lantern, a poisonous, saffron-colored mushroom. What is the structure of **X**?



The acetoacetic ester synthesis and direct enolate alkylation are two different methods that prepare similar ketones. 2-Butanone, for example, can be synthesized from acetone by direct enolate alkylation with CH_3I (Method [1]), or by alkylation of ethyl acetoacetate followed by hydrolysis and decarboxylation (Method [2]).

Method [1]
Direct enolate alkylation

The same product is formed by two different routes.


Method [2]
Acetoacetic ester synthesis

Why would you ever make 2-butanone from ethyl acetoacetate when you could make it in fewer steps from acetone? There are many factors to consider. First of all, synthetic organic chemists like to have a variety of methods to accomplish a single kind of reaction. Sometimes subtle changes in the structure of a starting material make one reaction work better than another.

In the chemical industry, moreover, cost is an important issue. Any reaction needed to make a large quantity of a useful drug or other consumer product must use cheap starting materials. Direct enolate alkylation usually requires a very strong base like LDA to be successful, whereas the acetoacetic ester synthesis utilizes NaOEt. NaOEt can be prepared from cheaper starting materials, and this makes the acetoacetic ester synthesis an attractive method, even though it involves more steps.

Thus, each method has its own advantages and disadvantages, depending on the starting material, the availability of reagents, the cost, and the occurrence of side reactions.

Problem 23.29 Nabumetone is a pain reliever and anti-inflammatory agent sold under the brand name of Relafen.




- Write out a synthesis of nabumetone from ethyl acetoacetate.
- What ketone and alkyl halide are needed to synthesize nabumetone by direct enolate alkylation?

KEY CONCEPTS


Substitution Reactions of Carbonyl Compounds at the α Carbon

Kinetic Versus Thermodynamic Enolates (23.4)



Kinetic enolate

- The less substituted enolate
- Favored by strong base, polar aprotic solvent, low temperature: LDA, THF, $-78\text{ }^\circ\text{C}$




Thermodynamic enolate

- The more substituted enolate
- Favored by strong base, protic solvent, higher temperature: NaOCH₂CH₃, CH₃CH₂OH, room temperature


Halogenation at the α Carbon

[1] Halogenation in acid (23.7A)




- The reaction occurs via enol intermediates.
- Monosubstitution of X for H occurs on the α carbon.

[2] Halogenation in base (23.7B)



- The reaction occurs via enolate intermediates.
- Polysubstitution of X for H occurs on the α carbon.


[3] Halogenation of *methyl* ketones in base—The haloform reaction (23.7B)



- The reaction occurs with methyl ketones and results in cleavage of a carbon–carbon σ bond.


Reactions of α -Halo Carbonyl Compounds (23.7C)

[1] Elimination to form α,β -unsaturated carbonyl compounds



- Elimination of the elements of Br and H forms a new π bond, giving an α,β -unsaturated carbonyl compound.

[2] Nucleophilic substitution



- The reaction follows an $\text{S}_{\text{N}}2$ mechanism, generating an α -substituted carbonyl compound.


Alkylation Reactions at the α Carbon

[1] Direct alkylation at the α carbon (23.8)




- The reaction forms a new C–C bond to the α carbon.
- LDA is a common base used to form an intermediate enolate.
- The alkylation in Step [2] follows an $\text{S}_{\text{N}}2$ mechanism.

[2] Malonic ester synthesis (23.9)



- The reaction is used to prepare carboxylic acids with one or two alkyl groups on the α carbon.
- The alkylation in Step [2] follows an $\text{S}_{\text{N}}2$ mechanism.

[3] Acetoacetic ester synthesis (23.10)









- The reaction is used to prepare ketones with one or two alkyl groups on the α carbon.
- The alkylation in Step [2] follows an $\text{S}_{\text{N}}2$ mechanism.


PROBLEMS

Enols, Enolates, and Acidic Protons


23.30 Draw enol tautomer(s) for each compound.



23.31 Both 2,4-pentanedione and ethyl acetoacetate have two carbonyl groups separated by a single carbon atom. Although an equilibrium mixture of 2,4-pentanedione tautomers contains 76% of the enol forms, an equilibrium mixture of ethyl acetoacetate tautomers contains only 8% of the enol forms. Suggest a reason for this difference.









2,4-pentanedione








ethyl acetoacetate







23.32 What hydrogen atoms in each compound have a $\text{p}K_a \leq 25$?



23.33 Rank the labeled protons in each compound in order of increasing acidity.




23.34 What is the major enolate (or carbanion) formed when each compound is treated with LDA?




23.35 How could IR spectroscopy be used to detect the presence of enol tautomers?


23.36 Why is the pK_a of the H_a protons in 1-acetylcyclohexene higher than the pK_a of the H_b protons?




23.37 Explain why 5,5-dimethyl-1,3-cyclohexanedione exists predominantly in its enol form, but 2,2-dimethyl-1,3-cyclohexanedione does not.



23.38 Explain why an optically active solution of (*R*)- α -methylbutyrophenone loses its optical activity when either dilute acid or base is added to the solution.




23.39 Although ibuprofen is sold as a racemic mixture, only the *S* enantiomer acts as an analgesic. In the body, however, some of the *R* enantiomer is converted to the *S* isomer by tautomerization to an enol and then protonation to regenerate the carbonyl compound. Write a stepwise mechanism for this isomerization.




23.40 Explain why the α protons of an ester are less acidic than the α protons of a ketone by ~ 5 pK_a units.

23.41 Explain why reactions that use LDA as base must be carried out under anhydrous conditions; that is, all traces of H_2O must be rigorously excluded.


23.42 Explain why 2,4-pentanedione forms two different alkylation products (**A** or **B**) when the number of equivalents of base is increased from one to two.



23.43 The cis ketone **A** is isomerized to the trans ketone **B** with aqueous NaOH. A similar isomerization reaction does not occur with the cis ketone **C**. Explain this difference in reactivity.




23.44 Treatment of α,β -unsaturated carbonyl compound **X** with base forms the diastereomer **Y**. Write a stepwise mechanism for this reaction. Explain why one stereogenic center changes configuration but the other does not.



Halogenation


23.45 Acid-catalyzed bromination of 2-pentanone ($\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_3$) forms two products: $\text{BrCH}_2\text{COCH}_2\text{CH}_2\text{CH}_3$ (**A**) and $\text{CH}_3\text{COCH}(\text{Br})\text{CH}_2\text{CH}_3$ (**B**). Explain why the major product is **B**, with the Br atom on the more substituted side of the carbonyl group.

23.46 Draw a stepwise mechanism for each reaction.




Malonic Ester Synthesis

23.47 What alkyl halides are needed to prepare each carboxylic acid using the malonic ester synthesis?




23.48 Use the malonic ester synthesis to prepare each carboxylic acid.



23.49 Devise a synthesis of valproic acid [$(\text{CH}_3\text{CH}_2\text{CH}_2)_2\text{CHCO}_2\text{H}$], a medicine used to treat epileptic seizures, using the malonic ester synthesis.

23.50 Synthesize each compound from diethyl malonate. You may use any other organic or inorganic reagents.




23.51 The enolate derived from diethyl malonate reacts with a variety of electrophiles (not just alkyl halides) to form new carbon-carbon bonds. With this in mind, draw the products formed when $\text{Na}^+\text{CH}(\text{COOEt})_2$ reacts with each electrophile, followed by treatment with H_2O .



Acetoacetic Ester Synthesis

23.52 What alkyl halides are needed to prepare each ketone using the acetoacetic ester synthesis?




23.53 Synthesize each compound from ethyl acetoacetate. You may use any other organic or inorganic reagents.




Reactions


23.54 Draw the organic products formed in each reaction.




23.55 Draw the products formed (including stereoisomers) in each reaction.




23.56 a. Identify intermediates **A–C** in the following stepwise conversion of *p*-isobutylbenzaldehyde to the analgesic ibuprofen.




b. Direct alkylation of **D** by treatment with one equivalent of LDA and CH_3I does not form ibuprofen. Identify the product of this reaction and explain how it is formed.




- 23.57 a. Clopidogrel is the generic name for Plavix, a drug used to prevent the formation of blood clots in patients that have a history of heart attacks or strokes. The racemic drug can be prepared from the racemic α -halo ester by the following reaction. What is the structure of clopidogrel?




- b. A single enantiomer of clopidogrel can be prepared in three steps from the chiral α -hydroxy acid **A**. Identify **B** and **C** in the following reaction sequence, and designate the configuration of the enantiomer formed by this route as *R* or *S*.



- 23.58 What reaction conditions—base, solvent, and temperature—are needed to convert ketone **A** to either **B** or **C** by an intramolecular alkylation reaction?




- 23.59 Explain why each of the following reactions will **not** proceed as written.




Mechanism


- 23.60 Draw a stepwise mechanism showing how two alkylation products are formed in the following reaction.




- 23.61 Draw a stepwise mechanism for the following reaction.



23.62 Draw stepwise mechanisms illustrating how each product is formed.




23.63 A key step in the synthesis of β -vetivone, a major constituent of vetiver, a perennial grass found in tropical and subtropical regions of the world, involved the reaction of compound **A** and dihalide **B** with two equivalents of LDA to form **C**. Draw a stepwise mechanism for this reaction. β -Vetivone contains a spiro ring system—that is, two rings that share a single carbon atom.




Synthesis


23.64 Convert acetophenone ($\text{C}_6\text{H}_5\text{COCH}_3$) into each of the following compounds. You may use any other organic compounds or required inorganic reagents. More than one step may be required.




23.65 Synthesize each compound from cyclohexanone and organic halides having ≤ 4 C's. You may use any other inorganic reagents.




23.66 Bupropion, sold under the trade name of Zyban, is an antidepressant that was approved to aid smoking cessation in 1997. Devise a synthesis of bupropion from benzene, organic compounds that have fewer than five carbons, and any required inorganic reagents.




23.67 Synthesize each product from ethyl acetoacetate ($\text{CH}_3\text{COCH}_2\text{CO}_2\text{Et}$) and the given starting material. You may also use any other organic compounds or required inorganic reagents.




23.68 Synthesize each compound from 3-pentanone [$(\text{CH}_3\text{CH}_2)_2\text{C}=\text{O}$]. You may also use benzene, organic alcohols having ≤ 3 C's, and any required inorganic reagents.



23.69 Treatment of ketone **A** with LDA followed by $\text{CH}_3\text{CH}_2\text{I}$ did not form the desired alkylation product **B**. What product was formed instead? Devise a multistep method to convert **A** to **B**, a synthetic intermediate used to prepare the anticancer drug tamoxifen (Section 23.8C and the chapter-opening molecule).





23.70 Capsaicin, the spicy component of hot peppers, can be prepared from amine **X** and acid chloride **Y**. Devise a synthesis of **Y** from (4*E*)-6-methyl-4-hepten-1-ol [$(\text{CH}_3)_2\text{CHCH}=\text{CH}(\text{CH}_2)_3\text{OH}$], $\text{CH}_2(\text{CO}_2\text{Et})_2$, and any required inorganic reagents.




Spectroscopy

- 23.71** Treatment of **W** with CH_3Li , followed by CH_3I , affords compound **Y** ($\text{C}_7\text{H}_{14}\text{O}$) as the major product. **Y** shows a strong absorption in its IR spectrum at 1713 cm^{-1} , and its $^1\text{H NMR}$ spectrum is given below. (a) Propose a structure for **Y**. (b) Draw a stepwise mechanism for the conversion of **W** to **Y**.




Challenge Problems


- 23.72** Explain why H_a is much less acidic than H_b . Then draw a mechanism for the following reaction.



- 23.73** The last step in the synthesis of β -vetivone (Problem 23.63) involves treatment of **C** with CH_3Li to form an intermediate **X**, which forms β -vetivone with aqueous acid. Identify the structure of **X** and draw a mechanism for converting **X** to β -vetivone.



- 23.74** Keeping in mind the mechanism for the dissolving metal reduction of alkynes to trans alkenes in Chapter 12, write a stepwise mechanism for the following reaction, which involves the conversion of an α,β -unsaturated carbonyl compound to a carbonyl compound with a new alkyl group on the α carbon.



24

Carbonyl Condensation Reactions

- 24.1 The aldol reaction
- 24.2 Crossed aldol reactions
- 24.3 Directed aldol reactions
- 24.4 Intramolecular aldol reactions
- 24.5 The Claisen reaction
- 24.6 The crossed Claisen and related reactions
- 24.7 The Dieckmann reaction
- 24.8 The Michael reaction
- 24.9 The Robinson annulation




Ibuprofen is the generic name for the pain reliever known by the trade names of Motrin and Advil. Like aspirin, ibuprofen acts as an anti-inflammatory agent by blocking the synthesis of prostaglandins from arachidonic acid. One step in a commercial synthesis of ibuprofen involves the reaction of a nucleophilic enolate with an electrophilic carbonyl group. In Chapter 24, we learn about the carbon-carbon bond-forming reactions of enolates with carbonyl electrophiles.

In Chapter 24, we examine **carbonyl condensations**—that is, reactions between two carbonyl compounds—a second type of reaction that occurs at the α carbon of a carbonyl group. Much of what is presented in Chapter 24 applies principles you have already learned. Many of the reactions may look more complicated than those in previous chapters, but they are fundamentally the same. Nucleophiles attack electrophilic carbonyl groups to form the products of nucleophilic addition or substitution, depending on the structure of the carbonyl starting material.

Every reaction in Chapter 24 forms a new carbon–carbon bond at the α carbon to a carbonyl group, so these reactions are extremely useful in the synthesis of complex natural products.

24.1 The Aldol Reaction

Chapter 24 concentrates on the second general reaction of enolates—**reaction with other carbonyl compounds**. In these reactions, one carbonyl component serves as the nucleophile and one serves as the electrophile, and a new carbon–carbon bond is formed.




The presence or absence of a leaving group on the electrophilic carbonyl carbon determines the structure of the product. Even though they appear somewhat more complicated, these reactions are often reminiscent of the nucleophilic addition and nucleophilic acyl substitution reactions of Chapters 21 and 22. Four types of reactions are examined:

- **Aldol reaction** (Sections 24.1–24.4)
- **Claisen reaction** (Sections 24.5–24.7)
- **Michael reaction** (Section 24.8)
- **Robinson annulation** (Section 24.9)

24.1A General Features of the Aldol Reaction

In the **aldol reaction**, two molecules of an aldehyde or ketone react with each other in the presence of base to form a **β -hydroxy carbonyl compound**. For example, treatment of acetaldehyde with aqueous OH^- forms 3-hydroxybutanal, a **β -hydroxy aldehyde**.

Many aldol products contain an **aldehyde** and an **alcohol**—hence the name **aldol**.




The mechanism of the aldol reaction has **three steps**, as shown in Mechanism 24.1. Carbon–carbon bond formation occurs in Step [2], when the nucleophilic enolate reacts with the electrophilic carbonyl carbon.




Mechanism 24.1 The Aldol Reaction

Step [1] Formation of a nucleophilic enolate



- In Step [1], the base removes a proton from the α carbon to form a **resonance-stabilized enolate**.

Steps [2]–[3] Nucleophilic addition and protonation



- In Step [2], the nucleophilic enolate attacks the electrophilic carbonyl carbon of another molecule of aldehyde, thus forming a new carbon–carbon bond. **This joins the α carbon of one aldehyde to the carbonyl carbon of a second aldehyde.**
- Protonation of the alkoxide in Step [3] forms the **β -hydroxy aldehyde**.

The aldol reaction is a reversible equilibrium, so the position of the equilibrium depends on the base and the carbonyl compound. OH^- is the base typically used in an aldol reaction. Recall from Section 23.3B that only a small amount of enolate forms with OH^- . In this case, that's appropriate because the starting aldehyde is needed to react with the enolate in the second step of the mechanism.

Aldol reactions can be carried out with either aldehydes or ketones. With aldehydes, the equilibrium usually favors the products, but with ketones the equilibrium favors the starting materials. There are ways of driving this equilibrium to the right, however, so we will write aldol products whether the substrate is an aldehyde or a ketone.


- The characteristic reaction of aldehydes and ketones is nucleophilic addition (Section 21.7). An aldol reaction is a nucleophilic addition in which an enolate is the nucleophile. See the comparison in Figure 24.1.

A **second example of an aldol** reaction is shown with propanal as starting material. The two molecules of the aldehyde that participate in the aldol reaction react in opposite ways.


- One molecule of propanal becomes an enolate—an electron-rich *nucleophile*.
- One molecule of propanal serves as the *electrophile* because its carbonyl carbon is electron deficient.

Figure 24.1


The aldol reaction—An example of nucleophilic addition




- Aldehydes and ketones react by nucleophilic addition. In an aldol reaction, an enolate is the nucleophile that adds to the carbonyl group.




These two examples illustrate the general features of the aldol reaction. **The α carbon of one carbonyl component becomes bonded to the carbonyl carbon of the other component.**



Problem 24.1 Draw the aldol product formed from each compound.




Problem 24.2 Which carbonyl compounds do *not* undergo an aldol reaction when treated with OH^- in H_2O ?




24.1B Dehydration of the Aldol Product

The β -hydroxy carbonyl compounds formed in the aldol reaction dehydrate more readily than other alcohols. In fact, under the basic reaction conditions, the initial aldol product is often not isolated. Instead, **it loses the elements of H_2O from the α and β carbons to form an α,β -unsaturated carbonyl compound.**

All alcohols—including β -hydroxy carbonyl compounds—dehydrate in the presence of acid. Only β -hydroxy carbonyl compounds dehydrate in the presence of base.



An aldol reaction is often called an **aldol condensation**, because the β -hydroxy carbonyl compound that is initially formed loses H_2O by dehydration. **A condensation reaction is one in which a small molecule, in this case H_2O , is eliminated during a reaction.**




It may or may not be possible to isolate the β -hydroxy carbonyl compound under the conditions of the aldol reaction. When the α,β -unsaturated carbonyl compound is further conjugated with a carbon-carbon double bond or a benzene ring, as in the case of Reaction [2], **elimination of H_2O is spontaneous** and the β -hydroxy carbonyl compound cannot be isolated.

The mechanism of dehydration consists of two steps: **deprotonation followed by loss of OH^-** , as shown in Mechanism 24.2.



Mechanism 24.2 Dehydration of β -Hydroxy Carbonyl Compounds with Base



- In Step [1], base removes a proton from the α carbon, thus forming a resonance-stabilized enolate.
- In Step [2], the electron pair of the enolate forms the π bond as OH^- is eliminated.


Like E1 elimination, E1cB requires two steps. Unlike E1, though, the intermediate in E1cB is a *carbanion*, not a carbocation. E1cB stands for **Elimination, unimolecular, conjugate base**.

This elimination mechanism, called the **E1cB mechanism**, differs from the two more general mechanisms of elimination, E1 and E2, which were discussed in Chapter 8. The E1cB mechanism involves two steps, and proceeds by way of an **anionic** intermediate.


Regular alcohols dehydrate only in the presence of acid but not base, because hydroxide is a poor leaving group. When the hydroxy group is β to a carbonyl group, however, loss of H and OH from the α and β carbons forms a **conjugated double bond**, and the stability of the conjugated system makes up for having such a poor leaving group.

Dehydration of the initial β -hydroxy carbonyl compound drives the equilibrium of an aldol reaction to the right, thus favoring product formation. Once the conjugated α,β -unsaturated carbonyl compound forms, it is *not* re-converted to the β -hydroxy carbonyl compound.

Problem 24.3 What unsaturated carbonyl compound is formed by dehydration of each β -hydroxy carbonyl compound?



Problem 24.4 Acid-catalyzed dehydration of β -hydroxy carbonyl compounds occurs by the mechanism discussed in Section 9.8. With this in mind, draw a stepwise mechanism for the following reaction.




24.1C Retrosynthetic Analysis

To utilize the aldol reaction in synthesis, you must be able to determine which aldehyde or ketone is needed to prepare a particular β -hydroxy carbonyl compound or α,β -unsaturated carbonyl compound—that is, you must be able to **work backwards, in the retrosynthetic direction**.

HOW TO Synthesize a Compound Using the Aldol Reaction

Example What starting material is needed to prepare each compound by an aldol reaction?




Step [1] Locate the α and β carbons of the carbonyl group.




- When a carbonyl group has two different α carbons, choose the side that contains the OH group (in a β -hydroxy carbonyl compound) or is part of the C=C (in an α,β -unsaturated carbonyl compound).

Step [2] Break the molecule into two components between the α and β carbons.

- The α carbon and all remaining atoms bonded to it belong to one carbonyl component. The β carbon and all remaining atoms bonded to it belong to the other carbonyl component. Both components are identical in all aldols we have thus far examined.



Problem 24.5 What aldehyde or ketone is needed to prepare each compound by an aldol reaction?




24.2 Crossed Aldol Reactions

In all of the aldol reactions discussed so far, the electrophilic carbonyl and the nucleophilic enolate have originated from the *same* aldehyde or ketone. Sometimes, though, it is possible to carry out an aldol reaction between two *different* carbonyl compounds.

- An aldol reaction between two different carbonyl compounds is called a *crossed aldol* or *mixed aldol reaction*.

24.2A A Crossed Aldol Reaction with Two Different Aldehydes, Both Having α H Atoms

When two different aldehydes, both having α H atoms, are combined in an aldol reaction, four different β -hydroxy carbonyl compounds are formed. Four products form, not one, because both aldehydes can lose an acidic α hydrogen atom and form an enolate in the presence of base. Both enolates can then react with both carbonyl compounds, as shown for acetaldehyde and propanal in the following reaction scheme.



- **Conclusion:** When two different aldehydes have α hydrogens, a crossed aldol reaction is *not* synthetically useful.


24.2B Synthetically Useful Crossed Aldol Reactions

Crossed aldols are synthetically useful in two different situations.

- A crossed aldol occurs when only one carbonyl component has α H atoms.


When one carbonyl compound has no α hydrogens, a crossed aldol reaction often leads to one product. Two common carbonyl compounds with no α hydrogens used for this purpose are **formaldehyde ($\text{CH}_2=\text{O}$)** and **benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$)**.

For example, reaction of $\text{C}_6\text{H}_5\text{CHO}$ (as the electrophile) with either acetaldehyde (CH_3CHO) or acetone [$(\text{CH}_3)_2\text{C}=\text{O}$] in the presence of base forms a single α,β -unsaturated carbonyl compound after dehydration.




The yield of a single crossed aldol product is increased further if the electrophilic carbonyl component is relatively unhindered (as is the case with most aldehydes), and if it is used in excess.

Problem 24.6 2-Pentylcinnamaldehyde, commonly called flosal, is a perfume ingredient with a jasmine-like odor. Flosal is an α,β -unsaturated aldehyde made by a crossed aldol reaction between benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$) and heptanal ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CHO}$), followed by dehydration. Draw a stepwise mechanism for the following reaction that prepares flosal.




Problem 24.7 Draw the products formed in each crossed aldol reaction.



- A crossed aldol occurs when one carbonyl component has especially acidic α H atoms.


A useful crossed aldol reaction takes place between an aldehyde or ketone and a β -dicarbonyl (or similar) compound.







As we learned in Section 23.3, the α hydrogens between two carbonyl groups are especially acidic, and so they are more readily removed than other α H atoms. As a result, **the β -dicarbonyl compound always becomes the enolate component of the aldol reaction.** Figure 24.2 shows the steps for the crossed aldol reaction between diethyl malonate and benzaldehyde. In this type of crossed aldol reaction, the initial β -hydroxy carbonyl compound *always* loses water to form the highly conjugated product.

β -Dicarbonyl compounds are sometimes called **active methylene compounds** because they are more reactive towards base than other carbonyl compounds. **1,3-Dinitriles** and **α -cyano carbonyl compounds** are also active methylene compounds.

Figure 24.2
Crossed aldol reaction
between benzaldehyde and
 $\text{CH}_2(\text{COOEt})_2$



Active methylene compounds



Problem 24.8 Draw the products formed in the crossed aldol reaction of phenylacetaldehyde ($\text{C}_6\text{H}_5\text{CH}_2\text{CHO}$) with each compound: (a) $\text{CH}_2(\text{COOEt})_2$; (b) $\text{CH}_2(\text{COCH}_3)_2$; (c) $\text{CH}_3\text{COCH}_2\text{CN}$.

24.2C Useful Transformations of Aldol Products

The aldol reaction is synthetically useful because it forms new carbon–carbon bonds, generating products with two functional groups. Moreover, the β-hydroxy carbonyl compounds formed in aldol reactions are readily transformed into a variety of other compounds. Figure 24.3 illustrates how the crossed aldol product obtained from cyclohexanone and formaldehyde ($\text{CH}_2=\text{O}$) can be converted to other compounds by reactions learned in earlier chapters.

Problem 24.9 What aldol product is formed when two molecules of butanal react together in the presence of base? What reagents are needed to convert this product to each of the following compounds?







Figure 24.3

Conversion of a β-hydroxy carbonyl compound into other compounds



- The β-hydroxy carbonyl compound formed from the crossed aldol reaction can be reduced with NaBH_4 , CH_3OH (Section 20.4A) to form a 1,3-diol (Reaction [1]) or dehydrated to form an α,β-unsaturated carbonyl compound (Reaction [2]).
- Reduction of the α,β-unsaturated carbonyl compound forms an allylic alcohol with NaBH_4 (Reaction [3]), or a ketone with H_2 and Pd-C (Reaction [4]); see Section 20.4C.
- Reaction of the α,β-unsaturated carbonyl compound with an organometallic reagent forms two different products depending on the choice of RM (Reaction [5]); see Section 20.15.

24.3 Directed Aldol Reactions


A **directed aldol reaction** is a variation of the crossed aldol reaction that clearly defines which carbonyl compound becomes the nucleophilic enolate and which reacts at the electrophilic carbonyl carbon. The strategy of a directed aldol reaction is as follows:

- [1] Prepare the enolate of one carbonyl component with LDA.
- [2] Add the second carbonyl compound (the electrophile) to this enolate.

Because the steps are done sequentially and a strong nonnucleophilic base is used to form the enolate of one carbonyl component only, a variety of carbonyl substrates can be used in the reaction. Both carbonyl components can have α hydrogens because only one enolate is prepared with LDA. Also, when an unsymmetrical ketone is used, LDA selectively forms the **less substituted, kinetic enolate**.

Sample Problem 24.1 illustrates the steps of a directed aldol reaction between a ketone and an aldehyde, both of which have α hydrogens.

Sample Problem 24.1 Draw the product of the following directed aldol reaction.



Solution

2-Methylcyclohexanone forms an enolate on the less substituted carbon, which then reacts with the electrophile, CH_3CHO .





Figure 24.4 illustrates how a directed aldol reaction was used in the synthesis of **periplanone B**, the sex pheromone of the female American cockroach.

Figure 24.4
A directed aldol reaction in the synthesis of periplanone B



- Periplanone B is an extremely active compound produced in small amounts by the female American cockroach. Its structure was determined using 200 μg of periplanone B from more than 75,000 female cockroaches. This structure was confirmed by synthesis in the laboratory in 1979.


To determine the needed carbonyl components for a directed aldol, follow the same strategy used for a regular aldol reaction in Section 24.1C, as shown in Sample Problem 24.2.

Sample Problem 24.2

What starting materials are needed to prepare *ar*-turmerone using a directed aldol reaction? *ar*-Turmerone is a principal component of the essential oil derived from turmeric root.




The dried and ground root of the turmeric plant, a tropical perennial in the ginger family, is an essential ingredient in curry powder.






Solution

When the desired product is an α,β -unsaturated carbonyl compound, identify the α and β carbons that are part of the C=C, and break the molecule into two components between these carbons.



Problem 24.10



What carbonyl starting materials are needed to prepare each compound using a directed aldol reaction?



Problem 24.11

A key step in the synthesis of donepezil (trade name Aricept) is a directed aldol reaction that forms α,β -unsaturated carbonyl compound **X**. What carbonyl starting materials are needed to prepare **X** using a directed aldol reaction? What reagents are needed to convert **X** to donepezil?


Donepezil (trade name Aricept) is a drug used to improve cognitive function in patients suffering from dementia and Alzheimer's disease.



24.4 Intramolecular Aldol Reactions

Aldol reactions with dicarbonyl compounds can be used to make five- and six-membered rings. The enolate formed from one carbonyl group is the nucleophile, and the carbonyl carbon of the other carbonyl group is the electrophile. For example, treatment of 2,5-hexanedione with base forms a five-membered ring.

2,5-Hexanedione is called a **1,4-dicarbonyl compound** to emphasize the relative positions of its carbonyl groups. 1,4-Dicarbonyl compounds are starting materials for synthesizing **five-membered rings**.




The steps in this process, shown in Mechanism 24.3, are no different from the general mechanisms of the aldol reaction and dehydration described previously in Section 24.1.




Mechanism 24.3 The Intramolecular Aldol Reaction

Step [1] Formation of an enolate




- Deprotonation of the CH₃ group with base forms a nucleophilic enolate, which is re-drawn to more clearly show the intramolecular reaction in Step [2].

Steps [2]–[4] Cyclization and dehydration



- In Step [2], the nucleophilic enolate attacks the electrophilic carbonyl carbon in the same molecule, forming a new carbon–carbon σ bond. **This generates the five-membered ring.**
- Protonation of the alkoxide in Step [3] and loss of H₂O by the two steps outlined in Mechanism 24.2 form a new C–C π bond.

When 2,5-hexanedione is treated with base in Step [1], two different enolates are possible—enolates **A** and **B**, formed by removal of H_a and H_b, respectively. Although enolate **A** goes on to form the five-membered ring, intramolecular cyclization using enolate **B** would lead to a strained three-membered ring.



Because the three-membered ring is much higher in energy than the enolate starting material, equilibrium greatly favors the starting materials and the **three-membered ring does not form**. Under the reaction conditions, enolate **B** is re-protonated to form 2,5-hexanedione, because all steps except dehydration are equilibria. **Thus, equilibrium favors formation of the more stable five-membered ring over the much less stable three-membered ring.**

In a similar fashion, six-membered rings can be formed from the intramolecular aldol reaction of **1,5-dicarbonyl compounds**.





Figure 24.5

The synthesis of progesterone using an intramolecular aldol reaction




- **Step [1]:** Oxidative cleavage of the alkene with O_3 , followed by Zn, H_2O (Section 12.10), gives the 1,5-dicarbonyl compound.
- **Step [2]:** Intramolecular aldol reaction of the 1,5-dicarbonyl compound with dilute OH^- in H_2O solution forms progesterone.
- **This two-step reaction sequence converts a five-membered ring into a six-membered ring.** Reactions that synthesize larger rings from smaller ones are called **ring expansion reactions**.


The synthesis of the female sex hormone **progesterone** by W. S. Johnson and co-workers at Stanford University is considered one of the classics in total synthesis. The last six-membered ring needed in the steroid skeleton was prepared by a two-step sequence using an intramolecular aldol reaction, as shown in Figure 24.5.

Problem 24.12 Draw a stepwise mechanism for the conversion of 2,6-heptanedione to 3-methyl-2-cyclohexenone with $NaOEt, EtOH$.

Problem 24.13 What cyclic product is formed when each 1,5-dicarbonyl compound is treated with aqueous OH^- ?




Problem 24.14 Following the two-step reaction sequence depicted in Figure 24.5, write out the steps needed to convert 1-methylcyclopentene to 2-cyclohexenone.



24.5 The Claisen Reaction

The **Claisen reaction** is the second general reaction of enolates with other carbonyl compounds. In the Claisen reaction, two molecules of an ester react with each other in the presence of an alkoxide base to form a **β -keto ester**. For example, treatment of ethyl acetate with $NaOEt$ forms ethyl acetoacetate after protonation with aqueous acid.

Unlike the aldol reaction, which is base-catalyzed, a full equivalent of base is needed to deprotonate the β -keto ester formed in Step [3] of the Claisen reaction.




The mechanism for the Claisen reaction (Mechanism 24.4) resembles the mechanism of an aldol reaction in that it involves nucleophilic addition of an enolate to an electrophilic carbonyl group. Because esters have a leaving group on the carbonyl carbon, however, loss of a leaving group occurs to form the product of **substitution, not addition**.




Mechanism 24.4 The Claisen Reaction

Step [1] Formation of a nucleophilic enolate




- In Step [1], the base removes a proton from the α carbon to form a **resonance-stabilized enolate**.

Steps [2]–[3] Nucleophilic addition and loss of the leaving group



- In Step [2], the nucleophilic enolate attacks the electrophilic carbonyl carbon of another molecule of ester, forming a new carbon–carbon bond. **This joins the α carbon of one ester to the carbonyl carbon of a second ester.**
- Elimination of the leaving group, EtO^- , forms a β -keto ester in Step [3]. Steps [1]–[3] are reversible equilibria.

Steps [4]–[5] Deprotonation and protonation




- Because the β -keto ester formed in Step [3] has especially acidic protons between its two carbonyl groups, a proton is removed under the basic reaction conditions to form an enolate (Step [4]). **The formation of this resonance-stabilized enolate drives the equilibrium in the Claisen reaction.**
- Protonation of this enolate with strong acid re-forms the neutral β -keto ester to complete the reaction.

Because the generation of a resonance-stabilized enolate from the product β -keto ester drives the Claisen reaction (Step [4] of Mechanism 24.4), only esters with two or three hydrogens on the α carbon undergo this reaction; that is, esters must have the general structure $\text{CH}_3\text{CO}_2\text{R}'$ or $\text{RCH}_2\text{CO}_2\text{R}'$.

- Keep in mind: The characteristic reaction of esters is nucleophilic substitution. A Claisen reaction is a nucleophilic substitution in which an enolate is the nucleophile.**

Figure 24.6 compares the general reaction for nucleophilic substitution of an ester with the Claisen reaction. Sample Problem 24.3 reinforces the basic features of the Claisen reaction.

Figure 24.6
The Claisen reaction—An example of nucleophilic substitution




- Esters react by nucleophilic substitution. In a Claisen reaction, an enolate is the nucleophile that adds to the carbonyl group.

Sample Problem 24.3 Draw the product of the following Claisen reaction.




Solution



To draw the product of any Claisen reaction, form a new carbon-carbon bond between the α carbon of one ester and the carbonyl carbon of another ester, with elimination of the leaving group ($^-\text{OCH}_3$ in this case).



Next, write out the steps of the reaction to verify this product.



Problem 24.15 What β -keto ester is formed when each ester is used in a Claisen reaction?



24.6 The Crossed Claisen and Related Reactions

Like the aldol reaction, it is sometimes possible to carry out a Claisen reaction with two different carbonyl components as starting materials.


- A Claisen reaction between two different carbonyl compounds is called a *crossed Claisen reaction*.

24.6A Two Useful Crossed Claisen Reactions

A crossed Claisen reaction is synthetically useful in two different instances.


- A crossed Claisen occurs between two different esters when only one has α hydrogens.

When one ester has no α hydrogens, a crossed Claisen reaction often leads to one product. Common esters with no α H atoms include ethyl formate (HCO_2Et) and ethyl benzoate ($\text{C}_6\text{H}_5\text{CO}_2\text{Et}$). For example, the reaction of ethyl benzoate (as the electrophile) with ethyl acetate (which forms the enolate) in the presence of base forms predominately one β -keto ester.



- A crossed Claisen occurs between a ketone and an ester.


The reaction of a ketone and an ester in the presence of base also forms the product of a crossed Claisen reaction. The enolate is generally formed from the ketone component, and the reaction works best when the ester has no α hydrogens. The product of this crossed Claisen reaction is a **β -dicarbonyl compound**, but *not* a β -keto ester.



Problem 24.16 What crossed Claisen product is formed from each pair of compounds?


- a. CH_3CH_2COOEt and HCO_2Et d. and
- b. $CH_3(CH_2)_5CO_2Et$ and HCO_2Et
- c. $(CH_3)_2C=O$ and CH_3CO_2Et

Problem 24.17




Sunscreen ingredients

Avobenzene is a conjugated compound that absorbs ultraviolet light with wavelengths in the 320–400 nm region, so it is a common ingredient in commercial sunscreens. Write out two different crossed Claisen reactions that form avobenzene.



24.6B Other Useful Variations of the Crossed Claisen Reaction


β -Dicarbonyl compounds are also prepared by reacting an enolate with **ethyl chloroformate** and **diethyl carbonate**.




These reactions resemble a Claisen reaction because they involve the same three steps:

- [1] Formation of an enolate
- [2] Nucleophilic addition to a carbonyl group
- [3] Elimination of a leaving group


For example, reaction of an ester enolate with diethyl carbonate yields a β -diester (Reaction [1]), whereas reaction of a ketone enolate with ethyl chloroformate forms a β -keto ester (Reaction [2]).




Reaction [2] is noteworthy because it provides easy access to **β -keto esters**, which are useful starting materials in the acetoacetic ester synthesis (Section 23.10). In this reaction, Cl^- is eliminated rather than ^-OEt in Step [3], because Cl^- is a better leaving group, as shown in the following steps.



Problem 24.18 Draw the products of each reaction.




Problem 24.19 Two steps in a synthesis of the analgesic ibuprofen, the chapter-opening molecule, include a carbonyl condensation reaction, followed by an alkylation reaction. Identify intermediates **A** and **B** in the synthesis of ibuprofen.




24.7 The Dieckmann Reaction

Intramolecular Claisen reactions of diesters form five- and six-membered rings. The enolate of one ester is the nucleophile, and the carbonyl carbon of the other is the electrophile. An intramolecular Claisen reaction is called a **Dieckmann reaction**. Two types of diesters give good yields of cyclic products.

- 1,6-Diesters yield five-membered rings by the Dieckmann reaction.




- 1,7-Diesters yield six-membered rings by the Dieckmann reaction.



The mechanism of the Dieckmann reaction is exactly the same as the mechanism of an intermolecular Claisen reaction. It is illustrated in Mechanism 24.5 for the formation of a six-membered ring.




Mechanism 24.5 The Dieckmann Reaction




- In Step [1], the base removes a proton to form an enolate, which attacks the carbonyl group of the second ester in Step [2], thus forming a new carbon–carbon bond.
- In Step [3], elimination of OEt^- forms the β -keto ester.
- To complete the reaction, the proton between the two carbonyl groups is removed with base, and then protonation of the enolate re-forms the β -keto ester (Steps [4] and [5]).

Problem 24.20 One synthesis of prostaglandin PGA_2 involved a Dieckmann reaction as a key step. What is the structure of **X** in the reactions drawn below?




Problem 24.21 What two β -keto esters are formed in the Dieckmann reaction of the following diester?




24.8 The Michael Reaction

Like the aldol and Claisen reactions, the **Michael reaction involves two carbonyl components—the enolate of one carbonyl compound and an α,β -unsaturated carbonyl compound.**

Two components of a Michael reaction




Recall from Section 20.15 that α,β -unsaturated carbonyl compounds are resonance stabilized and have **two electrophilic sites—the carbonyl carbon and the β carbon.**





- The Michael reaction involves the conjugate addition (1,4-addition) of a resonance-stabilized enolate to the β carbon of an α,β -unsaturated carbonyl system.

All conjugate additions add the **elements of H and Nu across the α and β carbons.** In the Michael reaction, the **nucleophile is an enolate.** Enolates of active methylene compounds are particularly common. The α,β -unsaturated carbonyl component is often called a **Michael acceptor.**

Conjugate addition



Michael reaction



Problem 24.22 Which of the following compounds can serve as Michael acceptors?




The Michael reaction always forms a new carbon–carbon bond on the β carbon of the Michael acceptor. Reaction [2] is used to illustrate the mechanism of the Michael reaction in Mechanism 24.6. The key step is nucleophilic addition of the enolate to the β carbon of the Michael acceptor in Step [2].




Mechanism 24.6 The Michael Reaction

Step [1] Enolate formation



- Base removes the acidic proton between the two carbonyl groups, **forming the enolate** in Step [1]. Only one of the three resonance structures is drawn.


Steps [2]–[3] Nucleophilic attack at the β carbon and protonation



- **The nucleophilic enolate adds to the β carbon** of the α,β-unsaturated carbonyl compound, forming a new carbon-carbon bond and a resonance-stabilized enolate.
- Protonation of the enolate forms the 1,4-addition product in Step [3].

When the product of a Michael reaction is also a β-keto ester, it can be hydrolyzed and decarboxylated by heating in aqueous acid, as discussed in Section 23.9. This forms a **1,5-dicarbonyl compound**. Figure 24.7 shows a Michael reaction that was a key step in the synthesis of **estrone**, a female sex hormone.

1,5-Dicarbonyl compounds are starting materials for intramolecular aldol reactions, as described in Section 24.4.



Problem 24.23 What product is formed when each pair of compounds is treated with NaOEt in ethanol?






Figure 24.7

Using a Michael reaction in the synthesis of the steroid estrone




Problem 24.24 What starting materials are needed to prepare each compound by the Michael reaction?




24.9 The Robinson Annulation

The word **annulation** comes from the Greek word *annulus* for ring. The Robinson annulation is named for English chemist Sir Robert Robinson, who was awarded the 1947 Nobel Prize in Chemistry.

The Robinson annulation is a ring-forming reaction that combines a Michael reaction with an intramolecular aldol reaction. Like the other reactions in Chapter 24, it involves enolates and it forms carbon–carbon bonds. The two starting materials for a Robinson annulation are an α,β -unsaturated carbonyl compound and an enolate.



The Robinson annulation forms a six-membered ring and three new carbon–carbon bonds—two σ bonds and one π bond. The product contains an α,β -unsaturated ketone in a cyclohexane ring—that is, a **2-cyclohexenone** ring. To generate the enolate component of the Robinson annulation, ^-OH in H_2O and ^-OEt in EtOH are typically used.




The mechanism of the Robinson annulation consists of two parts: a **Michael addition** to the α,β -unsaturated carbonyl compound to form a 1,5-dicarbonyl compound, followed by an **intramolecular aldol reaction** to form the six-membered ring. The mechanism is written out in two parts (Mechanisms 24.7 and 24.8) for Reaction [2] between methyl vinyl ketone and 2-methyl-1,3-cyclohexanedione.




Mechanism 24.7 The Robinson Annulation—Part [A] Michael Addition to Form a 1,5-Dicarbonyl Compound

Step [1] Enolate formation



- Base removes the most acidic proton—that is, the proton between the two carbonyl groups—forming the enolate in Step [1]. Only one of the three resonance structures is drawn.

Steps [2]–[3] Nucleophilic attack at the β carbon and protonation



- **Conjugate addition of the enolate to the β carbon** of the α,β -unsaturated carbonyl compound forms a new carbon-carbon bond and a resonance-stabilized enolate.
- Protonation of the enolate forms the 1,5-dicarbonyl compound.


Part [A] illustrates the three-step mechanism for the Michael addition that forms the first carbon-carbon σ bond, generating the 1,5-dicarbonyl compound. The first step always involves removal of the most acidic proton to form an enolate.

In Part [B] of the mechanism, an intramolecular aldol reaction followed by dehydration forms the six-membered ring.




Mechanism 24.8 The Robinson Annulation—Part [B] Intramolecular Aldol Reaction to Form a 2-Cyclohexenone

Steps [4]–[6] Intramolecular aldol reaction to form a β -hydroxy ketone



- The **intramolecular aldol reaction** consists of three steps: [4] **enolate formation**, [5] **nucleophilic attack**, and [6] **protonation**. This forms another carbon-carbon σ bond and a β -hydroxy carbonyl compound (compare Section 24.4).


Steps [7]–[8] Dehydration to form the α,β -unsaturated ketone



- Dehydration consists of two steps: **deprotonation** and **loss of OH^-** (Section 24.1B). This reaction forms the new π bond in the α,β -unsaturated ketone.


All of the parts of this mechanism have been discussed in previous sections of Chapter 24. However, the end result of the Robinson annulation—the formation of a 2-cyclohexenone ring—is new.

To draw the product of Robinson annulation without writing out the mechanism each time, place the α carbon of the compound that becomes the enolate next to the β carbon of the α,β -unsaturated carbonyl compound. Then, join the appropriate carbons together as shown. If you follow this method of drawing the starting materials, the double bond in the product always ends up in the same position in the six-membered ring.



Sample Problem 24.4

Draw the Robinson annulation product formed from the following starting materials.




Solution

Arrange the starting materials to put the reactive atoms next to each other. For example:


- Place the α,β -unsaturated carbonyl compound *to the left* of the carbonyl compound.
- Determine which α carbon will become the enolate. The most acidic H is always removed with base first, which in this case is the H on the α carbon between the two carbonyl groups. This α carbon is drawn adjacent to the β carbon of the α,β -unsaturated carbonyl compound.

Then draw the bonds to form the new six-membered ring.



Problem 24.25


Draw the products when each pair of compounds is treated with OH^- , H_2O in a Robinson annulation reaction.



To use the Robinson annulation in synthesis, you must be able to determine what starting materials are needed to prepare a given compound, by working in the retrosynthetic direction.


HOW TO Synthesize a Compound Using the Robinson Annulation

Example What starting materials are needed to synthesize the following compound using a Robinson annulation?




Step [1] Locate the 2-cyclohexenone ring and re-draw the target molecule if necessary.

- To most easily determine the starting materials, always arrange the α,β -unsaturated carbonyl system in the same location. The target compound may have to be flipped or rotated, and you must be careful not to move any bonds to the wrong location during this process.






Step [2] Break the 2-cyclohexenone ring into two components.

- Break the $C=C$. One half becomes the carbonyl group of the enolate component.
- Break the bond between the β carbon and the carbon to which it is bonded.



Problem 24.26 What starting materials are needed to synthesize each compound by a Robinson annulation?



KEY CONCEPTS

Carbonyl Condensation Reactions


The Four Major Carbonyl Condensation Reactions

Reaction type	Reaction [New C–C bonds are shown in red.]
[1] Aldol reaction (24.1)	$2 \text{ RCH}_2\text{C}(=\text{O})\text{H} \xrightleftharpoons{-\text{OH}, \text{H}_2\text{O}} \text{RCH}_2\text{C}(\text{OH})(\text{H})\text{CH}(\text{R})\text{CHO} \xrightarrow[-\text{OH}]{\text{or } \text{H}_3\text{O}^+} \text{RCH}_2\text{C}(\text{H})=\text{C}(\text{R})\text{CHO}$ <p>aldehyde (or ketone) β-hydroxy carbonyl compound (E and Z) α,β-unsaturated carbonyl compound</p>
[2] Claisen reaction (24.5)	$2 \text{ RCH}_2\text{C}(=\text{O})\text{OR}' \xrightarrow[\text{[2] H}_3\text{O}^+]{\text{[1] NaOR}' } \text{RCH}_2\text{C}(=\text{O})\text{CH}(\text{R})\text{C}(=\text{O})\text{OR}'$ <p>ester β-keto ester</p>
[3] Michael reaction (24.8)	$\text{R}-\text{C}(=\text{O})\text{CH}=\text{CH}_2 + \text{C}(=\text{O})\text{R}'' \xrightarrow[-\text{OH}]{-\text{OR}' \text{ or } \text{H}_2\text{O}} \text{R}-\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{C}(=\text{O})\text{R}''$ <p>α,β-unsaturated carbonyl compound carbonyl compound 1,5-dicarbonyl compound</p>
[4] Robinson annulation (24.9)	$\text{R}-\text{C}(=\text{O})\text{CH}=\text{CH}_2 + \text{Cyclohexanone} \xrightarrow[-\text{OH}]{\text{H}_2\text{O}} \text{2-cyclohexenone}$ <p>α,β-unsaturated carbonyl compound carbonyl compound 2-cyclohexenone</p>


Useful Variations

[New C–C bonds are shown in red.]

[1] Directed aldol reaction (24.3)




[2] Intramolecular aldol reaction (24.4)




[3] Dieckmann reaction (24.7)

a. With 1,6-diesters:



b. With 1,7-diesters:




PROBLEMS

The Aldol Reaction



24.27 Draw the product formed from an aldol reaction with the given starting material(s) using ⁻OH, H₂O.

- (CH₃)₂CHCHO only
- (CH₃)₂CHCHO + CH₂=O
- C₆H₅CHO + CH₃CH₂CH₂CHO
- (CH₃CH₂)₂C=O only
- (CH₃CH₂)₂C=O + CH₂=O






24.28 What four β-hydroxy aldehydes are formed by a crossed aldol reaction of CH₃CH₂CH₂CHO and C₆H₅CH₂CHO?






24.29 Draw the product formed in each directed aldol reaction.




24.30 Draw the product formed when each dicarbonyl compound undergoes an intramolecular aldol reaction followed by dehydration.







24.31 What starting materials are needed to synthesize each compound using an aldol or similar reaction?




24.32 A published synthesis of the analgesic nabumetone uses a crossed aldol reaction to form **X**. What is the structure of **X**? **X** is converted to nabumetone in one step by hydrogenation with H₂ and Pd-C. (See Problem 23.29 for another way to make nabumetone.)



24.33 What dicarbonyl compound is needed to prepare each compound by an intramolecular aldol reaction?





24.34 Identify the structures of **C** and **D** in the following reaction sequence.









The Claisen and Dieckmann Reactions

24.35 Draw the Claisen product formed from each ester.






24.36 What four compounds are formed from the crossed Claisen reaction of $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ and $\text{CH}_3\text{CH}_2\text{CO}_2\text{Et}$?


24.37 Draw the product formed from a Claisen reaction with the given starting materials using ^-OEt , EtOH .




24.38 What starting materials are needed to synthesize each compound by a crossed Claisen reaction?



24.39 The 1,3-diketone shown below can be prepared by two different Claisen reactions—namely, one that forms bond (a) and one that forms bond (b). What starting materials are needed for each of these reactions?




24.40 Even though **B** contains three ester groups, a single Dieckmann product results when **B** is treated with NaOCH_3 in CH_3OH , followed by H_3O^+ . Draw the structure and explain why it is the only product formed.




Michael Reaction


24.41 Draw the product formed from a Michael reaction with the given starting materials using ^-OEt , EtOH.



24.42 What starting materials are needed to prepare each compound using a Michael reaction?




24.43 β -Vetivone is isolated from vetiver, a perennial grass that yields a variety of compounds used in traditional eastern medicine, pest control, and fragrance. In one synthesis, ketone **A** is converted to β -vetivone by a two-step process: Michael reaction, followed by intramolecular aldol reaction. (a) What Michael acceptor is needed for the conjugate addition? (See Problem 23.63 for another method to form the bicyclic ring system of β -vetivone.) (b) Draw a stepwise mechanism for the aldol reaction, which forms the six-membered ring.




Robinson Annulation

24.44 Draw the product of each Robinson annulation from the given starting materials using ^-OH in H_2O solution.



24.45 What starting materials are needed to synthesize each compound using a Robinson annulation?











Reactions


24.46 Draw the organic products formed when butanal ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$) is treated with each reagent.

- | | | |
|--|---|---|
| a. OH^- , H_2O | f. NaBH_4 , CH_3OH | k. CrO_3 , H_2SO_4 |
| b. OH^- , $\text{CH}_2=\text{O}$, H_2O | g. H_2 , Pd-C | l. Br_2 , CH_3COOH |
| c. [1] LDA; [2] CH_3CHO ; [3] H_2O | h. $\text{HOCH}_2\text{CH}_2\text{OH}$, TsOH | m. $\text{Ph}_3\text{P}=\text{CH}_2$ |
| d. $\text{CH}_2(\text{CO}_2\text{Et})_2$, NaOEt, EtOH | i. CH_3NH_2 , mild acid | n. NaCN, HCl |
| e. [1] CH_3Li ; [2] H_2O | j. $(\text{CH}_3)_2\text{NH}$, mild acid | o. [1] LDA; [2] CH_3I |

24.47 Draw the organic products formed in each reaction.




24.48 Fill in the lettered reagents needed for each reaction.




24.49 Explain why vinyl halides such as $\text{CH}_2=\text{CHCl}$ undergo elimination by an E1cB mechanism more readily than alkyl halides such as $\text{CH}_3\text{CH}_2\text{Cl}$.

24.50 Identify lettered intermediates **A–D** in the following reaction sequence.




- 24.51** Identify compounds **A** and **B**, two synthetic intermediates in the 1979 synthesis of the plant growth hormone gibberellic acid by Corey and Smith. Gibberellic acid induces cell division and elongation, thus making plants tall and leaves large.




Mechanisms


- 24.52** When acetaldehyde (CH_3CHO) is treated with three equivalents of formaldehyde ($\text{CH}_2=\text{O}$) in the presence of aqueous Na_2CO_3 , $(\text{HOCH}_2)_3\text{CCHO}$ is formed as product. Draw a stepwise mechanism for this process.
- 24.53** In theory, the intramolecular aldol reaction of 6-oxoheptanal could yield the three compounds shown. It turns out, though, that 1-acetylcyclopentene is by far the major product. Why are the other two compounds formed in only minor amounts? Draw a stepwise mechanism to show how all three products are formed.




- 24.54** Draw a stepwise mechanism for each cyclization reaction.




- 24.55** Draw a stepwise mechanism for the following variation of the aldol reaction, often called a nitro aldol reaction.




- 24.56** Draw a stepwise mechanism for the following Robinson annulation. This reaction was a key step in a synthesis of the steroid cortisone by R. B. Woodward and co-workers at Harvard University in 1951.




24.63 Devise a synthesis of each compound from cyclopentanone, benzene, and organic alcohols having ≤ 3 C's. You may also use any required organic or inorganic reagents.




24.64 Devise a synthesis of each compound from $\text{CH}_3\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$, benzene, and alcohols having ≤ 2 C's. You may also use any required organic or inorganic reagents.



24.65 Devise a synthesis of each compound from cyclohexene. You may also use any required reagents.




24.66 Octinoxate is an unsaturated ester used as an active ingredient in sunscreens. (a) What carbonyl compounds are needed to synthesize this compound using a condensation reaction? (b) Devise a synthesis of octinoxate from the given organic starting materials and any other needed reagents.



General Problems


24.67 Four steps in the synthesis of helminthosporal, a toxin produced by a wheat plant fungus, are shown below.




- What compounds are needed to carry out Step [1]?
- What compounds are needed to carry out Step [2]?
- Write a detailed mechanism for Step [3].
- Write a detailed mechanism for Step [4]. What is unusual about the product of this reaction?
- Step [1] adds a formyl group ($\text{HCO}-$) and Step [3] removes it. Why was this apparently unnecessary process done?

Challenge Problems


24.68 Propose a stepwise mechanism for the following reaction of a β -keto ester. Suggest a reason why this rearrangement reaction occurs.




24.69 Isophorone is formed from three molecules of acetone [(CH₃)₂C=O] in the presence of base. Draw a mechanism for this process.



24.70 Draw a stepwise mechanism for the following reaction. [Hint: Two Michael reactions are needed.]




24.71 4-Methylpyridine reacts with benzaldehyde (C₆H₅CHO) in the presence of base to form **A**. (a) Draw a stepwise mechanism for this reaction. (b) Would you expect 2-methylpyridine or 3-methylpyridine to undergo a similar type of condensation reaction? Explain why or why not.



Amines

25

- 25.1 Introduction
- 25.2 Structure and bonding
- 25.3 Nomenclature
- 25.4 Physical properties
- 25.5 Spectroscopic properties
- 25.6 Interesting and useful amines
- 25.7 Preparation of amines
- 25.8 Reactions of amines—
General features
- 25.9 Amines as bases
- 25.10 Relative basicity of
amines and other
compounds
- 25.11 Amines as nucleophiles
- 25.12 Hofmann elimination
- 25.13 Reaction of amines with
nitrous acid
- 25.14 Substitution reactions of
aryl diazonium salts
- 25.15 Coupling reactions of aryl
diazonium salts
- 25.16 Application: Synthetic
dyes
- 25.17 Application: Sulfa drugs



Caffeine is a bitter-tasting compound found in coffee, tea, cola beverages, and chocolate. Caffeine is a mild stimulant, usually imparting a feeling of alertness after consumption. It also increases heart rate, dilates airways, and stimulates the secretion of stomach acid. Caffeine is an alkaloid, a naturally occurring amine derived from a plant source. In Chapter 25 we learn about the properties and reactions of amines.

We now leave the chemistry of carbonyl compounds to concentrate on amines, organic derivatives of ammonia (NH_3), formed by replacing one or more hydrogen atoms by alkyl or aryl groups. **Amines are stronger bases and better nucleophiles than other neutral organic compounds**, so much of Chapter 25 focuses on these properties.


Like that of alcohols, the chemistry of amines does not always fit neatly into one reaction class, and this can make learning the reactions of amines challenging. Many interesting natural products and widely used drugs are amines, so you also need to know how to introduce this functional group into organic molecules.

25.1 Introduction


Classifying amines as 1°, 2°, or 3° is reminiscent of classifying amides in

Chapter 22, but is *different* from classifying other atoms and functional groups as 1°, 2°, and 3°. Compare, for example, a 2° amine and a 2° alcohol. A 2° amine (R_2NH) has *two* C–N bonds. A 2° alcohol (R_2CHOH), on the other hand, has only *one* C–O bond, but *two* C–C bonds on the carbon bonded to oxygen.

Amines are organic nitrogen compounds, formed by replacing one or more hydrogen atoms of ammonia (NH_3) with alkyl groups. As discussed in Section 21.11, amines are classified as 1°, 2°, or 3° by the number of alkyl groups bonded to the *nitrogen* atom.




Like ammonia, **the amine nitrogen atom has a nonbonded electron pair**, making it both a base and a nucleophile. As a result, amines react with electrophiles to form **ammonium salts**—compounds with four bonds to nitrogen.



- The chemistry of amines is dominated by the nonbonded electron pair on the nitrogen atom.

Problem 25.1

Classify each amine in the following compounds as 1°, 2°, or 3°.



Problem 25.2

Draw the structure of a compound of molecular formula $\text{C}_4\text{H}_{11}\text{NO}$ that fits each description: (a) a compound that contains a 1° amine and a 3° alcohol; (b) a compound that contains a 3° amine and a 1° alcohol.

25.2 Structure and Bonding

An amine nitrogen atom is surrounded by three atoms and one nonbonded electron pair, making the N atom sp^3 hybridized and trigonal pyramidal, with bond angles of approximately 109.5° .





Figure 25.1
Electrostatic potential plots of
 CH_3NH_2 and $(\text{CH}_3)_3\text{N}$




- Both amines clearly show the electron-rich region (in red) at the N atom.


Because nitrogen is much more electronegative than carbon or hydrogen, **the C–N and N–H bonds are all polar**, with the N atom electron rich and the C and H atoms electron poor. The electrostatic potential maps in Figure 25.1 show the polar C–N and N–H bonds in CH_3NH_2 (methylamine) and $(\text{CH}_3)_3\text{N}$ (trimethylamine).

An amine nitrogen atom bonded to an electron pair and three different alkyl groups is technically a stereogenic center, so two nonsuperimposable trigonal pyramids can be drawn.

An amine with four different groups around N



This does not mean, however, that such an amine exists as two different enantiomers, because one is rapidly converted to the other at room temperature. The amine flips inside out, passing through a trigonal planar (achiral) transition state. **Because the two enantiomers interconvert, we can ignore the chirality of the amine nitrogen.**





In contrast, the chirality of an ammonium salt with four different groups on N cannot be ignored. Because there is no nonbonded electron pair on the nitrogen atom, interconversion cannot occur, and the N atom is just like a carbon atom with four different groups around it.

Two enantiomers of
an ammonium salt



- The N atom of an ammonium salt is a stereogenic center when N is surrounded by four different groups.

Problem 25.3 Label the stereogenic centers in each molecule.



Problem 25.4 The C–N bond length in CH_3NH_2 is 147 pm, whereas the C–N bond length in $\text{C}_6\text{H}_5\text{NH}_2$ is only 140 pm. Why is the C–N bond in the aromatic amine shorter?

25.3 Nomenclature

25.3A Primary Amines


Primary amines are named using either systematic or common names.

- To assign the **systematic name**, find the longest continuous carbon chain bonded to the amine nitrogen, and change the *-e* ending of the parent alkane to the suffix **-amine**. Then use the usual rules of nomenclature to number the chain and name the substituents.
- To assign a **common name**, name the alkyl group bonded to the nitrogen atom and add the suffix **-amine**, forming a single word.

Examples





Systematic name: **methanamine**
Common name: **methylamine**



Systematic name: **cyclohexanamine**
Common name: **cyclohexylamine**

25.3B Secondary and Tertiary Amines

Secondary and tertiary amines having identical alkyl groups are named by using the prefix **di-** or **tri-** with the name of the primary amine.




Secondary and tertiary amines having more than one kind of alkyl group are named as **N-substituted primary amines**, using the following procedure.


HOW TO Name 2° and 3° Amines with Different Alkyl Groups

Example Name the following 2° amine: $(\text{CH}_3)_2\text{CHNHCH}_3$.

Step [1] Designate the longest alkyl chain (or largest ring) bonded to the N atom as the parent amine and assign a common or systematic name.




Step [2] Name the other groups on the N atom as alkyl groups, alphabetize the names, and put the prefix **N-** before the name.




Sample Problem 25.1

Name each amine.




Solution

a. [1] A 1° amine: Find and name the longest chain containing the amine nitrogen.




[2] Number and name the substituents:




Answer: 4-methyl-1-pentanamine

b. For a 3° amine, one alkyl group on N is the principal R group and the others are substituents.

[1] Name the ring bonded to the N:




[2] Name the substituents:




Problem 25.5

Name each amine.




25.3C Aromatic Amines


Aromatic amines are named as derivatives of aniline.



aniline



N-ethylaniline



o-bromoaniline

25.3D Miscellaneous Nomenclature Facts

An NH_2 group named as a substituent is called an **amino group**.

There are many different **nitrogen heterocycles**, and each ring type is named differently depending on the number of N atoms in the ring, the ring size, and whether it is aromatic or not. The structures and names of four common nitrogen heterocycles are shown. In numbering these heterocycles, the **N atom is always placed at the "1" position**.



pyridine



pyrrole



piperidine




pyrrolidine

Problem 25.6 Draw a structure corresponding to each name.

- | | |
|--|-----------------------------------|
| a. 2,4-dimethyl-3-hexanamine | e. <i>N,N</i> -dimethylethylamine |
| b. <i>N</i> -methylpentylamine | f. 2-aminocyclohexanone |
| c. <i>N</i> -isopropyl- <i>p</i> -nitroaniline | g. <i>N</i> -methylaniline |
| d. <i>N</i> -methylpiperidine | h. <i>m</i> -ethylaniline |

25.4 Physical Properties

Amines exhibit dipole–dipole interactions because of the polar C–N and N–H bonds. **Primary and secondary amines are also capable of intermolecular hydrogen bonding**, because they contain N–H bonds. Because nitrogen is less electronegative than oxygen, however, intermolecular hydrogen bonds between N and H are *weaker* than those between O and H. How these factors affect the physical properties of amines is summarized in Table 25.1.



Problem 25.7 Arrange each group of compounds in order of increasing boiling point.





Table 25.1 Physical Properties of Amines

Property	Observation					
Boiling point and melting point	<ul style="list-style-type: none"> Primary (1°) and 2° amines have higher bp's than similar compounds (like ethers) incapable of hydrogen bonding, but lower bp's than alcohols that have stronger intermolecular hydrogen bonds. 					
	<table border="0" style="width: 100%; text-align: center;"> <tr> <td><chem>CH3CH2OCH2CH3</chem></td> <td><chem>CH3CH2CH2CH2NH2</chem></td> <td><chem>CH3CH2CH2CH2OH</chem></td> </tr> <tr> <td>MW = 74 bp 38 °C</td> <td>MW = 73 bp 78 °C</td> <td>MW = 74 bp 118 °C</td> </tr> </table> <p style="text-align: center;"> Increasing intermolecular forces Increasing boiling point </p>	<chem>CH3CH2OCH2CH3</chem>	<chem>CH3CH2CH2CH2NH2</chem>	<chem>CH3CH2CH2CH2OH</chem>	MW = 74 bp 38 °C	MW = 73 bp 78 °C
<chem>CH3CH2OCH2CH3</chem>	<chem>CH3CH2CH2CH2NH2</chem>	<chem>CH3CH2CH2CH2OH</chem>				
MW = 74 bp 38 °C	MW = 73 bp 78 °C	MW = 74 bp 118 °C				
Solubility	<ul style="list-style-type: none"> Tertiary (3°) amines have lower boiling points than 1° and 2° amines of comparable molecular weight, because they have no N–H bonds and are incapable of hydrogen bonding. 					
	<table border="0" style="width: 100%; text-align: center;"> <tr> <td> <div style="border: 1px solid black; padding: 2px; display: inline-block;">3° amine</div> <chem>CH3CH2N(CH3)2</chem> MW = 73 bp 38 °C no N–H bond </td> <td> <chem>CH3CH2-NH-CH2CH3</chem> ← <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">2° amine higher bp</div> MW = 73 bp 56 °C N–H bond </td> </tr> </table>	<div style="border: 1px solid black; padding: 2px; display: inline-block;">3° amine</div> <chem>CH3CH2N(CH3)2</chem> MW = 73 bp 38 °C no N–H bond	<chem>CH3CH2-NH-CH2CH3</chem> ← <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">2° amine higher bp</div> MW = 73 bp 56 °C N–H bond			
<div style="border: 1px solid black; padding: 2px; display: inline-block;">3° amine</div> <chem>CH3CH2N(CH3)2</chem> MW = 73 bp 38 °C no N–H bond	<chem>CH3CH2-NH-CH2CH3</chem> ← <div style="border: 1px solid black; padding: 2px; display: inline-block; margin-left: 10px;">2° amine higher bp</div> MW = 73 bp 56 °C N–H bond					

Figure 25.2
Mass spectrum of butylamine



- The molecular ion for $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ occurs at $m/z = 73$. This odd mass for a molecular ion is characteristic of an amine with an odd number of N atoms.

25.5 Spectroscopic Properties

Amines exhibit characteristic features in their mass spectra, IR spectra, and ^1H and ^{13}C NMR spectra.

25.5A Mass Spectra

Amines differ from compounds that contain only C, H, and O atoms, which always have a molecular ion with an *even* mass in their mass spectra. This is apparent in the mass spectrum of butylamine, which is shown in Figure 25.2.

The general molecular formula for an amine with one N atom is $\text{C}_n\text{H}_{2n+3}\text{N}$.

- Amines with an odd number of N atoms give an odd molecular ion in their mass spectra.


25.5B IR Spectra

Amines with N–H bonds show characteristic absorptions in their IR spectra.

- 1° Amines show *two* N–H absorptions at $3300\text{--}3500\text{ cm}^{-1}$.
- 2° Amines show *one* N–H absorption at $3300\text{--}3500\text{ cm}^{-1}$.

Because 3° amines have no N–H bonds, they do *not* absorb in this region in their IR spectra. The single bond region ($> 2500\text{ cm}^{-1}$) of the IR spectra for 1° , 2° , and 3° amines illustrates these features in Figure 25.3.

Figure 25.3 The single bond region of the IR spectra for a 1° , 2° , and 3° amine



Problem 25.8 Only one amine shows a molecular ion in its mass spectrum at $m/z = 59$ and has one peak in its IR spectrum at $\sim 3300\text{ cm}^{-1}$. What is its structure?

25.5C NMR Spectra

Amines exhibit the following characteristic ^1H NMR and ^{13}C NMR absorptions.

- **The NH signal appears between 0.5 and 5.0 ppm.** The exact location depends on the degree of hydrogen bonding and the concentration of the sample.
- The protons on the carbon bonded to the amine nitrogen are deshielded and typically absorb at **2.3–3.0 ppm.**
- In the ^{13}C NMR spectrum, the carbon bonded to the N atom is deshielded and typically absorbs at **30–50 ppm.**

Like the OH absorption of an alcohol, the **NH absorption is not split by adjacent protons, nor does it cause splitting of adjacent C–H absorptions in a ^1H NMR spectrum.** The NH peak of an amine is sometimes somewhat broader than other peaks in the spectrum. The ^1H NMR spectrum of *N*-methylaniline is shown in Figure 25.4.

Problem 25.9 What is the structure of an unknown compound with molecular formula $\text{C}_6\text{H}_{15}\text{N}$ that gives the following ^1H NMR absorptions: 0.9 (singlet, 1 H), 1.10 (triplet, 3 H), 1.15 (singlet, 9 H), and 2.6 (quartet, 2 H) ppm?


25.6 Interesting and Useful Amines

A great many simple and complex amines occur in nature, and others with biological activity have been synthesized in the lab.

25.6A Simple Amines and Alkaloids

Many low molecular weight amines have *very* foul odors. **Trimethylamine** $[(\text{CH}_3)_3\text{N}]$, formed when enzymes break down certain fish proteins, has the characteristic odor of rotting fish. **Putrescine** $(\text{NH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2)$ and **cadaverine** $(\text{NH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2)$ are

Figure 25.4
The ^1H NMR spectrum of *N*-methylaniline




- The CH_3 group appears as a singlet at 2.7 ppm because there is no splitting by the adjacent NH proton.
- The NH proton appears as a broad singlet at 3.6 ppm.
- The five H atoms of the aromatic ring appear as a complex pattern at 6.6–7.2 ppm.

The word **alkaloid** is derived from the word *alkali*, because aqueous solutions of alkaloids are slightly basic.

both poisonous diamines with putrid odors. They, too, are present in rotting fish, and are partly responsible for the odors of semen, urine, and bad breath.

Naturally occurring amines derived from plant sources are called **alkaloids**. Alkaloids previously encountered in the text include **quinine** (Problem 17.15), **morphine** (Section 22.9), and **cocaine** (Problem 3.42). Three other common alkaloids are **atropine**, **nicotine**, and **coniine**, illustrated in Figure 25.5.

25.6B Histamine and Antihistamines



Histamine, a rather simple triamine first discussed in Section 17.8, is responsible for a wide variety of physiological effects. Histamine is a vasodilator (it dilates capillaries), so it is released at the site of an injury or infection to increase blood flow. It is also responsible for the symptoms of allergies, including a runny nose and watery eyes. In the stomach, histamine stimulates the secretion of acid.

Understanding the central role of histamine in these biochemical processes has helped chemists design drugs to counteract some of its undesirable effects.





Figure 25.5
Three common alkaloids—
Atropine, nicotine, and coniine

atropine

nightshade

- **Atropine** is an alkaloid isolated from *Atropa belladonna*, the deadly nightshade plant. In the Renaissance, women used the juice of the berries of the nightshade to enlarge the pupils of their eyes for cosmetic reasons. Atropine causes an increase in heart rate, relaxes smooth muscles, and interferes with nerve impulses transmitted by acetylcholine. In higher doses atropine is poisonous, leading to convulsions, coma, and death.

nicotine

tobacco

- **Nicotine** is an addictive and highly toxic compound isolated from tobacco. In small doses it acts as a stimulant, but in large doses it causes depression, nausea, and even death. Nicotine is synthesized in plants as a defense against insect predators, and is used commercially as an insecticide.

coniine


hemlock

- **Coniine**, a poisonous alkaloid isolated from the seeds, leaves, and roots of hemlock (*Conium maculatum*), has been known since ancient times. Ingestion causes weakness, paralysis, and finally death. The Greek philosopher Socrates was executed by being forced to drink a potion prepared from hemlock in 339 B.C.


Antihistamines bind to the same active site of the enzyme that binds histamine in the cell, but they evoke a different response. An antihistamine like **fexofenadine** (trade name Allegra), for example, inhibits vasodilation, so it is used to treat the symptoms of the common cold and allergies. Unlike many antihistamines, fexofenadine does not cause drowsiness because it binds to histamine receptors but does not cross the blood–brain barrier, so it does not affect the central nervous system. **Cimetidine** (trade name Tagamet) is a histamine mimic that blocks the secretion of hydrochloric acid in the stomach, so it is used to treat individuals with ulcers.

25.6C Derivatives of 2-Phenylethylamine


A large number of physiologically active compounds are derived from **2-phenylethylamine**, $C_6H_5CH_2CH_2NH_2$. Some of these compounds are synthesized in cells and needed to maintain healthy mental function. Others are isolated from plant sources or are synthesized in the laboratory and have a profound effect on the brain because they interfere with normal neurochemistry. These compounds include **adrenaline**, **noradrenaline**, **methamphetamine**, and **mescaline**. Each contains a benzene ring bonded to a two-carbon unit with a nitrogen atom (shown in red).




a hormone secreted in response to stress
(Chapter 7, introductory molecule)



a neurotransmitter that increases heart rate
and dilates air passages



an addictive stimulant sold as
speed, meth, or crystal meth



a hallucinogen isolated from peyote, a cactus native
to the southwestern United States and Mexico



Cocaine, amphetamines, and several other addicting drugs increase the level of dopamine in the brain, which results in a pleasurable “high.” With time, the brain adapts to increased dopamine levels, so more drug is required for the same sensation.



Bufo toads from the Amazon jungle are the source of the hallucinogen bufotenin.


Another example, **dopamine**, is a neurotransmitter, a chemical messenger released by one nerve cell (neuron), which then binds to a receptor in a neighboring target cell (Figure 25.6). Dopamine affects brain processes that control movement and emotions, so proper dopamine levels are necessary to maintain an individual’s mental and physical health. For example, when dopamine-producing neurons die, the level of dopamine drops, resulting in the loss of motor control symptomatic of Parkinson’s disease.

Serotonin is a neurotransmitter that plays an important role in mood, sleep, perception, and temperature regulation. A deficiency of serotonin causes depression. Understanding the central role of serotonin in determining one’s mood has led to the development of a variety of drugs for the treatment of depression. The most widely used antidepressants today are selective serotonin reuptake inhibitors (SSRIs). These drugs act by inhibiting the reuptake of serotonin by the neurons that produce it, thus effectively increasing its concentration. Fluoxetine (trade name Prozac) is a common antidepressant that acts in this way.





Drugs that interfere with the metabolism of serotonin have a profound effect on mental state. For example, bufotenin, isolated from *Bufo* toads from the Amazon jungle, and psilocin, iso-



Figure 25.6 Dopamine—A neurotransmitter



lated from *Psilocybe* mushrooms, are very similar in structure to serotonin and both cause intense hallucinations.



Problem 25.10 LSD (a hallucinogen) and codeine (a narcotic) are structurally more complex derivatives of 2-phenylethylamine. Identify the atoms of 2-phenylethylamine in each of the following compounds.



25.7 Preparation of Amines

In the preparations of a given functional group, many different starting materials form a common product (amines, in this case).

Three types of reactions are used to prepare an amine:

- [1] **Nucleophilic substitution** using nitrogen nucleophiles
- [2] **Reduction** of other nitrogen-containing functional groups
- [3] **Reductive amination** of aldehydes and ketones


25.7A Nucleophilic Substitution Routes to Amines

Nucleophilic substitution is the key step in two different methods for synthesizing amines: direct nucleophilic substitution and the Gabriel synthesis of 1° amines.

Direct Nucleophilic Substitution

Conceptually, the simplest method to synthesize an amine is by S_N2 reaction of an alkyl halide with NH_3 or an amine. The method requires two steps:


- [1] **Nucleophilic attack** of the nitrogen nucleophile forms an ammonium salt.
- [2] **Removal of a proton** on N forms the amine.




The identity of the nitrogen nucleophile determines the type of amine or ammonium salt formed as product. One new carbon–nitrogen bond is formed in each reaction. Because the reaction follows an S_N2 mechanism, the alkyl halide must be unhindered—that is, CH_3X or RCH_2X .

Although this process seems straightforward, polyalkylation of the nitrogen nucleophile limits its usefulness. **Any amine formed by nucleophilic substitution still has a nonbonded electron pair, making it a nucleophile as well.** It will react with remaining alkyl halide to form a more substituted amine. Because of this, a mixture of 1°, 2°, and 3° amines often results. Only the final product—called a quaternary ammonium salt because it has four alkyl groups on N—cannot react further, and so the reaction stops.

As a result, this reaction is most useful for preparing 1° amines by using a very large excess of NH_3 (a relatively inexpensive starting material) and for preparing quaternary ammonium salts by alkylating any nitrogen nucleophile with one or more equivalents of alkyl halide.

Useful S_N2 substitutions


Problem 25.11 Draw the product of each reaction.



The Gabriel Synthesis of 1° Amines


To avoid polyalkylation, a nitrogen nucleophile can be used that reacts in a single nucleophilic substitution reaction—that is, the reaction forms a product that does not contain a nucleophilic nitrogen atom capable of reacting further.

The **Gabriel synthesis** consists of two steps and uses a resonance-stabilized nitrogen nucleophile to synthesize 1° amines via nucleophilic substitution. The Gabriel synthesis begins with **phthalimide**, one of a group of compounds called **imides**. The **N–H bond of an imide is especially acidic** because the resulting anion is resonance stabilized by the two flanking carbonyl groups.




An acid–base reaction forms a nucleophilic anion that can react with an unhindered alkyl halide—that is, CH_3X or RCH_2X —in an S_N2 reaction to form a substitution product. This alkylated imide is then hydrolyzed with aqueous base to give a 1° amine and a dicarboxylate. This reaction is similar to the hydrolysis of amides to afford carboxylate anions and amines, as discussed in Section 22.13. The overall result of this two-step sequence is **nucleophilic substitution of X by NH_2** , so the Gabriel synthesis can be used to prepare 1° amines only.


Steps in the Gabriel synthesis




- The Gabriel synthesis converts an alkyl halide into a 1° amine by a two-step process: nucleophilic substitution followed by hydrolysis.



Problem 25.12 What alkyl halide is needed to prepare each 1° amine by a Gabriel synthesis?



Problem 25.13 Which amines cannot be prepared by a Gabriel synthesis? Explain your choices.




25.7B Reduction of Other Functional Groups That Contain Nitrogen

Amines can be prepared by reduction of nitro compounds, nitriles, and amides. Because the details of these reactions have been discussed previously, they are presented here in summary form only.


[1] From nitro compounds (Section 18.14C)

Nitro groups are reduced to 1° amines using a variety of reducing agents.




[2] From nitriles (Section 22.18B)

Nitriles are reduced to 1° amines with LiAlH₄.




Because a cyano group is readily introduced by S_N2 substitution of alkyl halides with ⁻CN, this provides a **two-step method to convert an alkyl halide to a 1° amine with one more carbon atom**. The conversion of CH₃Br to CH₃CH₂NH₂ illustrates this two-step sequence.




[3] From amides (Section 20.7B)


Primary (1°), 2°, and 3° amides are reduced to 1°, 2°, and 3° amines, respectively, by using LiAlH₄.



Problem 25.14 What nitro compound, nitrile, and amide are reduced to each compound?




Problem 25.15 What amine is formed by reduction of each amide?



Problem 25.16 Explain why isopropylamine [(CH₃)₂CHNH₂] can be prepared by reduction of a nitro compound, but cannot be prepared by reduction of a nitrile, even though it is a 1° amine.

Problem 25.17 Which amines cannot be prepared by reduction of an amide?




25.7C Reductive Amination of Aldehydes and Ketones

Reductive amination is a two-step method that converts aldehydes and ketones into 1°, 2°, and 3° amines. Let's first examine this method using NH₃ to prepare 1° amines. There are two distinct parts in reductive amination:


- [1] **Nucleophilic attack of NH₃ on the carbonyl group forms an imine** (Section 21.11A), which is not isolated; then,
- [2] **Reduction of the imine forms an amine** (Section 20.7B).

**Reductive amination—
A two-step process**




- Reductive amination replaces a C=O by a C–H and C–N bond.


The most effective reducing agent for this reaction is sodium cyanoborohydride (NaBH_3CN). This hydride reagent is a derivative of sodium borohydride (NaBH_4), formed by replacing one H atom by CN.



Reductive amination combines two reactions we have already learned in a different way. Two examples are shown. The second reaction is noteworthy because the product is **amphetamine**, a potent central nervous system stimulant.



With a 1° or 2° amine as starting material, reductive amination is used to prepare 2° and 3° amines, respectively. Note the result: Reductive amination uses an aldehyde or ketone to replace one H atom on a nitrogen atom by an alkyl group, making a more substituted amine.



The synthesis of methamphetamine (Section 25.6C) by reductive amination is illustrated in Figure 25.7.

Problem 25.18

Draw the product of each reaction.





Figure 25.7


Synthesis of methamphetamine by reductive amination



- In reductive amination, one of the H atoms bonded to N is replaced by an alkyl group. As a result, a 1° amine is converted to a 2° amine and a 2° amine is converted to a 3° amine. In this reaction, CH_3NH_2 (a 1° amine) is converted to methamphetamine (a 2° amine).


To use reductive amination in synthesis, you must be able to determine what aldehyde or ketone and nitrogen compound are needed to prepare a given amine—that is, you must work backwards in the retrosynthetic direction. Keep in mind the following two points:

- One alkyl group on N comes from the carbonyl compound.
- The remainder of the molecule comes from NH_3 or an amine.



For example, 2-phenylethylamine is a 1° amine, so it has only one alkyl group bonded to N. This alkyl group must come from the carbonyl compound, and the rest of the molecule then comes from the nitrogen component. **For a 1° amine, the nitrogen component must be NH_3 .**


Retrosynthetic analysis for preparing 2-phenylethylamine:



There is usually more than one way to use reductive amination to synthesize 2° and 3° amines, as shown in Sample Problem 25.2 for a 2° amine.


Sample Problem 25.2

What aldehyde or ketone and nitrogen compound are needed to synthesize *N*-ethylcyclohexanamine by a reductive amination reaction?




Solution

Because *N*-ethylcyclohexanamine has two different alkyl groups bonded to the N atom, either R group can come from the carbonyl component and there are two different ways to form a C-N bond by reductive amination.




Because reductive amination adds one R group to a nitrogen atom, both routes to form the 2° amine begin with a 1° amine.

Problem 25.19 What starting materials are needed to prepare each drug using reductive amination? Give all possible pairs of compounds when more than one route is possible.



rimantadine
antiviral used to treat influenza




pseudoephedrine
nasal decongestant

Problem 25.20 (a) Explain why phentermine [$\text{PhCH}_2\text{C}(\text{CH}_3)_2\text{NH}_2$] can't be made by a reductive amination reaction. (b) Give a systematic name for phentermine, one of the components of the banned diet drug fen-phen.


25.8 Reactions of Amines—General Features

- The chemistry of amines is dominated by the lone pair of electrons on nitrogen.

Only three elements in the second row of the periodic table have nonbonded electron pairs in neutral organic compounds: nitrogen, oxygen, and fluorine. Because basicity and nucleophilicity decrease across the row, **nitrogen is the most basic and most nucleophilic** of these elements.




- Amines are stronger bases and nucleophiles than other neutral organic compounds.




- Amines react as bases with compounds that contain acidic protons.
- Amines react as nucleophiles with compounds that contain electrophilic carbons.

25.9 Amines as Bases

Amines react as bases with a variety of organic and inorganic acids.




What acids can be used to protonate an amine? Equilibrium favors the products of an acid–base reaction when the weaker acid and base are formed. Because the pK_a of many protonated amines is 10–11, the pK_a of the starting acid must be **less than 10** for equilibrium to favor the products. Amines are thus readily protonated by strong inorganic acids like HCl and H_2SO_4 , and by carboxylic acids as well.



The principles used in an extraction procedure were detailed in Section 19.12.

Because amines are protonated by aqueous acid, they can be separated from other organic compounds by extraction using a separatory funnel. **Extraction separates compounds based on solubility differences.** When an amine is protonated by aqueous acid, its solubility properties change.

For example, when cyclohexylamine is treated with aqueous HCl, it is protonated, forming an ammonium salt. Because the ammonium salt is ionic, it is soluble in water, but insoluble in organic solvents. A similar acid–base reaction does not occur with other organic compounds like alcohols, which are much less basic.




This difference in acid–base chemistry can be used to separate cyclohexylamine and cyclohexanol by the stepwise extraction procedure illustrated in Figure 25.8.

Figure 25.8 Separation of cyclohexylamine and cyclohexanol by an extraction procedure

Step [1] Dissolve cyclohexylamine and cyclohexanol in CH₂Cl₂.

Step [2] Add 10% HCl solution to form two layers.

Step [3] Separate the layers.



- Both compounds dissolve in the organic solvent CH₂Cl₂.

- Adding 10% aqueous HCl solution forms two layers. When the two layers are mixed, the HCl protonates the amine (RNH₂) to form RNH₃⁺Cl⁻, which dissolves in the aqueous layer.
- The cyclohexanol remains in the CH₂Cl₂ layer.


- Draining the lower layer out the bottom stopcock separates the two layers.
- Cyclohexanol (dissolved in CH₂Cl₂) is in one flask. The ammonium salt, RNH₃⁺Cl⁻ (dissolved in water), is in another flask.

- An amine can be separated from other organic compounds by converting it to a water-soluble ammonium salt by an acid–base reaction.

Thus, the water-soluble salt $C_6H_{11}NH_3^+Cl^-$ (obtained by protonation of $C_6H_{11}NH_2$) can be separated from water-insoluble cyclohexanol by an aqueous extraction procedure.


Problem 25.21

Draw the products of each acid–base reaction. Indicate whether equilibrium favors the reactants or products.




Many antihistamines and decongestants are sold as their hydrochloride salts.

Many water-insoluble amines with useful medicinal properties are sold as their water-soluble ammonium salts, which are more easily transported through the body in the aqueous medium of the blood. Benadryl, formed by treating diphenhydramine with HCl, is an over-the-counter antihistamine that is used to relieve the itch and irritation of skin rashes and hives.



Problem 25.22


Write out steps to show how each of the following pairs of compounds can be separated by an extraction procedure.



25.10 Relative Basicity of Amines and Other Compounds

The relative acidity of different compounds can be compared using their pK_a values. The relative *basicity* of different compounds (such as amines) can be compared using the pK_a values of their *conjugate acids*.

- The weaker the conjugate acid, the higher its pK_a and the stronger the base.




To compare the basicity of two compounds, keep in mind the following:

- Any factor that increases the electron density on the N atom increases an amine's basicity.
- Any factor that decreases the electron density on N decreases an amine's basicity.

25.10A Comparing an Amine and NH₃

Because alkyl groups are electron donating, they increase the electron density on nitrogen, which makes an amine like CH₃CH₂NH₂ more basic than NH₃. In fact, the pK_a of CH₃CH₂NH₃⁺ is higher than the pK_a of NH₄⁺, so **CH₃CH₂NH₂ is a stronger base than NH₃**.



The relative basicity of 1°, 2°, and 3° amines depends on additional factors, and will not be considered in this text.


- Primary (1°), 2°, and 3° alkylamines are more basic than NH₃ because of the electron-donating inductive effect of the R groups.

Problem 25.23


Which compound in each pair is more basic: (a) (CH₃)₂NH and NH₃; (b) CH₃CH₂NH₂ and ClCH₂CH₂NH₂?

25.10B Comparing an Alkylamine and an Arylamine

To compare an alkylamine (CH₃CH₂NH₂) and an arylamine (C₆H₅NH₂, aniline), we must look at the availability of the nonbonded electron pair on N. With CH₃CH₂NH₂, the electron pair is localized on the N atom. With an arylamine, however, the electron pair is now delocalized on the benzene ring. This *decreases* the electron density on N, and makes C₆H₅NH₂ less basic than CH₃CH₂NH₂.



Once again, pK_a values support this reasoning. Because the pK_a of CH₃CH₂NH₃⁺ is higher than the pK_a of C₆H₅NH₃⁺, **CH₃CH₂NH₂ is a stronger base than C₆H₅NH₂**.




- Arylamines are less basic than alkylamines because the electron pair on N is delocalized.

Substituted anilines are more or less basic than aniline depending on the nature of the substituent.

- Electron-donor groups add electron density to the benzene ring, making the arylamine more basic than aniline.


D = electron-donor group



D	
-NH ₂	
-OH	
-OR	
-NHCOR	
-R	

- Electron-withdrawing groups remove electron density from the benzene ring, making the arylamine less basic than aniline.

W = electron-withdrawing group




W	
-X	-CN
-CHO	-SO ₃ H
-COR	-NO ₂
-COOR	-NR ₃ ⁺
-COOH	

The effect of electron-donating and electron-withdrawing groups on the acidity of substituted benzoic acids was discussed in Section 19.11.


Whether a substituent donates or withdraws electron density depends on the balance of its inductive and resonance effects (Section 18.6 and Figure 18.7).

Sample Problem 25.3


Rank the following compounds in order of increasing basicity.



aniline




p-nitroaniline



p-methylaniline
(*p*-toluidine)


Solution

***p*-Nitroaniline:** NO₂ is an electron-withdrawing group, making the amine **less basic** than aniline.




The lone pair on N is delocalized on the O atom, decreasing the basicity of the amine.


***p*-Methylaniline:** CH₃ has an electron-donating inductive effect, making the amine **more basic** than aniline.




CH₃ inductively donates electron density, increasing the basicity of the amine.



p-nitroaniline



aniline




p-methylaniline
(*p*-toluidine)

Increasing basicity




Figure 25.9
Electrostatic potential plots
of substituted anilines



The NH_2 group gets more electron rich as the para substituent changes from $\text{NO}_2 \rightarrow \text{H} \rightarrow \text{CH}_3$. This is indicated by the color change around NH_2 (from green to yellow to red) in the electrostatic potential plot.

The electrostatic potential plots in Figure 25.9 demonstrate that the electron density of the nitrogen atoms in these anilines increases in the order shown.

Problem 25.24 Rank the compounds in each group in order of increasing basicity.



25.10C Comparing an Alkylamine and an Amide


To compare the basicity of an alkylamine (RNH_2) and an amide (RCONH_2), we must once again compare the availability of the nonbonded electron pair on nitrogen. With RNH_2 , the electron pair is localized on the N atom. With an amide, however, the electron pair is delocalized on the carbonyl oxygen by resonance. This *decreases* the electron density on N, making **an amide much less basic than an alkylamine**.




The electron pair on N is delocalized on O by resonance.

- Amides are much less basic than amines because the electron pair on N is delocalized.

In fact, amides are not much more basic than any carbonyl compound. When an amide is treated with acid, **protonation occurs at the carbonyl oxygen, not the nitrogen**, because the resulting cation is resonance stabilized. The product of protonation of the NH_2 group cannot be resonance stabilized.




Problem 25.25 Rank the following compounds in order of increasing basicity.



25.10D Heterocyclic Aromatic Amines


To determine the relative basicity of nitrogen heterocycles that are also aromatic, you must know whether the nitrogen lone pair is part of the aromatic π system.

For example, pyridine and pyrrole are both aromatic, but the nonbonded electron pair on the N atom in these compounds is located in different orbitals. Recall from Section 17.8C that the **lone pair of electrons in pyridine occupies an sp^2 hybridized orbital**, perpendicular to the plane of the molecule, so it is *not* part of the aromatic system, whereas that of pyrrole resides in a p orbital, making it part of the aromatic system. **The lone pair on pyrrole, therefore, is delocalized on all of the atoms of the five-membered ring**, making pyrrole a much weaker base than pyridine.



Protonation of pyrrole occurs at a ring carbon, not the N atom, as noted in Problem 17.51.

As a result, the pK_a of the conjugate acid of pyrrole is much less than that for the conjugate acid of pyridine.




- Pyrrole is much less basic than pyridine because its lone pair of electrons is part of the aromatic π system.

25.10E Hybridization Effects


The effect of hybridization on the acidity of an H–A bond was first discussed in Section 2.5D.

The hybridization of the orbital that contains an amine's lone pair also affects its basicity. This is illustrated by comparing the basicity of **piperidine** and **pyridine**, two nitrogen heterocycles. The lone pair in piperidine resides in an sp^3 hybrid orbital that has 25% s -character. The lone pair in pyridine resides in an sp^2 hybrid orbital that has 33% s -character.




- The higher the percent s -character of the orbital containing the lone pair, the more tightly the lone pair is held, and the weaker the base.

Pyridine is a weaker base than piperidine because its nonbonded pair of electrons resides in an sp^2 hybrid orbital. Although pyridine is an aromatic amine, its lone pair is *not* part of the delocalized π system, so its basicity is determined by the hybridization of its N atom. As a result, the pK_a value for the conjugate acid of pyridine is much lower than that for the conjugate acid of piperidine, making pyridine the weaker base.



Problem 25.26 Which nitrogen atom in each compound is more basic?



25.10F Summary

Acid–base chemistry is central to many processes in organic chemistry, so it has been a constant theme throughout this text. Tables 25.2 and 25.3 organize and summarize the acid–base principles

Table 25.2 Factors That Determine Amine Basicity








Factor	Example
[1] Inductive effects: Electron-donating groups bonded to N increase basicity.	• RNH_2 , R_2NH , and R_3N are more basic than NH_3 .
[2] Resonance effects: Delocalizing the lone pair on N decreases basicity.	• Arylamines ($C_6H_5NH_2$) are less basic than alkylamines (RNH_2). • Amides ($RCONH_2$) are much less basic than amines (RNH_2).
[3] Aromaticity: Having the lone pair on N as part of the aromatic π system decreases basicity.	• Pyrrole is less basic than pyridine.  less basic  more basic
[4] Hybridization effects: Increasing the percent s -character in the orbital with the lone pair decreases basicity.	• Pyridine is less basic than piperidine.  less basic  more basic

Table 25.3 Table of pK_a Values of Some Representative Organic Nitrogen Compounds


Compound	pK_a of the conjugate acid	Comment	
Ammonia	NH_3	9.3	
Alkylamines		11.1	Alkylamines have pK_a values of ~ 10–11.
	$(CH_3CH_2)_2NH$	11.1	
	$(CH_3CH_2)_3N$	11.0	
	$CH_3CH_2NH_2$	10.8	
Arylamines	$p\text{-}CH_3OC_6H_4NH_2$	5.3	The pK_a decreases as the electron density of the benzene ring decreases.
	$p\text{-}CH_3C_6H_4NH_2$	5.1	
	$C_6H_5NH_2$	4.6	
	$p\text{-}NO_2C_6H_4NH_2$	1.0	
Heterocyclic aromatic amines		5.3	The pK_a depends on whether the lone pair on N is localized or delocalized.
		0.4	
Amides	$RCONH_2$	-1	

discussed in Section 25.10. The principles in these tables can be used to determine the most basic site in a molecule that has more than one nitrogen atom, as shown in Sample Problem 25.4.

Sample Problem 25.4


 Which N atom in chloroquine is the strongest base?


Since 1945 chloroquine has been used to treat malaria, an infectious disease caused by a protozoan parasite that is spread by the *Anopheles* mosquito.




Solution

Examine the nitrogen atoms in chloroquine, labeled N1, N2, and N3, and recall that decreasing the electron density on N decreases basicity.



- N1 is bonded to an aromatic ring, so its lone pair is delocalized in the ring like aniline, decreasing basicity.
- The lone pair is localized on N2, but N2 is sp^2 hybridized. Increasing percent s -character decreases basicity.
- N3 has a localized lone pair and is sp^3 hybridized, making it the most basic site in the molecule.

Problem 25.27 Which N atom in each drug is more basic?



25.11 Amines as Nucleophiles


Amines react as nucleophiles with electrophilic carbon atoms. The details of these reactions have been described in Chapters 21 and 22, so they are summarized here only to emphasize the similar role that the amine nitrogen plays.

- Amines attack carbonyl groups to form products of nucleophilic addition or substitution.

The nature of the product depends on the carbonyl electrophile. These reactions are limited to 1° and 2° amines, because only these compounds yield neutral organic products.


[1] Reaction of 1° and 2° amines with aldehydes and ketones (Sections 21.11–21.12)

Aldehydes and ketones react with 1° amines to form **imines** and with 2° amines to form **enamines**. Both reactions involve nucleophilic addition of the amine to the carbonyl group to form a carbinolamine, which then loses water to form the final product.




[2] Reaction of NH₃ and 1° and 2° amines with acid chlorides and anhydrides (Sections 22.8–22.9)


Acid chlorides and anhydrides react with NH₃, 1° amines, and 2° amines to form 1°, 2°, and 3° **amides**, respectively. These reactions involve attack of the nitrogen nucleophile on the carbonyl group followed by elimination of a leaving group (Cl⁻ or RCOO⁻). The overall result of this reaction is substitution of the leaving group by the nitrogen nucleophile.



Problem 25.28 Draw the products formed when each carbonyl compound reacts with the following amines: [1] $\text{CH}_3\text{CH}_2\text{CH}_2\text{NH}_2$; [2] $(\text{CH}_3\text{CH}_2)_2\text{NH}$.



The conversion of amines to amides is useful in the synthesis of substituted anilines. For example, aniline itself does not undergo Friedel–Crafts reactions (Section 18.10B). Instead, its basic lone pair on N reacts with the Lewis acid (AlCl_3) to form a deactivated complex that does not undergo further reaction.



The N atom of an amide, however, is much less basic than the N atom of an amine, so it does not undergo a similar Lewis acid–base reaction with AlCl_3 . A three-step reaction sequence involving an intermediate amide can thus be used to form the products of the Friedel–Crafts reaction.

- [1] **Convert the amine (aniline) into an amide (acetanilide).**
- [2] **Carry out the Friedel–Crafts reaction.**
- [3] **Hydrolyze the amide** to generate the free amino group.

This three-step procedure is illustrated in Figure 25.10. In this way, **the amide serves as a protecting group for the NH_2 group**, in much the same way that *tert*-butyldimethylsilyl ethers and acetals are used to protect alcohols and carbonyls, respectively (Sections 20.12 and 21.15).

Problem 25.29 Devise a synthesis of each compound from aniline ($\text{C}_6\text{H}_5\text{NH}_2$).





Figure 25.10
An amide as a protecting group for an amine



A three-step sequence uses an amide as a protecting group.


[1] Treatment of aniline with acetyl chloride (CH_3COCl) forms an amide (acetanilide).

[2] Acetanilide, having a much less basic N atom compared to aniline, undergoes electrophilic aromatic substitution under Friedel-Crafts conditions, forming a mixture of ortho and para products.

[3] Hydrolysis of the amide forms the Friedel-Crafts substitution products.

25.12 Hofmann Elimination

Amines, like alcohols, contain a poor leaving group. To undergo a β elimination reaction, for example, a 1° amine would need to lose the elements of NH_3 across two adjacent atoms. The leaving group, NH_2^- , is such a strong base, however, that this reaction does not occur.




Problem: NH_2^- is a poor leaving group.


The only way around this obstacle is to convert NH_2^- into a better leaving group. The most common method to accomplish this is called a **Hofmann elimination**, which converts an amine into a quaternary ammonium salt prior to β elimination.

25.12A Details of the Hofmann Elimination

The **Hofmann elimination** converts an amine into an alkene.



The Hofmann elimination consists of three steps, as shown for the conversion of propylamine to propene.




- In Step [1], the amine reacts as a nucleophile in an S_N2 reaction with excess CH_3I to form a quaternary ammonium salt. **The $\text{N}(\text{CH}_3)_3$ group thus formed is a much better leaving group than NH_2 .**
- Step [2] converts one ammonium salt into another one with a different anion. The silver(I) oxide, Ag_2O , replaces the I^- anion with OH^- , a strong base.
- When the ammonium salt is heated in Step [3], **OH^- removes a proton from the β carbon atom**, forming the new π bond of the alkene. The mechanism of elimination is **E2**, so:

- All bonds are broken and formed in a single step.
- Elimination occurs through an anti periplanar geometry—that is, H and $\text{N}(\text{CH}_3)_3$ are oriented on opposite sides of the molecule.


The general E2 mechanism for the Hofmann elimination is shown in Mechanism 25.1.



Mechanism 25.1 The E2 Mechanism for the Hofmann Elimination




All Hofmann elimination reactions result in the formation of a new π bond between the α and β carbon atoms, as shown for cyclohexylamine and 2-phenylethylamine.






To help remember the reagents needed for the steps of the Hofmann elimination, keep in mind what happens in each step.

- **Step [1]** makes a good leaving group by forming a quaternary ammonium salt.
- **Step [2]** provides the strong base, OH^- , needed for elimination.
- **Step [3]** is the E2 elimination that forms the new π bond.

Problem 25.30 Draw an energy diagram for the following reaction. Label the axes, the starting materials, E_a , and ΔH° . Assume the reaction is exothermic. Draw a structure for the likely transition state.



Problem 25.31 Draw the product formed by treating each compound with excess CH_3I , followed by Ag_2O , and then heat.




25.12B Regioselectivity of the Hofmann Elimination

There is one major difference between a Hofmann elimination and other E2 eliminations.

- When constitutional isomers are possible, the major alkene has the *less* substituted double bond in a Hofmann elimination.


For example, Hofmann elimination of the elements of H and N(CH₃)₃ from 2-methylcyclopentanamine yields two constitutional isomers: the disubstituted alkene **A** (the major product) and the trisubstituted alkene **B** (the minor product).



This regioselectivity distinguishes a Hofmann elimination from other E2 eliminations, which form the more substituted double bond by the Zaitsev rule (Section 8.5). This result is sometimes explained by the size of the leaving group, N(CH₃)₃. **In a Hofmann elimination, the base removes a proton from the less substituted, more accessible β carbon atom, because of the bulky leaving group on the nearby α carbon.**

Sample Problem 25.5

Draw the major product formed from Hofmann elimination of the following amine.



Solution

The amine has three β carbons but two of them are identical, so two alkenes are possible. Draw elimination products by forming alkenes having a C=C between the α and β carbons. The major product has the **less substituted double bond**—that is, the alkene with the C=C between the α and β₁ carbons in this example.






Figure 25.11 contrasts the products formed by E2 elimination reactions using an alkyl halide and an amine as starting materials. Treatment of the alkyl halide (2-bromopentane) with base forms

Figure 25.11
A comparison of E2 elimination reactions using alkyl halides and amines




the more substituted alkene as the major product, following the **Zaitsev rule**. In contrast, the three-step Hofmann sequence of an amine (2-pentanamine) forms the less substituted alkene as major product.

Problem 25.32 Draw the major product formed by treating each amine with excess CH_3I , followed by Ag_2O , and then heat.




Problem 25.33 Draw the major product formed in each reaction.




25.13 Reaction of Amines with Nitrous Acid

Nitrous acid, HNO_2 , is a weak, unstable acid formed from NaNO_2 and a strong acid like HCl .




In the presence of acid, nitrous acid decomposes to ^+NO , the **nitrosonium ion**. This electrophile then goes on to react with the nucleophilic nitrogen atom of amines to form **diazonium salts** (RN_2^+Cl^-) from 1° amines and **N-nitrosamines** ($\text{R}_2\text{NN}=\text{O}$) from 2° amines.



25.13A Reaction of ^+NO with 1° Amines

Nitrous acid reacts with 1° alkylamines and arylamines to form **diazonium salts**. This reaction is called **diazotization**.


Preparation of diazonium salts







The mechanism for this reaction consists of many steps. It begins with nucleophilic attack of the amine on the nitrosonium ion, and it can conceptually be divided into two parts: formation of an **N-nitrosamine**, followed by loss of H_2O , as shown in Mechanism 25.2.



Mechanism 25.3 Formation of an *N*-Nitrosamine from a 2° Amine




Problem 25.34 Draw the product formed when each compound is treated with NaNO₂ and HCl.




25.14 Substitution Reactions of Aryl Diazonium Salts

Aryl diazonium salts undergo two general reactions:

- Substitution** of N₂ by an atom or a group of atoms **Z**.




- Coupling** of a diazonium salt with another benzene derivative to form an **azo compound**, a compound containing a nitrogen–nitrogen double bond.




Y = NH₂, NHR, NR₂, OH (a strong electron-donor group)

25.14A Specific Substitution Reactions


Aryl diazonium salts react with a variety of reagents to form products in which Z (an atom or group of atoms) replaces N₂, a very good leaving group. The mechanism of these reactions varies with the identity of Z, so we will concentrate on the products of the reactions, not the mechanisms.




[1] Substitution by OH—Synthesis of phenols




A diazonium salt reacts with H₂O to form a **phenol**.

[2] Substitution by Cl or Br—Synthesis of aryl chlorides and bromides


A diazonium salt reacts with copper(I) chloride or copper(I) bromide to form an **aryl chloride** or **aryl bromide**, respectively. This is called the **Sandmeyer reaction**. It provides an alternative to direct chlorination and bromination of an aromatic ring using Cl_2 or Br_2 and a Lewis acid catalyst.

[3] Substitution by F—Synthesis of aryl fluorides


A diazonium salt reacts with fluoroboric acid (HBF_4) to form an **aryl fluoride**. This is a useful reaction because aryl fluorides cannot be produced by direct fluorination with F_2 and a Lewis acid catalyst, as F_2 reacts too violently (Section 18.3).

[4] Substitution by I—Synthesis of aryl iodides

A diazonium salt reacts with sodium or potassium iodide to form an aryl iodide. This, too, is a useful reaction because aryl iodides cannot be produced by direct iodination with I_2 and a Lewis acid catalyst, as I_2 reacts too slowly (Section 18.3).

[5] Substitution by CN—Synthesis of benzonitriles


A diazonium salt reacts with copper(I) cyanide to form a benzonitrile. Because a cyano group can be hydrolyzed to a carboxylic acid, reduced to an amine or aldehyde, or converted to a ketone with organometallic reagents, this reaction provides easy access to a wide variety of benzene derivatives using chemistry described in Section 22.18.

[6] Substitution by H—Synthesis of benzene

A diazonium salt reacts with hypophosphorus acid (H_3PO_2) to form benzene. This reaction has limited utility because it reduces the functionality of the benzene ring by replacing N_2 with a hydrogen atom. Nonetheless, this reaction is useful in synthesizing compounds that have substitution patterns that are not available by other means.


Figure 25.12

The synthesis of 1,3,5-tribromobenzene from benzene




- Nitration followed by reduction forms aniline ($C_6H_5NH_2$) from benzene (Steps [1] and [2]).
- Bromination of aniline yields the tribromo derivative in Step [3].
- The NH_2 group is removed by a two-step process: diazotization with $NaNO_2$ and HCl (Step [4]), followed by substitution of the diazonium ion by H with H_3PO_2 .

For example, it is not possible to synthesize 1,3,5-tribromobenzene from benzene by direct bromination. Because Br is an ortho, para director, bromination with Br_2 and $FeBr_3$ will not add Br substituents meta to each other on the ring.




It is possible, however, to add three Br atoms meta to each other when aniline is the starting material. Because an NH_2 group is a very powerful ortho, para director, three Br atoms are introduced in a single step on halogenation (Section 18.10A). Then, the NH_2 group can be removed by diazotization and reaction with H_3PO_2 .



The complete synthesis of 1,3,5-tribromobenzene from benzene is outlined in Figure 25.12.


25.14B Using Diazonium Salts in Synthesis

Diazonium salts provide easy access to many different benzene derivatives. Keep in mind the following four-step sequence, because it will be used to synthesize many substituted benzenes.



Sample Problems 25.6 and 25.7 apply these principles to two different multistep syntheses.


Sample Problem 25.6 Synthesize *m*-chlorophenol from benzene.



Solution

Both OH and Cl are ortho, para directors, but they are located meta to each other. The OH group must be formed from a diazonium salt, which can be made from an NO₂ group by a stepwise method.


Retrosynthetic Analysis



Working backwards:


- [1] Form the OH group from NO₂ by a three-step procedure using a diazonium salt.
- [2] Introduce Cl meta to NO₂ by halogenation.
- [3] Add the NO₂ group by nitration.

Synthesis



- Nitration followed by chlorination meta to the NO₂ group forms the meta disubstituted benzene (Steps [1]–[2]).
- Reduction of the nitro group followed by diazotization forms the diazonium salt in Step [4], which is then converted to the desired phenol by treatment with H₂O (Step [5]).


Sample Problem 25.7 Synthesize *p*-bromobenzaldehyde from benzene.



Solution

Because the two groups are located para to each other and Br is an ortho, para director, Br should be added to the ring first. To add the CHO group, recall that it can be formed from CN by reduction.


Retrosynthetic Analysis



Working backwards:


- [1] Form the CHO group by reduction of CN.
- [2] Prepare the CN group from an NO₂ group by a three-step sequence using a diazonium salt.
- [3] Introduce the NO₂ group by nitration, para to the Br atom.
- [4] Introduce Br by bromination with Br₂ and FeBr₃.

Synthesis




- Bromination followed by nitration forms a disubstituted benzene with two para substituents (Steps [1]–[2]), which can be separated from its undesired ortho isomer.
- Reduction of the NO₂ group followed by diazotization forms the diazonium salt in Step [4], which is converted to a nitrile by reaction with CuCN (Step [5]).
- Reduction of the CN group with DIBAL-H (a mild reducing agent) forms the CHO group and completes the synthesis.

Problem 25.35 Draw the product formed in each reaction.




Problem 25.36 Devise a synthesis of each compound from benzene.




25.15 Coupling Reactions of Aryl Diazonium Salts

The second general reaction of diazonium salts is **coupling**. When a diazonium salt is treated with an aromatic compound that contains a strong electron-donor group, the two rings join together to form an **azo compound**, a compound with a nitrogen–nitrogen double bond.



Synthetic dyes are described in more detail in Section 25.16.

Azo compounds are highly conjugated, rendering them colored (Section 16.15). Many of these compounds, such as the azo compound “butter yellow,” are synthetic dyes. Butter yellow was once used to color margarine.




This reaction is another example of **electrophilic aromatic substitution**, with the **diazonium salt acting as the electrophile**. Like all electrophilic substitutions (Section 18.2), the mechanism has two steps: addition of the electrophile (the diazonium ion) to form a resonance-stabilized carbocation, followed by deprotonation, as shown in Mechanism 25.4.




Mechanism 25.4 Azo Coupling

Step [1] Addition of the diazonium ion to form a carbocation



- **Step [1]** The electrophilic diazonium ion reacts with the electron-rich benzene ring to form a resonance-stabilized carbocation. (Only one resonance structure is drawn.)

Step [2] Loss of a proton to re-form the aromatic ring




- **Step [2]** Loss of a proton regenerates the aromatic ring.

Because a diazonium salt is weakly electrophilic, the reaction occurs only when the benzene ring has a **strong electron-donor group Y**, where **Y = NH₂, NHR, NR₂, or OH**. Although these groups activate both the ortho and para positions, para substitution occurs unless the para position already has another substituent present.


To determine what starting materials are needed to synthesize a particular azo compound, always divide the molecule into two components: **one has a benzene ring with a diazonium ion, and one has a benzene ring with a very strong electron-donor group**.

Break the molecule into two components here.



Sample Problem 25.8


What starting materials are needed to synthesize the following azo compound?




Solution

Both benzene rings in methyl orange have a substituent, but only one group, N(CH₃)₂, is a strong electron donor. In determining the two starting materials, the **diazonium ion must be bonded to the ring that is not bonded to N(CH₃)₂**.


Break the molecule into two components here.



Problem 25.37 Draw the product formed when $C_6H_5N_2^+Cl^-$ reacts with each compound.



Problem 25.38 What starting materials are needed to synthesize each azo compound?




25.16 Application: Synthetic Dyes

Azo compounds have two important applications: as dyes and as sulfa drugs, the first synthetic antibiotics (Section 25.17).

25.16A Natural and Synthetic Dyes

Until 1856, all dyes were natural in origin, obtained from plants, animals, or minerals. Three natural dyes known for centuries are **indigo**, **tyrian purple**, and **alizarin**.




The blue dye **indigo**, derived from the plant *Indigofera tinctoria*, has been used in India for thousands of years. Traders introduced it to the Mediterranean area and then to Europe. **Tyrian purple**, a natural dark purple dye obtained from the mucous gland of a Mediterranean snail of the genus *Murex*, was a symbol of royalty before the collapse of the Roman empire. **Alizarin**, a bright red dye obtained from madder root (*Rubia tinctorum*), a plant native to India and north-eastern Asia, has been found in cloth entombed with Egyptian mummies.

Because all three of these dyes were derived from natural sources, they were difficult to obtain, making them expensive and available only to the privileged. This all changed when William Henry Perkin, an 18-year-old student with a makeshift home laboratory, serendipitously prepared a purple dye, which would later be called mauveine, during his failed attempt to synthesize the antimalarial drug quinine. Mauveine is a mixture of two compounds that differ in the presence of only one methyl group on one of the aromatic rings.




A purple shawl dyed with Perkin's mauveine



Perkin's discovery marked the beginning of the chemical industry. He patented the dye and went on to build a factory to commercially produce it on a large scale. This event began the surge of research in organic chemistry, not just in the synthesis of dyes, but in the production of perfumes, anesthetics, inks, and drugs as well. Perkin was a wealthy man when he retired at the age of 36 to devote the rest of his life to basic chemical research. The most prestigious award given by the American Chemical Society is named the Perkin Medal in his honor.

Many common synthetic dyes, such as alizarine yellow R, para red, and Congo red, are **azo compounds**, prepared by the diazonium coupling reaction described in Section 25.15.



Although natural and synthetic dyes are quite varied in structure, **all of them are colored because they are highly conjugated**. A molecule with eight or more π bonds in conjugation absorbs light in the visible region of the electromagnetic spectrum (Section 16.15A), taking on the color from the visible spectrum that it does *not* absorb.


Problem 25.39

(a) What two components are needed to prepare para red by azo coupling? (b) What two components are needed to prepare alizarine yellow R?


25.16B How Dyes Bind to Fabric

To be classified as a dye, a compound must be colored *and* it must bind to fabric. There are many ways for this binding to occur. Compounds that bind to fabric by some type of attractive forces are called **direct dyes**. These attractive forces may involve electrostatic interactions, van der Waals forces, hydrogen bonding, or sometimes, even covalent bonding. The type of interaction depends on the structure of the dye and the fiber. Thus, a compound that is good for dyeing wool or silk, both polyamides, may be poor for dyeing cotton, a carbohydrate (Figure 22.4).

Wool and silk contain charged functional groups, such as NH_3^+ and COO^- . Because of this, they bind to ionic dyes by electrostatic interactions. For example, positively charged NH_3^+ groups bonded to the protein backbone are electrostatically attracted to anionic groups in a dye like methyl orange.



Cotton, on the other hand, binds dyes by hydrogen bonding interactions with its many OH groups. Thus, Congo red is bound to the cellulose backbone by hydrogen bonds.



Problem 25.40

Explain why Dacron, a polyester first discussed in Section 22.16B, does not bind well with an anionic dye such as methyl orange.


25.17 Application: Sulfa Drugs

Although they may seem quite unrelated, the synthesis of colored dyes led to the development of the first synthetic antibiotics. Much of the early effort in this field was done by the German chemist Paul Ehrlich, who worked with synthetic dyes and used them to stain tissues. This led him on a search for dyes that were lethal to bacteria without affecting other tissue cells, hoping that these dyes could treat bacterial infections. For many years this effort was unsuccessful.

Then, in 1935, Gerhard Domagk, a German physician working for a dye manufacturer, first used a synthetic dye as a drug to kill bacteria. His daughter had contracted a streptococcal infection, and as she neared death, he gave her **prontosil**, an azo dye that inhibited the growth of certain bacteria in mice. His daughter recovered, and the modern era of synthetic antibiotics was initiated. For his pioneering work, Domagk was awarded the Nobel Prize in Physiology or Medicine in 1939.



Kalaupapa is a remote and inaccessible peninsula on the Hawaiian island of Molokai that has served as home for individuals suffering from Hansen's disease, commonly called leprosy. Once thought to be very contagious, Hansen's disease is now known to be a treatable bacterial infection completely cured with sulfa drugs.



Prontosil and other sulfur-containing antibiotics are collectively called **sulfa drugs**. Prontosil is not the active agent itself. In cells, it is metabolized to **sulfanilamide**, the active drug. To understand how sulfanilamide functions as an antibacterial agent we must examine **folic acid**, which microorganisms synthesize from *p*-aminobenzoic acid.






Figure 25.13

Two common sulfa drugs



- Sulfamethoxazole is the sulfa drug in Bactrim, and sulfisoxazole is sold as Gantrisin. Both drugs are commonly used in the treatment of ear and urinary tract infections.

Sulfanilamide and *p*-aminobenzoic acid are similar in size and shape and have related functional groups. Thus, when sulfanilamide is administered, bacteria attempt to use it in place of *p*-aminobenzoic acid to synthesize folic acid. Derailing folic acid synthesis means that the bacteria cannot grow and reproduce. Sulfanilamide only affects bacterial cells, though, because humans do not synthesize folic acid, and must obtain it from their diets.



These compounds are similar in size and shape.

Many other compounds of similar structure have been prepared and are still widely used as antibiotics. The structures of two other sulfa drugs are shown in Figure 25.13.

KEY CONCEPTS

Amines

General Facts

- Amines are organic nitrogen compounds having the general structure RNH_2 , R_2NH , or R_3N , with a lone pair of electrons on N (25.1).
- Amines are named using the suffix *-amine* (25.3).
- All amines have polar C–N bonds. Primary (1°) and 2° amines have polar N–H bonds and are capable of intermolecular hydrogen bonding (25.4).
- The lone pair on N makes amines strong organic bases and nucleophiles (25.8).



Summary of Spectroscopic Absorptions (25.5)

Mass spectra	Molecular ion	Amines with an odd number of N atoms give an odd molecular ion.
IR absorptions	N–H	$3300\text{--}3500\text{ cm}^{-1}$ (two peaks for RNH_2 , one peak for R_2NH)
^1H NMR absorptions	NH	0.5–5 ppm (no splitting with adjacent protons)
	CH–N	2.3–3.0 ppm (deshielded $\text{C}_{\text{sp}^3}\text{--H}$)
^{13}C NMR absorption	C–N	30–50 ppm

Comparing the Basicity of Amines and Other Compounds (25.10)


- Alkylamines (RNH_2 , R_2NH , and R_3N) are more basic than NH_3 because of the electron-donating R groups (25.10A).
- Alkylamines (RNH_2) are more basic than arylamines ($\text{C}_6\text{H}_5\text{NH}_2$), which have a delocalized lone pair from the N atom (25.10B).
- Arylamines with electron-donor groups are more basic than arylamines with electron-withdrawing groups (25.10B).
- Alkylamines (RNH_2) are more basic than amides (RCONH_2), which have a delocalized lone pair from the N atom (25.10C).
- Aromatic heterocycles with a localized electron pair on N are more basic than those with a delocalized lone pair from the N atom (25.10D).
- Alkylamines with a lone pair in an sp^3 hybrid orbital are more basic than those with a lone pair in an sp^2 hybrid orbital (25.10E).

Preparation of Amines (25.7)

[1] Direct nucleophilic substitution with NH_3 and amines (25.7A)




- The mechanism is $\text{S}_{\text{N}}2$.
- The reaction works best for CH_3X or RCH_2X .
- The reaction works best to prepare 1° amines and quaternary ammonium salts.

[2] Gabriel synthesis (25.7A)




- The mechanism is $\text{S}_{\text{N}}2$.
- The reaction works best for CH_3X or RCH_2X .
- Only 1° amines can be prepared.

[3] Reduction methods (25.7B)




[4] Reductive amination (25.7C)



- Reductive amination adds one alkyl group (from an aldehyde or ketone) to a nitrogen nucleophile.
- Primary (1°), 2°, and 3° amines can be prepared.


Reactions of Amines

[1] Reaction as a base (25.9)




[2] Nucleophilic addition to aldehydes and ketones (25.11)


With 1° amines:



With 2° amines:



[3] Nucleophilic substitution with acid chlorides and anhydrides (25.11)




[4] Hofmann elimination (25.12)




[5] Reaction with nitrous acid (25.13)

With 1° amines:




With 2° amines:




Reactions of Diazonium Salts

[1] Substitution reactions (25.14)















[2] Coupling to form azo compounds (25.15)









PROBLEMS

Nomenclature

25.41 Give a systematic or common name for each compound.





25.42 Draw the structure that corresponds to each name.



25.43 Draw all constitutional isomers of molecular formula $\text{C}_4\text{H}_{11}\text{N}$, and give an acceptable name for each amine.




Chiral Compounds

25.44 How many stereogenic centers are present in each compound? Draw all possible stereoisomers.







Basicity

25.45 Which compound in each pair is the stronger base?




25.46 Rank the compounds in each group in order of increasing basicity.




25.47 How does the $\text{p}K_a$ of the conjugate acid of benzylamine ($\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$) compare to the $\text{p}K_a$'s of the conjugate acids of cyclohexylamine (10.7) and aniline (4.6)? Explain your choice.


- 25.48** Decide which N atom in each molecule is most basic and draw the product formed when each compound is treated with $\text{CH}_3\text{CO}_2\text{H}$. Benazepril (trade name Lotensin) is a β blocker used to treat high blood pressure and congestive heart failure. Varenicline (trade name Chantix) is used to help smokers quit their habit.



- 25.49** Rank the nitrogen atoms in each compound in order of increasing basicity. Isoniazid is a drug used to treat tuberculosis, whereas histamine (Section 25.6B) causes the runny nose and watery eyes associated with allergies.




- 25.50** Abilify is the trade name for aripiprazole, a drug used to treat depression, schizophrenia, and bipolar disorders. (a) Rank the N atoms in aripiprazole in order of increasing basicity. (b) What product is formed when aripiprazole is treated with HCl ?




- 25.51** Explain why *m*-nitroaniline is a stronger base than *p*-nitroaniline.

- 25.52** Explain the observed difference in the $\text{p}K_a$ values of the conjugate acids of amines **A** and **B**.




- 25.53** Why is pyrrole more acidic than pyrrolidine?




Preparation of Amines


- 25.54** How would you prepare 3-phenyl-1-propanamine ($\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$) from each compound?







- 25.55** What amide(s) can be used to prepare each amine by reduction?



- 25.56** What carbonyl and nitrogen compounds are needed to make each compound by reductive amination? When more than one set of starting materials is possible, give all possible methods.



25.57 Draw the product of each reductive amination reaction.



25.58 How would you prepare benzylamine ($C_6H_5CH_2NH_2$) from each compound? In some cases, more than one step is required.

- a. $C_6H_5CH_2Br$ c. $C_6H_5CONH_2$ e. $C_6H_5CH_3$ g. $C_6H_5NH_2$
 b. C_6H_5CN d. C_6H_5CHO f. C_6H_5COOH h. benzene

Extraction

25.59 How would you separate toluene ($C_6H_5CH_3$), benzoic acid (C_6H_5COOH), and aniline ($C_6H_5NH_2$) by an extraction procedure?

Reactions

25.60 What products are formed when *N*-ethylaniline ($C_6H_5NHCH_2CH_3$) is treated with each reagent?

- a. HCl e. CH_3I (excess) h. The product in (g), then HNO_3 , H_2SO_4
 b. CH_3COOH f. CH_3I (excess), followed by Ag_2O and Δ i. The product in (g), then [1] $LiAlH_4$; [2] H_2O
 c. $(CH_3)_2C=O$ g. CH_3CH_2COCl j. The product in (h), then H_2 , Pd-C
 d. CH_2O , $NaBH_3CN$





25.61 Draw the products formed when *p*-methylaniline ($p-CH_3C_6H_4NH_2$) is treated with each reagent.

- a. HCl e. $(CH_3)_2C=O$ h. $NaNO_2$, HCl
 b. CH_3COCl f. CH_3COCl , $AlCl_3$ i. Step (b), then CH_3COCl , $AlCl_3$
 c. $(CH_3CO)_2O$ g. CH_3COOH j. CH_3CHO , $NaBH_3CN$
 d. excess CH_3I







25.62 How would you convert $CH_3CH_2CH_2CH_2NH_2$ into each compound?

- a. $CH_3CH_2CH_2CH_2NHCO_2C_6H_5$ d. $CH_3CH_2CH_2CH_2NHCH_2C_6H_5$
 b. $CH_3CH_2CH_2CH_2N=C(CH_2CH_3)_2$ e. $CH_3CH_2CH_2CH_2NHCH_2CH_3$
 c. $CH_3CH_2CH=CH_2$ f. $[CH_3CH_2CH_2CH_2N(CH_3)_3]^+I^-$


25.63 Draw the products formed when each amine is treated with [1] CH_3I (excess); [2] Ag_2O ; [3] Δ . Indicate the major product when a mixture results.



25.64 What reagents are needed to convert acetophenone ($C_6H_5COCH_3$) into each compound? In some cases, more than one step is required.













25.65 Answer the following questions about benzphetamine, a habit-forming diet pill sold under the trade name Didrex.






- a. Label the stereogenic center(s).
 b. What amide(s) can be reduced to form benzphetamine?
 c. What carbonyl compound and amine can be used to make benzphetamine by reductive amination? Draw all possible methods.
 d. What products are formed by Hofmann elimination from benzphetamine? Label the major product.


25.66 Draw the organic products formed in each reaction.



25.67 What is the major Hofmann elimination product formed from each amine?



25.68 Draw the product formed when **A** is treated with each reagent.




- a. H_2O
b. H_3PO_2
c. CuCl
d. CuBr

- e. CuCN
f. HBF_4
g. NaI

- h. $\text{C}_6\text{H}_5\text{NH}_2$
i. $\text{C}_6\text{H}_5\text{OH}$
j. KI



25.69 Explain why so much meta product is formed when aniline is nitrated with HNO_3 and H_2SO_4 . For this reason, nitration of aniline is **not** a useful reaction to prepare either *o*- or *p*-nitroaniline.




25.70 A chiral amine **A** having the *R* configuration undergoes Hofmann elimination to form an alkene **B** as the major product. **B** is oxidatively cleaved with ozone, followed by CH_3SCH_3 , to form $\text{CH}_2=\text{O}$ and $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}$. What are the structures of **A** and **B**?

Mechanism

25.71 Draw a stepwise mechanism for each reaction.




25.72 Draw a stepwise mechanism for the following reaction.




25.73 Propose a reason why aryl diazonium salts are more stable than alkyl diazonium salts.

25.74 Alkyl diazonium salts are unstable even at low temperature. They decompose to form carbocations, which go on to form products of substitution, elimination, and (sometimes) rearrangement. Keeping this in mind, draw a stepwise mechanism that forms all of the following products.




25.75 Tertiary (3°) aromatic amines react with NaNO_2 and HCl to afford products of electrophilic aromatic substitution. Draw a stepwise mechanism for this nitrosation reaction and explain why it occurs only on benzene rings with strong ortho, para activating groups.




Synthesis

25.76 Devise a stepwise reaction sequence to convert 4-phenyl-2-butanone ($\text{PhCH}_2\text{CH}_2\text{COCH}_3$) into each alkene: (a) $\text{PhCH}_2\text{CH}_2\text{CH}=\text{CH}_2$; (b) $\text{PhCH}_2\text{CH}=\text{CHCH}_3$.

25.77 Devise a synthesis of each compound from benzene. You may use any other organic or inorganic reagents.




25.78 Devise a synthesis of each compound from aniline ($\text{C}_6\text{H}_5\text{NH}_2$) as starting material.







25.79 Devise at least three different methods to prepare *N*-methylbenzylamine ($\text{PhCH}_2\text{NHCH}_3$) from benzene, any one-carbon organic compounds, and any required reagents.






25.80 Safrole, which is isolated from sassafras (Problem 21.36), can be converted to the illegal stimulant MDMA (3,4-methylenedioxyamphetamine, "Ecstasy") by a variety of methods. (a) Devise a synthesis that begins with safrole and uses a nucleophilic substitution reaction to introduce the amine. (b) Devise a synthesis that begins with safrole and uses reductive amination to introduce the amine.






25.81 Devise a synthesis of the hallucinogen mescaline (Section 25.6) from each starting material.









25.82 Synthesize each compound from benzene. Use a diazonium salt as one of the synthetic intermediates.



25.83 Devise a synthesis of each biologically active compound from benzene.



25.84 Devise a synthesis of each compound from benzene, any organic alcohols having three carbons or fewer, and any required reagents.




Spectroscopy


25.85 Draw the structures of the eight isomeric amines that have a molecular ion in the mass spectrum at $m/z = 87$ and show two peaks in their IR spectra at $3300\text{--}3500\text{ cm}^{-1}$.

25.86 Three isomeric compounds, **A**, **B**, and **C**, all have molecular formula $C_8H_{11}N$. The 1H NMR and IR spectral data of **A**, **B**, and **C** are given below. What are their structures?


Compound **A**: IR peak at 3400 cm^{-1}





Compound **B**: IR peak at 3310 cm^{-1}



Compound **C**: IR peaks at 3430 and 3350 cm^{-1}




- 25.87** Treatment of compound **D** with LiAlH_4 followed by H_2O forms compound **E**. **D** shows a molecular ion in its mass spectrum at $m/z = 71$ and IR absorptions at $3600\text{--}3200$ and 2263 cm^{-1} . **E** shows a molecular ion in its mass spectrum at $m/z = 75$ and IR absorptions at 3636 and $3600\text{--}3200\text{ cm}^{-1}$. Propose structures for **D** and **E** from these data and the given $^1\text{H NMR}$ spectra.




Challenge Problems



- 25.88** The $\text{p}K_a$ of the conjugate acid of guanidine is 13.6, making it one of the strongest neutral organic bases. Offer an explanation.



- 25.89** Draw the product **Y** of the following reaction sequence. **Y** was an intermediate in the remarkable synthesis of cyclooctatetraene by Wilstatter in 1911.



- 25.90** Devise a synthesis of each compound from the given starting material(s). Albuterol is a bronchodilator and proparacaine is a local anesthetic.



26

Carbon–Carbon Bond-Forming Reactions in Organic Synthesis

- 26.1 Coupling reactions of organocuprate reagents
- 26.2 Suzuki reaction
- 26.3 Heck reaction
- 26.4 Carbenes and cyclopropane synthesis
- 26.5 Simmons–Smith reaction
- 26.6 Metathesis



Bombykol is a sex pheromone secreted by the female silkworm moth *Bombyx mori* to attract mates. Bombykol's structure was elucidated in 1959 using 6.4 mg of material obtained from 500,000 silkworm moths. One step in an efficient synthesis of bombykol is the Suzuki reaction, a stereospecific carbon–carbon bond-forming reaction between a vinylborane and a vinyl halide to form a conjugated diene. In Chapter 26, we learn how to prepare a variety of substrates using novel carbon–carbon bond-forming reactions.

To form the carbon skeletons of complex molecules, organic chemists need an extensive repertoire of carbon–carbon bond-forming reactions. In Chapter 20, for example, we learned about the reactions of organometallic reagents—organolithium reagents, Grignard reagents, and organocuprates—with carbonyl substrates. In Chapters 23 and 24, we studied the reactions of nucleophilic enolates that form new carbon–carbon bonds.

Chapter 26 presents more carbon–carbon bond-forming reactions that are especially useful tools in organic synthesis. While previous chapters have concentrated on the reactions of one or two functional groups, the reactions in this chapter utilize a variety of starting materials and conceptually different reactions that form many types of products. All follow one central theme—they form new carbon–carbon bonds under mild conditions, making them versatile synthetic methods.


26.1 Coupling Reactions of Organocuprate Reagents

Several carbon–carbon bond-forming reactions involve the coupling of an organic halide ($R'X$) with an organometallic reagent or alkene. Three useful reactions are discussed in Sections 26.1–26.3:


[1] Reaction of an organic halide with an organocuprate reagent (Section 26.1)



[2] Suzuki reaction: Reaction of an organic halide with an organoboron reagent in the presence of a palladium catalyst (Section 26.2)




[3] Heck reaction: Reaction of an organic halide with an alkene in the presence of a palladium catalyst (Section 26.3)



A complete list of reactions that form C–C bonds appears in Appendix D.



26.1A General Features of Organocuprate Coupling Reactions

In addition to their reactions with acid chlorides, epoxides, and α,β -unsaturated carbonyl compounds (Sections 20.13–20.15), organocuprate reagents (R_2CuLi) also react with organic halides $R'-X$ to form coupling products $R-R'$ that contain a new C–C bond. Only one R group of the organocuprate is transferred to form the product, while the other becomes part of RCu , a reaction by-product.







A variety of organic halides can be used, including methyl and 1° alkyl halides, as well as vinyl and aryl halides that contain X bonded to an sp^2 hybridized carbon. Some cyclic 2° alkyl halides give reasonable yields of product, but 3° alkyl halides are too sterically hindered. The halogen X in $R'X$ may be Cl, Br, or I.

Examples




Coupling reactions with vinyl halides are **stereospecific**. For example, reaction of *trans*-1-bromo-1-hexene with $(\text{CH}_3)_2\text{CuLi}$ forms *trans*-2-heptene as the only stereoisomer (Equation [3]).

Problem 26.1 Draw the product of each coupling reaction.




Problem 26.2 Identify reagents **A** and **B** in the following reaction scheme. This synthetic sequence was used to prepare the C_{18} juvenile hormone, the opening molecule in Chapter 20.



26.1B Using Organocuprate Couplings to Synthesize Hydrocarbons

Since organocuprate reagents (R_2CuLi) are prepared in two steps from alkyl halides (RX), this method ultimately converts two organic halides (RX and $\text{R}'\text{X}$) into a hydrocarbon $\text{R-R}'$ with a new carbon–carbon bond. A hydrocarbon can often be made by two different routes, as shown in Sample Problem 26.1.




A general ligand bonded to a metal is often designated as **L**. Pd bonded to four ligands is denoted as PdL₄.

Ac is the abbreviation for an acetyl group, **CH₃C=O**, so OAc (or ⁻OAc) is the abbreviation for acetate, **CH₃CO₂⁻**.


Organopalladium compounds—compounds that contain a carbon–palladium bond—are generally prepared in situ during the course of a reaction, from another palladium reagent such as Pd(OAc)₂ or Pd(PPh₃)₄. In most useful reactions only a catalytic amount of palladium reagent is utilized.

Two common processes, called **oxidative addition** and **reductive elimination**, dominate many reactions of palladium compounds.

- **Oxidative addition** is the addition of a reagent (such as RX) to a metal, often increasing the number of groups around the metal by two.




- **Reductive elimination** is the elimination of two groups that surround the metal, often forming new C–H or C–C bonds.



Reaction mechanisms with palladium compounds are often multistep. During the course of a reaction, the identity of some groups bonded to Pd will be known with certainty, while the identity of other ligands might not be known. Consequently, only the crucial reacting groups around a metal are usually drawn and the other ligands are not specified.


26.2B Details of the Suzuki Reaction

The **Suzuki reaction** is a palladium-catalyzed coupling of an organic halide (R'X) with an organoborane (R–BY₂) to form a product (R–R') with a new C–C bond. Pd(PPh₃)₄ is the typical palladium catalyst, and the reaction is carried out in the presence of a base such as NaOH or NaOCH₂CH₃.




Vinyl halides and aryl halides, both of which contain a halogen X bonded directly to an *sp*² hybridized carbon, are most often used, and the halogen is usually Br or I. The Suzuki reaction is completely **stereospecific**, as shown in Example [3]; a *cis* vinyl halide and a *trans* vinylborane form a *cis,trans*-1,3-diene.

Examples




The organoboranes used in the Suzuki reaction are prepared from two sources.


- **Vinylboranes**, which have a boron atom bonded to a carbon-carbon double bond, are prepared by hydroboration of an alkyne using catecholborane, a commercially available reagent. **Hydroboration adds the elements of H and B in a syn fashion to form a trans vinylborane.** With terminal alkynes, hydroboration always places the boron atom on the *less substituted* terminal carbon.



- **Arylboranes**, which have a boron atom bonded to a benzene ring, are prepared from organolithium reagents by reaction with trimethyl borate $[\text{B}(\text{OCH}_3)_3]$.



Problem 26.4 Draw the product of each reaction.




The mechanism of the Suzuki reaction can be conceptually divided into three parts: oxidative addition of $\text{R}'-\text{X}$ to the palladium catalyst, transfer of an alkyl group from the organoborane to palladium, and reductive elimination of $\text{R}-\text{R}'$, forming a new carbon-carbon bond. A general halide $\text{R}'-\text{X}$ and organoborane $\text{R}-\text{BY}_2$ are used to illustrate this process in Mechanism 26.1. Since the palladium reagent is regenerated during reductive elimination, only a catalytic amount of palladium is needed.




Mechanism 26.1 Suzuki Reaction

Part [1] Formation of an organopalladium compound by oxidative addition




- Loss of two triphenylphosphine ligands followed by oxidative addition forms an organopalladium compound.
- The organopalladium compound $(\text{Ph}_3\text{P})_2\text{Pd}(\text{R}')\text{X}$ bears no net charge, and by convention, organometallic chemists generally omit formal charges on the phosphorus ligands and the metal.

Part [2] Transfer of an alkyl group from boron to palladium



- Reaction of the organoborane $\text{R}-\text{BY}_2$ with ^-OH forms a nucleophilic boron intermediate that transfers an alkyl group from boron to palladium in Step [3].

Part [3] Reductive elimination




- Reductive elimination of $\text{R}-\text{R}'$ forms the coupling product.
- The palladium catalyst $\text{Pd}(\text{PPh}_3)_2$ is regenerated so the cycle can repeat.

The Suzuki reaction was a key step in the synthesis of **bombykol**, the sex pheromone of the female silkworm moth and the chapter-opening molecule, and **humulene**, a lipid isolated from hops, as shown in Figure 26.1. The synthesis of humulene illustrates that an intramolecular Suzuki reaction can form a ring. Sample Problem 26.2 shows how a conjugated diene can be prepared from an alkyne and vinyl halide using a Suzuki reaction.


Figure 26.1

Synthesis of two natural products using the Suzuki reaction




Sample Problem 26.2

Devise a synthesis of (1*Z*,3*E*)-1-phenyl-1,3-octadiene from 1-hexyne and (*Z*)-2-bromostyrene using a Suzuki coupling.




Solution

This synthesis can be accomplished in two steps. Hydroboration of 1-hexyne with catecholborane forms a vinylborane. Coupling of this vinylborane with (*Z*)-2-bromostyrene gives the desired 1,3-diene. The *E* configuration of the vinylborane and the *Z* configuration of the vinyl bromide are both retained in the product.




Problem 26.5

Synthesize each compound from the given starting materials.






26.3 Heck Reaction

The **Heck reaction** is a **palladium-catalyzed coupling of a vinyl or aryl halide with an alkene to form a more highly substituted alkene with a new C–C bond**. Palladium(II) acetate $[\text{Pd}(\text{OAc})_2]$ in the presence of a triarylphosphine $[\text{P}(o\text{-tolyl})_3]$ is the typical catalyst, and the reaction is carried out in the presence of a base such as triethylamine. The Heck reaction is a **substitution reaction** in which one H atom of the alkene starting material is replaced by the R' group of the vinyl or aryl halide.







The alkene component is typically ethylene or a monosubstituted alkene ($\text{CH}_2=\text{CHZ}$), and the halogen X is usually Br or I. When $\text{Z} = \text{Ph}$, COOR , or CN in a monosubstituted alkene, **the new C–C bond is formed on the less substituted carbon to afford a trans alkene**. When a vinyl halide is used as the organic halide, the reaction is **stereospecific**, as shown in Example [3]; the trans stereochemistry of the vinyl iodide is retained in the product.


Examples





Problem 26.6 Draw the coupling product formed when each pair of compounds is treated with $\text{Pd}(\text{OAc})_2$, $\text{P}(\text{o-tolyl})_3$, and $(\text{CH}_3\text{CH}_2)_3\text{N}$.



To use the Heck reaction in synthesis, you must determine what alkene and what organic halide are needed to prepare a given compound. To work backwards, locate the double bond with the aryl, COOR , or CN substituent, and break the molecule into two components at the end of the $\text{C}=\text{C}$ not bonded to one of these substituents. Sample Problem 26.3 illustrates this retrosynthetic analysis.




Sample Problem 26.3 What starting materials are needed to prepare each alkene using a Heck reaction?




Solution

To prepare an alkene of general formula $R'CH=CHZ$ by the Heck reaction, two starting materials are needed—an alkene ($CH_2=CHZ$) and a vinyl or aryl halide ($R'X$).



Problem 26.7 What starting materials are needed to prepare each compound using a Heck reaction?




The actual palladium catalyst in the Heck reaction is thought to contain a palladium atom bonded to two tri(*o*-tolyl)phosphine ligands, abbreviated as $Pd(PAr_3)_2$. In this way it resembles the divalent palladium catalyst used in the Suzuki reaction. The mechanism of the Heck reaction conceptually consists of three parts: oxidative addition of the halide $R'X$ to the palladium catalyst, addition of the resulting organopalladium reagent to the alkene, and two successive eliminations. A general organic halide $R'X$ and alkene $CH_2=CHZ$ are used to illustrate the process in Mechanism 26.2.




Mechanism 26.2 Heck Reaction

Step [1] Formation of an organopalladium compound by oxidative addition




- Oxidative addition of $R'X$ forms an organopalladium compound.

Step [2] Addition of R' and Pd to the double bond




- Addition of R' and Pd to the π bond places the Pd on the carbon with the Z substituent.

Steps [3]–[4] Two successive eliminations




- Elimination of H and Pd forms the alkene in Step [3] and transfers a hydrogen to Pd.
- Reductive elimination of HX regenerates the palladium catalyst $Pd(PAr_3)_2$ in Step [4].


The **trans** methyl groups in *trans*-2-butene become **trans** substituents in the cyclopropane. Addition from either side of the alkene yields an equal amount of two enantiomers—a **racemic mixture**.



Problem 26.8 Draw all stereoisomers formed when each alkene is treated with CHCl_3 and $\text{KOC}(\text{CH}_3)_3$.



Finally, dihalo cyclopropanes can be converted to dialkyl cyclopropanes by reaction with organocuprates (Section 26.1). For example, cyclohexene can be converted to a bicyclic product having four new C–C bonds by the following two-step sequence: cyclopropanation with dibromocarbene ($:\text{CBr}_2$) and reaction with lithium dimethylcuprate, $\text{LiCu}(\text{CH}_3)_2$.




Problem 26.9 What reagents are needed to convert 2-methylpropene $[(\text{CH}_3)_2\text{C}=\text{CH}_2]$ to each compound? More than one step may be required.





26.5 Simmons–Smith Reaction

Although the reaction of dihalocarbenes with alkenes gives good yields of halogenated cyclopropanes, this is not usually the case with **methylene**, $:\text{CH}_2$, the simplest carbene. Methylene is readily formed by heating diazomethane, CH_2N_2 , which decomposes and loses N_2 , but the reaction of $:\text{CH}_2$ with alkenes often affords a complex mixture of products. Thus, this reaction cannot be reliably used for cyclopropane synthesis.



Nonhalogenated cyclopropanes can be prepared by the reaction of an alkene with diiodomethane, CH_2I_2 , in the presence of a copper-activated zinc reagent called zinc–copper couple $[\text{Zn}(\text{Cu})]$. This process, the **Simmons–Smith reaction**, is named for H. E. Simmons and R. D. Smith, DuPont chemists who discovered the reaction in 1959.




The Simmons–Smith reaction does not involve a free carbene. Rather, the reaction of CH_2I_2 with $\text{Zn}(\text{Cu})$ forms (iodomethyl)zinc iodide, which transfers a CH_2 group to an alkene, as shown in Mechanism 26.5.




Mechanism 26.5 Simmons–Smith Reaction

Step [1] Formation of the Simmons–Smith reagent




- Reaction of CH_2I_2 with zinc–copper couple forms ICH_2ZnI , the **Simmons–Smith reagent**. Since the CH_2 group is bonded to the metal and does not exist as a free carbene, this intermediate is called a **carbenoid**.

Step [2] Formation of the cyclopropane ring




- The Simmons–Smith reagent transfers a CH_2 group to the alkene, forming two new C–C bonds in a single step.

The Simmons–Smith reaction is stereospecific. The relative position of substituents in the alkene reactant is *retained* in the cyclopropane product, as shown for the conversion of *cis*-3-hexene to *cis*-1,2-diethylcyclopropane.



Problem 26.10 What product is formed when each alkene is treated with CH_2I_2 and $\text{Zn}(\text{Cu})$?




Problem 26.11 What stereoisomers are formed when *trans*-3-hexene is treated with CH_2I_2 and $\text{Zn}(\text{Cu})$?

26.6 Metathesis


Recall from Section 10.1 that **olefin** is another name for an **alkene**.

The word *metathesis* is derived from the Greek words *meta* (change) and *thesis* (position). The 2005 Nobel Prize in Chemistry was awarded to Robert Grubbs of the California Institute of Technology, Yves Chauvin of the Institut Français du Pétrole, and Richard Schrock of the Massachusetts Institute of Technology for their work on olefin metathesis.

Alkene metathesis, more commonly called **olefin metathesis**, is a reaction between two alkene molecules that results in the interchange of the carbons of their double bonds. Two σ and two π bonds are broken and two new σ and two new π bonds are formed.





Olefin metathesis occurs in the presence of a complex transition metal catalyst that contains a carbon–metal double bond. The metal is typically ruthenium (Ru), tungsten (W), or molybdenum (Mo). In a widely used catalyst, called **Grubbs catalyst**, the metal is Ru.



Grubbs catalyst

Olefin metathesis is an equilibrium process and, with many alkene substrates, a mixture of starting material and two or more alkene products is present at equilibrium, making the reaction useless for preparative purposes. With terminal alkenes, however, one metathesis product is $\text{CH}_2=\text{CH}_2$ (a gas), which escapes from the reaction mixture and drives the equilibrium to the right. As a result, monosubstituted alkenes ($\text{RCH}=\text{CH}_2$) and 2,2-disubstituted alkenes ($\text{R}_2\text{C}=\text{CH}_2$) are excellent metathesis substrates because high yields of a single alkene product are obtained, as shown in Equations [1] and [2].



To draw the products of any metathesis reaction:

- [1] Arrange two molecules of the starting alkene adjacent to each other as in Figure 26.2 where styrene ($\text{PhCH}=\text{CH}_2$) is used as the starting material.
- [2] Then, break the double bonds in the starting material and form two new double bonds using carbon atoms that were *not* previously bonded to each other in the starting alkenes.

There are always two ways to arrange the starting alkenes (Pathways [1] and [2] in Figure 26.2). In this example, the two products of the reaction, $\text{PhCH}=\text{CHPh}$ and $\text{CH}_2=\text{CH}_2$ are formed in the first reaction pathway (Pathway [1]), while starting material is re-formed in the second pathway (Pathway [2]). Whenever the starting alkene is regenerated, it can go on to form product when the catalytic cycle is repeated.

Problem 26.12 Draw the products formed when each alkene is treated with Grubbs catalyst.






Figure 26.2
Drawing the products of olefin metathesis using styrene ($\text{PhCH}=\text{CH}_2$) as starting material



- Overall reaction: $2 \text{PhCH}=\text{CH}_2 \rightarrow \text{PhCH}=\text{CHPh} + \text{CH}_2=\text{CH}_2$.
- There are always two ways to join the C's of a single alkene together to form metathesis products (Pathways [1] and [2]).
- When like C's of the alkene substrate are joined in the first reaction (Pathway [1]), $\text{PhCH}=\text{CHPh}$ (in a cis and trans mixture) and $\text{CH}_2=\text{CH}_2$ are formed. Since $\text{CH}_2=\text{CH}_2$ escapes as a gas from the reaction mixture, only $\text{PhCH}=\text{CHPh}$ is isolated as product.
- When unlike C's of $\text{PhCH}=\text{CH}_2$ are joined in the second reaction (Pathway [2]), starting material is formed, which can re-enter the catalytic cycle to form product by the first pathway.
- In this way, a single constitutional isomer, $\text{PhCH}=\text{CHPh}$, is isolated with two new C–C bonds.

Problem 26.13 What products are formed when *cis*-2-pentene undergoes metathesis? Use this reaction to explain why metathesis of a 1,2-disubstituted alkene ($\text{RCH}=\text{CHR}'$) is generally not a practical method for alkene synthesis.


The mechanism for olefin metathesis is complex, and involves **metal–carbene intermediates—intermediates that contain a metal–carbon double bond**. The mechanism is drawn for the reaction of a terminal alkene ($\text{RCH}=\text{CH}_2$) with Grubbs catalyst, abbreviated as $\text{Ru}=\text{CHPh}$, to form $\text{RCH}=\text{CHR}$ and $\text{CH}_2=\text{CH}_2$. To begin metathesis, Grubbs catalyst reacts with the alkene substrate to form two new metal–carbenes **A** and **B** by a two-step process: addition of $\text{Ru}=\text{CHPh}$ to the alkene to yield two different metallocyclobutanes (Step [1]), followed by elimination to form **A** and **B** (Steps [2a] and [2b]). The alkene by-products formed in this process ($\text{RCH}=\text{CHPh}$ and $\text{PhCH}=\text{CH}_2$) are present in only a small amount since Grubbs reagent is used catalytically.



Each of these metal–carbene intermediates **A** and **B** then reacts with more starting alkene to form metathesis products, as shown in Mechanism 26.6. This mechanism is often written in a circle to emphasize the catalytic cycle. The mechanism demonstrates how two molecules of $\text{RCH}=\text{CH}_2$ are converted to $\text{RCH}=\text{CHR}$ and $\text{CH}_2=\text{CH}_2$.



Mechanism 26.6 Olefin Metathesis: $2 \text{RCH}=\text{CH}_2 \rightarrow \text{RCH}=\text{CHR} + \text{CH}_2=\text{CH}_2$




- Reaction of $\text{Ru}=\text{CH}_2$ with the alkene $\text{RCH}=\text{CH}_2$ forms a metallocyclobutane in Step [1], which eliminates $\text{CH}_2=\text{CH}_2$, a metathesis product, in Step [2]. Ru can bond to either the more or less substituted end of the alkene ($\text{RCH}=\text{CH}_2$), but product is formed only when Ru bonds to the *more* substituted C, as shown.
- Metal–carbene complex **B**, also formed in Step [2], adds to alkene $\text{RCH}=\text{CH}_2$ to form another metallocyclobutane, which decomposes to $\text{RCH}=\text{CHR}$, the second metathesis product, in Step [4]. Ru can bond to either the more or less substituted end of the alkene, but product is formed only when Ru bonds to the *less* substituted C, as shown.
- Since **A** is also formed in Step [4], the catalyst is regenerated and the cycle can begin again.
- The mechanism can be written beginning with reagent **A** or **B**, and all steps are equilibria.

When a diene is used as starting material, ring closure occurs. These reactions are typically run in very dilute solution so that the two reactive ends of the *same* molecule have a higher


A metathesis reaction that forms a ring is called **ring-closing metathesis (RCM)**.

probability of finding each other for reaction than two functional groups in *different* molecules. These high-dilution conditions thus favor **intramolecular** rather than *intermolecular* metathesis.

Ring-closing metathesis




Examples



Because metathesis catalysts are compatible with the presence of many functional groups (such as OH, OR, and C=O) and because virtually any ring size can be prepared, metathesis has been used to prepare many complex natural products (Figure 26.3).


Figure 26.3

Ring-closing metathesis in the synthesis of epothilone A and Sch38516




- **Epothilone A**, a promising anticancer agent, was first isolated from soil bacteria collected from the banks of the Zambezi River in South Africa.
- **Sch38516** is an antiviral agent active against influenza A.
- The new C–C bonds formed during metathesis are indicated in red. In both metathesis reactions, $\text{CH}_2=\text{CH}_2$ is also formed.


Problem 26.14 Draw the product formed from ring-closing metathesis of each compound.



Problem 26.15 When diene **X** is treated with Grubbs catalyst under high-dilution conditions, compound **Y** (molecular formula $C_{50}H_{90}O_3$) is isolated. When **X** is treated with Grubbs catalyst at usual reaction concentrations, three compounds **Z**, **Z'**, and **Z''** (all having molecular formula $C_{102}H_{184}O_6$) are isolated. Draw structures for **Y**, **Z**, **Z'**, and **Z''** and explain why this difference is observed.




Sample Problem 26.5 What starting material is needed to synthesize each compound by a ring-closing metathesis reaction?




Solution


To work in the retrosynthetic direction, cleave the C=C in the product, and bond each carbon of the original alkene to a CH_2 group using a double bond.



The resulting compound has a carbon chain with two terminal alkenes.



Problem 26.16 What starting material is needed to synthesize each compound by a ring-closing metathesis reaction?



KEY CONCEPTS

Carbon–Carbon Bond-Forming Reactions in Organic Synthesis


Coupling Reactions

[1] Coupling reactions of organocuprate reagents (26.1)




- R'X can be CH₃X, RCH₂X, 2° cyclic halides, vinyl halides, and aryl halides.
- X may be Cl, Br, or I.
- With vinyl halides, coupling is stereospecific.

[2] Suzuki reaction (26.2)



- R'X is most often a vinyl halide or aryl halide.
- With vinyl halides, coupling is stereospecific.


[3] Heck reaction (26.3)



- R'X is a vinyl halide or aryl halide.
- Z = H, Ph, COOR, or CN.
- With vinyl halides, coupling is stereospecific.
- The reaction forms trans alkenes.


Cyclopropane Synthesis

[1] Addition of dihalocarbenes to alkenes (26.4)



- The reaction occurs with syn addition.
- The position of substituents in the alkene is retained in the cyclopropane.

[2] Simmons–Smith reaction (26.5)



- The reaction occurs with syn addition.
- The position of substituents in the alkene is retained in the cyclopropane.


Metathesis (26.6)

[1] Intermolecular reaction



- Metathesis works best when CH₂=CH₂, a gas that escapes from the reaction mixture, is one of the products formed.

[2] Intramolecular reaction











- Ring-closing metathesis forms rings of any size from diene starting materials.





PROBLEMS

Coupling Reactions






26.17 Draw the products formed in each reaction.






26.18 What organic halide is needed to convert lithium divinylcuprate $[(CH_2=CH)_2CuLi]$ to each compound?






26.19 How can you convert ethynylcyclohexane to dienes **A–C** using a Suzuki reaction? You may use any other organic compounds and inorganic reagents. Is it possible to synthesize diene **D** using a Suzuki reaction? Explain why or why not.



26.20 What compound is needed to convert styrene ($C_6H_5CH=CH_2$) to each product using a Heck reaction?









26.21 In addition to organic halides, alkyl tosylates ($R'OTs$, Section 9.13) also react with organocuprates (R_2CuLi) to form coupling products $R-R'$. When 2° alkyl tosylates are used as starting materials (R_2CHOTs), inversion of the configuration at a stereogenic center results. Keeping this in mind, draw the product formed when each compound is treated with $(CH_3)_2CuLi$.



26.22 What steps are needed to convert 1-butene ($CH_3CH_2CH=CH_2$) to octane [$CH_3(CH_2)_6CH_3$] using a coupling reaction with an organocuprate reagent? All carbon atoms in octane must come from 1-butene.

Cyclopropanes




26.23 Draw the products (including stereoisomers) formed in each reaction.






26.24 Treatment of cyclohexene with $C_6H_5CHI_2$ and $Zn(Cu)$ forms two stereoisomers of molecular formula $C_{13}H_{16}$. Draw their structures and explain why two compounds are formed.

Metathesis




26.25 What ring-closing metathesis product is formed when each substrate is treated with Grubbs catalyst under high-dilution conditions?




26.26 What starting material is needed to prepare each compound by a ring-closing metathesis reaction?




26.27 Metathesis reactions can be carried out with two *different* alkene substrates in one reaction mixture. Depending on the substitution pattern around the $C=C$, the reaction may lead to one major product or a mixture of many products. For each pair of alkene substrates, draw all metathesis products formed. (Disregard any starting materials that may also be present at equilibrium.) With reference to the three examples, discuss when alkene metathesis with two different alkenes is a synthetically useful reaction.





26.28 Draw the structure of the two products of molecular formula $C_{15}H_{26}O_2$ formed when **M** is treated with Grubbs catalyst under high-dilution conditions.



26.29 When certain cycloalkenes are used in metathesis reactions, ring-opening metathesis polymerization (ROMP) occurs to form a high molecular weight polymer, as shown with cyclopentene as the starting material. The reaction is driven to completion by relief of strain in the cycloalkene.











What products are formed by ring-opening metathesis polymerization of each alkene?




General Reactions

26.30 Draw the products formed in each reaction.




Mechanisms


26.31 In addition to using CHX_3 and base to synthesize dihalocarbenes (Section 26.4), dichlorocarbene ($:\text{CCl}_2$) can be prepared by heating sodium trichloroacetate. Draw a stepwise mechanism for this reaction.




26.32 Draw a stepwise mechanism for the following reaction.




26.33 Sulfur ylides, like the phosphorus ylides of Chapter 21, are useful intermediates in organic synthesis, as shown in Problem 21.79. Methyl *trans*-chrysanthemate, an intermediate in the synthesis of the insecticide pyrethrin I (Section 26.4), can be prepared from diene **A** and a sulfur ylide. Draw a stepwise mechanism for this reaction.



- 26.34** Although diazomethane (CH_2N_2) is often not a useful reagent for preparing cyclopropanes, other diazo compounds give good yields of more complex cyclopropanes. Draw a stepwise mechanism for the conversion of a diazo compound **A** into **B**, an intermediate in the synthesis of sirenin, the sperm attractant produced by the female gametes of the water mold *Allomyces*.




- 26.35** The reaction of cyclohexene with iodobenzene under Heck conditions forms **E**, a coupling product with the new phenyl group on the allylic carbon, but none of the “expected” coupling product **F** with the phenyl group bonded directly to the carbon–carbon double bond.



- Draw a stepwise mechanism that illustrates how **E** is formed.
- Step [2] in Mechanism 26.2 proceeds with syn addition of Pd and R^1 to the double bond. What does the formation of **E** suggest about the stereochemistry of the elimination reaction depicted in Step [3] of Mechanism 26.2?


Synthesis

- 26.36** Devise a synthesis of diene **A** from (*Z*)-2-bromostyrene as the only organic starting material. Use a Suzuki reaction in one step of the synthesis.




- 26.37** Devise a synthesis of (*E*)-1-phenyl-1-hexene ($\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CHPh}$) using hydrocarbons having ≤ 6 C's and a Suzuki reaction as one of the steps.


- 26.38** Devise a synthesis of the given trans vinylborane, which can be used for bombykol synthesis (Figure 26.1). All of the carbon atoms in the vinylborane must come from acetylene, 1,9-nonanediol, and catecholborane.




- 26.39** Devise a synthesis of each compound using a Heck reaction as one step. You may use benzene, $\text{CH}_2=\text{CHCO}_2\text{Et}$, organic alcohols having two carbons or fewer, and any required inorganic reagents.




- 26.40** Devise a synthesis of each compound from cyclohexene and any required organic or inorganic reagents.




26.41 Devise a synthesis of each compound from benzene. You may also use any organic compounds having four carbons or fewer, and any required inorganic reagents.




26.42 Devise a synthesis of each substituted cyclopropane. Use acetylene ($\text{HC}\equiv\text{CH}$) as a starting material in parts (a) and (b), and cyclohexanone as a starting material in parts (c) and (d). You may use any other organic compounds and any needed reagents.




26.43 Biaryls, compounds containing two aromatic rings joined by a C–C bond, can often be efficiently made by two different Suzuki couplings; that is, either aromatic ring can be used to form the organoborane needed for coupling. In some cases, however, only one route is possible. With this in mind, synthesize each of the following biaryls using benzene as the starting material for each aromatic ring. When more than one route is possible, draw both of them. You may use any required organic or inorganic reagents.




26.44 Draw the product formed from the ring-closing metathesis of each compound. Then, devise a synthesis of each metathesis starting material using any of the following compounds: $\text{CH}_2(\text{CO}_2\text{Et})_2$, alcohols with less than five carbons, and any needed organic and inorganic reagents.




26.45 Draw the product formed from the ring-closing metathesis of each compound. Then, devise a synthesis of each metathesis starting material from benzene, alcohols with less than five carbons, and any needed organic and inorganic reagents.





26.46 What reagents are needed to carry out transformations [1]–[3] in the synthesis of aldehyde **A**? **A** can be converted to the antitumor agent maytansine in several steps.




- 26.47** Devise a synthesis of each of the following compounds. Besides inorganic reagents, you may use hydrocarbons and halides having ≤ 6 C's, and $\text{CH}_2=\text{CHCOOCH}_3$ as starting materials. Each synthesis must use at least one of the carbon–carbon bond-forming reactions in this chapter.





(two enantiomers)



(+ enantiomer)





(two enantiomers)




(two enantiomers)

Challenge Problems


- 26.48** Many variations of ring-closing metathesis have now been reported. For example, tandem ring-opening–ring-closing metathesis can occur with cyclic alkenes that contain two additional carbon–carbon double bonds. In this reaction, the cycloalkene is cleaved, and two new rings are formed. [1] What compounds are formed in this tandem reaction with the following substrates? [2] Devise a synthesis of the substrate in part (b) that uses a Diels–Alder reaction with diethyl maleate as the dienophile.



- 26.49** Suzuki coupling of aryl iodide **A** and vinylborane **B** affords compound **C**, which is converted to **D** in the presence of aqueous acid. Identify compounds **C** and **D** and draw a stepwise mechanism for the conversion of **C** to **D**.



- 26.50** Dimethyl cyclopropanes can be prepared by the reaction of an α,β -unsaturated carbonyl compound **X** with two equivalents of a Wittig reagent **Y**. Draw a stepwise mechanism for this reaction.



Carbohydrates

27

- 27.1 Introduction
- 27.2 Monosaccharides
- 27.3 The family of D-aldoses
- 27.4 The family of D-ketoses
- 27.5 Physical properties of monosaccharides
- 27.6 The cyclic forms of monosaccharides
- 27.7 Glycosides
- 27.8 Reactions of monosaccharides at the OH groups
- 27.9 Reactions at the carbonyl group—Oxidation and reduction
- 27.10 Reactions at the carbonyl group—Adding or removing one carbon atom
- 27.11 The Fischer proof of the structure of glucose
- 27.12 Disaccharides
- 27.13 Polysaccharides
- 27.14 Other important sugars and their derivatives



Lactose, a carbohydrate formed from two simple sugars, glucose and galactose, is the principal sugar in dairy products. Many individuals, mainly of Asian and African descent, lack adequate amounts of the enzyme lactase necessary to digest and absorb lactose. This condition, lactose intolerance, is associated with abdominal cramping and recurrent diarrhea, and is precipitated by the ingestion of milk and dairy products. Individuals who are lactose intolerant can drink lactose-free milk. Tablets that contain the lactase enzyme can also be taken when ice cream or other milk products are ingested. In Chapter 27, we learn about the structure, synthesis, and properties of carbohydrates like lactose.

Chapters 27, 28, and 29 discuss *biomolecules, organic compounds found in biological systems*. You have already learned many facts about these compounds in previous chapters while you studied other organic compounds having similar properties. In Chapter 10 (Alkenes), for example, you learned that the presence of double bonds determines whether a fatty acid is part of a fat or an oil. In Chapter 19 (Carboxylic Acids and the Acidity of the O–H Bond), you learned that amino acids are the building blocks of proteins.

Chapter 27 focuses on carbohydrates, the largest group of biomolecules in nature, comprising ~50% of the earth's biomass. Chapter 28 concentrates on proteins (and the amino acids that compose them), whereas Chapter 29 explores lipids. These compounds are all organic molecules, so many of the same principles and chemical reactions that you have already studied will be examined once again. But, as you will see, each class of compound has its own unique features that we must learn as well.

27.1 Introduction


Carbohydrates were given their name because their molecular formulas could be written as $C_n(H_2O)_n$, making them **hydrates of carbon**.

Carbohydrates such as glucose and cellulose were discussed in Sections 5.1, 6.4, and 21.17.

Although the metabolism of lipids provides more energy per gram than the metabolism of carbohydrates, glucose is the preferred source when a burst of energy is needed during exercise. Glucose is water soluble, so it can be quickly and easily transported through the bloodstream to the tissues.

Carbohydrates, commonly referred to as sugars and starches, are polyhydroxy aldehydes and ketones, or compounds that can be hydrolyzed to them. The cellulose in plant stems and tree trunks and the chitin in the exoskeletons of arthropods and mollusks are both complex carbohydrates. Four examples are shown in Figure 27.1. They include not only glucose and cellulose, but also doxorubicin (an anticancer drug) and 2'-deoxyadenosine 5'-monophosphate (a nucleotide base from DNA), both of which have a carbohydrate moiety as part of a larger molecule.


Carbohydrates are storehouses of chemical energy. They are synthesized in green plants and algae by **photosynthesis**, a process that uses the energy from the sun to convert carbon dioxide and water into glucose and oxygen. This energy is released when glucose is metabolized. The oxidation of glucose is a multistep process that forms carbon dioxide, water, and a great deal of energy (Section 6.4).



27.2 Monosaccharides


The word *saccharide* comes from the Latin word *saccharum* meaning "sugar."

The simplest carbohydrates are called **monosaccharides** or **simple sugars**. **Monosaccharides have three to seven carbon atoms** in a chain, with a **carbonyl group** at either the terminal carbon (C1) or the carbon adjacent to it (C2). In most carbohydrates, each of the remaining carbon atoms has a **hydroxy group**. Monosaccharides are usually drawn vertically, with the carbonyl group at the top.



- Monosaccharides with an aldehyde carbonyl group at C1 are called aldoses.
- Monosaccharides with a ketone carbonyl group at C2 are called ketoses.

Figure 27.1
Some examples of carbohydrates



These compounds illustrate the structural diversity of carbohydrates. **Glucose** is the most common simple sugar, whereas **cellulose**, which comprises wood, plant stems, and grass, is the most common carbohydrate in the plant world. **Doxorubicin**, an anticancer drug that has a carbohydrate ring as part of its structure, has been used in the treatment of leukemia, Hodgkin's disease, and cancers of the breast, bladder, and ovaries. **2'-Deoxyadenosine 5'-monophosphate** is one of the four nucleotides that form DNA.




D-Fructose is almost twice as sweet as normal table sugar (sucrose) with about the same number of calories per gram. "Lite" food products use only half as much fructose as sucrose for the same level of sweetness, and so they have fewer calories.



Dihydroxyacetone is the active ingredient in many artificial tanning agents.

Several examples of simple carbohydrates are shown. D-Glyceraldehyde and dihydroxyacetone have the same molecular formula, so they are **constitutional isomers**, as are D-glucose and D-fructose.



All carbohydrates have common names. The simplest aldehyde, glyceraldehyde, and the simplest ketone, dihydroxyacetone, are the only monosaccharides whose names do not end in the suffix **-ose**. (The prefix "D-" is explained in Section 27.2C.)

A monosaccharide is called:

- a **triose** if it has 3 C's;
- a **tetrose** if it has 4 C's;
- a **pentose** if it has 5 C's;
- a **hexose** if it has 6 C's, and so forth.

These terms are then combined with the words *aldose* and *ketose* to indicate both the number of carbon atoms in the monosaccharide and whether it contains an aldehyde or ketone. Thus, glyceraldehyde is an **aldotriose** (three C atoms and an aldehyde), glucose is an **aldohexose** (six C atoms and an aldehyde), and fructose is a **ketohexose** (six C atoms and a ketone).


Problem 27.1 Draw the structure of (a) a ketotetrose; (b) an aldopentose; (c) an aldotetrose.

27.2A Fischer Projection Formulas

A striking feature of carbohydrate structure is the presence of stereogenic centers. **All carbohydrates except for dihydroxyacetone contain one or more stereogenic centers.**

The simplest aldehyde, glyceraldehyde, has one stereogenic center, so there are two possible **enantiomers**. Only the enantiomer with the *R* configuration occurs naturally.


Two different representations for each enantiomer of glyceraldehyde



The stereogenic centers in sugars are often depicted following a different convention than is usually seen for other stereogenic centers. Instead of drawing a tetrahedron with two bonds in the plane, one in front of the plane, and one behind it, the **tetrahedron is tipped so that horizontal bonds come forward (drawn on wedges) and vertical bonds go behind (on dashed lines)**. This structure is then abbreviated by a **cross formula**, also called a **Fischer projection formula**. In a Fischer projection formula:

- A carbon atom is located at the intersection of the two lines of the cross.
- The horizontal bonds come forward, on wedges.
- The vertical bonds go back, on dashed lines.
- In a carbohydrate, the aldehyde or ketone carbonyl is put at or near the top.


Using a Fischer projection formula, (*R*)-glyceraldehyde becomes:



Do not rotate a Fischer projection formula in the plane of the page, because you might inadvertently convert a compound into its enantiomer. When using Fischer projections it is usually best to convert them to structures with wedges and dashes, and then manipulate them. Although a Fischer projection formula can be used for the stereogenic center in any compound, it is most commonly used for monosaccharides.


Sample Problem 27.1

Convert each compound to a Fischer projection formula.




Solution

Rotate and re-draw each molecule to place the horizontal bonds in front of the plane and the vertical bonds behind the plane. Then use a cross to represent the stereogenic center.



Problem 27.2 Draw each stereogenic center using a Fischer projection formula.




R,S designations can be assigned to any stereogenic center drawn as a Fischer projection formula in the following manner:

- [1] Assign priorities (1 → 4) to the four groups bonded to the stereogenic center using the rules detailed in Section 5.6.
- [2] When the lowest priority group occupies a vertical bond—that is, it projects *behind* the plane on a dashed line—tracing a circle in the clockwise direction (from priority group 1 → 2 → 3) gives the *R* configuration. Tracing a circle in the counterclockwise direction gives the *S* configuration.
- [3] When the lowest priority group occupies a horizontal bond—that is, it projects *in front* of the plane on a wedge—reverse the answer obtained in Step [2] to designate the configuration.


Sample Problem 27.2


Re-draw each Fischer projection formula using wedges and dashes for the stereogenic center, and label the center as *R* or *S*.

**Solution**





For each molecule:

- [1] Convert the Fischer projection formula to a representation with wedges and dashes.
- [2] Assign priorities (Section 5.6).
- [3] Determine *R* or *S* in the usual manner. Reverse the answer if priority group [4] is oriented forward (on a wedge).






Problem 27.3 Label each stereogenic center as *R* or *S*.



27.2B Monosaccharides with More Than One Stereogenic Center


The number of possible stereoisomers of a monosaccharide increases exponentially with the number of stereogenic centers present. **An aldohexose has four stereogenic centers, and so it has $2^4 = 16$ possible stereoisomers**, or eight pairs of enantiomers.

General structure of an aldohexose



[* denotes a stereogenic center.]

Fischer projection formulas are also used for compounds like aldohexoses that contain several stereogenic centers. In this case, the molecule is drawn with a vertical carbon skeleton and the stereogenic centers are stacked one above another. Using this convention, **all horizontal bonds project forward (on wedges)**.




Although Fischer projections are commonly used to depict monosaccharides with many stereogenic centers, care must be exercised in using them since they do not give a true picture of the three-dimensional structures they represent. Because each stereogenic center is drawn in the less stable eclipsed conformation, the Fischer projection of glucose really represents the molecule in a cylindrical conformation, as shown in Figure 27.2.

Problem 27.4 Assign *R,S* designations to each stereogenic center in glucose.

Figure 27.2


A Fischer projection and the 3-D structure of glucose



27.2C D and L Monosaccharides

Although the prefixes *R* and *S* can be used to designate the configuration of stereogenic centers in monosaccharides, an older system of nomenclature uses the prefixes *D*- and *L*-, instead. Naturally occurring glyceraldehyde with the *R* configuration is called the **D-isomer**. Its enantiomer, (*S*)-glyceraldehyde, is called the **L-isomer**.


Fischer projections for the enantiomers of glyceraldehyde




The letters **D** and **L** are used to label all monosaccharides, even those with multiple stereogenic centers. **The configuration of the stereogenic center farthest from the carbonyl group determines whether a monosaccharide is D- or L-**.

The two designations, *D* and *d*, refer to very different phenomena. The “*D*” designates the configuration around a stereogenic center. In a *D* monosaccharide, the OH group on the stereogenic center farthest from the carbonyl group is on the right in a Fischer projection. The “*d*,” on the other hand, is an abbreviation for “dextrorotatory;” that is, a *d*-compound rotates the plane of polarized light in the clockwise direction. A *D*-sugar may be dextrorotatory or it may be levorotatory. There is no direct correlation between *D* and *d* or *L* and *l*.

- A *D*-sugar has the OH group on the stereogenic center farthest from the carbonyl on the right in a Fischer projection (like *D*-glyceraldehyde).
- An *L*-sugar has the OH group on the stereogenic center farthest from the carbonyl on the left in a Fischer projection (like *L*-glyceraldehyde).




Glucose and all other naturally occurring sugars are D-sugars. L-Glucose, a compound that does not occur in nature, is the enantiomer of *D*-glucose. L-Glucose has the opposite configuration at *every* stereogenic center.



Problem 27.5 How many stereogenic centers are present in each type of monosaccharide: (a) an aldotetrose; (b) a ketohexose?


Problem 27.6 (a) Label compounds **A**, **B**, and **C** as D- or L-sugars. (b) How are compounds **A** and **B** related? **A** and **C**? **B** and **C**? Choose from enantiomers, diastereomers, or constitutional isomers.



27.3 The Family of D-Aldoses

Beginning with D-glyceraldehyde, one may formulate other D-aldoses having four, five, or six carbon atoms by adding carbon atoms (each bonded to H and OH), one at a time, between C1 and C2. Two D-aldotetroses can be formed from D-glyceraldehyde, one with the new OH group on the right and one with the new OH group on the left. Their names are D-erythrose and D-threose. They are two diastereomers, each with two stereogenic centers.

The common name of each monosaccharide indicates both the number of atoms it contains and the configuration at each of the stereogenic centers. Because the common names are firmly entrenched in the chemical literature, no systematic method has ever been established to name these compounds.



Because each aldotetrose has two stereogenic centers, there are 2^2 or four possible stereoisomers. D-Erythrose and D-threose are two of them. The other two are their enantiomers, called L-erythrose and L-threose, respectively. The configuration around each stereogenic center is exactly the opposite in its enantiomer. All four stereoisomers of the D-aldotetroses are shown in Figure 27.3.

To continue forming the family of D-aldoses, we must add another carbon atom (bonded to H and OH) just below the carbonyl of either tetrose. Because there are two D-aldotetroses to begin with, and there are two ways to place the new OH (right or left), there are now four D-aldopentoses: D-ribose, D-arabinose, D-xylose, and D-lyxose. Each aldopentose now has three stereogenic centers, so there are $2^3 = 8$ possible stereoisomers, or four pairs of enantiomers. The D-enantiomer of each pair is shown in Figure 27.4.

Finally, to form the D-aldohexoses, we must add another carbon atom (bonded to H and OH) just below the carbonyl of all the aldopentoses. Because there are four D-aldopentoses to begin with, and there are two ways to place the new OH (right or left), there are now eight D-aldohexoses. Each aldohexose now has four stereogenic centers, so there are $2^4 = 16$ possible stereoisomers, or eight pairs of enantiomers. Only the D-enantiomer of each pair is shown in Figure 27.4.

The tree of D-aldoses (Figure 27.4) is arranged in pairs of compounds that are bracketed together. Each pair of compounds, such as D-glucose and D-mannose, has the same configuration around all of its stereogenic centers except for one.

D-Ribose, D-arabinose, and D-xylose are all common aldopentoses in nature. D-Ribose is the carbohydrate component of RNA, the polymer that translates the genetic information of DNA for protein synthesis.

Figure 27.3

The four stereoisomeric aldotetroses






Figure 27.4

The family of D-aldoses having three to six carbon atoms



Of the D-aldohexoses, only D-glucose and D-galactose are common in nature. **D-Glucose is by far the most abundant of all D-aldoses.** D-Glucose comes from the hydrolysis of starch and cellulose, and D-galactose comes from the hydrolysis of fruit pectins.

- Two diastereomers that differ in the configuration around one stereogenic center only are called *epimers*.



Problem 27.7 How is it possible that D-glucose is dextrorotatory but D-fructose is levorotatory?

Problem 27.8 How many different aldoheptoses are there? How many are D-sugars? Draw all D-aldohexoses having the *R* configuration at C2 and C3.


Problem 27.9 Draw two possible epimers of D-erythrose. Name each of these compounds using Figure 27.4.

27.4 The Family of D-Ketoses

The family of D-ketoses, shown in Figure 27.5, is formed from dihydroxyacetone by adding a new carbon (bonded to H and OH) between C2 and C3. Having a carbonyl group at C2 decreases the number of stereogenic centers in these monosaccharides, so that there are only four D-ketohexoses. The most common naturally occurring ketose is D-fructose.

Figure 27.5

The family of D-ketoses having three to six carbon atoms



- Problem 27.10** Referring to the structures in Figures 27.4 and 27.5, classify each pair of compounds as enantiomers, epimers, diastereomers but not epimers, or constitutional isomers of each other.
- D-allose and L-allose
 - D-altrose and D-gulose
 - D-galactose and D-talose
 - D-mannose and D-fructose
 - D-fructose and D-sorbose
 - L-sorbose and L-tagatose

- Problem 27.11**
- Draw the enantiomer of D-fructose.
 - Draw an epimer of D-fructose at C4. What is the name of this compound?
 - Draw an epimer of D-fructose at C5. What is the name of this compound?

- Problem 27.12** Referring to Figure 27.5, which D-ketohexoses have the S configuration at C3?


27.5 Physical Properties of Monosaccharides

Monosaccharides have the following physical properties:

- They are all **sweet tasting**, but their relative sweetness varies a great deal.
- They are polar compounds with **high melting points**.
- The presence of so many polar functional groups capable of hydrogen bonding makes them **water soluble**.
- Unlike most other organic compounds, monosaccharides are so polar that they are **insoluble in organic solvents like diethyl ether**.

27.6 The Cyclic Forms of Monosaccharides


Although the monosaccharides in Figures 27.4 and 27.5 are drawn as acyclic carbonyl compounds containing several hydroxy groups, the hydroxy and carbonyl groups of monosaccharides can undergo intramolecular cyclization reactions to form **hemiacetals** having either five or six atoms in the ring. This process was first discussed in Section 21.16.



- A six-membered ring containing an O atom is called a *pyranose ring*.
- A five-membered ring containing an O atom is called a *furanose ring*.

Cyclization of a hydroxy carbonyl compound always forms a stereogenic center at the hemiacetal carbon, called the **anomeric carbon**. The two hemiacetals are called **anomers**.


- Anomers are stereoisomers of a cyclic monosaccharide that differ in the position of the OH group at the hemiacetal carbon.



Cyclization forms the more stable ring size in a given molecule. **The most common monosaccharides, the aldohexoses like glucose, typically form a pyranose ring**, so our discussion begins with forming a cyclic hemiacetal from D-glucose.


27.6A Drawing Glucose as a Cyclic Hemiacetal

Which of the five OH groups in glucose is at the right distance from the carbonyl group to form a six-membered ring? The **O atom on the stereogenic center farthest from the carbonyl (C5)** is six atoms from the carbonyl carbon, placing it in the proper position for cyclization to form a pyranose ring.




To translate the acyclic form of glucose into a cyclic hemiacetal, we must draw the hydroxy aldehyde in a way that suggests the position of the atoms in the new ring, and then draw the ring. **By convention the O atom in the new pyranose ring is drawn in the upper right-hand corner of the six-membered ring.**

Rotating the groups on the bottom stereogenic center in **A** places all six atoms needed for the ring (including the OH) in a vertical line (**B**). Re-drawing this representation as a Fischer projection makes the structure appear less cluttered (**C**). Twisting this structure and rotating it 90° forms **D**. Structures **A–D** are four different ways of drawing the same acyclic structure of D-glucose.



We are now set to draw the cyclic hemiacetal formed by nucleophilic attack of the OH group on C5 on the aldehyde carbonyl. Because cyclization creates a new stereogenic center, there are **two cyclic forms of D-glucose**, an **α anomer** and a **β anomer**. All the original stereogenic centers maintain their configuration in both of the products formed.

- The **α anomer** of a D monosaccharide has the OH group drawn **down**, **trans** to the CH₂OH group at C5. The **α anomer** of D-glucose is called **α -D-glucose**, or **α -D-glucopyranose** (to emphasize the six-membered ring).
- The **β anomer** of a D monosaccharide has the OH group drawn **up**, **cis** to the CH₂OH group at C5. The **β anomer** is called **β -D-glucose**, or **β -D-glucopyranose** (to emphasize the six-membered ring).




The **α anomer** in any monosaccharide has the anomeric OH group and the CH₂OH group **trans**. The **β anomer** has the anomeric OH group and the CH₂OH group **cis**.

These flat, six-membered rings used to represent the cyclic hemiacetals of glucose and other sugars are called **Haworth projections**. The cyclic forms of glucose now have **five stereogenic centers**, **the four from the starting hydroxy aldehyde and the new anomeric carbon**. **α -D-Glucose** and **β -D-glucose** are **diastereomers**, because only the anomeric carbon has a different configuration.

The mechanism for this transformation is exactly the same as the mechanism that converts a hydroxy aldehyde to a cyclic hemiacetal (Mechanism 21.11). The acyclic aldehyde and two cyclic hemiac-

Figure 27.6

The three forms of glucose




etals are all in equilibrium. Each cyclic hemiacetal can be isolated and crystallized separately, but when any one compound is placed in solution, an equilibrium mixture of all three forms results. This process is called **mutarotation**. At equilibrium, the mixture has 37% of the α anomer, 63% of the β anomer, and only trace amounts of the acyclic hydroxy aldehyde, as shown in Figure 27.6.

27.6B Haworth Projections

To convert an acyclic monosaccharide to a Haworth projection, follow a stepwise procedure.


HOW TO Draw a Haworth Projection from an Acyclic Aldohexose

Example Convert D-mannose to a Haworth projection.




Step [1] Place the O atom in the upper right corner of a hexagon, and add the CH_2OH group on the first carbon counterclockwise from the O atom.

- For **D-sugars**, the CH_2OH group is drawn **up**. For **L-sugars**, the CH_2OH group is drawn **down**.



Step [2] Place the anomeric carbon on the first carbon clockwise from the O atom.


- For an α **anomer**, the **OH** is drawn **down** in a D-sugar.
- For a β **anomer**, the **OH** is drawn **up** in a D-sugar.



- Remember: The carbonyl carbon becomes the anomeric carbon (a new stereogenic center).




Step [3] Add the substituents at the three remaining stereogenic centers clockwise around the ring.

- The substituents on the **right side** of the Fischer projection are drawn **down**.
- The substituents on the **left** are drawn **up**.




Problem 27.13 Convert each aldohexose to the indicated anomer using a Haworth projection.

- a. Draw the α anomer of: b. Draw the α anomer of: c. Draw the β anomer of:



Sample Problem 27.3 shows how to convert a Haworth projection back to the acyclic form of a monosaccharide. It doesn't matter whether the hemiacetal is the α or β anomer, because both anomers give the *same* hydroxy aldehyde.


Sample Problem 27.3 Convert the following Haworth projection to the acyclic form of the aldohexose.




Solution

To convert the substituents to the acyclic form, start at the pyranose O atom, and work in a **counterclockwise** fashion around the ring, and from bottom-to-top along the chain.

- [1] Draw the carbon skeleton, placing the CHO on the top and the CH₂OH on the bottom.




Proceed in a **counterclockwise** fashion around the ring.



- [2] Classify the sugar as D- or L-.


- The CH₂OH is drawn **up**, so it is a **D-sugar**.
- A D-sugar has the OH group on the bottom stereogenic center on the **right**.




- [3] Add the three other stereogenic centers.

- **Up** substituents go on the **left**.
- **Down** substituents go on the **right**.

The anomeric C becomes the C=O.



Problem 27.14 Convert each Haworth projection to its acyclic form.




27.6C Three-Dimensional Representations for D-Glucose

Because the chair form of a six-membered ring gives the truest picture of its three-dimensional shape, we must learn to convert Haworth projections into chair forms.

To convert a Haworth projection to a chair form:

- Draw the pyranose ring with the O atom as an “up” atom.
- The “up” substituents in a Haworth projection become the “up” bonds (either axial or equatorial) on a given carbon atom on a puckered six-membered ring.
- The “down” substituents in a Haworth projection become the “down” bonds (either axial or equatorial) on a given carbon atom on a puckered six-membered ring.

As a result, the three-dimensional chair form of β -D-glucose is drawn in the following manner:




Glucose has all substituents larger than a hydrogen atom in the more roomy equatorial positions, making it the most stable and thus most prevalent monosaccharide. The β anomer is the major isomer at equilibrium, moreover, because the hemiacetal OH group is in the equatorial position, too. Figure 27.7 shows both anomers of D-glucose drawn as chair conformations.

Problem 27.15 Convert each Haworth projection in Problem 27.14 to a three-dimensional representation using a chair pyranose ring.

27.6D Furanoses

Certain monosaccharides—notably aldopentoses and ketohexoses—form furanose rings, not pyranose rings, in solution. The same principles apply to drawing these structures as for drawing pyranose rings, except the ring size is one atom smaller.

Figure 27.7 Three-dimensional representations for both anomers of D-glucose







Honey was the first and most popular sweetening agent until it was replaced by sugar (from sugarcane) in modern times. Honey is a mixture consisting largely of D-fructose and D-glucose.

- Cyclization always forms a new stereogenic center at the anomeric carbon, so two different anomers are possible. For a D-sugar, the OH group is drawn down in the α anomer and up in the β anomer.
- Use the same drawing conventions for adding substituents to the five-membered ring. With D-sugars, the CH_2OH group is drawn up.

With D-ribose, the OH group used to form the five-membered furanose ring is located on C4. Cyclization yields two anomers at the new stereogenic center, which are called α -D-ribofuranose and β -D-ribofuranose.



The same procedure can be used to draw the furanose form of D-fructose, the most common ketohexose. Because the carbonyl group is at C2 (instead of C1, as in the aldoses), the OH group at C5 reacts to form the hemiacetal in the five-membered ring. Two anomers are formed.



Problem 27.16 Aldotetroses exist in the furanose form. Draw both anomers of D-erythrose.

27.7 Glycosides

Keep in mind the difference between a hemiacetal and an acetal:



hemiacetal

acetal

- one OH group
- one OR group
- two OR groups

Because monosaccharides exist in solution in an equilibrium between acyclic and cyclic forms, they undergo three types of reactions:

- **Reaction of the hemiacetal**
- **Reaction of the hydroxy groups**
- **Reaction of the carbonyl group**


Even though the acyclic form of a monosaccharide may be present in only trace amounts, the equilibrium can be tipped in its favor by Le Châtelier's principle (Section 9.8). Suppose, for example, that the carbonyl group of the acyclic form reacts with a reagent, thus depleting its

equilibrium concentration. The equilibrium will then shift to compensate for the loss, thus producing more of the acyclic form, which can react further.

Note, too, that monosaccharides have two different types of OH groups. Most are “regular” alcohols, and as such, undergo reactions characteristic of alcohols. The anomeric OH group, on the other hand, is part of a hemiacetal, giving it added reactivity.

27.7A Glycoside Formation

Treatment of a monosaccharide with an alcohol and HCl converts the hemiacetal into an acetal called a **glycoside**. For example, treatment of α -D-glucose with CH_3OH and HCl forms two glycosides that are diastereomers at the acetal carbon. The α and β labels are assigned in the same way as anomers: with a D-sugar, an α glycoside has the new OR group (OCH_3 group in this example) down, and a β glycoside has the new OR group up.




Mechanism 27.1 explains why a single anomer forms two glycosides. The reaction proceeds by way of a **planar carbocation**, which undergoes nucleophilic attack from two different directions to give a mixture of diastereomers. Because both α - and β -D-glucose form the same planar carbocation, each yields the same mixture of two glycosides.




Mechanism 27.1 Glycoside Formation

Steps [1]–[2] Protonation and loss of the leaving group



- Protonation of the hemiacetal OH group, followed by **loss of H_2O** , forms a resonance-stabilized cation (Steps [1] and [2]).

Steps [3]–[4] Nucleophilic attack and deprotonation



- **Nucleophilic attack of CH_3OH** on the planar carbocation occurs from both sides to yield α and β glycosides after loss of a proton (Steps [3] and [4]).

The mechanism also explains why only the hemiacetal OH group reacts. Protonation of the hemiacetal OH, followed by loss of H_2O , forms a resonance-stabilized carbocation in Step [2]. A resonance-stabilized carbocation is not formed by loss of H_2O from any other OH group.


Unlike cyclic hemiacetals, **glycosides are acetals, and so they do not undergo mutarotation.** When a single glycoside is dissolved in H_2O , it is *not* converted to an equilibrium mixture of α and β glycosides.

- Glycosides are acetals with an alkoxy group (OR) bonded to the anomeric carbon.

Problem 27.17 What glycosides are formed when each monosaccharide is treated with $\text{CH}_3\text{CH}_2\text{OH}$, HCl :
(a) β -D-mannose; (b) α -D-gulose; (c) β -D-fructose?

27.7B Glycoside Hydrolysis

Because glycosides are acetals, **they are hydrolyzed with acid and water to cyclic hemiacetals and a molecule of alcohol.** A mixture of two anomers is formed from a single glycoside. For example, treatment of methyl α -D-glucopyranoside with aqueous acid forms a mixture of α - and β -D-glucose and methanol.




The mechanism for glycoside hydrolysis is just the reverse of glycoside formation. It involves two parts: **formation of a planar carbocation**, followed by **nucleophilic attack of H_2O** to form anomeric hemiacetals, as shown in Mechanism 27.2.




Mechanism 27.2 Glycoside Hydrolysis

Steps [1]–[2] Protonation and loss of the leaving group




- Protonation of the acetal OCH_3 group, followed by **loss of CH_3OH** , forms a resonance-stabilized cation (Steps [1] and [2]).

Steps [3]–[4] Nucleophilic attack and deprotonation



- **Nucleophilic attack of H_2O** on the planar carbocation occurs from both sides to yield α and β anomers after loss of a proton (Steps [3] and [4]).

Problem 27.18 Draw a stepwise mechanism for the following reaction.




27.7C Naturally Occurring Glycosides



The berries of the black nightshade plant (*Solanum nigrum*) are a source of the poisonous alkaloid solanine.

Salicin and **solanine** are two naturally occurring compounds that contain glycoside bonds as part of their structure. Salicin is an analgesic isolated from willow bark, and solanine is a poisonous compound isolated from the berries of the deadly nightshade plant. Solanine is also produced in the leaves, stem, and green spots on the skin of potatoes as a defense against insects and predators. It is believed that the role of the sugar rings in both salicin and solanine is to increase their water solubility.




Glycosides are common in nature. All disaccharides and polysaccharides are formed by joining monosaccharides together with glycosidic linkages. These compounds are discussed in detail beginning in Section 27.12.

Problem 27.19

(a) Label all the O atoms that are part of a glycoside in rebaudioside A. Rebaudioside A, marketed under the trade name Truvia, is a sweet glycoside obtained from the stevia plant, which has been used for centuries in Paraguay to sweeten foods. (b) The alcohol or phenol formed from the hydrolysis of a glycoside is called an **aglycon**. What aglycon and monosaccharides are formed by the hydrolysis of rebaudioside A?




Rebaudioside A, a naturally occurring glycoside about 400 times sweeter than table sugar, is obtained from the leaves of the stevia plant, a shrub native to Central and South America.



27.8 Reactions of Monosaccharides at the OH Groups

Because monosaccharides contain OH groups, they undergo reactions typical of alcohols—that is, they are converted to **ethers** and **esters**. Because the cyclic hemiacetal form of a monosaccharide contains an OH group, this form of a monosaccharide must be drawn as the starting material for any reaction that occurs at an OH group.


All OH groups of a cyclic monosaccharide are converted to ethers by treatment with base and an alkyl halide. For example, α -D-glucose reacts with silver(I) oxide (Ag_2O , a base) and excess CH_3I to form pentamethyl ether.



Ag_2O removes a proton from each alcohol, forming an alkoxide (RO^-), which then reacts with CH_3I in an $\text{S}_{\text{N}}2$ reaction. Because no C–O bonds are broken, the configuration of all substituents in the starting material is **retained**, forming a single product.


The product contains two different types of ether bonds. There are four “regular” ethers formed from the “regular” hydroxyls. The new ether from the hemiacetal is now part of an **acetal**—that is, a **glycoside**.

The four ether bonds that are *not* part of the acetal do not react with any reagents except strong acids like HBr and HI (Section 9.14). **The acetal ether, on the other hand, is hydrolyzed with aqueous acid** (Section 27.7B). Aqueous hydrolysis of a single glycoside (like the pentamethyl ether of α -D-glucose) yields both anomers of the product monosaccharide.



- “Regular” ethers are shown in blue.
- The acetal ether is shown in red.

The OH groups of monosaccharides can also be converted to esters. For example, treatment of β -D-glucose with either acetic anhydride or acetyl chloride in the presence of pyridine (a base) converts all OH groups into acetate esters.



Monosaccharides are so polar that they are insoluble in common organic solvents, making them difficult to isolate and use in organic reactions. Monosaccharide derivatives that have five ether or ester groups in place of the OH groups, however, are readily soluble in organic solvents.

Problem 27.20 Draw the products formed when β -D-galactose is treated with each reagent.

- $\text{Ag}_2\text{O} + \text{CH}_3\text{I}$
- $\text{NaH} + \text{C}_6\text{H}_5\text{CH}_2\text{Cl}$
- The product in (b), then H_3O^+
- $\text{Ac}_2\text{O} + \text{pyridine}$
- $\text{C}_6\text{H}_5\text{COCl} + \text{pyridine}$
- The product in (c), then $\text{C}_6\text{H}_5\text{COCl} + \text{pyridine}$


27.9 Reactions at the Carbonyl Group—Oxidation and Reduction

Oxidation and reduction reactions occur at the carbonyl group of monosaccharides, so they all begin with the monosaccharide drawn in the acyclic form. We will confine our discussion to aldoses as starting materials.

27.9A Reduction of the Carbonyl Group

Like other aldehydes, the **carbonyl group of an aldose is reduced to a 1° alcohol using NaBH_4** . This alcohol is called an **alditol**. For example, reduction of D-glucose with NaBH_4 in CH_3OH yields glucitol (also called sorbitol).


Glucitol occurs naturally in some fruits and berries. It is sometimes used as a substitute for sucrose (table sugar). With six polar OH groups capable of hydrogen bonding, glucitol is readily hydrated. It is used as an additive to prevent certain foods from drying out.



Problem 27.21 A 2-ketohexose is reduced with NaBH_4 in CH_3OH to form a mixture of D-galactitol and D-talitol. What is the structure of the 2-ketohexose?

27.9B Oxidation of Aldoses


Aldoses contain 1° and 2° alcohols and an aldehyde, all of which are oxidizable functional groups. Two different types of oxidation reactions are particularly useful—oxidation of the aldehyde to a carboxylic acid (an **aldonic acid**) and oxidation of both the aldehyde and the 1° alcohol to a diacid (an **aldaric acid**).



[1] Oxidation of the aldehyde to a carboxylic acid


The aldehyde carbonyl is the most easily oxidized functional group in an aldose, and so a variety of reagents oxidize it to a carboxy group, forming an **aldonic acid**.

Figure 27.8
Examples of reducing and nonreducing sugars



- Carbohydrates containing a hemiacetal are in equilibrium with an acyclic aldehyde, making them reducing sugars.
- Glycosides are acetals, so they are *not* in equilibrium with any acyclic aldehyde, making them nonreducing sugars.


Three reagents used for this process produce a characteristic color change because the oxidizing agent is reduced to a colored product that is easily visible. As described in Section 20.8, **Tollens reagent** oxidizes aldehydes to carboxylic acids using Ag_2O in NH_4OH , and forms a mirror of Ag as a by-product. **Benedict's** and **Fehling's reagents** use a blue Cu^{2+} salt as an oxidizing agent, which is reduced to Cu_2O , a brick-red solid. Unfortunately, none of these reagents gives a high yield of aldonic acid. When the aldonic acid is needed to carry on to other reactions, $\text{Br}_2 + \text{H}_2\text{O}$ is used as the oxidizing agent.



- Any carbohydrate that exists as a *hemiacetal* is in equilibrium with a small amount of acyclic aldehyde, so it is oxidized to an aldonic acid.
- Glycosides are acetals, not hemiacetals, so they are *not* oxidized to aldonic acids.


Carbohydrates that can be oxidized with Tollens, Benedict's, or Fehling's reagent are called **reducing sugars**. Those that do not react with these reagents are called **nonreducing sugars**. Figure 27.8 shows examples of reducing and nonreducing sugars.

Problem 27.22 Classify each compound as a reducing or nonreducing sugar.




[2] Oxidation of both the aldehyde and 1° alcohol to a diacid

Both the aldehyde and 1° alcohol of an aldose are oxidized to carboxy groups by treatment with warm nitric acid, forming an **aldaric acid**. Under these conditions, D-glucose is converted to D-glucaric acid.



Because aldaric acids have identical functional groups on both terminal carbons, some aldaric acids contain a plane of symmetry, making them achiral molecules. For example, oxidation of D-allose forms an achiral, optically inactive aldaric acid. This contrasts with D-glucaric acid formed from glucose, which has no plane of symmetry, and is thus still optically active.




Problem 27.23 Draw the products formed when D-arabinose is treated with each reagent: (a) Ag_2O , NH_4OH ; (b) Br_2 , H_2O ; (c) HNO_3 , H_2O .

Problem 27.24 Which aldoses are oxidized to optically inactive aldaric acids: (a) D-erythrose; (b) D-lyxose; (c) D-galactose?

27.10 Reactions at the Carbonyl Group—Adding or Removing One Carbon Atom


Two common procedures in carbohydrate chemistry result in adding or removing one carbon atom from the skeleton of an aldose. The **Wohl degradation** shortens an aldose chain by one carbon, whereas the **Kiliani–Fischer synthesis** lengthens it by one. Both reactions involve cyanohydrins as intermediates. Recall from Section 21.9 that cyanohydrins are formed from aldehydes by addition of the elements of HCN. Cyanohydrins can also be re-converted to carbonyl compounds by treatment with base.




- Forming a cyanohydrin adds one carbon to a carbonyl group.
- Re-converting a cyanohydrin to a carbonyl compound removes one carbon.

27.10A The Wohl Degradation

The Wohl degradation is a stepwise procedure that shortens the length of an aldose chain by cleavage of the C1–C2 bond. As a result, an aldohexose is converted to an aldopentose having the same configuration at its bottom three stereogenic centers (C3–C5). For example, the Wohl degradation converts D-glucose into D-arabinose.




The Wohl degradation consists of three steps, illustrated here beginning with D-glucose.



- [1] Treatment of D-glucose with hydroxylamine (NH_2OH) forms an **oxime** by nucleophilic addition. This reaction is analogous to the formation of imines discussed in Section 21.11.
- [2] Dehydration of the oxime to a nitrile occurs with acetic anhydride and sodium acetate. The nitrile product is a cyanohydrin.
- [3] **Treatment of the cyanohydrin with base results in loss of the elements of HCN to form an aldehyde having one fewer carbon.**


The Wohl degradation converts a stereogenic center at C2 in the original aldose to an sp^2 hybridized C=O. As a result, a pair of aldoses that are epimeric at C2, such as D-galactose and D-talose, yield the same aldose (D-lyxose, in this case) upon Wohl degradation.




Problem 27.25 What two aldoses yield D-xylose on Wohl degradation?

27.10B The Kiliani–Fischer Synthesis

The Kiliani–Fischer synthesis lengthens a carbohydrate chain by adding one carbon to the aldehyde end of an aldose, thus forming a new stereogenic center at C2 of the product. The product consists of epimers that differ only in their configuration about the one new stereogenic center. For example, the Kiliani–Fischer synthesis converts D-arabinose into a mixture of D-glucose and D-mannose.



The Kiliani–Fischer synthesis, shown here beginning with D-arabinose, consists of three steps. “Squiggly” lines are meant to indicate that two different stereoisomers are formed at the new stereogenic center. As with the Wohl degradation, the key intermediate is a cyanohydrin.



- [1] Treating an aldose with NaCN and HCl adds the elements of HCN to the carbonyl group, forming a **cyanohydrin** and a new carbon–carbon bond. Because the sp^2 hybridized carbonyl carbon is converted to an sp^3 hybridized carbon with four different groups, a new stereogenic center is formed in this step.
- [2] Reduction of the nitrile with H_2 and Pd-BaSO₄, a poisoned Pd catalyst, forms an **imine**.
- [3] Hydrolysis of the imine with aqueous acid forms an aldehyde that has one more carbon than the aldose that began the sequence.

Note that the **Wohl degradation** and the **Kiliani–Fischer synthesis** are conceptually opposite transformations.

- The **Wohl degradation** *removes* a carbon atom from the aldehyde end of an aldose. Two aldoses that are epimers at C2 form the same product.
- The **Kiliani–Fischer synthesis** *adds* a carbon to the aldehyde end of an aldose, forming two epimers at C2.

Problem 27.26

What aldoses are formed when the following aldoses are subjected to the Kiliani–Fischer synthesis: (a) D-threose; (b) D-ribose; (c) D-galactose?

27.10C Determining the Structure of an Unknown Monosaccharide

The reactions in Sections 27.9–27.10 can be used to determine the structure of an unknown monosaccharide, as shown in Sample Problem 27.4.

Sample Problem 27.4


A D-aldopentose **A** is oxidized to an optically inactive aldaric acid with HNO_3 . **A** is formed by the Kiliani–Fischer synthesis of a D-aldotetrose **B**, which is also oxidized to an optically inactive aldaric acid with HNO_3 . What are the structures of **A** and **B**?

Solution

Use each fact to determine the relative orientation of the OH groups in the D-aldopentose.


Fact [1] A D-aldopentose **A** is oxidized to an optically inactive aldaric acid with HNO_3 .

An optically inactive aldaric acid must contain a **plane of symmetry**. There are only two ways to arrange the OH groups in a five-carbon D-aldaric acid, for this to be the case. Thus, only two structures are possible for **A**, labeled **A'** and **A''**.




Fact [2] **A** is formed by the Kiliani–Fischer synthesis from a D-aldotetrose **B**.

A' and **A''** are each prepared from a D-aldotetrose (**B'** and **B''**) that has the same configuration at the bottom two stereogenic centers.



Fact [3] The D-aldotetrose is oxidized to an optically inactive aldaric acid upon treatment with HNO_3 .

Only the aldaric acid from **B'** has a plane of symmetry, making it optically inactive. Thus, **B'** is the correct structure for the D-aldotetrose **B**, and therefore **A'** is the structure of the D-aldopentose **A**.



Problem 27.27 D-Aldopentose **A** is oxidized to an optically inactive aldaric acid. On Wohl degradation, **A** forms an aldotetrose **B** that is oxidized to an optically active aldaric acid. What are the structures of **A** and **B**?

27.11 The Fischer Proof of the Structure of Glucose

The Fischer proof is remarkable because it was done at a time when determining melting points and optical rotations were the most sophisticated techniques available to the chemist.

In 1951, the technique of X-ray crystallography confirmed that (+)-glucose had the **D** configuration, as assumed by Fischer more than 50 years earlier.

Both Fischer projections and the Kiliani–Fischer synthesis are named after **Emil Fischer**, a noted chemist of the late nineteenth and early twentieth centuries, who received the Nobel Prize in Chemistry in 1902 for his work in carbohydrate chemistry. Fischer’s most elegant work is the subject of Section 27.11.

In 1891, only 10 years after the tetrahedral structure of carbon was proposed, Fischer determined the *relative* configuration of the four stereogenic centers in naturally occurring (+)-glucose. This body of work is called the **Fischer proof** of the structure of glucose.

Because glucose has four stereogenic centers, there are $2^4 = 16$ possible stereoisomers, or eight pairs of enantiomers. In 1891, there was no way to determine the **absolute configuration** of (+)-glucose—that is, the *exact* three-dimensional arrangement of the four stereogenic centers. Because there was no way to distinguish between enantiomers, Fischer could only determine the **relative arrangement of the OH groups to each other**.

Because of this, Fischer began with an assumption. He assumed that naturally occurring glucose had the **D configuration**—namely, that the OH group on the stereogenic center farthest from the aldehyde was oriented on the *right* in a Fischer projection. He then set out to determine the orientation of all other OH groups relative to it. Thus, the Fischer proof determined which of the eight D-aldohexoses was (+)-glucose.

The strategy used by Fischer is similar to that used in Sample Problem 27.4, in which the structure of an aldopentose is determined by piecing together different facts. The Fischer proof is much more complicated, though, because the relative orientation of more stereogenic centers had to be determined.

The reasoning behind the Fischer proof is easier to follow if the eight possible D-aldohexoses are arranged in pairs of epimers at C2. These compounds are labeled **1–8** in Figure 27.9. When organized in this way, each pair of epimers would also be formed as the products of a Kiliani–Fischer synthesis beginning with a particular D-aldopentose (lettered **A–D** in Figure 27.9).


To follow the steps in the Fischer proof, we must determine what information can be obtained from each experimental result.

Fact [1] Kiliani–Fischer synthesis of arabinose, an aldopentose, forms glucose and mannose.


Because the Kiliani–Fischer synthesis forms two epimers at C2, glucose and mannose have the same configurations at three stereogenic centers (C3–C5), but opposite configurations at C2. Thus, glucose and mannose are either **1** and **2**, **3** and **4**, **5** and **6**, or **7** and **8**.

Figure 27.9

The D-aldopentoses and D-aldohexoses needed to illustrate the Fischer proof




- The aldohexoses (**1–8**) are arranged in pairs of epimers at C2.
- Kiliani–Fischer synthesis using aldopentoses **A–D** forms each pair of epimers.



Fact [2] Glucose and mannose are both oxidized to optically active aldaric acids.

Because an optically active aldaric acid does not have a plane of symmetry, any aldohexose that forms an aldaric acid with a plane of symmetry can be eliminated.




- Thus, aldohexoses **1** (and therefore its epimer **2**) and **7** (and its epimer **8**) can be eliminated, so that glucose and mannose are one of two pairs of epimers: either **3** and **4**, or **5** and **6**.

Fact [3] Arabinose is oxidized to an optically active aldaric acid.

We can now narrow down the possible structures for arabinose, the aldopentose that forms glucose and mannose from the Kiliani–Fischer synthesis (Fact [1]).


- Because glucose and mannose are epimers **3** and **4** or **5** and **6** (Fact [2]), arabinose must be either **B** or **C**.
- Because oxidation of **C** forms an optically *inactive* aldaric acid, arabinose must have structure **B**, an aldopentose that gives an optically active aldaric acid.




- Thus, glucose and mannose are structures **3** and **4**, but, with the given information, it is not possible to decide which is glucose and which is mannose.

Fact [4] When the functional groups on the two end carbons of glucose are interchanged, glucose is converted to a different aldohexose.

To determine whether glucose had structure **3** or **4**, a method was devised to interchange the two functional groups at the ends of an aldohexose. The CHO at C1 was converted to CH₂OH, and the CH₂OH at C6 was converted to CHO.



The results of this process are different for compounds **3** and **4**. Compound **4** gives a compound that is identical to itself, whereas compound **3** gives a compound that is different from itself.



- Because glucose gives a different aldohexose after the two end groups are interchanged, it must have structure **3**, and mannose must have structure **4**. **The proof is complete.**


Problem 27.28 Besides D-mannose, only one other D-aldohexose yields itself when the CHO and CH₂OH groups on the end carbon atoms are interchanged. What is the name and structure of this D-aldohexose?

Problem 27.29 A D-aldohexose **A** is formed from an aldopentose **B** by the Kiliani–Fischer synthesis. Reduction of **A** with NaBH₄ forms an optically inactive alditol. Oxidation of **B** forms an optically active aldaric acid. What are the structures of **A** and **B**?


27.12 Disaccharides

Disaccharides contain two monosaccharides joined together by a glycosidic linkage. The **general features of a disaccharide** include the following:

Some features of a disaccharide



glycosidic linkage
This bond may be α or β .



a 1 \rightarrow 4- β -glycoside

- [1] Two monosaccharide rings may be five- or six-membered, but six-membered rings are much more common. The two rings are connected by an O atom that is part of an acetal, called a **glycosidic linkage**, which may be oriented α or β .
- [2] The **glycoside is formed from the anomeric carbon of one monosaccharide and any OH group on the other monosaccharide**. All disaccharides have **one acetal**, together with either a hemiacetal or another acetal.
- [3] With pyranose rings, the carbon atoms in each ring are numbered beginning with the anomeric carbon. The most common disaccharides contain two monosaccharides in which the hemiacetal carbon of one ring (C1) is joined to C4 of the other ring.


The three most abundant disaccharides are **maltose**, **lactose**, and **sucrose**.

27.12A Maltose




Maltose gets its name from malt, the liquid obtained from barley and other cereal grains.

Maltose, a disaccharide formed by the hydrolysis of starch, is found in germinated grains such as barley. Maltose contains two glucose units joined together by a 1→4- α -glycoside bond.



Because one glucose ring of maltose still contains a hemiacetal, it exists as a mixture of α and β anomers. Only the β anomer is shown. Maltose exhibits two properties of all carbohydrates that contain a hemiacetal: it undergoes **mutarotation**, and it reacts with oxidizing agents, making it a **reducing sugar**.

Hydrolysis of maltose forms two molecules of glucose. The C1–O bond is cleaved in this process, and a mixture of glucose anomers forms. The mechanism for this hydrolysis is exactly the same as the mechanism for glycoside hydrolysis in Section 27.7B.



Problem 27.30 Draw a stepwise mechanism for the acid-catalyzed hydrolysis of maltose to two molecules of glucose.


Problem 27.31 Draw the α anomer of maltose. What products are formed on hydrolysis of this form of maltose?

27.12B Lactose



Milk contains the disaccharide lactose.

As noted in the chapter opener, **lactose** is the principal disaccharide found in milk from both humans and cows. Unlike many mono- and disaccharides, lactose is not appreciably sweet. Lactose consists of **one galactose** and **one glucose unit**, joined by a 1→4- β -glycoside bond from the anomeric carbon of galactose to C4 of glucose.



Like maltose, lactose also contains a hemiacetal, so it exists as a mixture of α and β anomers. The β anomer is drawn. Lactose undergoes **mutarotation**, and it reacts with oxidizing agents, making it a **reducing sugar**.

Lactose is digested in the body by first cleaving the 1→4-β-glycoside bond using the enzyme *lactase*. Individuals who are lactose intolerant no longer produce lactase, and so they are unable to digest and absorb lactose.

Problem 27.32 Cellobiose, a disaccharide obtained by the hydrolysis of cellulose, is composed of two glucose units joined together in a 1→4-β-glycoside bond. What is the structure of cellobiose?

27.12C Sucrose

Sucrose, the disaccharide found in sugarcane and used as table sugar (Figure 27.10), is the most common disaccharide in nature. It contains **one glucose unit** and **one fructose unit**.

The structure of sucrose has several features that make it different from maltose and lactose. First of all, sucrose contains one six-membered ring (glucose) and one five-membered ring (fructose), whereas both maltose and lactose contain two six-membered rings. In sucrose the six-membered glucose ring is joined by an α-glycosidic bond to C2 of a fructofuranose ring. The numbering in a fructofuranose is different from the numbering in a pyranose ring. The anomeric carbon is now designated as C2, so the anomeric carbons of the glucose and fructose rings are both used to form the glycosidic linkage.

As a result, **sucrose contains two acetals but no hemiacetal**. Sucrose, therefore, is a **nonreducing sugar** and **it does not undergo mutarotation**.

Sucrose's pleasant sweetness has made it a widely used ingredient in baked goods, cereals, bread, and many other products. It is estimated that the average American ingests 100 lb of sucrose annually. Like other carbohydrates, however, sucrose contains many calories. To reduce caloric intake while maintaining sweetness, a variety of artificial sweeteners have been developed. These include sucralose, aspartame, and saccharin (Figure 27.11). These compounds are much sweeter than sucrose so only a small amount of each compound is needed to achieve the same level of perceived sweetness.

Figure 27.10

Sucrose





Figure 27.11
Artificial sweeteners




27.13 Polysaccharides

Polysaccharides contain three or more monosaccharides joined together. Three prevalent polysaccharides in nature are **cellulose**, **starch**, and **glycogen**, each of which consists of repeating glucose units joined by different glycosidic bonds.

27.13A Cellulose

The structure of cellulose was previously discussed in Section 5.1.

Cellulose is found in the cell walls of nearly all plants, where it gives support and rigidity to wood and plant stems. Cotton is essentially pure cellulose.




The 1→4-β-glycosidic bonds are shown in red.


Ball-and-stick models showing the three-dimensional structures of cellulose and starch were given in Figure 5.2.

Cellulose is an unbranched polymer composed of repeating glucose units joined in a 1→4-β-glycosidic linkage. The β-glycosidic linkage creates long linear chains of cellulose molecules that stack in sheets, creating an extensive three-dimensional array. A network of intermolecular hydrogen bonds between the chains and sheets means that only the few OH groups on the surface are available to hydrogen bond to water, making this very polar compound water insoluble.

Cellulose acetate, a cellulose derivative, is made by treating cellulose with acetic anhydride and sulfuric acid. The resulting product has acetate esters in place of every OH group. Cellulose acetate is spun into fibers that are used for fabrics called *acetates*, which have a deep luster and satin appearance.



Cellulose can be hydrolyzed to glucose by cleaving all of the β -glycosidic bonds, yielding both anomers of glucose.




A **β -glycosidase** is the general name of an enzyme that hydrolyzes a β -glycoside linkage.

In cells, the hydrolysis of cellulose is accomplished by an enzyme called a **β -glucosidase**, which cleaves all the β -glycoside bonds formed from glucose. Humans do not possess this enzyme, and therefore cannot digest cellulose. Ruminant animals, on the other hand, such as cattle, deer, and camels, have bacteria containing a β -glucosidase in their digestive systems, so they can derive nutritional benefit from eating grass and leaves.

27.13B Starch

Starch is the main carbohydrate found in the seeds and roots of plants. Corn, rice, wheat, and potatoes are common foods that contain a great deal of starch.

Starch is a polymer composed of repeating glucose units joined in α -glycosidic linkages. Both starch and cellulose are polymers of glucose, but starch contains α glycoside bonds, whereas cellulose contains β glycoside bonds. The two common forms of starch are **amylose** and **amylopectin**.



Amylose, which comprises about 20% of starch molecules, has an unbranched skeleton of glucose molecules with **1 \rightarrow 4- α -glycoside bonds**. Because of this linkage, an amylose chain adopts

a helical arrangement, giving it a very different three-dimensional shape from the linear chains of cellulose. Amylose was first described in Section 5.1.

Amylopectin, which comprises about 80% of starch molecules, likewise consists of a backbone of glucose units joined in **α -glycosidic bonds**, but it also contains considerable branching along the chain. The linear linkages of amylopectin are formed by **1 \rightarrow 4- α -glycoside bonds**, similar to amylose. The branches are linked to the chain with **1 \rightarrow 6- α -glycosidic linkages**.

Both forms of starch are water soluble. Because the OH groups in these starch molecules are not buried in a three-dimensional network, they are more available for hydrogen bonding with water molecules, leading to greater water solubility than cellulose has.

The ability of amylopectin to form branched polymers is a unique feature of carbohydrates. Other types of polymers in the cell, such as the proteins discussed in Chapter 28, occur in nature only as linear molecules.

α -Glycosidase is the general name of an enzyme that hydrolyzes an α -glycoside linkage.

Both amylose and amylopectin are hydrolyzed to glucose with cleavage of the glycosidic bonds. The human digestive system has the necessary **α -glucosidase** enzymes needed to catalyze this process. Bread and pasta made from wheat flour, rice, and corn tortillas are all sources of starch that are readily digested.

27.13C Glycogen

Glycogen is the major form in which polysaccharides are stored in animals. Glycogen, a polymer of glucose containing **α -glycosidic bonds**, has a branched structure similar to amylopectin, but the branching is much more extensive.

Glycogen is stored principally in the liver and muscle. When glucose is needed for energy in the cell, glucose units are hydrolyzed from the ends of the glycogen polymer, and then further metabolized with the release of energy. Because glycogen has a highly branched structure, there are many glucose units at the ends of the branches that can be cleaved whenever the body needs them.

Problem 27.33


Draw the structure of: (a) a polysaccharide formed by joining D-mannose units in 1 \rightarrow 4- β -glycosidic linkages; (b) a polysaccharide formed by joining D-glucose units in 1 \rightarrow 6- α -glycosidic linkages. The polysaccharide in (b) is dextran, a component of dental plaque.

27.14 Other Important Sugars and Their Derivatives

Many other examples of simple and complex carbohydrates with useful properties exist in the biological world. In Section 27.14, we examine some carbohydrates that contain nitrogen atoms.


27.14A Amino Sugars and Related Compounds

Amino sugars contain an NH_2 group instead of an OH group at a non-anomeric carbon. The most common amino sugar in nature, **D-glucosamine**, is formally derived from D-glucose by replacing the OH at C2 with NH_2 . Although it is not classified as a drug, and therefore not regulated by the Food and Drug Administration, glucosamine is available in many over-the-counter treatments for osteoarthritis. Glucosamine is thought to promote the repair of deteriorating cartilage in joints.



Acetylation of glucosamine with acetyl CoA (Section 22.17) forms **N-acetyl-D-glucosamine**, abbreviated as **NAG**. **Chitin**, the second most abundant carbohydrate polymer, is a polysaccharide


Problem 27.35 Draw the products of each reaction.



Problem 27.36 Draw a stepwise mechanism for the conversion of β -D-glucose to both anomers of *N*-ethyl glucopyranoside, the equation written in Reaction [1].


The prefix **deoxy** means “without oxygen.”

The *N*-glycosides of two sugars, **D-ribose** and **2-deoxy-D-ribose**, are especially noteworthy, because they form the building blocks of RNA and DNA, respectively. 2-Deoxyribose is so named because it lacks an OH group at C2 of ribose.




- Reaction of D-ribose with certain amine heterocycles forms *N*-glycosides called ribonucleosides.
- This same reaction of 2-deoxy-D-ribose forms deoxyribonucleosides.

An example of a **ribonucleoside** and a **deoxyribonucleoside** are drawn. These *N*-glycosides have the β orientation. Numbering in the sugar ring begins at the anomeric carbon (1'), and proceeds in a clockwise fashion around the ring.




Only five common nitrogen heterocycles are used to form these nucleosides. Three compounds have one ring, and are derived from a nitrogen heterocycle called **pyrimidine**. Two are bicyclic, and are derived from a nitrogen heterocycle called **purine**. These five amines are referred to as **bases**. Each base is designated by a one-letter abbreviation, as shown in the names and structures drawn. Note that uracil (U) occurs only in ribonucleosides and thymine (T) occurs only in deoxyribonucleosides.

- Each nucleoside has two parts, a sugar and a base, joined together by a β *N*-glycosidic linkage.




When one OH group of the sugar nucleus is bonded to a phosphate, the derivatives are called **ribonucleotides** and **deoxyribonucleotides**.



- Ribonucleotides are the building blocks of the polymer ribonucleic acid, or RNA, the messenger molecules that convert genetic information into proteins.
- Deoxyribonucleotides are the building blocks of the polymer deoxyribonucleic acid, or DNA, the molecules that are responsible for the storage of all genetic information.

Short segments of both RNA and DNA are shown in Figure 27.12. Note the central role of the sugar moiety in both RNA and DNA. The sugar residues are bonded to two phosphate groups, thus connecting the chain of RNA or DNA together. The sugar residues are also bonded to the nitrogen base via the anomeric carbon.



DNA is composed of two polynucleotide strands that wind around each other to form a double helix, resembling a spiral ladder. The sides of the ladder are composed of the sugar–phosphate backbone of the polymer and the rungs are composed of the bases, as shown in Figure 27.13.

The nitrogen bases on one strand of DNA hydrogen bond to nitrogen bases on the other strand. A purine base on one strand hydrogen bonds with a pyrimidine base on the other strand. Two types of bases, called **base pairs**, hydrogen bond to each other: adenine hydrogen bonds with thymine (A–T) and cytosine hydrogen bonds with guanine (C–G). Although much more can be said about the intricate structure of DNA, our discussion will end here since the focus of this chapter is carbohydrates, not nucleic acids.





Figure 27.12
Short segments of RNA
and DNA

Problem 27.37 Draw the structures of the nucleosides formed from each of the following components:
(a) ribose + uracil; (b) 2-deoxyribose + guanine

Problem 27.38 (a) Why can't two purine bases (A and G) form a base pair and hydrogen bond to each other on two strands of DNA in the double helix? (b) Why is hydrogen bonding between guanine and cytosine more favorable than hydrogen bonding between guanine and thymine?

Figure 27.13 DNA—A double helix



- Two polynucleotide strands form the double helix of DNA. The backbone of each polymer strand is composed of sugar–phosphate residues. Hydrogen bonding of base pairs (A–T and C–G) holds the two strands of DNA together.


KEY CONCEPTS

Carbohydrates

Important Terms


- **Aldose** A monosaccharide containing an aldehyde (27.2)
- **Ketose** A monosaccharide containing a ketone (27.2)
- **D-Sugar** A monosaccharide with the OH bonded to the stereogenic center farthest from the carbonyl group drawn on the right in the Fischer projection (27.2C)
- **Epimers** Two diastereomers that differ in configuration around one stereogenic center only (27.3)
- **Anomers** Monosaccharides that differ in configuration at the hemiacetal OH group (27.6)
- **Glycoside** An acetal derived from a monosaccharide hemiacetal (27.7)

Acyclic, Haworth, and 3-D Representations for D-Glucose (27.6)




Reactions of Monosaccharides Involving the Hemiacetal

[1] Glycoside formation (27.7A)



- Only the hemiacetal OH reacts.
- A mixture of α and β glycosides forms.


[2] Glycoside hydrolysis (27.7B)



- A mixture of α and β anomers forms.


Reactions of Monosaccharides at the OH Groups

[1] Ether formation (27.8)



- All OH groups react.
- The stereochemistry at all stereogenic centers is retained.


[2] Ester formation (27.8)



- All OH groups react.
- The stereochemistry at all stereogenic centers is retained.


Reactions of Monosaccharides at the Carbonyl Group

[1] Oxidation of aldoes (27.9B)




- Aldonic acids are formed using:
 - Ag_2O , NH_4OH
 - Cu^{2+}
 - Br_2 , H_2O
- Aldaric acids are formed with HNO_3 , H_2O .

[2] Reduction of aldoes to alditols (27.9A)




[3] Wohl degradation (27.10A)



- The C1-C2 bond is cleaved to shorten an aldose chain by one carbon.
- The stereochemistry at all other stereogenic centers is retained.
- Two epimers at C2 form the same product.



[4] Kiliani-Fischer synthesis (27.10B)



- One carbon is added to the aldehyde end of an aldose.
- Two epimers at C2 are formed.



Other Reactions

[1] Hydrolysis of disaccharides (27.12)

[2] Formation of *N*-glycosides (27.14B)

PROBLEMS

Fischer Projections


27.39 Classify each compound as identical to **A** or its enantiomer.27.40 Convert each compound to a Fischer projection and label each stereogenic center as *R* or *S*.

Monosaccharide Structure and Stereochemistry

27.41 Draw the C4 epimer of D-xylose and name the monosaccharide.

27.42 For D-arabinose:


- Draw its enantiomer.
- Draw an epimer at C3.
- Draw a diastereomer that is not an epimer.
- Draw a constitutional isomer that still contains a carbonyl group.

27.43 Consider the following six compounds (**A–F**).

How are the two compounds in each pair related? Choose from enantiomers, epimers, diastereomers but not epimers, constitutional isomers, and identical compounds.

- A** and **B**
- A** and **C**
- B** and **C**
- A** and **D**
- E** and **F**

27.44 Consider the monosaccharide aldopentoses **A** and **B**, drawn below.




- Which of the following terms describe **A** and **B**: epimers, anomers, enantiomers, diastereomers, and reducing sugars?
- Draw the acyclic form of both **A** and **B**, and name each compound.


27.45 Draw a Haworth projection for each compound using the structures in Figures 27.4 and 27.5.

- β -D-talopyranose
- β -D-mannopyranose
- α -D-galactopyranose
- α -D-ribofuranose
- α -D-tagatofuranose

27.46 Draw both pyranose anomers of each aldohexose using a three-dimensional representation with a chair pyranose. Label each anomer as α or β .



27.47 Convert each cyclic monosaccharide into its acyclic form.




27.48 D-Arabinose can exist in both pyranose and furanose forms.

- Draw the α and β anomers of D-arabinofuranose.
- Draw the α and β anomers of D-arabinopyranose.

27.49 The most stable conformation of the pyranose ring of most D-aldohexoses places the largest group, CH_2OH , in the equatorial position. An exception to this is the aldohexose D-idose. Draw the two possible chair conformations of either the α or β anomer of D-idose. Explain why the more stable conformation has the CH_2OH group in the axial position.


Monosaccharide Reactions

27.50 Draw the products formed when α -D-gulose is treated with each reagent.



- CH_3I , Ag_2O
- CH_3OH , HCl
- $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$, Ag_2O
- $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$, HCl
- Ac_2O , pyridine
- $\text{C}_6\text{H}_5\text{COCl}$, pyridine
- The product in (a), then H_3O^+
- The product in (b), then Ac_2O , pyridine
- The product in (g), then $\text{C}_6\text{H}_5\text{CH}_2\text{Cl}$, Ag_2O
- The product in (d), then CH_3I , Ag_2O

27.51 Draw the products formed when D-altrose is treated with each reagent.




- CH_3OH , HCl
- $(\text{CH}_3)_2\text{CHOH}$, HCl
- NaBH_4 , CH_3OH
- Br_2 , H_2O
- HNO_3 , H_2O
- $[\text{1}] \text{NH}_2\text{OH}$; $[\text{2}] (\text{CH}_3\text{CO})_2\text{O}$, NaOCOCH_3 ; $[\text{3}] \text{NaOCH}_3$
- $[\text{1}] \text{NaCN}$, HCl ; $[\text{2}] \text{H}_2$, Pd-BaSO_4 ; $[\text{3}] \text{H}_3\text{O}^+$
- CH_3I , Ag_2O
- Ac_2O , pyridine
- $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$, mild H^+

27.52 Answer Problem 27.51 using D-xylose as the starting material.


27.53 What aglycon and monosaccharides are formed when salicin and solanine (Section 27.7C) are each hydrolyzed with aqueous acid?

27.54 What two aldohexoses yield D-arabinose upon Wohl degradation?

27.55 What products are formed when each compound is subjected to a Kiliani-Fischer synthesis?




27.56 How would you convert D-glucose into each compound? More than one step is required.




27.57 Which D-aldopentoses are reduced to optically inactive alditols using NaBH_4 , CH_3OH ?

27.58 What products are formed when each compound is treated with aqueous acid?




Mechanisms


27.59 Draw a stepwise mechanism for the acid-catalyzed interconversion of two glucose anomers by mutarotation.




27.60 Draw a stepwise mechanism for the following reaction.




27.61 Draw a stepwise mechanism for the following hydrolysis.



- 27.62** In the oxidation of D-allose to D-allonic acid, a lactone having the general structure **A** is isolated. Draw a stepwise mechanism to account for the formation of **A**. Use wedges and dashes to indicate the stereochemistry of all stereogenic centers in **A**.



- 27.63** The following isomerization reaction, drawn using D-glucose as starting material, occurs with all aldohexoses in the presence of base. Draw a stepwise mechanism that illustrates how each compound is formed.




Identifying Monosaccharides

- 27.64** Which D-aldopentose is oxidized to an optically active aldaric acid and undergoes the Wohl degradation to yield a D-aldotetrose that is oxidized to an optically active aldaric acid?
- 27.65** What other D-aldopentose forms the same alditol as D-arabinose when reduced with NaBH_4 in CH_3OH ?
- 27.66** Identify compounds **A–D**. A D-aldopentose **A** is oxidized with HNO_3 to an optically inactive aldaric acid **B**. **A** undergoes the Kiliani–Fischer synthesis to yield **C** and **D**. **C** is oxidized to an optically active aldaric acid. **D** is oxidized to an optically inactive aldaric acid.
- 27.67** A D-aldopentose **A** is reduced to an optically active alditol. Upon Kiliani–Fischer synthesis, **A** is converted to two D-aldohexoses, **B** and **C**. **B** is oxidized to an optically inactive aldaric acid. **C** is oxidized to an optically active aldaric acid. What are the structures of **A–C**?
- 27.68** A D-aldohexose **A** is reduced to an optically active alditol **B** using NaBH_4 in CH_3OH . **A** is converted by Wohl degradation to an aldopentose **C**, which is reduced to an optically inactive alditol **D**. **C** is converted by Wohl degradation to aldotetrose **E**, which is oxidized to an optically active aldaric acid **F**. When the two ends of aldohexose **A** are interconverted, a different aldohexose **G** is obtained. What are the structures of **A–G**?

Disaccharides and Polysaccharides


- 27.69** Draw the structure of a disaccharide formed from two galactose units joined by a 1→4-β-glycosidic linkage.
- 27.70** Draw the structure of a disaccharide formed from two mannose units joined by a 1→4-α-glycosidic linkage.
- 27.71** Identify the lettered compounds in the following reactions.



27.72 For each disaccharide in Problem 27.71:

- Identify the glycosidic linkage.
- Classify the glycosidic bond as α or β and use numbers to designate its location.
- Classify each disaccharide as reducing or nonreducing.


27.73 Consider the tetrasaccharide stachyose drawn below. Stachyose is found in white jasmine, soybeans, and lentils. Because humans cannot digest it, its consumption causes flatulence.



- Label all glycoside bonds.
- Classify each glycosidic linkage as α or β and use numbers to designate its location between two rings (e.g., 1 \rightarrow 4- β).
- What products are formed when stachyose is hydrolyzed with H_3O^+ ?
- Is stachyose a reducing sugar?
- What product is formed when stachyose is treated with excess CH_3I , Ag_2O ?
- What products are formed when the product in (e) is treated with H_3O^+ ?


27.74 Deduce the structure of the disaccharide isomaltose from the following data.

- Hydrolysis yields D-glucose exclusively.
- Isomaltose is cleaved with α -glycosidase enzymes.
- Isomaltose is a reducing sugar.
- Methylation with excess CH_3I , Ag_2O and then hydrolysis with H_3O^+ forms two products:



27.75 Deduce the structure of the disaccharide trehalose from the following data. Trehalose is the "blood sugar" of the insect world. It is found in bacterial spores, fungi, and many insects whose natural environment has large variations in temperature.

- Hydrolysis yields D-glucose exclusively.
- Trehalose is hydrolyzed by α -glycosidase enzymes.
- Trehalose is a nonreducing sugar.
- Methylation with excess CH_3I , Ag_2O , followed by hydrolysis with H_3O^+ , forms only one product:




27.76 Draw the structure of each of the following compounds:

- A polysaccharide formed by joining D-glucosamine in 1 \rightarrow 6- α -glycosidic linkages.
- A disaccharide formed by joining D-mannose and D-glucose in a 1 \rightarrow 4- β -glycosidic linkage using mannose's anomeric carbon.
- An α -N-glycoside formed from D-arabinose and $\text{C}_6\text{H}_5\text{CH}_2\text{NH}_2$.
- A ribonucleoside formed from D-ribose and thymine.


Challenge Problems

27.77 Draw a stepwise mechanism for the following reaction.



27.78 Deduce the structure of the trisaccharide (**X**) from the given information.

[1] Methylation with excess CH_3I , Ag_2O and then hydrolysis with H_3O^+ forms three products.



(Both anomers of each compound are formed.)


[2] **X** is cleaved with a β -glycosidase enzyme to give a disaccharide and D-galactose.

[3] **X** is cleaved with an α -glycosidase enzyme to give a disaccharide and D-fructose.

28

Amino Acids and Proteins

- 28.1 Amino acids
- 28.2 Synthesis of amino acids
- 28.3 Separation of amino acids
- 28.4 Enantioselective synthesis of amino acids
- 28.5 Peptides
- 28.6 Peptide sequencing
- 28.7 Peptide synthesis
- 28.8 Automated peptide synthesis
- 28.9 Protein structure
- 28.10 Important proteins



Myoglobin is a globular protein that contains 153 amino acids joined together, as well as a non-protein portion called a heme unit. The heme group consists of a large nitrogen heterocycle complexed with the Fe^{2+} cation. The Fe^{2+} cation binds oxygen in the blood and stores it in tissues. Whales have a particularly high myoglobin concentration in their muscles. It serves as an oxygen reservoir for the whale while it is submerged for long periods of time. In Chapter 28, we discuss the properties of proteins and the amino acids from which they are synthesized.

Of the four major groups of biomolecules—lipids, carbohydrates, nucleic acids, and proteins—proteins have the widest array of functions. **Keratin** and **collagen**, for example, are part of a large group of structural proteins that form long insoluble fibers, giving strength and support to tissues. Hair, horns, hooves, and fingernails are all made up of keratin. **Collagen** is found in bone, connective tissue, tendons, and cartilage. **Enzymes** are proteins that catalyze and regulate all aspects of cellular function. **Membrane proteins** transport small organic molecules and ions across cell membranes. **Insulin**, the hormone that regulates blood glucose levels, **fibrinogen** and **thrombin**, which form blood clots, and **hemoglobin**, which transports oxygen from the lungs to tissues, are all proteins.


In Chapter 28 we discuss proteins and their primary components, the amino acids.

28.1 Amino Acids

Amino acids were first discussed in Section 19.14.

Naturally occurring amino acids have an amino group (NH_2) bonded to the α carbon of a carboxy group (COOH), and so they are called **α -amino acids**.

- All proteins are polyamides formed by joining amino acids together.



28.1A General Features of α -Amino Acids

The 20 amino acids that occur naturally in proteins differ in the identity of the R group bonded to the α carbon. The R group is called the **side chain** of the amino acid.


The simplest amino acid, called glycine, has $\text{R} = \text{H}$. **All other amino acids ($\text{R} \neq \text{H}$) have a stereogenic center on the α carbon.** As is true for monosaccharides, the prefixes **D** and **L** are used to designate the configuration at the stereogenic center of amino acids. Common, naturally occurring amino acids are called **L-amino acids**. Their enantiomers, D-amino acids, are rarely found in nature. These general structures are shown in Figure 28.1. According to *R,S* designations, all L-amino acids except cysteine have the **S configuration**.

All amino acids have common names. These names can be represented by either a one-letter or a three-letter abbreviation. Figure 28.2 is a listing of the 20 naturally occurring amino acids, together with their abbreviations. Note the variability in the R groups. A side chain can be a simple alkyl group, or it can have additional functional groups such as OH, SH, COOH, or NH_2 .


- Amino acids with an additional COOH group in the side chain are called acidic amino acids.
- Those with an additional basic N atom in the side chain are called basic amino acids.
- All others are neutral amino acids.

Figure 28.1
The general features of an α -amino acid

Simplest amino acid, $\text{R} = \text{H}$


















Two possible enantiomers when $\text{R} \neq \text{H}$





Only this isomer occurs in proteins.

Figure 28.2 The 20 naturally occurring amino acids




Neutral amino acids

Name	Structure	Abbreviations	Name	Structure	Abbreviations
Alanine		Ala A	Phenylalanine*		Phe F
Asparagine		Asn N	Proline		Pro P
Cysteine		Cys C	Serine		Ser S
Glutamine		Gln Q	Threonine*		Thr T
Glycine		Gly G	Tryptophan*		Trp W
Isoleucine*		Ile I	Tyrosine		Tyr Y
Leucine*		Leu L	Valine*		Val V
Methionine*		Met M			

Acidic amino acids

Name	Structure	Abbreviations
Aspartic acid		Asp D
Glutamic acid		Glu E


Basic amino acids

Name	Structure	Abbreviations
Arginine*		Arg R
Histidine*		His H
Lysine*		Lys K

Essential amino acids are labeled with an asterisk (*).

Look closely at the structures of proline, isoleucine, and threonine.

- **All amino acids are 1° amines except for proline**, which has its N atom in a five-membered ring, making it a 2° amine.
- **Isoleucine and threonine** contain an additional stereogenic center at the β carbon, so there are four possible stereoisomers, only one of which is naturally occurring.




Humans can synthesize only 10 of these 20 amino acids. The remaining 10 are called **essential amino acids** because they must be obtained from the diet. These are labeled with an asterisk in Figure 28.2.

Problem 28.1 Draw the other three stereoisomers of L-isoleucine, and label the stereogenic centers as *R* or *S*.

28.1B Acid–Base Behavior

Recall from Section 19.14B that an amino acid has both an acidic and a basic functional group, so proton transfer forms a salt called a **zwitterion**.



The zwitterion is neutral.

This neutral form of an amino acid does **not** really exist.

This salt is the neutral form of an amino acid.

This form exists at **pH \approx 6**.

- **Amino acids do not exist to any appreciable extent as uncharged neutral compounds. They exist as salts, giving them high melting points and making them water soluble.**

Amino acids exist in different charged forms, as shown in Figure 28.3, depending on the pH of the aqueous solution in which they are dissolved. For neutral amino acids, the overall charge is +1, 0, or -1. Only at pH \sim 6 does the zwitterionic form exist.

The $-\text{COOH}$ and $-\text{NH}_3^+$ groups of an amino acid are ionizable, because they can lose a proton in aqueous solution. As a result, they have different $\text{p}K_a$ values. The $\text{p}K_a$ of the $-\text{COOH}$ group is typically \sim 2, whereas that of the $-\text{NH}_3^+$ group is \sim 9, as shown in Table 28.1.

Some amino acids, such as aspartic acid and lysine, have acidic or basic side chains. These additional ionizable groups complicate somewhat the acid–base behavior of these amino acids. Table 28.1 lists the $\text{p}K_a$ values for these acidic and basic side chains as well.

Figure 28.3
How the charge of a neutral amino acid depends on the pH




Table 28.1 pK_a Values for the Ionizable Functional Groups of an α -Amino Acid

Amino acid	α -COOH	α -NH ₃ ⁺	Side chain	pI
Alanine	2.35	9.87	—	6.11
Arginine	2.01	9.04	12.48	10.76
Asparagine	2.02	8.80	—	5.41
Aspartic acid	2.10	9.82	3.86	2.98
Cysteine	2.05	10.25	8.00	5.02
Glutamic acid	2.10	9.47	4.07	3.08
Glutamine	2.17	9.13	—	5.65
Glycine	2.35	9.78	—	6.06
Histidine	1.77	9.18	6.10	7.64
Isoleucine	2.32	9.76	—	6.04
Leucine	2.33	9.74	—	6.04
Lysine	2.18	8.95	10.53	9.74
Methionine	2.28	9.21	—	5.74
Phenylalanine	2.58	9.24	—	5.91
Proline	2.00	10.00	—	6.30
Serine	2.21	9.15	—	5.68
Threonine	2.09	9.10	—	5.60
Tryptophan	2.38	9.39	—	5.88
Tyrosine	2.20	9.11	10.07	5.63
Valine	2.29	9.72	—	6.00

Table 28.1 also lists the isoelectric points (pI) for all of the amino acids. Recall from Section 19.14C that the **isoelectric point is the pH at which an amino acid exists primarily in its neutral form**, and that it can be calculated from the average of the pK_a values of the α -COOH and α -NH₃⁺ groups (for neutral amino acids only).

Problem 28.2 What form exists at the isoelectric point of each of the following amino acids: (a) valine; (b) leucine; (c) proline; (d) glutamic acid?



Problem 28.3 Explain why the pK_a of the $-\text{NH}_3^+$ group of an α -amino acid is lower than the pK_a of the ammonium ion derived from a 1° amine (RNH_3^+). For example the pK_a of the $-\text{NH}_3^+$ group of alanine is 9.7 but the pK_a of CH_3NH_3^+ is 10.63.

28.2 Synthesis of Amino Acids

Amino acids can be prepared in a variety of ways in the laboratory. Three methods are described, each of which is based on reactions learned in previous chapters.

28.2A S_N2 Reaction of α -Halo Acids with NH_3

The most direct way to synthesize an α -amino acid is by S_N2 reaction of an α -halo carboxylic acid with a large excess of NH_3 .




Although the alkylation of ammonia with simple alkyl halides does not generally afford high yields of 1° amines (Section 25.7A), this reaction using α -halo carboxylic acids does form the desired amino acids in good yields. In this case, the amino group in the product is both less basic and more sterically crowded than other 1° amines, so that a single alkylation occurs and the desired amino acid is obtained.


Problem 28.4 What α -halo carbonyl compound is needed to synthesize each amino acid: (a) glycine; (b) isoleucine; (c) phenylalanine?

28.2B Alkylation of a Diethyl Malonate Derivative


The second method for preparing amino acids is based on the malonic ester synthesis. Recall from Section 23.9 that this synthesis converts diethyl malonate to a carboxylic acid with a new alkyl group on its α carbon atom.



This reaction can be adapted to the synthesis of α -amino acids by using a commercially available derivative of diethyl malonate as starting material. This compound, **diethyl acetamidomalonate**, has a nitrogen atom on the α carbon, which ultimately becomes the NH_2 group on the α carbon of the amino acid.




The malonic ester synthesis consists of three steps, and so does this variation to prepare an amino acid.



- [1] **Deprotonation** of diethyl acetamidomalonate with NaOEt forms an enolate by removal of the acidic proton between the two carbonyl groups.
- [2] **Alkylation** of the enolate with an unhindered alkyl halide (usually CH_3X or RCH_2X) forms a substitution product with a new R group on the α carbon.
- [3] Heating the alkylation product with aqueous acid results in **hydrolysis** of both esters and the amide, followed by **decarboxylation** to form the amino acid.

Phenylalanine, for example, can be synthesized as follows:




Problem 28.5 The enolate derived from diethyl acetamidomalonate is treated with each of the following alkyl halides. After hydrolysis and decarboxylation, what amino acid is formed: (a) CH_3I ; (b) $(\text{CH}_3)_2\text{CHCH}_2\text{Cl}$; (c) $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)\text{Br}$?


Problem 28.6 What amino acid is formed when $\text{CH}_3\text{CONHCH}(\text{CO}_2\text{Et})_2$ is treated with the following series of reagents: [1] NaOEt; [2] $\text{CH}_2=\text{O}$; [3] H_3O^+ , Δ ?

28.2C Strecker Synthesis

The third method, the **Strecker amino acid synthesis**, converts an aldehyde into an amino acid by a two-step sequence that adds one carbon atom to the aldehyde carbonyl. Treating an aldehyde with NH_4Cl and NaCN first forms an α -amino nitrile, which can then be hydrolyzed in aqueous acid to an amino acid.



The Strecker synthesis of alanine, for example, is as follows:




Mechanism 28.1 for the formation of the α -amino nitrile from an aldehyde (the first step in the Strecker synthesis) consists of two parts: **nucleophilic addition of NH_3** to form an imine, followed by **addition of cyanide** to the $\text{C}=\text{N}$ bond. Both parts are related to earlier mechanisms involving imines (Section 21.11) and cyanohydrins (Section 21.9).




Mechanism 28.1 Formation of an α -Amino Nitrile

Part [1] Nucleophilic attack of NH_3 to form an imine



• **Part [1]** Nucleophilic attack of NH_3 followed by proton transfer and loss of H_2O forms an imine. Loss of H_2O occurs by the same three-step process outlined in Mechanism 21.5.


Part [2] Nucleophilic attack of CN^- to form an α -amino nitrile



• **Part [2]** Protonation of the imine followed by nucleophilic attack of CN^- gives the α -amino nitrile.

Figure 28.4

The synthesis of methionine by three different methods



Three methods of amino acid synthesis:

[1] S_N2 reaction using an α-halo carboxylic acid

[2] Alkylation of diethyl acetamidomalonate

[3] Strecker synthesis

The details of the second step of the Strecker synthesis, the hydrolysis of a nitrile (RCN) to a carboxylic acid (RCOOH), have already been presented in Section 22.18A.

Figure 28.4 shows how the amino acid methionine can be prepared by all three methods in Section 28.2.

Problem 28.7 What aldehyde is needed to synthesize each amino acid by the Strecker synthesis: (a) valine; (b) leucine; (c) phenylalanine?

Problem 28.8 Draw the products of each reaction.



28.3 Separation of Amino Acids

No matter which of the preceding methods is used to synthesize an amino acid, all three yield a racemic mixture. Naturally occurring amino acids exist as a single enantiomer, however, so the two enantiomers obtained must be separated if they are to be used in biological applications. This is not an easy task. Two enantiomers have the same physical properties, so they cannot be separated by common physical methods, such as distillation or chromatography. Moreover, they react in the same way with achiral reagents, so they cannot be separated by chemical reactions either.

Nonetheless, strategies have been devised to separate two enantiomers using physical separation techniques and chemical reactions. We examine two different strategies in Section 28.3. Then, in Section 28.4, we will discuss a method that affords optically active amino acids without the need for separation.


- The separation of a racemic mixture into its component enantiomers is called *resolution*. Thus, a racemic mixture is *resolved* into its component enantiomers.

28.3A Resolution of Amino Acids

The oldest, and perhaps still the most widely used method to separate enantiomers exploits the following fact: **enantiomers have the same physical properties, but diastereomers have**

Figure 28.5

Resolution of a racemic mixture by converting it to a mixture of diastereomers



Enantiomers A and B can be separated by reaction with a single enantiomer of a chiral reagent, Y. The process of resolution requires three steps:


- [1] Reaction of enantiomers **A** and **B** with **Y** forms two diastereomers, **AY** and **BY**.
- [2] Diastereomers **AY** and **BY** have different physical properties, so they can be separated by physical methods such as fractional distillation or crystallization.
- [3] **AY** and **BY** are then re-converted to **A** and **B** by a chemical reaction. The two enantiomers **A** and **B** are now separated from each other, and resolution is complete.

different physical properties. Thus, a racemic mixture can be resolved using the following general strategy.

- [1] **Convert a pair of enantiomers into a pair of diastereomers**, which are now separable because they have different melting points and boiling points.
- [2] **Separate the diastereomers.**
- [3] **Re-convert each diastereomer into the original enantiomer**, now separated from the other.


This general three-step process is illustrated in Figure 28.5.

To resolve a racemic mixture of amino acids such as (*R*)- and (*S*)-alanine, the racemate is first treated with acetic anhydride to form ***N*-acetyl amino acids**. Each of these amides contains one stereogenic center and they are still *enantiomers*, so they are *still inseparable*.




HOW TO Use (*R*)- α -Methylbenzylamine to Resolve a Racemic Mixture of Amino Acids

Step [1] React both enantiomers with the *R* isomer of the chiral amine.




These salts have the *same* configuration around one stereogenic center, but the *opposite* configuration about the other stereogenic center.

Step [2] Separate the diastereomers.



Step [3] Regenerate the amino acid by hydrolysis of the amide.




Step [1] is just an acid–base reaction in which the racemic mixture of *N*-acetyl alanines reacts with the same enantiomer of the resolving agent, in this case (*R*)- α -methylbenzylamine. The salts that form are **diastereomers, not enantiomers**, because they have the same configuration about one stereogenic center, but the opposite configuration about the other stereogenic center.

In **Step [2]**, the diastereomers are separated by some physical technique, such as crystallization or distillation.

In **Step [3]**, the amides can be hydrolyzed with aqueous base to regenerate the amino acids. The amino acids are now separated from each other. The optical activity of the amino acids can be measured and compared to their known rotations to determine the purity of each enantiomer.

Problem 28.9 Which of the following amines can be used to resolve a racemic mixture of amino acids?




Problem 28.10 Write out a stepwise sequence that shows how a racemic mixture of leucine enantiomers can be resolved into optically active amino acids using (*R*)- α -methylbenzylamine.

28.3B Kinetic Resolution of Amino Acids Using Enzymes


A second strategy used to separate amino acids is based on the fact that two enantiomers react differently with chiral reagents. An **enzyme** is typically used as the chiral reagent.

To illustrate this strategy, we begin again with the two enantiomers of *N*-acetyl alanine, which were prepared by treating a racemic mixture of (*R*)- and (*S*)-alanine with acetic anhydride (Section 28.3A). Enzymes called **acylases** hydrolyze amide bonds, such as those found in *N*-acetyl alanine, but only for amides of L-amino acids. Thus, when a racemic mixture of *N*-acetyl alanines is treated with an acylase, only the amide of L-alanine (the *S* stereoisomer) is hydrolyzed to generate L-alanine, whereas the amide of D-alanine (the *R* stereoisomer) is untouched. The reaction mixture now consists of one amino acid and one *N*-acetyl amino acid. Because they have different functional groups with different physical properties, they can be physically separated.



- Separation of two enantiomers by a chemical reaction that selectively occurs for only one of the enantiomers is called *kinetic resolution*.

Problem 28.11 Draw the organic products formed in the following reaction.




28.4 Enantioselective Synthesis of Amino Acids


Although the two methods introduced in Section 28.3 for resolving racemic mixtures of amino acids make enantiomerically pure amino acids available for further research, half of the reaction product is useless because it has the undesired configuration. Moreover, each of these procedures is costly and time-consuming.

If we use a chiral reagent to synthesize an amino acid, however, it is possible to favor the formation of the desired enantiomer over the other, without having to resort to a resolution. For example, single enantiomers of amino acids have been prepared by using **enantioselective (or asymmetric) hydrogenation reactions**. The success of this approach depends on finding a chiral catalyst, in much the same way that a chiral catalyst is used for the Sharpless asymmetric epoxidation (Section 12.15).

The necessary starting material is an alkene. Addition of H₂ to the double bond forms an *N*-acetyl amino acid with a new stereogenic center on the α carbon to the carboxy group. With proper choice of a chiral catalyst, the naturally occurring *S* configuration can be obtained as product.




Several chiral catalysts with complex structures have now been developed for this purpose. Many contain **rhodium** as the metal, complexed to a chiral molecule containing one or more phosphorus atoms. One example, abbreviated simply as **Rh***, is drawn below.



Ryoji Noyori shared the 2001 Nobel Prize in Chemistry for developing methods for asymmetric hydrogenation reactions using the chiral BINAP catalyst.

This catalyst is synthesized from a rhodium salt and a phosphorus compound, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (**BINAP**). It is the BINAP moiety (Figure 28.6) that makes the catalyst chiral.


Figure 28.6
The structure of BINAP



Twistoflex and helicene (Section 17.5) are two more aromatic compounds whose shape makes them chiral.

BINAP is one of a small number of molecules that is chiral even though it has no tetrahedral stereogenic centers. Its shape makes it a chiral molecule. The two naphthalene rings of the BINAP molecule are oriented at almost 90° to each other to minimize steric interactions between the hydrogen atoms on adjacent rings. This rigid three-dimensional shape makes BINAP non-superimposable on its mirror image, and thus it is a chiral compound.

The following graphic shows how enantioselective hydrogenation can be used to synthesize a single stereoisomer of phenylalanine. Treating achiral alkene **A** with H_2 and the chiral rhodium catalyst Rh^* forms the *S* isomer of *N*-acetyl phenylalanine in 100% *ee*. Hydrolysis of the acetyl group on nitrogen then yields a single enantiomer of phenylalanine.




Problem 28.12 What alkene is needed to synthesize each amino acid by an enantioselective hydrogenation reaction using H_2 and Rh^* : (a) alanine; (b) leucine; (c) glutamine?

28.5 Peptides

When amino acids are joined together by amide bonds, they form larger molecules called **peptides** and **proteins**.

- A *dipeptide* has two amino acids joined together by *one* amide bond.
- A *tripeptide* has three amino acids joined together by *two* amide bonds.




Polypeptides and **proteins** both have many amino acids joined together in long linear chains, but the term **protein** is usually reserved for polymers of more than 40 amino acids.

- The amide bonds in peptides and proteins are called *peptide bonds*.
- The individual amino acids are called *amino acid residues*.


28.5A Simple Peptides

To form a dipeptide, the amino group of one amino acid forms an amide bond with the carboxy group of another amino acid. Because each amino acid has both an amino group and a carboxy group, **two different dipeptides can be formed**. This is illustrated with alanine and cysteine.

[1] The COOH group of alanine can combine with the NH₂ group of cysteine.



[2] The COOH group of cysteine can combine with the NH₂ group of alanine.



These compounds are **constitutional isomers** of each other. Both have a free amino group at one end of their chains and a free carboxy group at the other.

- The amino acid with the free amino group is called the **N-terminal amino acid**.
- The amino acid with the free carboxy group is called the **C-terminal amino acid**.


By convention, **the N-terminal amino acid is always written at the left end of the chain and the C-terminal amino acid at the right**. The peptide can be abbreviated by writing the one- or three-letter symbols for the amino acids in the chain from the N-terminal to the C-terminal end. Thus, Ala-Cys has alanine at the N-terminal end and cysteine at the C-terminal end, whereas Cys-Ala has cysteine at the N-terminal end and alanine at the C-terminal end. Sample Problem 28.1 shows how this convention applies to a tripeptide.

Sample Problem 28.1

Draw the structure of the following tripeptide, and label its N-terminal and C-terminal amino acids: Ala-Gly-Ser.

Solution

Draw the structures of the amino acids in order from **left to right**, placing the COOH of one amino acid *next* to the NH₂ group of the adjacent amino acid. Always draw the **NH₂ group on the left** and the **COOH group on the right**. Then, join adjacent COOH and NH₂ groups together in amide bonds to form the tripeptide.



The N-terminal amino acid is **alanine**, and the C-terminal amino acid is **serine**.


The tripeptide in Sample Problem 28.1 has one N-terminal amino acid, one C-terminal amino acid, and two peptide bonds.

- No matter how many amino acid residues are present, there is only *one* N-terminal amino acid and *one* C-terminal amino acid.
- For n amino acids in the chain, the number of amide bonds is $n - 1$.

Problem 28.13 Draw the structure of each peptide. Label the N-terminal and C-terminal amino acids and all amide bonds.

- a. Val-Glu b. Gly-His-Leu c. M-A-T-T


Problem 28.14 Name each peptide using both the one-letter and the three-letter abbreviations for the names of the component amino acids.



Problem 28.15 How many different tripeptides can be formed from three different amino acids?


28.5B The Peptide Bond

The carbonyl carbon of an amide is sp^2 hybridized and has **trigonal planar** geometry. A second resonance structure can be drawn that delocalizes the nonbonded electron pair on the N atom. Amides are more resonance stabilized than other acyl compounds, so the resonance structure having the $C=N$ makes a significant contribution to the hybrid.



Resonance stabilization has important consequences. Rotation about the C–N bond is restricted because it has partial double bond character. As a result, there are two possible conformations.

Two conformations of the peptide bond




Recall from Section 16.6 that 1,3-butadiene can also exist as *s-cis* and *s-trans* conformations. In 1,3-butadiene, the *s-cis* conformation has the two double bonds on the same side of the single bond (dihedral angle = 0°), whereas the *s-trans* conformation has them on opposite sides (dihedral angle = 180°).

- The *s-trans* conformation has the two R groups oriented on *opposite* sides of the C–N bond.
- The *s-cis* conformation has the two R groups oriented on the *same* side of the C–N bond.
- The *s-trans* conformation of a peptide bond is typically more stable than the *s-cis*, because the *s-trans* has the two bulky R groups located farther from each other.

A second consequence of resonance stabilization is that **all six atoms involved in the peptide bond lie in the same plane**. All bond angles are $\sim 120^\circ$ and the C=O and N–H bonds are oriented 180° from each other.

The planar geometry of the peptide bond is analogous to the planar geometry of ethylene (or any other alkene), where the double bond between sp^2 hybridized carbon atoms makes all of the bond angles $\sim 120^\circ$ and puts all six atoms in the same plane.




These six atoms lie in a plane.

The structure of a tetrapeptide illustrates the results of these effects in a long peptide chain.

- The *s-trans* arrangement makes a long chain with a zigzag arrangement.
- In each peptide bond, the N–H and C=O bonds lie parallel and at 180° with respect to each other.

A tetrapeptide



Problem 28.16 Draw the *s-cis* and *s-trans* conformations for the dipeptide formed from two glycine molecules.


28.5C Interesting Peptides

Even relatively simple peptides can have important biological functions. **Bradykinin**, for example, is a peptide hormone composed of nine amino acids. It stimulates smooth muscle contraction, dilates blood vessels, and causes pain. Bradykinin is a component of bee venom.



bradykinin

Oxytocin and **vasopressin** are nonapeptide hormones, too. Their sequences are identical except for two amino acids, yet this is enough to give them very different biological activities. Oxytocin induces labor by stimulating the contraction of uterine muscles, and it stimulates the flow of milk in nursing mothers. Vasopressin, on the other hand, controls blood pressure by regulating smooth muscle contraction. The N-terminal amino acid in both hormones is a cysteine residue, and the C-terminal residue is glycine. Instead of a free carboxy group, both peptides have an NH_2 group in place of OH, so this is indicated with the additional NH_2 group drawn at the end of the chain.



The structure of both peptides includes a **disulfide bond**, a form of covalent bonding in which the $-\text{SH}$ groups from two cysteine residues are oxidized to form a sulfur–sulfur bond. In oxytocin and vasopressin, the disulfide bonds make the peptides cyclic. Three-dimensional structures of oxytocin and vasopressin are shown in Figure 28.7.






Figure 28.7

Three-dimensional structures of vasopressin and oxytocin




The artificial sweetener **aspartame** (Figure 27.11) is the methyl ester of the dipeptide Asp–Phe. This synthetic peptide is 180 times sweeter (on a gram-for-gram basis) than sucrose (common table sugar). Both of the amino acids in aspartame have the naturally occurring L-configuration. If the D-amino acid is substituted for either Asp or Phe, the resulting compound tastes bitter.



aspartame
the methyl ester of Asp–Phe
a synthetic artificial sweetener

Problem 28.17 Draw the structure of leu-enkephalin, a pentapeptide that acts as an analgesic and opiate, and has the following sequence: Tyr–Gly–Gly–Phe–Leu. (The structure of a related peptide, met-enkephalin, appeared in Section 22.6B.)

Problem 28.18 Glutathione, a powerful antioxidant that destroys harmful oxidizing agents in cells, is composed of glutamic acid, cysteine, and glycine, and has the following structure:



glutathione

- What product is formed when glutathione reacts with an oxidizing agent?
- What is unusual about the peptide bond between glutamic acid and cysteine?

28.6 Peptide Sequencing

To determine the structure of a peptide, we must know not only what amino acids comprise it, but also the sequence of the amino acids in the peptide chain. Although mass spectrometry has become an increasingly powerful method for the analysis of high molecular weight proteins (Section 13.4), chemical methods to determine peptide structure are still widely used and presented in this section.

28.6A Amino Acid Analysis


The structure determination of a peptide begins by analyzing the total amino acid composition. The amide bonds are first hydrolyzed by heating with hydrochloric acid for 24 h to form the individual amino acids. The resulting mixture is then separated using high-performance liquid chromatography (HPLC), a technique in which a solution of amino acids is placed on a column and individual amino acids move through the column at characteristic rates, often dependent on polarity.

This process determines both the identity of the individual amino acids and the amount of each present, but it tells nothing about the order of the amino acids in the peptide. For example, complete hydrolysis and HPLC analysis of the tetrapeptide Gly–Gly–Phe–Tyr would indicate the presence of three amino acids—glycine, phenylalanine, and tyrosine—and show that there are twice as many glycine residues as phenylalanine or tyrosine residues. The exact order of the amino acids in the peptide chain must then be determined by additional methods

28.6B Identifying the N-Terminal Amino Acid—The Edman Degradation

To determine the sequence of amino acids in a peptide chain, a variety of procedures are often combined. One especially useful technique is to identify the N-terminal amino acid using the **Edman degradation**. In the Edman degradation, amino acids are cleaved one at a time from the N-terminal end, the identity of the amino acid determined, and the process repeated until the entire sequence is known. Automated sequencers using this methodology are now available to sequence peptides containing up to about 50 amino acids.

The Edman degradation is based on the reaction of the nucleophilic NH_2 group of the N-terminal amino acid with the electrophilic carbon of phenyl isothiocyanate, $\text{C}_6\text{H}_5\text{N}=\text{C}=\text{S}$. When the N-terminal amino acid is removed from the peptide chain, two products are formed: **an N-phenylthiohydantoin (PTH) and a new peptide with one fewer amino acid.**




The *N*-phenylthiohydantoin derivative contains the atoms of the N-terminal amino acid. This product identifies the N-terminal amino acid in the peptide because the PTH derivatives of all 20 naturally occurring amino acids are known and characterized. The new peptide formed in the Edman degradation has one amino acid fewer than the original peptide. Moreover, it contains a new N-terminal amino acid, so the process can be repeated.

Mechanism 28.2 illustrates some of the key steps of the Edman degradation. The nucleophilic N-terminal NH_2 group adds to the electrophilic carbon of phenyl isothiocyanate to form an *N*-phenylthiourea, the product of nucleophilic addition (Part [1]). Intramolecular cyclization followed by elimination results in cleavage of the terminal amide bond in Part [2] to form a new peptide with one fewer amino acid. A sulfur heterocycle, called a thiazolinone, is also formed, which rearranges by a multistep pathway (Part [3]) to form an *N*-phenylthiohydantoin. The R group in this product identifies the amino acid located at the N-terminal end.




Mechanism 28.2 Edman Degradation

Part [1] Formation of an *N*-phenylthiourea




- Addition of the free amino group from the *N*-terminal amino acid to the electrophilic carbon of phenyl isothiocyanate followed by proton transfer forms an *N*-phenylthiourea.

Part [2] Cleavage of the amide bond from the *N*-terminal amino acid



- Nucleophilic addition in Step [3] followed by loss of the amino group in Step [4] forms two products: a five-membered thiazolinone ring and a peptide chain that contains one fewer amino acid than the original peptide.

Part [3] Rearrangement to form an *N*-phenylthiohydantoin (PTH)



- Under the conditions of the reaction, the thiazolinone rearranges by a multistep pathway to form an *N*-phenylthiohydantoin (PTH). This product contains the original *N*-terminal amino acid.

In theory a protein of any length can be sequenced using the Edman degradation, but in practice, the accumulation of small quantities of unwanted by-products limits sequencing to proteins having fewer than approximately 50 amino acids.

Problem 28.19 Draw the structure of the *N*-phenylthiohydantoin formed by initial Edman degradation of each peptide: (a) Ala–Gly–Phe–Phe; (b) Val–Ile–Tyr.

28.6C Partial Hydrolysis of a Peptide

Additional structural information can be obtained by cleaving some, but not all, of the amide bonds in a peptide. Partial hydrolysis of a peptide with acid forms smaller fragments in a random fashion. Sequencing these peptides and identifying sites of overlap can be used to determine the sequence of the complete peptide, as shown in Sample Problem 28.2.

Sample Problem 28.3

Deduce the sequence of a pentapeptide that contains the amino acids Ala, Glu, Gly, Ser, and Tyr, from the following experimental data. Edman degradation cleaves Gly from the pentapeptide, and carboxypeptidase forms Ala and a tetrapeptide. Treatment of the pentapeptide with chymotrypsin forms a dipeptide and a tripeptide. Partial hydrolysis forms Gly, Ser, and the tripeptide Tyr–Glu–Ala.

Solution

Use each result to determine the location of an amino acid in the pentapeptide.


Experiment	Result
• Edman degradation identifies the N-terminal amino acid—in this case, Gly.	→ Gly- _ - _ - _ - _
• Carboxypeptidase identifies the C-terminal amino acid (Ala) when it is cleaved from the end of the chain.	→ Gly- _ - _ - _ -Ala
• Chymotrypsin cleaves amide bonds that contain a carbonyl group from an aromatic amino acid—Tyr in this case. Because a dipeptide and tripeptide are obtained after treatment with chymotrypsin, Tyr must be the C-terminal amino acid of either the di- or tripeptide. As a result, Tyr must be either the second or third amino acid in the pentapeptide chain.	→ Gly-Tyr- _ - _ -Ala or Gly- _ -Tyr- _ -Ala
• Partial hydrolysis forms the tripeptide Tyr–Glu–Ala. Because Ala is the C-terminal amino acid, this result identifies the last three amino acids in the chain.	→ Gly- _ -Tyr–Glu–Ala
• The last amino acid, Ser, must be located at the only remaining position, the second amino acid in the pentapeptide, and the complete sequence is determined.	→ Gly–Ser–Tyr–Glu–Ala

Problem 28.22

Deduce the sequence of a heptapeptide that contains the amino acids Ala, Arg, Glu, Gly, Leu, Phe, and Ser, from the following experimental data. Edman degradation cleaves Leu from the heptapeptide, and carboxypeptidase forms Glu and a hexapeptide. Treatment of the heptapeptide with chymotrypsin forms a hexapeptide and a single amino acid. Treatment of the heptapeptide with trypsin forms a pentapeptide and a dipeptide. Partial hydrolysis forms Glu, Leu, Phe, and the tripeptides Gly–Ala–Ser and Ala–Ser–Arg.

28.7 Peptide Synthesis

The synthesis of a specific dipeptide, such as Ala–Gly from alanine and glycine, is complicated because both amino acids have two functional groups. As a result, four products—namely, Ala–Ala, Ala–Gly, Gly–Gly, and Gly–Ala—are possible.




How do we selectively join the COOH group of alanine with the NH₂ group of glycine?


- Protect the functional groups that we don't want to react, and then form the amide bond.

HOW TO Synthesize a Dipeptide from Two Amino Acids


Example




Step [1] Protect the NH₂ group of alanine.




Step [2] Protect the COOH group of glycine.




Step [3] Form the amide bond with DCC.




Dicyclohexylcarbodiimide (**DCC**) is a reagent commonly used to form amide bonds (see Section 22.10D). DCC makes the OH group of the carboxylic acid a better leaving group, thus **activating the carboxy group toward nucleophilic attack**.



Step [4] Remove one or both protecting groups.





Two widely used amino protecting groups convert an amine into a **carbamate**, a functional group having a carbonyl bonded to both an oxygen and a nitrogen atom. Since the N atom of the carbamate is bonded to a carbonyl group, the protected amino group is no longer nucleophilic.





For example, the ***tert*-butoxycarbonyl protecting group**, abbreviated as **Boc**, is formed by reacting the amino acid with di-*tert*-butyl dicarbonate in a nucleophilic acyl substitution reaction.

Adding a Boc protecting group




To be a useful protecting group, the Boc group must be removed under reaction conditions that do not affect other functional groups in the molecule. It can be removed with an acid such as **trifluoroacetic acid, HCl, or HBr**.

Removing a Boc protecting group




A second amino protecting group, the **9-fluorenylmethoxycarbonyl protecting group**, abbreviated as **Fmoc**, is formed by reacting the amino acid with 9-fluorenylmethyl chloroformate in a nucleophilic acyl substitution reaction.

Adding an Fmoc protecting group




While the Fmoc protecting group is stable to most acids, it can be removed by treatment with base (NH_3 or an amine), to regenerate the free amino group.


Removing an Fmoc protecting group




The carboxy group is usually protected as a **methyl** or **benzyl ester** by reaction with an alcohol and an acid.



These esters are usually removed by hydrolysis with aqueous base.




One advantage of using a benzyl ester for protection is that it can also be removed with H_2 in the presence of a Pd catalyst. This process is called **hydrogenolysis**. These conditions are especially mild, because they avoid the use of either acid or base. Benzyl esters can also be removed with HBr in acetic acid.



The specific reactions needed to synthesize the dipeptide Ala-Gly are illustrated in Sample Problem 28.4.


Sample Problem 28.4

Draw out the steps in the synthesis of the dipeptide Ala-Gly.




Solution


Step [1] Protect the NH_2 group of alanine using a Boc group.



Step [2] Protect the COOH group of glycine as a benzyl ester.




Step [3] Form the amide bond with DCC.




Step [4] Remove one or both protecting groups.

The protecting groups can be removed in a stepwise fashion, or in a single reaction.



This method can be applied to the synthesis of tripeptides and even larger polypeptides. After the protected dipeptide is prepared in Step [3], only one of the protecting groups is removed, and this dipeptide is coupled to a third amino acid with one of its functional groups protected, as illustrated in the following equations.



Problem 28.23

Devise a synthesis of each peptide from amino acid starting materials: (a) Leu-Val; (b) Ala-Ile-Gly; (c) Ala-Gly-Ala-Gly.


28.8 Automated Peptide Synthesis

The method described in Section 28.7 works well for the synthesis of small peptides. It is extremely time-consuming to synthesize larger proteins by this strategy, however, because each step requires isolation and purification of the product. The synthesis of larger polypeptides is usually accomplished by using the **solid phase technique** originally developed by R. Bruce Merrifield of Rockefeller University.


Development of the solid phase technique earned Merrifield the 1984 Nobel Prize in Chemistry and has made possible the synthesis of many polypeptides and proteins.

In the **Merrifield method** an amino acid is attached to an **insoluble polymer**. Amino acids are sequentially added, one at a time, thereby forming successive peptide bonds. Because impurities and by-products are not attached to the polymer chain, they are removed simply by washing them away with a solvent at each stage of the synthesis.

A commonly used polymer is a **polystyrene derivative** that contains $-\text{CH}_2\text{Cl}$ groups bonded to some of the benzene rings in the polymer chain. The Cl atoms serve as handles that allow attachment of amino acids to the chain.




An Fmoc-protected amino acid is attached to the polymer at its carboxy group by an $\text{S}_{\text{N}}2$ reaction.



Once the first amino acid is bound to the polymer, additional amino acids can be added sequentially. The steps of the solid phase peptide synthesis technique are illustrated in the accompanying scheme. In the last step, HF cleaves the polypeptide chain from the polymer.

HOW TO Synthesize a Peptide Using the Merrifield Solid Phase Technique

Step [1]
Attach an Fmoc-protected amino acid to the polymer.




—Continued


HOW TO, continued . . .

Step [2]


Remove the protecting group.

**Step [3]**


Form the amide bond with DCC.

**Step [4]**

Repeat Steps [2] and [3].

**Step [5]**

Remove the protecting group and detach the peptide from the polymer.



The Merrifield method has now been completely automated, so it is possible to purchase peptide synthesizers that automatically carry out all of the above operations and form polypeptides in high yield in a matter of hours, days, or weeks, depending on the length of the chain of the desired product. The instrument is pictured in Figure 28.8. For example, the protein ribonuclease, which contains 128 amino acids, has been prepared by this technique in an overall yield of 17%. This remarkable synthesis involved 369 separate reactions, and thus the yield of each individual reaction was > 99%.

Problem 28.24 Outline the steps needed to synthesize the tetrapeptide Ala–Leu–Ile–Gly using the Merrifield technique.

Figure 28.8
Automated peptide synthesizer




28.9 Protein Structure

Now that you have learned some of the chemistry of amino acids, it's time to study proteins, the large polymers of amino acids that are responsible for so much of the structure and function of all living cells. We begin with a discussion of the **primary, secondary, tertiary, and quaternary structure** of proteins.

28.9A Primary Structure

The **primary structure** of proteins is the particular sequence of amino acids that is joined together by **peptide bonds**. The most important element of this primary structure is the **amide bond**.


- Rotation around the amide C–N bond is *restricted* because of electron delocalization, and the *s-trans* conformation is the more stable arrangement.
- In each peptide bond, the N–H and C=O bonds are directed 180° from each other.



Although rotation about the amide bonds is restricted, **rotation about the other σ bonds in the protein backbone is not**. As a result, the peptide chain can twist and bend into a variety of different arrangements that constitute the secondary structure of the protein.

28.9B Secondary Structure

The **three-dimensional conformations of localized regions of a protein are called its secondary structure**. These regions arise due to hydrogen bonding between the N–H proton of one amide and C=O oxygen of another. Two arrangements that are particularly stable are called the **α -helix** and the **β -pleated sheet**.



α -Helix

The α -helix forms when a peptide chain twists into a right-handed or clockwise spiral, as shown in Figure 28.9. Four important features of the α -helix are as follows:

- [1] **Each turn of the helix has 3.6 amino acids.**
- [2] **The N–H and C=O bonds point along the axis of the helix.** All C=O bonds point in one direction, and all N–H bonds point in the opposite direction.
- [3] **The C=O group of one amino acid is hydrogen bonded to an N–H group four amino acid residues farther along the chain.** Thus, hydrogen bonding occurs between two amino acids *in the same chain*. Note, too, that the hydrogen bonds are parallel to the axis of the helix.
- [4] **The R groups of the amino acids extend outward** from the core of the helix.

An α -helix can form only if there is rotation about the bonds at the α carbon of the amide carbonyl group, and not all amino acids can do this. For example, proline, the amino acid whose nitrogen atom forms part of a five-membered ring, is more rigid than other amino acids, and its C_{α} –N bond cannot rotate the necessary amount. Additionally, it has no N–H proton with which to form an intramolecular hydrogen bond to stabilize the helix. Thus, proline cannot be part of an α -helix.

Both the myosin in muscle and α -keratin in hair are proteins composed almost entirely of α -helices.

β -Pleated Sheet


The β -pleated sheet secondary structure forms when two or more peptide chains, called **strands**, line up side-by-side, as shown in Figure 28.10. All β -pleated sheets have the following characteristics:

- [1] **The C=O and N–H bonds lie in the plane of the sheet.**
- [2] **Hydrogen bonding often occurs between the N–H and C=O groups of nearby amino acid residues.**

Figure 28.9


Two different illustrations of the α -helix

a. The right-handed α -helix




All atoms of the α -helix are drawn in this representation. All C=O bonds are pointing up and all N–H bonds are pointing down.

b. The backbone of the α -helix



Only the peptide backbone is drawn in this representation. The hydrogen bonds between the C=O and N–H of amino acids four residues away from each other are shown.

Figure 28.10
Three-dimensional structure
of the β -pleated sheet



- The β -pleated sheet consists of extended strands of the peptide chains held together by hydrogen bonding. The C=O and N-H bonds lie in the plane of the sheet, and the R groups (shown as orange balls) alternate above and below the plane.

[3] The **R groups are oriented above and below the plane** of the sheet, and alternate from one side to the other along a given strand.


The β -pleated sheet arrangement most commonly occurs with amino acids with small R groups, like alanine and glycine. With larger R groups steric interactions prevent the chains from getting close together and so the sheet cannot be stabilized by hydrogen bonding.

The peptide strands of β -pleated sheets can actually be oriented in two different ways, as shown in Figure 28.11.

- In a *parallel* β -pleated sheet, the strands run in the *same* direction from the N- to C-terminal amino acid.
- In an *antiparallel* β -pleated sheet, the strands run in the *opposite* direction.

Most proteins have regions of α -helix and β -pleated sheet, in addition to other regions that cannot be characterized by either of these arrangements. Shorthand symbols are often used to indicate regions of a protein that have α -helix or β -pleated sheet. A **flat helical ribbon** is used for

Figure 28.11 The parallel and antiparallel forms of the β -pleated sheet



The two peptide chains are arranged in the same direction. Hydrogen bonds occur between N-H and C=O bonds in adjacent chains.


[Note: R groups on the carbon chain are omitted for clarity.]

The two peptide chains are arranged in opposite directions. Hydrogen bonding between the N-H and C=O groups still holds the two chains together.

the α -helix, and a **flat wide arrow** is used for the β -pleated sheet. These representations are often used in **ribbon diagrams** to illustrate protein structure.



α -helix shorthand



β -pleated sheet shorthand

Proteins are drawn in a variety of ways to illustrate different aspects of their structure. Figure 28.12 illustrates three different representations of the protein lysozyme, an enzyme found in both plants and animals. Lysozyme catalyzes the hydrolysis of bonds in bacterial cell walls, weakening them, often causing the bacteria to burst.

Spider dragline silk is a strong yet elastic protein because it has regions of β -pleated sheet and regions of α -helix (Figure 28.13). α -Helical regions impart elasticity to the silk because the peptide chain is twisted (not fully extended), so it can stretch. β -Pleated sheet regions are almost fully extended, so they can't be stretched further, but their highly ordered three-dimensional structure imparts strength to the silk. Thus, spider silk suits the spider by comprising both types of secondary structure with beneficial properties.

Problem 28.25 Suggest a reason why antiparallel β -pleated sheets are generally more stable than parallel β -pleated sheets.


Problem 28.26 Consider two molecules of a tetrapeptide composed of only alanine residues. Draw the hydrogen bonding interactions that result when these two peptides adopt a parallel β -pleated sheet arrangement. Answer this same question for the antiparallel β -pleated sheet arrangement.

28.9C Tertiary and Quaternary Structure


The three-dimensional shape adopted by the entire peptide chain is called its tertiary structure. A peptide generally folds into a conformation that maximizes its stability. In the aqueous environment of the cell, proteins often fold in such a way as to place a large number of polar and charged groups on their outer surface, to maximize the dipole–dipole and hydrogen bonding interactions with water. This generally places most of the nonpolar side chains in the interior of the protein, where van der Waals interactions between these hydrophobic groups help stabilize the molecule, too.

Figure 28.12 Lysozyme


a. Ball-and-stick model



b. Space-filling model




c. Ribbon diagram



(a) The ball-and-stick model of lysozyme shows the protein backbone with color-coded C, N, O, and S atoms. Individual amino acids are most clearly located using this representation. (b) The space-filling model uses color-coded balls for each atom in the backbone of the enzyme and illustrates how the atoms fill the space they occupy. (c) The ribbon diagram shows regions of α -helix and β -sheet that are not clearly in evidence in the other two representations.

Figure 28.13
Different regions of secondary structure in spider silk




Spider silk has regions of α -helix and β -pleated sheet that make it both strong and elastic. The green coils represent the α -helical regions, and the purple arrows represent the β -pleated sheet regions. The yellow lines represent other areas of the protein that are neither α -helix nor β -pleated sheet.

In addition, polar functional groups hydrogen bond with each other (not just water), and amino acids with charged side chains like $-\text{COO}^-$ and $-\text{NH}_3^+$ can stabilize tertiary structure by electrostatic interactions.


Finally, **disulfide bonds are the only covalent bonds that stabilize tertiary structure.** As previously mentioned, these strong bonds form by oxidation of two cysteine residues either on the same polypeptide chain or another polypeptide chain of the same protein.

Disulfide bonds can form in two different ways.

Between two SH groups on the same chain.



Between two SH groups on different chains.



The nonapeptides **oxytocin** and **vasopressin** (Section 28.5C) contain intramolecular disulfide bonds. **Insulin**, on the other hand, consists of two separate polypeptide chains (**A** and **B**) that are covalently linked by two intermolecular disulfide bonds, as shown in Figure 28.14. The **A** chain, which also has an intramolecular disulfide bond, has 21 amino acid residues, whereas the **B** chain has 30.


Figure 28.15 schematically illustrates the many different kinds of intramolecular forces that stabilize the secondary and tertiary structures of polypeptide chains.

The shape adopted when two or more folded polypeptide chains aggregate into one protein complex is called the **quaternary structure** of the protein. Each individual polypeptide chain is called a **subunit** of the overall protein. **Hemoglobin**, for example, consists of two α and two β subunits held together by intermolecular forces in a compact three-dimensional shape. The unique function of hemoglobin is possible only when all four subunits are together.


The four levels of protein structure are summarized in Figure 28.16.

Figure 28.14

Insulin



Insulin is a small protein consisting of two polypeptide chains (designated as the **A** and **B** chains) held together by two disulfide bonds. An additional disulfide bond joins two cysteine residues within the **A** chain.



Synthesized by groups of cells in the pancreas called the islets of Langerhans, insulin is the protein that regulates the levels of glucose in the blood. Insufficiency of insulin results in diabetes. Many of the abnormalities associated with this disease can be controlled by the injection of insulin. Until the availability of human insulin through genetic engineering techniques, all insulin used by diabetics was obtained from pigs and cattle. The amino acid sequences of these insulin proteins is slightly different from that of human insulin. Pig insulin differs in one amino acid only, whereas bovine insulin has three different amino acids. This is shown in the accompanying table.

Position of residue →	Chain A			Chain B
	8	9	10	30
Human insulin	Thr	Ser	Ile	Thr
Pig insulin	Thr	Ser	Ile	Ala
Bovine insulin	Ala	Ser	Val	Ala

Problem 28.27 What types of stabilizing interactions exist between each of the following pairs of amino acids?
 a. Ser and Tyr b. Val and Leu c. Two Phe residues

Problem 28.28 The fibroin proteins found in silk fibers consist of large regions of β -pleated sheets stacked one on top of another. (a) Explain why having a glycine at every other residue allows the β -pleated sheets to stack on top of each other. (b) Why are silk fibers insoluble in water?


28.10 Important Proteins

Proteins are generally classified according to their three-dimensional shapes.

- **Fibrous proteins** are composed of long linear polypeptide chains that are bundled together to form rods or sheets. These proteins are insoluble in water and serve structural roles, giving strength and protection to tissues and cells.
- **Globular proteins** are coiled into compact shapes with hydrophilic outer surfaces that make them water soluble. Enzymes and transport proteins are globular to make them soluble in the blood and other aqueous environments in cells.

Figure 28.17

Anatomy of a hair—
It begins with α -keratin.



a **supercoil** or **superhelix**. These, in turn, form larger and larger bundles of fibers, ultimately forming a strand of hair, as shown schematically in Figure 28.17.

α -Keratins also have a number of cysteine residues, and because of this, disulfide bonds are formed between adjacent helices. The number of disulfide bridges determines the strength of the material. Claws, horns, and fingernails have extensive networks of disulfide bonds, making them extremely hard.


Straight hair can be made curly by cleaving the disulfide bonds in α -keratin, and then rearranging and re-forming them, as shown schematically in Figure 28.18. First, the disulfide bonds in the straight hair are reduced to thiol groups, so the bundles of α -keratin chains are no longer held in their specific “straight” orientation. Then, the hair is wrapped around curlers and treated with an oxidizing agent that converts the thiol groups back to disulfide bonds, now with twists and turns in the keratin backbone. This makes the hair look curly and is the chemical basis for a “permanent.”

28.10B Collagen

Collagen, the most abundant protein in vertebrates, is found in connective tissues such as bone, cartilage, tendons, teeth, and blood vessels. Glycine and proline account for a large fraction of its amino acid residues, whereas cysteine accounts for very little. Because of the high proline content, it cannot form a right-handed α -helix. Instead, it forms an elongated left-handed helix, and then three of these helices wind around each other to form a right-handed **superhelix** or **triple helix**. The side chain of glycine is only a hydrogen atom, so the high glycine content allows the

Figure 28.18


The chemistry of a
“permanent”—Making
straight hair curly



To make straight hair curly, the disulfide bonds holding the α -helical chains together are cleaved by reduction. This forms free thiol groups ($-\text{SH}$). The hair is turned around curlers and then an oxidizing agent is applied. This re-forms the disulfide bonds in the hair, but between different thiol groups, now giving it a curly appearance.

Figure 28.19

Two different representations for the triple helix of collagen



- In collagen, three polypeptide chains having an unusual left-handed helix wind around each other in a right-handed triple helix. The high content of small glycine residues allows the chains to lie close to each other, permitting hydrogen bonding between the chains.

collagen superhelices to lie compactly next to each other, thus stabilizing the superhelices via hydrogen bonding. Two views of the collagen superhelix are shown in Figure 28.19.


28.10C Hemoglobin and Myoglobin

Hemoglobin and **myoglobin**, two globular proteins, are called **conjugated proteins** because they are composed of a protein unit and a nonprotein molecule called a **prosthetic group**. The prosthetic group in hemoglobin and myoglobin is **heme**, a complex organic compound containing the Fe^{2+} ion complexed with a nitrogen heterocycle called a **porphyrin**. The Fe^{2+} ion of hemoglobin and myoglobin binds oxygen in the blood. Hemoglobin, which is present in red blood cells, transports oxygen to wherever it is needed in the body, whereas myoglobin stores oxygen in tissues. Ribbon diagrams for myoglobin and hemoglobin are shown in Figure 28.20.

Figure 28.20


Protein ribbon diagrams for myoglobin and hemoglobin

a. Myoglobin




Myoglobin consists of a single polypeptide chain with a heme unit shown in a ball-and-stick model.

b. Hemoglobin



Hemoglobin consists of two α and two β chains shown in red and blue, respectively, and four heme units shown in ball-and-stick models.




Myoglobin, the chapter-opening molecule, has 153 amino acid residues in a single polypeptide chain. It has eight separate α -helical sections that fold back on one another, with the prosthetic heme group held in a cavity inside the polypeptide. Most of the polar residues are found on the outside of the protein so that they can interact with the water solvent. Spaces in the interior of the protein are filled with nonpolar amino acids. Myoglobin gives cardiac muscle its characteristic red color.

Hemoglobin consists of four polypeptide chains (two α subunits and two β subunits), each of which carries a heme unit. Hemoglobin has more nonpolar amino acids than myoglobin. When each subunit is folded, some of these remain on the surface. The van der Waals attraction between these hydrophobic groups is what stabilizes the quaternary structure of the four subunits.

Carbon monoxide is poisonous because it binds to the Fe^{2+} of hemoglobin more strongly than does oxygen. Hemoglobin complexed with CO cannot carry O_2 from the lungs to the tissues. Without O_2 in the tissues for metabolism, cells cannot function, so they die.

The properties of all proteins depend on their three-dimensional shape, and their shape depends on their primary structure—that is, their amino acid sequence. This is particularly well exemplified by comparing normal hemoglobin with **sickle cell hemoglobin**, a mutant variation in which a single amino acid of both β subunits is changed from glutamic acid to valine. The replacement of one acidic amino acid (Glu) with one nonpolar amino acid (Val) changes the shape of hemoglobin, which has profound effects on its function. Deoxygenated red blood cells with sickle cell hemoglobin become elongated and crescent shaped, and they are unusually fragile. As a result, they do not flow easily through capillaries, causing pain and inflammation, and they break open easily, leading to severe anemia and organ damage. The end result is often a painful and premature death.

This disease, called **sickle cell anemia**, is found almost exclusively among people originating from central and western Africa, where malaria is an enormous health problem. Sickle cell hemoglobin results from a genetic mutation in the DNA sequence that is responsible for the synthesis of hemoglobin. Individuals who inherit this mutation from both parents develop sickle cell anemia, whereas those who inherit it from only one parent are said to have the sickle cell trait. They do not develop sickle cell anemia and they are more resistant to malaria than individuals without the mutation. This apparently accounts for this detrimental gene being passed on from generation to generation.




When red blood cells take on a “sickled” shape in persons with sickle cell disease, they occlude capillaries (causing organ injury) and they break easily (leading to profound anemia). This devastating illness results from the change of a single amino acid in hemoglobin. Note the single sickled cell surrounded by three red cells with normal morphology.

KEY CONCEPTS


Amino Acids and Proteins

Synthesis of Amino Acids (28.2)

[1] From α -halo carboxylic acids by S_N2 reaction




[2] By alkylation of diethyl acetamidomalonate



- Alkylation works best with unhindered alkyl halides—that is, CH_3X and RCH_2X .

[3] Strecker synthesis




Preparation of Optically Active Amino Acids


[1] Resolution of enantiomers by forming diastereomers (28.3A)

- Convert a racemic mixture of amino acids into a racemic mixture of *N*-acetyl amino acids [(*S*)- and (*R*)- $\text{CH}_3\text{CONHCH(R)COOH}$].
- Treat the enantiomers with a chiral amine to form a mixture of diastereomers.
- Separate the diastereomers.
- Regenerate the amino acids by protonation of the carboxylate salt and hydrolysis of the *N*-acetyl group.

[2] Kinetic resolution using enzymes (28.3B)



[3] By enantioselective hydrogenation (28.4)




Summary of Methods Used for Peptide Sequencing (28.6)


- Complete hydrolysis of all amide bonds in a peptide gives the identity and amount of the individual amino acids.
- Edman degradation identifies the N-terminal amino acid. Repeated Edman degradations can be used to sequence a peptide from the N-terminal end.
- Cleavage with carboxypeptidase identifies the C-terminal amino acid.
- Partial hydrolysis of a peptide forms smaller fragments that can be sequenced. Amino acid sequences common to smaller fragments can be used to determine the sequence of the complete peptide.
- Selective cleavage of a peptide occurs with trypsin and chymotrypsin to identify the location of specific amino acids (Table 28.2).

Adding and Removing Protecting Groups for Amino Acids (28.7)


[1] Protection of an amino group as a Boc derivative




[2] Deprotection of a Boc-protected amino acid




[3] Protection of an amino group as an Fmoc derivative




[4] Deprotection of an Fmoc-protected amino acid



[5] Protection of a carboxy group as an ester




[6] Deprotection of an ester group



Synthesis of Dipeptides (28.7)

[1] Amide formation with DCC



[2] Four steps are needed to synthesize a dipeptide:

- Protect** the amino group of one amino acid with a Boc or Fmoc group.
- Protect** the carboxy group of the second amino acid as an ester.
- Form the amide bond with **DCC**.
- Remove both protecting groups** in one or two reactions.

Summary of the Merrifield Method of Peptide Synthesis (28.8)


- Attach an Fmoc-protected amino acid to a polymer derived from polystyrene.
- Remove the Fmoc protecting group.
- Form the amide bond with a second Fmoc-protected amino acid by using DCC.
- Repeat steps [2] and [3].
- Remove the protecting group and detach the peptide from the polymer.

PROBLEMS

Amino Acids

28.29 Explain why L-alanine has the *S* configuration but L-cysteine has the *R* configuration.

28.30



- (*S*)-Penicillamine, an amino acid that does not occur in proteins, is used as a copper chelating agent to treat Wilson's disease, an inherited defect in copper metabolism. (*R*)-Penicillamine is toxic, sometimes causing blindness. Draw the structures of (*R*)- and (*S*)-penicillamine.
- What disulfide is formed from oxidation of L-penicillamine?

28.31 Explain why amino acids are insoluble in diethyl ether but *N*-acetyl amino acids are soluble.

28.32 Histidine is classified as a basic amino acid because one of the N atoms in its five-membered ring is readily protonated by acid. Which N atom in histidine is protonated and why?

28.33 Tryptophan is not classified as a basic amino acid even though it has a heterocycle containing a nitrogen atom. Why is the N atom in the five-membered ring of tryptophan not readily protonated by acid?

28.34 What is the structure of each amino acid at its isoelectric point: (a) alanine; (b) methionine; (c) aspartic acid; (d) lysine?

28.35 To calculate the isoelectric point of amino acids having other ionizable functional groups, we must also take into account the pK_a of the additional functional group in the side chain.

For an acidic amino acid (one with an additional acidic OH group):

$$pI = \frac{pK_a(\alpha\text{-COOH}) + pK_a(\text{second COOH})}{2}$$

For a basic amino acid (one with an additional basic NH group):

$$pI = \frac{pK_a(\alpha\text{-NH}_3^+) + pK_a(\text{side chain NH})}{2}$$

- Indicate which pK_a values must be used to calculate the *pI* of each of the following amino acids: [1] glutamic acid; [2] lysine; [3] arginine.
 - In general, how does the *pI* of an acidic amino acid compare to that of a neutral amino acid?
 - In general, how does the *pI* of a basic amino acid compare to the *pI* of a neutral amino acid?
- 28.36** What is the predominant form of each of the following amino acids at pH = 1? What is the overall charge on the amino acid at this pH? (a) threonine; (b) methionine; (c) aspartic acid; (d) arginine
- 28.37** What is the predominant form of each of the following amino acids at pH = 11? What is the overall charge on the amino acid? (a) valine; (b) proline; (c) glutamic acid; (d) lysine

- 28.38 a. Draw the structure of the tripeptide A–A–A, and label the two ionizable functional groups.
 b. What is the predominant form of A–A–A at pH = 1?
 c. The pK_a values for the two ionizable functional groups (3.39 and 8.03) differ considerably from the pK_a values of alanine (2.35 and 9.87; see Table 28.1). Account for the observed pK_a differences.


Synthesis and Reactions of Amino Acids

28.39 Draw the organic product formed when the amino acid leucine is treated with each reagent.

- a. CH_3OH , H^+
 b. CH_3COCl , pyridine
 c. $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$, H^+
 d. Ac_2O , pyridine
 e. HCl (1 equiv)
 f. NaOH (1 equiv)
 g. $\text{C}_6\text{H}_5\text{COCl}$, pyridine
 h. $[(\text{CH}_3)_3\text{COCO}]_2\text{O}$, $(\text{CH}_3\text{CH}_2)_3\text{N}$
 i. The product in (d), then $\text{NH}_2\text{CH}_2\text{COOCH}_3 + \text{DCC}$
 j. The product in (h), then $\text{NH}_2\text{CH}_2\text{COOCH}_3 + \text{DCC}$
 k. Fmoc-Cl , Na_2CO_3 , H_2O
 l. $\text{C}_6\text{H}_5\text{N}=\text{C}=\text{S}$

28.40 Answer Problem 28.39 using phenylalanine as a starting material.

28.41 Draw the organic products formed in each reaction.




28.42 What alkyl halide is needed to synthesize each amino acid from diethyl acetamidomalonate: (a) Asn; (b) His; (c) Trp?


28.43 Devise a synthesis of threonine from diethyl acetamidomalonate.

28.44 Devise a synthesis of each amino acid from acetaldehyde (CH_3CHO): (a) glycine; (b) alanine.

28.45 Identify the lettered intermediates in the following reaction scheme. This is an alternative method to synthesize amino acids, based on the Gabriel synthesis of 1° amines (Section 25.7A).




28.46 Glutamic acid is synthesized by the following reaction sequence. Draw a stepwise mechanism for Steps [1]–[3].




Resolution; The Synthesis of Chiral Amino Acids

28.47 Write out a scheme for the resolution of the two enantiomers of racemic lactic acid [$\text{CH}_3\text{CH}(\text{OH})\text{COOH}$] using (*R*)- α -methylbenzylamine as resolving agent.




- 28.48** Another strategy used to resolve amino acids involves converting the carboxy group to an ester and then using a *chiral carboxylic acid* to carry out an acid–base reaction at the free amino group. The general plan is drawn below using (*R*)-mandelic acid as resolving agent. Using a racemic mixture of alanine enantiomers and (*R*)-mandelic acid as resolving agent, write out the steps showing how a resolution process would occur.




- 28.49** Brucine is a poisonous alkaloid obtained from *Strychnos nux vomica*, a tree that grows in India, Sri Lanka, and northern Australia. Write out a resolution scheme similar to the one given in Section 28.3A, which shows how a racemic mixture of phenylalanine can be resolved using brucine.



- 28.50** Draw the organic products formed in each reaction.



- 28.51** What two steps are needed to convert **A** to L-dopa, an uncommon amino acid that is effective in treating Parkinson's disease? These two steps are the key reactions in the first commercial asymmetric synthesis using a chiral transition metal catalyst. This process was developed at Monsanto in 1974.



Peptide Structure and Sequencing


- 28.52** Draw the structure for each peptide: (a) Phe–Ala; (b) Gly–Gln; (c) Lys–Gly; (d) R–H.

- 28.53** For each tetrapeptide [1] Ala–Gln–Cys–Ser; [2] Asp–Arg–Val–Tyr:

- a. Name the peptide using one-letter abbreviations.
b. Draw the structure.

- c. Label all amide bonds.
d. Label the N-terminal and C-terminal amino acids.

28.54 Name each peptide using both the three-letter and one-letter abbreviations of the component amino acids.



28.55 Explain why a peptide C–N bond is stronger than an ester C–O bond.

28.56 Draw the *s*-trans and *s*-cis conformations of the peptide bond in the dipeptide Ala–Ala.

28.57 Draw the amino acids and peptide fragments formed when the decapeptide A–P–F–L–K–W–S–G–R–G is treated with each reagent or enzyme: (a) chymotrypsin; (b) trypsin; (c) carboxypeptidase; (d) $C_6H_5N=C=S$.

28.58 Give the amino acid sequence of each peptide using the fragments obtained by partial hydrolysis of the peptide with acid.

- A tetrapeptide that contains Ala, Gly, His, and Tyr, which is hydrolyzed to the dipeptides His–Tyr, Gly–Ala, and Ala–His.
- A pentapeptide that contains Glu, Gly, His, Lys, and Phe, which is hydrolyzed to His–Gly–Glu, Gly–Glu–Phe, and Lys–His.


28.59 Angiotensin is an octapeptide that narrows blood vessels, thereby increasing blood pressure. ACE inhibitors are a group of drugs used to treat high blood pressure by blocking the synthesis of angiotensin in the body. Angiotensin contains the amino acids Arg (2 equiv), His, Ile, Phe, Pro, Tyr, and Val. Determine the sequence of angiotensin using the following fragments obtained by partial hydrolysis with acid: Ile–His–Pro–Phe, Arg–Arg–Val, Tyr–Ile–His, and Val–Tyr.

28.60 Use the given experimental data to deduce the sequence of an octapeptide that contains the following amino acids: Ala, Gly (2 equiv), His (2 equiv), Ile, Leu, and Phe. Edman degradation cleaves Gly from the octapeptide, and carboxypeptidase forms Leu and a heptapeptide. Partial hydrolysis forms the following fragments: Ile–His–Leu, Gly, Gly–Ala–Phe–His, and Phe–His–Ile.









28.61 An octapeptide contains the following amino acids: Arg, Glu, His, Ile, Leu, Phe, Tyr, and Val. Carboxypeptidase treatment of the octapeptide forms Phe and a heptapeptide. Treatment of the octapeptide with chymotrypsin forms two tetrapeptides, **A** and **B**. Treatment of **A** with trypsin yields two dipeptides, **C** and **D**. Edman degradation cleaves the following amino acids from each peptide: Glu (octapeptide), Glu (**A**), Ile (**B**), Glu (**C**), and Val (**D**). Partial hydrolysis of tetrapeptide **B** forms Ile–Leu in addition to other products. Deduce the structure of the octapeptide and fragments **A–D**.

Peptide Synthesis


28.62 Draw all the products formed in the following reaction.




28.63 Draw the organic products formed in each reaction.



- 28.64** Draw all the steps in the synthesis of each peptide from individual amino acids: (a) Gly-Ala; (b) Phe-Leu; (c) Ile-Ala-Phe.
- 28.65** Write out the steps for the synthesis of each peptide using the Merrifield method: (a) Ala-Leu-Phe-Phe; (b) Phe-Gly-Ala-Ile.
- 28.66** An amino acid [RCH(NH₂)COOH] can readily be converted to an *N*-acetyl amino acid [RCH(NHCOCH₃)COOH] using acetic anhydride. Why can't this acetyl group be used as an amino protecting group, in place of the Boc group, for peptide synthesis?
- 28.67** Another method to form a peptide bond involves a two-step process:
- [1] Conversion of a Boc-protected amino acid to a *p*-nitrophenyl ester.
 - [2] Reaction of the *p*-nitrophenyl ester with an amino acid ester.




- a. Why does a *p*-nitrophenyl ester “activate” the carboxy group of the first amino acid to amide formation?
- b. Would a *p*-methoxyphenyl ester perform the same function? Why or why not?




- 28.68** In addition to forming an Fmoc-protected amino acid using Fmoc-Cl, an Fmoc protecting group can also be added to an amino group using reagent **A**.


- a. Draw the mechanism for the following reaction that adds an Fmoc group to an amino acid.



- b. Draw the mechanism for the reaction that removes an Fmoc group from an amino acid under the following conditions:




- 28.69** Many different insoluble polymers, called resins, are currently available for automated peptide synthesis. For example, the Wang resin contains benzene rings substituted with $-\text{CH}_2\text{OH}$ groups that serve as sites of attachment for amino acids. Propose reaction conditions that would bind an Fmoc-protected amino acid to a Wang resin. What reaction conditions could be used to remove the polypeptide from the resin after the synthesis is complete?




Proteins

- 28.70** Which of the following amino acids are typically found in the interior of a globular protein, and which are typically found on the surface: (a) phenylalanine; (b) aspartic acid; (c) lysine; (d) isoleucine; (e) arginine; (f) glutamic acid?
- 28.71** After the peptide chain of collagen has been formed, many of the proline residues are hydroxylated on one of the ring carbon atoms. Why is this process important for the triple helix of collagen?



Challenge Problems

- 28.72** Devise a stepwise synthesis of the tripeptide Val–Leu–Val from 3-methylbutanal $[(\text{CH}_3)_2\text{CHCH}_2\text{CHO}]$ as the only organic starting material. You may also use any required inorganic or organic reagents.
- 28.73** Besides asymmetric hydrogenation (Section 28.4), several other methods are now available for the synthesis of optically active amino acids. How might a reaction like the Strecker synthesis be adapted to the preparation of chiral amino acids?
- 28.74** As shown in Mechanism 28.2, the final steps in the Edman degradation result in rearrangement of a thiazolinone to an *N*-phenylthiohydantoin. Draw a stepwise mechanism for this acid-catalyzed reaction.



Lipids

29

- 29.1 Introduction
- 29.2 Waxes
- 29.3 Triacylglycerols
- 29.4 Phospholipids
- 29.5 Fat-soluble vitamins
- 29.6 Eicosanoids
- 29.7 Terpenes
- 29.8 Steroids



Cholesterol is the most prominent member of the steroid family, a group of organic lipids that contains a tetracyclic structure. Cholesterol is synthesized in the liver and is found in almost all body tissues. It is a vital component for healthy cell membranes and serves as the starting material for the synthesis of all other steroids. But, as the general public now knows well, elevated cholesterol levels can lead to coronary artery disease. For this reason, consumer products are now labeled with their cholesterol content. In Chapter 29, we learn about the properties of cholesterol and other lipids.

We conclude the discussion of the organic molecules in biological systems by turning our attention to **lipids**, biomolecules that are soluble in organic solvents. Unlike the carbohydrates in Chapter 27 and the amino acids and proteins in Chapter 28, lipids contain many carbon–carbon and carbon–hydrogen bonds and few functional groups.

Since lipids are the biomolecules that most closely resemble the hydrocarbons we studied in Chapters 4 and 10, we have already learned many facts that directly explain their properties. Since there is no one functional group that is present in all lipids, however, the chemistry of lipids draws upon knowledge learned in many prior chapters.

29.1 Introduction

The word *lipid* comes from the Greek word *lipos* for “fat.”

- **Lipids are biomolecules that are soluble in organic solvents.**

Lipids are unique among organic molecules because their identity is defined on the basis of a *physical property* and not by the presence of a particular functional group. Because of this, lipids come in a wide variety of structures and they have many different functions in the cell. Three examples are given in Figure 29.1.




The large number of **carbon–carbon and carbon–hydrogen σ bonds in lipids makes them very soluble in organic solvents and insoluble in water.** Monosaccharides (from which carbohydrates are formed) and amino acids (from which proteins are formed), on the other hand, are very polar, so they tend to be water soluble. Because lipids share many properties with hydrocarbons, several features of lipid structure and properties have already been discussed. Table 29.1 summarizes sections of the text where aspects of lipid chemistry were covered previously.

Table 29.1 Summary of Lipid Chemistry Discussed Prior to Chapter 29

Topic	Section	Topic	Section
• Vitamin A	3.5	• Lipid oxidation	15.11
• Soap	3.6	• Vitamin E	15.12
• Phospholipids, the cell membrane	3.7	• Steroid synthesis	16.14
• Lipids Part 1	4.15	• Prostaglandins	19.6
• Leukotrienes	9.16	• Lipid hydrolysis	22.12A
• Fats and oils	10.6	• Soap	22.12B
• Oral contraceptives	11.4	• Cholesteryl esters	22.17
• Hydrogenation of oils	12.4	• Steroid synthesis	24.8

Figure 29.1


Three examples of lipids




All lipids have many C–C and C–H bonds, but there is no one functional group common to all lipids.

Lipids can be categorized as hydrolyzable or nonhydrolyzable.

- [1] **Hydrolyzable lipids can be cleaved into smaller molecules by hydrolysis with water.** Most hydrolyzable lipids contain an ester unit. We will examine three subgroups: waxes, triacylglycerols, and phospholipids.



- [2] **Nonhydrolyzable lipids cannot be cleaved into smaller units by aqueous hydrolysis.** Nonhydrolyzable lipids tend to be more varied in structure. We will examine four different types: fat-soluble vitamins, eicosanoids, terpenes, and steroids.





29.2 Waxes



Water beads up on the surface of a leaf because of the leaf's waxy coating.

Waxes are the simplest hydrolyzable lipids. **Waxes are esters (RCOOR') formed from a high molecular weight alcohol (R'OH) and a fatty acid (RCOOH).**

Because of their long hydrocarbon chains, **waxes are very hydrophobic.** They form a protective coating on the feathers of birds to make them water repellent, and on leaves to prevent water evaporation. **Lanolin**, a wax composed of a complex mixture of high molecular weight esters, coats the wool fibers of sheep. **Spermaceti wax**, isolated from the heads of sperm whales, is largely $\text{CH}_3(\text{CH}_2)_{14}\text{COO}(\text{CH}_2)_{15}\text{CH}_3$. The three-dimensional structure of this compound shows how small the ester group is compared to the long hydrocarbon chains.



Problem 29.1


Carnauba wax, a wax that coats the leaves of the Brazilian palm tree, is used for hard, high-gloss finishes for floors, boats, and automobiles. (a) Draw the structure of one component of carnauba wax, formed from an unbranched 32-carbon carboxylic acid and a straight chain 34-carbon alcohol. (b) Draw the structure of a second component of carnauba wax, formed by the polymerization of $\text{HO}(\text{CH}_2)_{17}\text{COOH}$.

29.3 Triacylglycerols

Triacylglycerols, or triglycerides, are the most abundant lipids, and for this reason we have already discussed many of their properties in earlier sections of this text.

- Triacylglycerols are triesters that produce glycerol and three molecules of fatty acid upon hydrolysis.

Line structures of stearic, oleic, linoleic, and linolenic acids can be found in Table 10.2. Ball-and-stick models of these fatty acids are shown in Figure 10.6.



Simple triacylglycerols are composed of three identical fatty acid side chains, whereas **mixed triacylglycerols** have two or three different fatty acids. Table 29.2 lists the most common fatty acids used to form triacylglycerols.

What are the characteristics of these fatty acids?

- All fatty acid chains are unbranched, but they may be saturated or unsaturated.
- Naturally occurring fatty acids have an even number of carbon atoms.
- Double bonds in naturally occurring fatty acids generally have the *Z* configuration.
- The melting point of a fatty acid depends on the degree of unsaturation.

Fats and oils are triacylglycerols; that is, they are triesters of glycerol and these fatty acids.

- Fats have higher melting points, making them solids at room temperature.
- Oils have lower melting points, making them liquids at room temperature.

This melting point difference correlates with the number of degrees of unsaturation present in the fatty acid side chains. **As the number of double bonds increases, the melting point decreases, as it does for the constituent fatty acids as well.**

Table 29.2 The Most Common Fatty Acids in Triacylglycerols


Number of C atoms	Number of C=C bonds	Structure	Name	Mp (°C)
Saturated fatty acids				
12	0	CH ₃ (CH ₂) ₁₀ COOH	lauric acid	44
14	0	CH ₃ (CH ₂) ₁₂ COOH	myristic acid	58
16	0	CH ₃ (CH ₂) ₁₄ COOH	palmitic acid	63
18	0	CH ₃ (CH ₂) ₁₆ COOH	stearic acid	69
20	0	CH ₃ (CH ₂) ₁₈ COOH	arachidic acid	77
Unsaturated fatty acids				
16	1	CH ₃ (CH ₂) ₅ CH=CH(CH ₂) ₇ COOH	palmitoleic acid	1
18	1	CH ₃ (CH ₂) ₇ CH=CH(CH ₂) ₇ COOH	oleic acid	4
18	2	CH ₃ (CH ₂) ₄ (CH=CHCH ₂) ₂ (CH ₂) ₆ COOH	linoleic acid	-5
18	3	CH ₃ CH ₂ (CH=CHCH ₂) ₃ (CH ₂) ₆ COOH	linolenic acid	-11
20	4	CH ₃ (CH ₂) ₄ (CH=CHCH ₂) ₄ (CH ₂) ₂ COOH	arachidonic acid	-49

The most common saturated fatty acids are palmitic and stearic acids. The most common unsaturated fatty acid is oleic acid.

Linoleic and linolenic acids are called **essential fatty acids** because we cannot synthesize them and must acquire them in our diets.

Figure 29.2

Three-dimensional structures of a saturated and unsaturated triacylglycerol



- Three saturated side chains lie parallel to each other, making a compact lipid.
- One Z double bond in a fatty acid side chain produces a twist so the lipid is no longer so compact.



Unlike other vegetable oils, oils from palm and coconut trees are very high in saturated fats. Considerable evidence currently suggests that diets high in saturated fats lead to a greater risk of heart disease. For this reason, the demand for coconut and palm oils has decreased considerably in recent years, and many coconut plantations previously farmed in the South Pacific are no longer in commercial operation.



Three-dimensional structures of a saturated and unsaturated triacylglycerol are shown in Figure 29.2. With no double bonds, the three side chains of the saturated lipid lie parallel to each other, making it possible for this compound to pack relatively efficiently in a crystalline lattice, thus leading to a high melting point. In the unsaturated lipid, however, a single Z double bond places a kink in the side chain, making it more difficult to pack efficiently in the solid state, thus leading to a lower melting point.

Solid fats have a relatively high percentage of saturated fatty acids and are generally of animal origin. Liquid oils have a higher percentage of unsaturated fatty acids and are generally of vegetable origin. Table 29.3 lists the fatty acid composition of some common fats and oils.

Table 29.3 Fatty Acid Composition of Some Fats and Oils

Source	% Saturated fatty acids	% Oleic acid	% Linoleic acid
beef	49–62	37–43	2–3
milk	37	33	3
coconut	86	7	—
corn	11–16	19–49	34–62
olive	11	84	4
palm	43	40	8
safflower	9	13	78
soybean	15	20	52


Data from *Merck Index*, 10th ed. Rahway, NJ: Merck and Co.; and Wilson, et al., 1967, *Principles of Nutrition*, 2nd ed. New York: Wiley.

Fish oils, such as cod liver and herring oils, are very rich in polyunsaturated triacylglycerols. These triacylglycerols pack so poorly that they have very low melting points; thus, they remain liquids even in the cold water inhabited by these fish.



The hydrolysis, hydrogenation, and oxidation of triacylglycerols—reactions originally discussed in Chapters 12, 15, and 22—are summarized here for your reference.


[1] Hydrolysis of triacylglycerols (Section 22.12A)



Three ester units are cleaved.


Hydrolysis of a triacylglycerol with water in the presence of either acid, base, or an enzyme yields glycerol and three fatty acids. This cleavage reaction follows the same mechanism as any other ester hydrolysis (Section 22.11). This reaction is the first step in triacylglycerol metabolism.

[2] Hydrogenation of unsaturated fatty acids (Section 12.4)



The double bonds of an unsaturated fatty acid can be hydrogenated by using H₂ in the presence of a transition metal catalyst. Hydrogenation converts a liquid oil to a solid fat. This process, sometimes called **hardening**, is used to prepare margarine from vegetable oils.

[3] Oxidation of unsaturated fatty acids (Section 15.11)




Allylic C–H bonds are weaker than other C–H bonds and are thus susceptible to oxidation with molecular oxygen by a radical process. The hydroperoxide formed by this process is unstable, and it undergoes further oxidation to products that often have a disagreeable odor. This oxidation process turns an oil rancid.

In the cell, the principal function of triacylglycerols is energy storage. Complete metabolism of a triacylglycerol yields CO_2 and H_2O , and a great deal of energy. This overall reaction is reminiscent of the combustion of alkanes in fossil fuels, a process that also yields CO_2 and H_2O and provides energy to heat homes and power automobiles (Section 4.14B). Fundamentally both processes convert C–C and C–H bonds to C–O bonds, a highly exothermic reaction.

Metabolism of a lipid

The average body fat content of men and women is ~20% and ~25%, respectively. (For elite athletes, however, the averages are more like <10% for men and <15% for women.) This stored fat can fill the body's energy needs for two or three months.



Carbohydrates provide an energy boost, but only for the short term, such as during strenuous exercise. Our long-term energy needs are met by triacylglycerols, because they store ~38 kJ/g, whereas carbohydrates and proteins store only ~16 kJ/g.

Because triacylglycerols release heat on combustion, they can in principle be used as fuels for vehicles. In fact, coconut oil was used as a fuel during both World War I and World War II, when gasoline and diesel supplies ran short. Since coconut oil is more viscous than petroleum products and freezes at 24°C , engines must be modified to use it and it can't be used in cold climates. Nonetheless, a limited number of trucks and boats can now use vegetable oils, sometimes blended with diesel, as a fuel source. When the price of crude oil is high, the use of these **biofuels** becomes economically attractive.




Problem 29.2

How would you expect the melting point of eicosapentaenoic acid [$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_5(\text{CH}_2)_2\text{COOH}$] to compare with the melting points of the fatty acids listed in Table 29.2?

Problem 29.3

Draw the products formed when triacylglycerol **A** is treated with each reagent. Rank compounds **A**, **B**, and **C** in order of increasing melting point.



Problem 29.4

The main fatty acid component of the triacylglycerols in coconut oil is lauric acid, $\text{CH}_3(\text{CH}_2)_{10}\text{COOH}$. Explain why coconut oil is a liquid at room temperature even though it contains a large fraction of this saturated fatty acid.

Problem 29.5

Unlike many fats and oils, the cocoa butter used to make chocolate is remarkably uniform in composition. All triacylglycerols contain oleic acid esterified to the 2° OH group of glycerol, and either palmitic acid or stearic acid esterified to the 1° OH groups. Draw the structures of two possible triacylglycerols that compose cocoa butter.




29.4 Phospholipids

Phospholipids are hydrolyzable lipids that contain a phosphorus atom. There are two common types of phospholipids: **phosphoacylglycerols** and **sphingomyelins**. Both classes are found almost exclusively in the cell membranes of plants and animals, as discussed in Section 3.7.


The phosphorus atom in a phosphodiester shares 10 electrons. Recall that third-row elements (such as P and S) can be surrounded by more than eight electrons.

Phospholipids are organic derivatives of phosphoric acid, formed by replacing two of the H atoms by R groups. This type of functional group is called a **phosphodiester**, or a **phosphoric acid diester**. These compounds are phosphorus analogues of carboxylic esters. In cells, the remaining OH group on phosphorus loses its proton, giving the phosphodiester a net negative charge.




29.4A Phosphoacylglycerols

Phosphoacylglycerols (or phosphoglycerides) are the second most abundant type of lipid. They form the principal lipid component of most cell membranes. Their structure resembles the triacylglycerols of Section 29.3 with one important difference. In phosphoacylglycerols, only two of the hydroxy groups of glycerol are esterified with fatty acids. The third OH group is part of a phosphodiester, which is also bonded to another low molecular weight alcohol.



There are two prominent types of phosphoacylglycerols. They differ in the identity of the R'' group in the phosphodiester.

- When R'' = CH₂CH₂NH₃⁺, the compound is called a **phosphatidylethanolamine** or **cephalin**.
- When R'' = CH₂CH₂N(CH₃)₃⁺, the compound is called a **phosphatidylcholine**, or **lecithin**.




The middle carbon of the glycerol backbone of all of these compounds is a stereogenic center, usually with the *R* configuration.

The phosphorus side chain of a phosphoacylglycerol makes it different from a triacylglycerol. The two fatty acid side chains form two nonpolar “tails” that lie parallel to each other, while the phosphodiester end of the molecule is a charged or polar “head.” A three-dimensional structure of a phosphoacylglycerol is shown in Figure 29.3.

As discussed in Section 3.7, when these phospholipids are mixed with water, they assemble in an arrangement called a **lipid bilayer**. The ionic heads of the phospholipid are oriented on the outside

Figure 29.3
Three-dimensional structure
of a phosphoacylglycerol



- A phosphoacylglycerol has two distinct regions: two nonpolar tails due to the long-chain fatty acids, and a very polar head from the charged phosphodiester.

and the nonpolar tails on the inside. The identity of the fatty acids in the phospholipid determines the rigidity of this bilayer. When the fatty acids are saturated, they pack well in the interior of the lipid bilayer, and the membrane is quite rigid. When there are many unsaturated fatty acids, the nonpolar tails cannot pack as well and the bilayer is more fluid. Thus, important characteristics of this lipid bilayer are determined by the three-dimensional structure of the molecules that comprise it.

Cell membranes are composed of these lipid bilayers (see Figure 3.7). Proteins and cholesterol are embedded in the membranes as well, but the phospholipid bilayer forms the main fabric of the insoluble barrier that protects the cell.

Problem 29.6 Draw the structure of a lecithin containing oleic acid and palmitic acid as the fatty acid side chains.

Problem 29.7 Phosphoacylglycerols should remind you of soaps (Section 3.6). In what ways are these compounds similar?

29.4B Sphingomyelins

Sphingomyelins, the second major class of phospholipids, are derivatives of the amino alcohol **sphingosine**, in much the same way that triacylglycerols and phosphoacylglycerols are derivatives of glycerol. Other notable features of a sphingomyelin include:

- **A phosphodiester at C1.**
- **An amide formed with a fatty acid at C2.**





Figure 29.4

A comparison of a triacylglycerol, a phosphoacylglycerol, and a sphingomyelin



- A triacylglycerol has three nonpolar side chains.
- The three OH groups of glycerol are esterified with three fatty acids.
- A phosphoacylglycerol has two nonpolar side chain tails and one ionic head.
- Two OH groups of glycerol are esterified with fatty acids.
- A phosphodiester is located on a terminal carbon.
- A sphingomyelin has two nonpolar side chain tails and one ionic head.
- A sphingomyelin is formed from sphingosine, not glycerol. One of the nonpolar tails is an amide.
- A phosphodiester is located on a terminal carbon.

Like phosphoacylglycerols, **sphingomyelins are also a component of the lipid bilayer of cell membranes.** The coating that surrounds and insulates nerve cells, the **myelin sheath**, is particularly rich in sphingomyelins, and is vital for proper nerve function. Deterioration of the myelin sheath as seen in multiple sclerosis leads to disabling neurological problems.

Figure 29.4 compares the structural features of the most common hydrolyzable lipids: a triacylglycerol, a phosphoacylglycerol, and a sphingomyelin.

Problem 29.8 Why are phospholipids, but not triacylglycerols, found in cell membranes?


29.5 Fat-Soluble Vitamins

Vitamins are organic compounds required in small quantities for normal metabolism (Section 3.5). Because our cells cannot synthesize these compounds, they must be obtained in the diet. Vitamins can be categorized as fat soluble or water soluble. **The fat-soluble vitamins are lipids.**


The four fat-soluble vitamins—**A, D, E, and K**—are found in fruits and vegetables, fish, liver, and dairy products. Although fat-soluble vitamins must be obtained from the diet, they do not have to be ingested every day. Excess vitamins are stored in fat cells, and then used when needed. Figure 29.5 shows the structure of these vitamins and summarizes their functions.

Electrostatic potential plots of vitamins A and E (Figure 29.6) show that the electron density is virtually uniform in these compounds. The large regions of nonpolar C–C and C–H bonds tend to obscure small dipoles that occur in the one or two polar bonds, making these vitamins nonpolar and hydrophobic.


Problem 29.9 Explain why regularly ingesting a large excess of a fat-soluble vitamin can lead to severe health problems, whereas ingesting a large excess of a water-soluble vitamin often causes no major health problems.

Figure 29.5 The fat-soluble vitamins


- **Vitamin A** (retinol, Section 3.5) is obtained from fish liver oils and dairy products, and is synthesized from β -carotene, the orange pigment in carrots.
- In the body, vitamin A is converted to 11-*cis*-retinal, the light-sensitive compound responsible for vision in all vertebrates (Section 21.11B). It is also needed for healthy mucous membranes.
- A deficiency of vitamin A causes night blindness, as well as dry eyes and skin.



- **Vitamin D₃** is the most abundant of the D vitamins. Strictly speaking, it is not a vitamin because it can be synthesized in the body from cholesterol. Nevertheless, it is classified as such, and many foods (particularly milk) are fortified with vitamin D₃ so that we get enough of this vital nutrient.
- Vitamin D helps regulate both calcium and phosphorus metabolism.
- A deficiency of vitamin D causes rickets, a bone disease characterized by knock-knees, spinal curvature, and other deformities.



- The term **vitamin E** refers to a group of structurally similar compounds, the most potent being α -tocopherol (Section 15.12).
- Vitamin E is an antioxidant, so it protects unsaturated side chains in fatty acids from oxidation.
- A deficiency of vitamin E causes numerous neurologic problems.



- **Vitamin K** (phylloquinone) regulates the synthesis of prothrombin and other proteins needed for blood to clot.
- A deficiency of vitamin K leads to excessive and sometimes fatal bleeding because of inadequate blood clotting.


29.6 Eicosanoids

The word **eicosanoid** is derived from the Greek word *eikosi*, meaning 20.


The **eicosanoids** are a group of biologically active compounds containing 20 carbon atoms derived from arachidonic acid. The **prostaglandins** (Section 19.6) and the **leukotrienes** (Section 9.16) are two types of eicosanoids. Two others are the **thromboxanes** and **prostacyclins**.

Figure 29.6

Electrostatic potential plots of vitamins A and E



- The electron density is distributed fairly evenly among the carbon atoms of these vitamins due to their many nonpolar C–C and C–H bonds.




All eicosanoids are very potent compounds present in low concentration in cells. They are **local mediators**, meaning that they perform their function in the environment in which they are synthesized. This distinguishes them from **hormones**, which are first synthesized and then transported in the bloodstream to their site of action. Eicosanoids are not stored in cells; rather, they are synthesized from arachidonic acid in response to an external stimulus.

The synthesis of prostaglandins, thromboxanes, and prostacyclins begins with the oxidation of arachidonic acid with O_2 by a **cyclooxygenase** enzyme, which forms an unstable cyclic intermediate, PGG_2 . PGG_2 is then converted via different pathways to these three classes of compounds. Leukotrienes are formed by a different pathway, using an enzyme called a **lipoxygenase**. These four paths for arachidonic acid are summarized in Figure 29.7.

Each eicosanoid is associated with specific types of biological activity (Table 29.4). In some cases, the effects oppose one another. For example, thromboxanes are vasoconstrictors that trigger blood platelet aggregation, whereas prostacyclins are vasodilators that inhibit platelet aggregation. The levels of these two eicosanoids must be in the right balance for cells to function properly.

Because of their wide range of biological functions, prostaglandins and their analogues have found several clinical uses. For example, **dinoprostone**, the generic name for PGE_2 , is administered to relax the smooth muscles of the uterus when labor is induced, and to terminate pregnancies in the early stages.




Other details of the biosynthesis of leukotrienes and prostaglandins were given in Sections 9.16 and 19.6, respectively.

Table 29.4 Biological Activity of the Eicosanoids


Eicosanoid	Effect	Eicosanoid	Effect
Prostaglandins	• Lower blood pressure	Thromboxanes	• Constrict blood vessels
	• Inhibit blood platelet aggregation		• Trigger blood platelet aggregation
	• Control inflammation	Prostacyclins	• Dilate blood vessels
	• Lower gastric secretions		• Inhibit blood platelet aggregation
	• Stimulate uterine contractions		Leukotrienes
• Relax smooth muscles of the uterus			

Figure 29.7

The conversion of arachidonic acid to prostaglandins, thromboxanes, prostacyclins, and leukotrienes



Because prostaglandins themselves are unstable in the body, often having half-lives of only minutes, more stable analogues have been developed that retain their important biological activity longer. For example, **misoprostol**, an analogue of PGE₁, is sold as a mixture of stereoisomers. Misoprostol is administered to prevent gastric ulcers in patients who are at high risk of developing them.




Studying the biosynthesis of eicosanoids has led to other discoveries as well. For example, aspirin and other nonsteroidal anti-inflammatory drugs (**NSAIDs**) inactivate the cyclooxygenase enzyme needed for prostaglandin synthesis. In this way, NSAIDs block the synthesis of the prostaglandins that cause inflammation (Section 19.6).


More recently, it has been discovered that two *different* cyclooxygenase enzymes, called **COX-1** and **COX-2**, are responsible for prostaglandin synthesis. COX-1 is involved with the usual production of prostaglandins, but COX-2 is responsible for the synthesis of additional prostaglandins in inflammatory diseases like arthritis. **NSAIDs like aspirin and ibuprofen inactivate both the**

COX-1 and COX-2 enzymes. This activity also results in an increase in gastric secretions, making an individual more susceptible to ulcer formation.


A group of anti-inflammatory drugs that block only the COX-2 enzyme was developed in the 1990s. These drugs—**rofecoxib**, **valdecoxib**, and **celecoxib**—do not cause an increase in gastric secretions, and thus were touted as especially effective NSAIDs for patients with arthritis, who need daily doses of these medications. Unfortunately, both rofecoxib and valdecoxib have now been removed from the market, since their use has been associated with an increased risk of heart attack and stroke.



Generic name: rofecoxib
Trade name: Vioxx



Generic name: valdecoxib
Trade name: Bextra



Generic name: celecoxib
Trade name: Celebrex


A COX-3 enzyme was also reported in 2002. Its activity is inhibited by acetaminophen, the active ingredient in the pain reliever Tylenol.

The discovery of drugs that block prostaglandin synthesis illustrates how basic research in organic chemistry can lead to important practical applications. Elucidating the structure and biosynthesis of prostaglandins began as a project in basic research. It has now resulted in a number of applications that benefit many individuals with various illnesses.


Problem 29.10 How are the two isomers of misoprostol related?

29.7 Terpenes

Terpenes are lipids composed of repeating five-carbon units called isoprene units. An **isoprene unit** has five carbons: four in a row, with a one-carbon branch on a middle carbon.



Terpenes have a wide variety of structures. They can be acyclic or have one or more rings. They may have only carbon and hydrogen atoms, or they may have heteroatoms as well. The most common heteroatom in terpenes is oxygen. Many **essential oils**, a group of compounds isolated from plant sources by distillation, are terpenes. Examples include myrcene and menthol.



29.7A Locating Isoprene Units in Terpenes

How do we identify the isoprene units in these molecules? Start at one end of the molecule near a branch point. Then **look for a four-carbon chain with a one-carbon branch**. This forms one isoprene unit. Continue along the chain or around the ring until all the carbons are part of an isoprene unit. Keep in mind the following:




bayberry plant
(source of myrcene)



peppermint plant
(source of menthol)

- An isoprene unit may be composed of C–C σ bonds only, or there may be π bonds at any position.
- Isoprene units are always connected by one or more carbon–carbon bonds.
- Each carbon atom is part of one isoprene unit only.
- Every isoprene unit has five carbon atoms. Heteroatoms may be present but their presence is ignored in locating isoprene units.

Myrcene and menthol, for example, each have 10 carbon atoms, so they are composed of two isoprene units.




Terpenes are classified by the number of isoprene units they contain. A **monoterpene contains 10 carbons** and has two isoprene units, a **sesquiterpene contains 15 carbons** and has three isoprene units, and so forth. The different terpene classes are summarized in Table 29.5.

Several examples, with the isoprene units labeled in red, are given in Figure 29.8.

Table 29.5 Classes of Terpenes





Name	Number of C atoms	Number of isoprene units
Monoterpene	10	2
Sesquiterpene	15	3
Diterpene	20	4
Sesterterpene	25	5
Triterpene	30	6
Tetraterpene	40	8

Figure 29.8
Examples of some
common terpenes




- Isoprene units are labeled in red, with C–C bonds (in black) joining two units.
- The source of each terpene is given in parentheses.

Problem 29.11 Locate the isoprene units in each compound.



Problem 29.12 Manoalide, a sesterterpene isolated from the Pacific marine sponge *Luffariella variabilis* by Scheuer and co-workers at the University of Hawai'i at Mānoa, has anti-inflammatory, analgesic, and antifungal properties. Find the isoprene units in manoalide.




29.7B The Biosynthesis of Terpenes


Terpene biosynthesis is an excellent example of how syntheses in nature occur with high efficiency. There are two ways this is accomplished.

- [1] **The same reaction is used over and over again to prepare progressively more complex compounds.**
- [2] **Key intermediates along the way serve as the starting materials for a wide variety of other compounds.**

All terpenes are synthesized from dimethylallyl diphosphate and isopentenyl diphosphate. Both of these five-carbon compounds are synthesized, in turn, in a multistep process from three molecules of acetyl CoA (Section 22.17).



Diphosphate, abbreviated as **OPP**, is often used as a leaving group in biological systems. **It is a good leaving group because it is a weak, resonance-stabilized base.**




The overall strategy of terpene biosynthesis from dimethylallyl diphosphate and isopentenyl diphosphate is summarized in Figure 29.9.

There are three basic parts:

- [1] The two C₅ diphosphates are converted to **geranyl diphosphate**, a C₁₀ monoterpene. Geranyl diphosphate is the starting material for all other monoterpenes.

Figure 29.9
An outline of terpene biosynthesis



[2] Geranyl diphosphate is converted to **farnesyl diphosphate**, a C_{15} **sesquiterpene**, by addition of a five-carbon unit. Farnesyl diphosphate is the starting material for all sesquiterpenes and diterpenes.


[3] Two molecules of farnesyl diphosphate are converted to **squalene**, a C_{30} **triterpene**. Squalene is the starting material for all triterpenes and steroids.

The biological formation of geranyl diphosphate from the two five-carbon diphosphates involves three steps: **loss of the leaving group**, **nucleophilic attack**, and **loss of a proton**, as shown in Mechanism 29.1.




Mechanism 29.1 Biological Formation of Geranyl Diphosphate

Steps [1]–[2] Loss of the leaving group and nucleophilic attack to form a new C–C bond



- Loss of the diphosphate leaving group forms a resonance-stabilized carbocation in Step [1], which reacts with the nucleophilic double bond of the 1° diphosphate to form a new C–C bond and a 3° carbocation in Step [2].
- Steps [1] and [2] are analogous to an S_N1 mechanism because the leaving group (^-OPP) is lost before the nucleophile (a $C=C$) attacks.

Step [3] Loss of a proton




- Loss of a proton forms geranyl diphosphate in Step [3].

The biological conversion of geranyl diphosphate to farnesyl diphosphate involves the same three steps, as shown in Mechanism 29.2.




Mechanism 29.2 Biological Formation of Farnesyl Diphosphate

Steps [1]–[2] Loss of the leaving group and nucleophilic attack to form a new C–C bond




- Loss of the diphosphate leaving group forms a resonance-stabilized carbocation in Step [1], which reacts with isopentenyl diphosphate to form a new C–C bond and a 3° carbocation in Step [2].

Step [3] Loss of a proton




- Loss of a proton forms a new π bond and farnesyl diphosphate in Step [3].

Two molecules of farnesyl diphosphate react to form squalene, from which all other triterpenes and steroids are synthesized.



Aqueous hydrolysis of geranyl and farnesyl diphosphates forms the monoterpene geraniol and the sesquiterpene farnesol, respectively.




All other terpenes are biologically derived from geranyl and farnesyl diphosphates by a series of reactions. Cyclic compounds are formed by intramolecular reactions involving nucleophilic attack of π bonds on intermediate carbocations. To form some cyclic compounds, the *E*

double bond in geranyl diphosphate must first isomerize to an isomeric diphosphate with a *Z* double bond, neryl diphosphate, by the process illustrated in Mechanism 29.3. Isomerization forms a substrate with a leaving group and nucleophilic double bond in close proximity, so that an intramolecular reaction can occur.




Mechanism 29.3 Isomerization of Geranyl Diphosphate to Neryl Diphosphate

Steps [1]–[2] Isomerization of geranyl diphosphate to linalyl diphosphate




- Loss of the diphosphate leaving group forms a resonance-stabilized carbocation in Step [1], which reacts with the diphosphate anion to form linalyl diphosphate, a constitutional isomer.
- The original *E* double bond is now a single bond that can rotate freely.

Steps [3]–[4] Isomerization of linalyl diphosphate to neryl diphosphate




- Loss of the diphosphate leaving group forms a resonance-stabilized carbocation in Step [3]. The only difference in the products of Steps [1] and [3] is the geometry around the internal carbon–carbon double bond.
- Nucleophilic attack with diphosphate forms neryl diphosphate, a stereoisomer of geranyl diphosphate. The diphosphate leaving group of neryl diphosphate is now in closer proximity to the double bond at the other end of the chain, so that intramolecular cyclization can occur.


In the synthesis of α -terpineol or limonene, for example, geranyl diphosphate isomerizes to form neryl diphosphate (Step [1] in the following reaction sequence). Neryl diphosphate then cyclizes to a 3° carbocation by intramolecular attack (Steps [2]–[3]). Nucleophilic attack of water on this carbocation yields α -terpineol (Step [4]) or loss of a proton yields limonene (Step [5]). Both products are cyclic monoterpenes.



Problem 29.13 Write a stepwise mechanism for the following reaction.



Problem 29.14 Draw a stepwise mechanism for the conversion of geranyl diphosphate to α -terpinene.




29.8 Steroids

The steroids are a group of tetracyclic lipids, many of which are biologically active.


29.8A Steroid Structure

Steroids are composed of three six-membered rings and one five-membered ring, joined together as drawn. Many steroids also contain two methyl groups, called **angular methyl groups**, at the two ring junctions indicated. The steroid rings are lettered **A**, **B**, **C**, and **D**, and the 17 ring carbons are numbered as shown. The two angular methyl groups are numbered C18 and C19.


General steroid skeleton



Numbering the steroid skeleton





Whenever two rings are fused together, the substituents at the ring fusion can be arranged *cis* or *trans*. To see more easily why this is true, consider **decalin**, which consists of two six-membered rings fused together. ***trans*-Decalin** has the two hydrogen atoms at the ring fusion on opposite sides, whereas ***cis*-decalin** has them on the same side.



Three-dimensional structures of these molecules show how different these two possible arrangements actually are. The two rings of *trans*-decalin lie roughly in the same plane, whereas the two rings of *cis*-decalin are almost perpendicular to each other. **The *trans* arrangement is lower in energy and therefore more stable.**

Figure 29.10

The three-dimensional structure of the steroid nucleus



In steroids, each ring fusion could theoretically have the cis or trans configuration, but, by far the most common arrangement is all trans. Because of this, **all four rings of the steroid skeleton lie in the same plane**, and the ring system is fairly rigid. The two angular methyl groups are oriented perpendicular to the plane of the molecule. These methyl groups make one side of the steroid skeleton significantly more hindered than the other, as shown in Figure 29.10.

Although steroids have the same fused-ring arrangement of carbon atoms, they differ in the identity and location of the substituents attached to that skeleton.


Problem 29.15

(a) Draw a skeletal structure of the anabolic steroid 4-androstene-3,17-dione, also called “andro,” from the following description. Andro contains the tetracyclic steroid skeleton with carbonyl groups at C3 and C17, a double bond between C4 and C5, and methyl groups bonded to C10 and C13. (b) Add wedges and dashes for all stereogenic centers with the following information: the configuration at C10 is *R*, the configuration at C13 is *S*, and all substituents at ring fusions are trans to each other.

29.8B Cholesterol

Cholesterol, the chapter-opening molecule, has the tetracyclic carbon skeleton characteristic of steroids. It also has eight stereogenic carbons (seven on rings and one on a side chain), so there are $2^8 = 256$ possible stereoisomers. In nature, however, only the following stereoisomer exists:

Cholesterol has also been discussed in Sections 3.4C and 4.15. The role of cholesterol in plaque formation and atherosclerosis was discussed in Section 22.17.



Konrad Bloch and Feodor Lynen shared the 1964 Nobel Prize in Physiology or Medicine for unraveling the complex transformation of squalene to cholesterol.

Cholesterol is essential to life because it forms an important component of cell membranes and is the starting material for the synthesis of all other steroids. Humans do not have to ingest cholesterol, because it is synthesized in the liver and then transported to other tissues through the bloodstream. Because cholesterol has only one polar OH group and many nonpolar C–C and C–H bonds, it is insoluble in water (and, thus, in the aqueous medium of the blood).

Cholesterol is synthesized in the body from squalene, a C₃₀ triterpene that is itself prepared from smaller terpenes, as discussed in Section 29.7B. Because the biosynthesis of all terpenes begins with acetyl CoA, every one of the 27 carbon atoms of cholesterol comes from the same two-carbon precursor. The major steps in the conversion of squalene to cholesterol are given in Figure 29.11.

The conversion of squalene to cholesterol consists of five different parts:

- [1] **Epoxidation** of squalene with an enzyme, squalene epoxidase, gives squalene oxide, which contains a single epoxide on one of the six double bonds.
- [2] **Cyclization** of squalene oxide yields a carbocation, called the protosterol cation. This reaction results in the formation of four new C–C bonds and the tetracyclic ring system.
- [3] **The protosterol carbocation rearranges** by a series of 1,2-shifts of either a hydrogen or methyl group to form another 3° carbocation.
- [4] **Loss of a proton** gives an alkene called **lanosterol**. Although lanosterol has seven stereogenic centers, a single stereoisomer is formed.
- [5] Lanosterol is then converted to cholesterol by a multistep process that results in removal of three methyl groups.

Several drugs are now available to reduce the level of cholesterol in the bloodstream. These compounds act by blocking the biosynthesis of cholesterol at its very early stages. Two examples include atorvastatin (Lipitor) and simvastatin (Zocor), whose structures appear in Figure 29.12.

Problem 29.16

Draw the enantiomer and any two diastereomers of cholesterol. Does the OH group of cholesterol occupy an axial or equatorial position?

Figure 29.11

The biosynthesis of cholesterol






Figure 29.12
Two cholesterol-lowering drugs




Generic name: atorvastatin
Trade name: Lipitor



Generic name: simvastatin
Trade name: Zocor

Problem 29.17 Treatment of cholesterol with mCPBA results in formation of a single epoxide **A**, with the stereochemistry drawn. Why isn't the isomeric epoxide **B** formed to any extent?



29.8C Other Steroids


Many other important steroids are hormones secreted by the endocrine glands. Two classes are the **sex hormones** and the **adrenal cortical steroids**.

There are two types of female sex hormones, **estrogens** and **progestins**. The male sex hormones are called **androgens**. The most important members of each hormone type are given in Table 29.6.

Table 29.6 The Female and Male Sex Hormones

Structure	Properties
<p>estradiol</p>	<ul style="list-style-type: none"> • Estradiol and estrone are estrogens synthesized in the ovaries. They control the development of secondary sex characteristics in females and regulate the menstrual cycle.
<p>estrone</p>	
<p>progesterone</p>	<ul style="list-style-type: none"> • Progesterone is often called the “pregnancy hormone.” It is responsible for the preparation of the uterus for implantation of a fertilized egg.
<p>testosterone</p>	
<p>androsterone</p>	

Synthetic analogues of these steroids have found important uses, such as in oral contraceptives, first mentioned in Section 11.4.




Synthetic androgen analogues, called **anabolic steroids**, promote muscle growth. They were first developed to help individuals whose muscles had atrophied from lack of use following surgery. They have since come to be used by athletes and body builders, although their use is not permitted in competitive sports. Many physical and psychological problems result from their prolonged use.




Some body builders use anabolic steroids to increase muscle mass. Long-term or excessive use can cause many health problems, including high blood pressure, liver damage, and cardiovascular disease.

Anabolic steroids, such as stanozolol, nandrolone, and tetrahydrogestrinone have the same effect on the body as testosterone, but they are more stable, so they are not metabolized as quickly. Tetrahydrogestrinone (also called THG or The Clear), the performance-enhancing drug used by track star Marion Jones during the 2000 Sydney Olympics, was considered a “designer steroid” because it was initially undetected in urine tests for doping. After its chemical structure and properties were determined, it was added to the list of banned anabolic steroids in 2004.



A second group of steroid hormones includes the **adrenal cortical steroids**. Three examples of these hormones are **cortisone**, **cortisol**, and **aldosterone**. All of these compounds are synthesized in the outer layer of the adrenal gland. Cortisone and cortisol serve as anti-inflammatory agents and they also regulate carbohydrate metabolism. Aldosterone regulates blood pressure and volume by controlling the concentration of Na^+ and K^+ in body fluids.




KEY CONCEPTS


Lipids

Hydrolyzable Lipids

[1] **Waxes** (29.2)—Esters formed from a long-chain alcohol and a long-chain carboxylic acid.




[2] **Triacylglycerols** (29.3)—Triesters of glycerol with three fatty acids.




[3] **Phospholipids** (29.4)

a. Phosphatidylethanolamine (cephalin)




R, R' = long carbon chain

b. Phosphatidylcholine (lecithin)



R, R' = long carbon chain

c. Sphingomyelin



R = long carbon chain
R' = H or CH₃

Nonhydrolyzable Lipids


[1] **Fat-soluble vitamins** (29.5)—Vitamins A, D, E, and K.

[2] **Eicosanoids** (29.6)—Compounds containing 20 C's derived from arachidonic acid. There are four types: prostaglandins, thromboxanes, prostacyclins, and leukotrienes.

[3] **Terpenes** (29.7)—Lipids composed of repeating 5 C units called isoprene units.

Isoprene unit	Types of terpenes			
	[1] monoterpene	10 C's	[4] sesterterpene	25 C's
	[2] sesquiterpene	15 C's	[5] triterpene	30 C's
	[3] diterpene	20 C's	[6] tetraterpene	40 C's


[4] **Steroids** (29.8)—Tetracyclic lipids composed of three six-membered and one five-membered ring.



PROBLEMS

Waxes, Triacylglycerols, and Phospholipids


- 29.18** One component of lanolin, the wax that coats sheep's wool, is derived from cholesterol and stearic acid. Draw its structure, including the correct stereochemistry at all stereogenic centers.
- 29.19** Draw all possible constitutional isomers of a triacylglycerol formed from one mole each of palmitic, oleic, and linoleic acids. Locate the tetrahedral stereogenic centers in each constitutional isomer.
- 29.20** What is the structure of an optically inactive triacylglycerol that yields two moles of oleic acid and one mole of palmitic acid when hydrolyzed in aqueous acid?
- 29.21** Triacylglycerol **L** yields compound **M** when treated with excess H_2 , Pd-C. Ozonolysis of **L** ($[1] \text{O}_3$; $[2] (\text{CH}_3)_2\text{S}$) affords compounds **N–P**. What is the structure of **L**?



- 29.22** Draw the structure of the following phospholipids:
- A cephalin formed from two molecules of stearic acid.
 - A sphingomyelin formed from palmitic acid.


Prostaglandins

- 29.23** A difficult problem in the synthesis of $\text{PGF}_{2\alpha}$ is the introduction of the OH group at C15 in the desired configuration.
- Label this stereogenic center as *R* or *S*.
 - A well known synthesis of $\text{PGF}_{2\alpha}$ involves reaction of **A** with $\text{Zn}(\text{BH}_4)_2$, a metal hydride reagent similar in reactivity to NaBH_4 , to form two isomeric products, **B** and **C**. Draw their structures and indicate their stereochemical relationship.
 - Suggest a reagent to convert **A** to the single stereoisomer **X**.




Terpenes

29.24 Locate the isoprene units in each compound.




29.25 Classify each terpene in Problem 29.24 (e.g., as a monoterpene, sesquiterpene, etc.).

29.26 An isoprene unit can be thought of as having a head and a tail. The “head” of the isoprene unit is located at the end of the chain nearest the branch point, and the “tail” is located at the end of the carbon chain farthest from the branch point. Most isoprene units are connected together in a “head-to-tail” fashion, as illustrated. For both lycopene (Problem 29.24), and squalene (Figure 29.9), decide which isoprene units are connected in a head-to-tail fashion and which are not.




These two isoprene units are connected in a head-to-tail fashion.

29.27 Draw a stepwise mechanism for the conversion of neryl diphosphate to α -pinene. α -Pinene is a component of pine oil and rosemary oil.




- 29.28** Flexibilene is a terpene isolated from *Sinularia flexibilis*, a soft coral found in the Indian Ocean. Draw a stepwise mechanism for the formation of flexibilene from farnesyl diphosphate and isopentenyl diphosphate. What is unusual about the cyclization that forms the 15-membered ring of flexibilene?



flexibilene


- 29.29** The biosynthesis of lanosterol from squalene has intrigued chemists since its discovery. It is now possible, for example, to synthesize polycyclic compounds from acyclic or monocyclic precursors by reactions that form several C–C bonds in a single reaction mixture.

- a. Draw a stepwise mechanism for the following reaction.
b. Show how **X** can be converted to 16,17-dehydroprogesterone. (*Hint*: See Figure 24.5 for a related conversion.)




Steroids


- 29.30** Draw three-dimensional structures for each decalin derivative.



- 29.31** Draw three-dimensional structures for each alcohol. Label the OH groups as occupying axial or equatorial positions.




- 29.32** Axial alcohols are oxidized faster than equatorial alcohols by PCC and other Cr^{6+} oxidants. Which OH group in each compound is oxidized faster?




- 29.33** (a) Draw a skeletal structure of the anabolic steroid methenolone from the following description. Methenolone contains the tetracyclic steroid skeleton with a carbonyl group at C3, a hydroxyl at C17, a double bond between C1 and C2, and methyl groups bonded to C1, C10, and C13. (b) Add wedges and dashes for all stereogenic centers with the following information: the configuration at C10 is *R*, the configuration at C13 is *S*, the configuration at C17 is *S*, and all substituents at ring fusions are trans to each other. (c) Draw the structure of Primobolan, the product formed when methenolone is treated with $\text{CH}_3(\text{CH}_2)_5\text{COCl}$ and pyridine. Primobolan is an anabolic steroid that can be taken orally or by injection and has been used illegally by well-known Major League Baseball players.

29.34 Draw a three-dimensional representation for androsterone.




- 29.35 a. Draw a three-dimensional structure for the following steroid.
 b. What is the structure of the single stereoisomer formed by reduction of this ketone with H_2 , Pd-C? Explain why only one stereoisomer is formed.




- 29.36 Draw the products formed when cholesterol is treated with each reagent. Indicate the stereochemistry around any stereogenic centers in the product.
- a. CH_3COCl c. PCC e. $[\text{1}] \text{BH}_3 \cdot \text{THF}$; $[\text{2}] \text{H}_2\text{O}_2, ^-\text{OH}$
 b. H_2 , Pd-C d. oleic acid, H^+

Challenge Problems


29.37 Draw a stepwise mechanism for the following conversion, which forms camphene. Camphene is a component of camphor and citronella oils.



29.38 Draw a stepwise mechanism for the following reaction.



29.39 Farnesyl diphosphate is cyclized to sesquiterpene **A**, which is then converted to the bicyclic product epi-aristolochene. Write a stepwise mechanism for both reactions.



30

Synthetic Polymers

- 30.1 Introduction
- 30.2 Chain-growth polymers—
Addition polymers
- 30.3 Anionic polymerization of epoxides
- 30.4 Ziegler–Natta catalysts and polymer stereochemistry
- 30.5 Natural and synthetic rubbers
- 30.6 Step-growth polymers—
Condensation polymers
- 30.7 Polymer structure and properties
- 30.8 Green polymer synthesis
- 30.9 Polymer recycling and disposal



Polyethylene terephthalate (PET) is a synthetic polymer formed by the reaction of ethylene glycol ($\text{HOCH}_2\text{CH}_2\text{OH}$) and terephthalic acid. Because PET is lightweight and impervious to air and moisture, it is commonly used for transparent soft drink containers. PET is also used to produce synthetic fibers, sold under the trade name of Dacron. Of the six most common synthetic polymers, PET is the most easily recycled, in part because beverage bottles that bear the recycling code "1" are composed almost entirely of PET. Recycled polyethylene terephthalate is used for fleece clothing and carpeting. In Chapter 30, we learn about the preparation and properties of synthetic polymers like polyethylene terephthalate.

Chapter 30 discusses polymers, large organic molecules composed of repeating units—called **monomers**—that are covalently bonded together. Polymers occur naturally, as in the polysaccharides and proteins of Chapters 27 and 28, respectively, or they are synthesized in the laboratory.

This chapter concentrates on **synthetic polymers**, and expands on the material already presented in Chapters 15 and 22. Thousands of synthetic polymers have now been prepared. While some exhibit properties that mimic naturally occurring compounds, many others have unique properties. Although all polymers are large molecules, the size and branching of the polymer chain and the identity of the functional groups all contribute to determining an individual polymer's properties, thus making it suited for a particular product.

30.1 Introduction

A **polymer** is a large organic molecule composed of repeating units—called **monomers**—that are covalently bonded together. The word *polymer* is derived from the Greek words *poly* + *meros* meaning “many parts.”

Polymerization is the joining together of monomers to make polymers.

Synthetic polymers are perhaps more vital to the fabric of modern society than any other group of compounds prepared in the laboratory. Nylon backpacks and polyester clothing, car bumpers and CD cases, milk jugs and grocery bags, artificial heart valves and condoms—all these products and innumerable others are made of synthetic polymers. Since 1976, the U.S. production of synthetic polymers has exceeded its steel production. Figure 30.1 illustrates several consumer products and the polymers from which they are made.

Synthetic polymers can be classified as chain-growth or step-growth polymers.

- **Chain-growth polymers, also called addition polymers, are prepared by chain reactions.** These compounds are formed by adding monomers to the growing end of a polymer chain. The conversion of vinyl chloride to poly(vinyl chloride) is an example of chain-growth polymerization. These reactions were introduced in Section 15.14.






Figure 30.1 Polymers in some common consumer products



- We are surrounded by synthetic polymers in our daily lives. This cyclist rides on synthetic rubber tires, drinks from a polyethylene water bottle, wears a protective Lexan helmet and goggles, and uses a lightweight nylon backpack.

- **Step-growth polymers, also called condensation polymers, are formed when monomers containing two functional groups come together and lose a small molecule such as H₂O or HCl.** In this method, any two reactive molecules can combine, so the monomer is not necessarily added to the end of a growing chain. Step-growth polymerization is used to prepare polyamides and polyesters, as discussed in Section 22.16.




In contrast to many of the organic molecules encountered in Chapters 1–26, which have molecular weights much less than 1000 grams per mole (g/mol), polymers generally have high molecular weights, ranging from 10,000 to 1,000,000 grams per mole (g/mol). Synthetic polymers are really mixtures of individual polymer chains of varying lengths, so the reported molecular weight is an average value based on the average size of the polymer chain.

By convention, we often simplify the structure of a polymer by placing brackets around the repeating unit that forms the chain, as shown in Figure 30.2.

Problem 30.1 Give the shorthand structures of poly(vinyl chloride) and nylon 6,6 in Section 30.1.

30.2 Chain-Growth Polymers—Addition Polymers


Chain-growth polymerization is a chain reaction that converts an organic starting material, usually an alkene, to a polymer via a reactive intermediate—a radical, cation, or anion.



- The alkene can be ethylene (CH₂=CH₂) or a derivative of ethylene (CH₂=CHZ or CH₂=CZ₂).
- The substituent Z (in part) determines whether radicals, cations, or anions are formed as intermediates.
- An initiator—a radical, cation, or anion—is needed to begin polymerization.
- Since chain-growth polymerization is a chain reaction, the mechanism involves initiation, propagation, and termination (Section 15.4).





Figure 30.2

Drawing a polymer in a shorthand representation



In most chain-growth polymerizations, an initiator adds to the carbon–carbon double bond of one monomer to form a reactive intermediate, which then reacts with another molecule of monomer to build the chain. Polymerization of $\text{CH}_2=\text{CHZ}$ results in a carbon chain having the Z substituents on every other carbon atom.

Problem 30.2 What polymer is formed by chain-growth polymerization of each monomer?




30.2A Radical Polymerization

Radical polymerization of alkenes was first discussed in Section 15.14, and is included here to emphasize its relationship to other methods of chain-growth polymerization. The initiator is often a peroxy radical ($\text{RO}\cdot$), formed by cleavage of the weak O–O bond in an organic peroxide, ROOR . Mechanism 30.1 is written with styrene ($\text{CH}_2=\text{CHPh}$) as the starting material.




Mechanism 30.1 Radical Polymerization of $\text{CH}_2=\text{CHPh}$

Part [1] Initiation: Formation of a carbon radical in two steps




- Homolysis of the weak O–O bond of the peroxide forms $\text{RO}\cdot$, which then adds to a molecule of monomer to form a carbon radical.

Part [2] Propagation: Growth of the polymer chain by C–C bond formation



- In Step [3], the carbon radical formed during initiation adds to another alkene molecule to form a new C–C bond and another carbon radical. Addition gives the more substituted carbon radical—that is, the **unpaired electron is always located on the carbon atom having the phenyl substituent**.
- Step [3] occurs repeatedly, thus growing the polymer chain.

Part [3] Termination: Removal of radicals by formation of a σ bond




- To terminate the chain, two radicals can combine to form a stable bond, thus ending the polymerization process.

Radical polymerization of $\text{CH}_2=\text{CHZ}$ is favored by Z substituents that stabilize a radical by electron delocalization. Each addition step occurs to put the intermediate radical on the carbon bearing the Z substituent. With styrene as the starting material, the intermediate radical is benzylic and highly resonance stabilized. Figure 30.3 shows several monomers used in radical polymerization reactions.

Figure 30.3


Monomers used in radical polymerization reactions




ethylene




vinyl chloride



styrene



vinyl acetate




Problem 30.3 What polymer is formed by the radical polymerization of each monomer?

- a. $\text{CH}_2 = \text{C}(\text{CH}_3)\text{CO}_2\text{CH}_3$ b. $\text{CH}_2 = \text{C}(\text{CH}_3)\text{CN}$

Problem 30.4 Draw the mechanism for the radical polymerization of vinyl acetate ($\text{CH}_2 = \text{CHOCOCH}_3$) using $(\text{CH}_3)_3\text{CO}-\text{OC}(\text{CH}_3)_3$ as the initiator.


Chain termination can occur by radical coupling, as shown in Mechanism 30.1. Chain termination can also occur by **disproportionation**, a process in which a hydrogen atom is transferred from one polymer radical to another, forming a new C–H bond on one polymer chain, and a double bond on the other.



30.2B Chain Branching

The choice of reaction parameters greatly affects the properties of a synthetic polymer. In Section 15.14, we learned that there are two common types of polyethylene—**high-density polyethylene (HDPE)** and **low-density polyethylene (LDPE)**. High-density polyethylene, which consists of long chains of CH_2 groups joined together in a linear fashion, is strong and hard because the linear chains pack well, resulting in strong van der Waals interactions. Low-density polyethylene, on the other hand, consists of long carbon chains with many branches along the chain. Branching prohibits the chains from packing well, so LDPE has weaker intermolecular interactions, making it a much softer, pliable material.

HDPE is used in milk containers and water jugs, whereas LDPE is used in plastic bags and insulation.




Branching occurs when a radical on one growing polyethylene chain abstracts a hydrogen atom from a CH₂ group in another polymer chain, as shown in Mechanism 30.2. The new 2° radical then continues chain propagation by adding to another molecule of ethylene, thus forming a branch point.




Mechanism 30.2 Forming Branched Polyethylene During Radical Polymerization

Step [1] Abstraction of a hydrogen atom




- Abstraction of a hydrogen atom from an existing polymer chain forms a 2° radical in the middle of the polymer chain.

Step [2] Growth of the polymer chain forming a branch point



- Addition of the radical to another molecule of ethylene forms a new radical and a branch point along the polymer chain. Step [2] occurs repeatedly, and a long branch grows off the original polymer chain.

Problem 30.5 Explain why radical polymerization of styrene forms branched chains with 4° carbons as in **A**, but none with 3° carbons as in **B**.



30.2C Ionic Polymerization

Chain-growth polymerization can also occur by way of cationic or anionic intermediates. **Cationic polymerization is an example of electrophilic addition to an alkene involving carbocations.** Cationic polymerization occurs with alkene monomers that have substituents capable of stabilizing intermediate carbocations, such as alkyl groups or other electron-donor groups. The initiator is an electrophile such as a proton source or Lewis acid.

Mechanism 30.3 illustrates cationic polymerization of the general monomer CH₂=CHZ using BF₃·H₂O, the Lewis acid–base complex formed from BF₃ and H₂O, as the initiator.


Since cationic polymerization involves carbocations, addition follows Markovnikov's rule to form the more stable, more substituted carbocation. Chain termination can occur by a variety of pathways, such as loss of a proton to form an alkene. Examples of alkene monomers that undergo cationic polymerization are shown in Figure 30.4.

Problem 30.6 Explain why cationic polymerization is an effective method of polymerizing CH₂=C(CH₃)₂ but not CH₂=CH₂.




Mechanism 30.3 Cationic Polymerization of $\text{CH}_2=\text{CHZ}$

Part [1] Initiation: Formation of a carbocation




- Electrophilic addition of H^+ from $\text{BF}_3 \cdot \text{H}_2\text{O}$ forms a carbocation.

Part [2] Propagation: Growth of the polymer chain by C–C bond formation



- In Step [3], the carbocation adds to another alkene molecule to form a new C–C bond. Addition always forms a carbocation stabilized by an electron-donating Z substituent.
- Step [3] occurs repeatedly, thus growing the polymer chain.

Part [3] Termination: Loss of a proton




- Loss of a proton forms a new π bond to terminate the chain.

Although alkenes readily react with electron-deficient radicals and electrophiles, alkenes do not generally react with anions and other nucleophiles. Consequently, **anionic polymerization takes place only with alkene monomers that contain electron-withdrawing groups** such as COR, COOR, or CN, which can stabilize an intermediate negative charge. The initiator is a strong nucleophile, such as an organolithium reagent, RLi. Mechanism 30.4 illustrates anionic polymerization of the general monomer $\text{CH}_2=\text{CHZ}$.




Mechanism 30.4 Anionic Polymerization of $\text{CH}_2=\text{CHZ}$

Part [1] Initiation: Formation of a carbanion




- Nucleophilic addition of RLi forms a carbanion stabilized by an electron-withdrawing group Z.

Part [2] Propagation: Growth of the polymer chain by C–C bond formation



- In Step [2], the carbanion adds to another alkene molecule to form a new C–C bond. Addition always forms a carbanion adjacent to the Z substituent.
- Step [2] occurs repeatedly, thus growing the polymer chain.

Part [3] Termination: Addition of a proton source to stop the chain




- Addition of H_2O (or another electrophile) terminates the chain by a Brønsted–Lowry acid–base reaction.

In contrast to other types of chain-growth polymerization, there are no efficient methods of terminating the chain mechanism in anionic polymerization. The reaction continues until all the initiator and monomer have been consumed, so that the end of each polymer chain contains

Figure 30.4

Common polymers formed by ionic chain-growth polymerization



- A chain-growth polymer is named by adding the prefix *poly* to the name of the monomer from which it is made. When the name of the monomer contains two words, this name is enclosed in parentheses and preceded by the prefix *poly*.

a carbanion (Step [2] in Mechanism 30.4). Anionic polymerization is often called **living polymerization** because polymerization will begin again if more monomer is added at this stage. To terminate anionic polymerization an electrophile such as H_2O or CO_2 must be added. Examples of alkene monomers that undergo anionic polymerization are shown in Figure 30.4.

Problem 30.7 Which method of ionic polymerization—cationic or anionic—is preferred for each monomer? Explain your choices.


- a. $\text{CH}_2=\text{C}(\text{CH}_3)\text{COOCH}_3$ b. $\text{CH}_2=\text{CHCH}_3$ c. $\text{CH}_2=\text{CHOC}(\text{CH}_3)_3$ d. $\text{CH}_2=\text{CHCOCH}_3$

Problem 30.8 Draw a stepwise mechanism for the conversion of acrylonitrile ($\text{CH}_2=\text{CHC}\equiv\text{N}$) to polyacrylonitrile, $-\text{[CH}_2\text{CHC}\equiv\text{N}]_n-$, using butyllithium (BuLi) as the initiator and CO_2 as the electrophile to terminate the chain.

Problem 30.9 Explain why styrene ($\text{CH}_2=\text{CHPh}$) can be polymerized to polystyrene by all three methods of chain-growth polymerization.

30.2D Copolymers


All polymers discussed thus far are **homopolymers**, because they have been prepared by the polymerization of a single monomer. **Copolymers, on the other hand, are polymers prepared by joining two or more monomers (X and Y) together.**



- An *alternating copolymer* is formed when X and Y alternate regularly along the chain.
- A *random copolymer* is formed when X and Y are randomly distributed along the chain.

The structure of the copolymer depends on the relative amount and reactivity of **X** and **Y**, as well as the conditions used for polymerization.

Several copolymers are commercially important and used in a wide range of consumer products. For example, the copolymer of vinyl chloride and vinylidene chloride forms **Saran**, the film used in the well-known plastic food wrap. Copolymerization of 1,3-butadiene and styrene forms **styrene-butadiene rubber (SBR)**, the polymer used almost exclusively in automobile tires.



Problem 30.10 Draw the alternating copolymer formed from each set of monomers.

- a. $\text{CH}_2=\text{CHPh}$ and $\text{CH}_2=\text{CHCN}$ b. $\text{F}_2\text{C}=\text{CFCl}$ and $\text{CH}_2=\text{CF}_2$


Problem 30.11 Draw the mechanism for the radical copolymerization of $\text{CH}_2=\text{CHCH}=\text{CH}_2$ and $\text{CH}_2=\text{CHPh}$ to form styrene-butadiene rubber, $-\text{[CH}_2\text{CH}=\text{CHCH}_2\text{CH}_2\text{CHPh}]_n-$.

30.3 Anionic Polymerization of Epoxides

Alkene monomers are the most common starting materials in chain-growth polymerizations, but epoxides can also serve as starting materials, forming **polyethers**. The strained three-membered ring of an epoxide is readily opened with a nucleophile (such as OH^- or OR^-) to form an alkoxide, which can then ring open another epoxide monomer to build the polymer chain. Unlike the other methods of chain-growth polymerization that join monomers together with C–C bonds, this process forms **new C–O bonds** in the polymer backbone.


For example, the ring opening of ethylene oxide with a OH^- initiator affords an alkoxide nucleophile, which propagates the chain by reacting with more ethylene oxide. This process yields

poly(ethylene glycol), PEG, a polymer used in lotions and creams. The many C–O bonds in these polymers make them highly water soluble.




The ring opening of epoxides with nucleophiles was first discussed in Section 9.15.

Under anionic conditions, the ring opening follows an $\text{S}_{\text{N}}2$ mechanism. Thus, the ring opening of an unsymmetrical epoxide occurs at the more accessible, less substituted carbon.




Problem 30.12 What polymer is formed by anionic polymerization of each monomer?



30.4 Ziegler–Natta Catalysts and Polymer Stereochemistry

Polymers prepared from monosubstituted alkene monomers ($\text{CH}_2=\text{CHZ}$) can exist in three different configurations, called **isotactic**, **syndiotactic**, and **atactic**:



- An *isotactic* polymer has all Z groups on the same side of the carbon backbone.
- A *syndiotactic* polymer has the Z groups alternating from one side of the carbon chain to the other.
- An *atactic* polymer has the Z groups oriented randomly along the polymer chain.

The more regular arrangement of the Z substituents in isotactic and syndiotactic polymers allows them to pack together better, making the polymer stronger and more rigid. In contrast, the chains of an atactic polymer tend to pack less closely together, resulting in a lower melting, softer polymer. Radical polymerization often affords an atactic polymer, but the particular reaction conditions can greatly affect the stereochemistry of the polymer formed.

In 1953, Karl Ziegler and Giulio Natta developed a new method of polymerizing alkene monomers using a metal catalyst to promote chain-growth polymerization. These catalysts, now called **Ziegler–Natta catalysts**, offer two advantages over other methods of chain-growth polymerization.

- The stereochemistry of the polymer is easily controlled. Polymerization affords isotactic, syndiotactic, or atactic polymers depending on the catalyst.
- Long, linear chains of polymer are prepared without significant branching. Radicals are not formed as reactive intermediates, so intermolecular hydrogen abstraction, which leads to chain branching, does not occur.


Ziegler and Natta received the 1963 Nobel Prize in Chemistry for their pioneering work on polymerization catalysts.

Many different Ziegler–Natta catalysts are used for polymerization, but most consist of an organoaluminum compound such as $(\text{CH}_3\text{CH}_2)_2\text{AlCl}$ and TiCl_4 , a Lewis acid. The active catalyst is thought to be an alkyl titanium compound, formed by transfer of an ethyl group from $(\text{CH}_3\text{CH}_2)_2\text{AlCl}$ to TiCl_4 , although many mechanistic details are not known with certainty. It is generally agreed that the alkene monomer coordinates to an alkyl titanium complex, and then inserts into the Ti–C bond to form a new carbon–carbon bond, as shown in Mechanism 30.5.




Mechanism 30.5 Ziegler–Natta Polymerization of $\text{CH}_2=\text{CH}_2$

Step [1] Formation of the catalyst



- Reaction of the organoaluminum compound with TiCl_4 forms a Ziegler–Natta catalyst with a Ti–C bond.

Steps [2] and [3] Growth of the polymer chain by C–C bond formation



- An alkene monomer coordinates with the titanium catalyst in Step [2].
- Insertion of $\text{CH}_2=\text{CH}_2$ into the Ti–C bond forms a new C–C bond in Step [3].
- Repeating Steps [2] and [3] again and again yields the long polymer chain.

The Ziegler–Natta polymerization of ethylene forms **high-density polyethylene, HDPE**, composed of long linear carbon chains that pack closely together, forming a rigid polymer. By using specialized manufacturing techniques that force the polymer chains to pack closely in the solid phase as a set of linear extended chains, this material is converted to ultra high-density polyethylene, a synthetic organic material stronger than steel.

Recently developed Ziegler–Natta polymerizations utilize zirconium complexes that are soluble in the reaction solvents typically used, and so they are **homogeneous catalysts**. Reactions that use these soluble catalysts are called **coordination polymerizations**.

30.5 Natural and Synthetic Rubbers




Dyneema, the strongest fabric known, is made of ultra high-density polyethylene (Section 30.4), and is used for ropes, nets, bulletproof vests, and crash helmets.

Locating isoprene units in terpenes was discussed in Section 29.7.




Natural rubber is obtained from latex that oozes from cuts made to the bark of the rubber tree. Waterproof latex is the rubber tree's natural protection, exuded in response to an injury. Although rubber was produced exclusively in Brazil until the late 1800s, today most of the world's rubber comes from plantations in Southeast Asia, Sri Lanka, and Indonesia.

Natural rubber is a terpene composed of repeating isoprene units, in which all the double bonds have the Z configuration. Because natural rubber is a hydrocarbon, it is water insoluble, and thus useful for waterproofing. The Z double bonds cause bends and kinks in the polymer chain, making it a soft material.




The polymerization of isoprene under radical conditions forms a stereoisomer of natural rubber called **gutta-percha**, in which all the double bonds have the E configuration. Gutta-percha is also a naturally occurring polymer, although considerably less common than its Z stereoisomer. Polymerization of isoprene with a Ziegler–Natta catalyst forms natural rubber with all the double bonds having the desired Z configuration.



Natural rubber is too soft to be a useful material for most applications. Moreover, when natural rubber is stretched, the chains become elongated and slide past each other until the material pulls apart. In 1839, Charles Goodyear discovered that mixing hot rubber with sulfur produced a stronger and more elastic material. This process, called **vulcanization**, results in cross-linking of the hydrocarbon chains by disulfide bonds, as shown in Figure 30.5. When the polymer is stretched, the chains no longer can slide past each other and tearing does not occur. Vulcanized rubber is an **elastomer**, a polymer that stretches when stressed but then returns to its original shape when the stress is alleviated.

Figure 30.5
Vulcanized rubber




- Vulcanized rubber contains many disulfide bonds that cross-link the hydrocarbon chains together.

Gutta-percha, a much harder material than natural rubber obtained from latex, is used in golf ball casings.


The degree of cross-linking affects the rubber's properties. Harder rubber used for automobile tires has more cross-linking than the softer rubber used for rubber bands.

Other synthetic rubbers can be prepared by the polymerization of different 1,3-dienes using Ziegler–Natta catalysts. For example, the polymerization of 1,3-butadiene affords (*Z*)-poly(1,3-butadiene), and the polymerization of 2-chloro-1,3-butadiene yields neoprene, a polymer used in wet suits and tires.



Problem 30.13 Assign the *E* or *Z* configuration to the double bonds in neoprene. Draw a stereoisomer of neoprene in which all the double bonds have the opposite configuration.

Problem 30.14 The polymerization of $\text{CH}_2=\text{CHCH}=\text{CH}_2$ under radical conditions affords products **A** and **B**. Draw a mechanism that accounts for their formation.



30.6 Step-Growth Polymers—Condensation Polymers


Step-growth polymers, the second major class of polymers, are formed when monomers containing two functional groups come together and lose a small molecule such as H_2O or HCl . Commercially important step-growth polymers include:

- Polyamides
- Polyesters
- Polyurethanes
- Polycarbonates
- Epoxy resins


30.6A Polyamides

Nylon 6,6 is used in many products including parachutes and clothing.

Nylons are polyamides formed by step-growth polymerization. In Section 22.16A, we learned that **nylon 6,6** can be prepared by the reaction of a diacid chloride and a diamine. Nylon 6,6 can also be prepared by heating adipic acid and 1,6-diaminohexane. A Brønsted–Lowry acid–base reaction forms a diammonium salt, which loses H_2O at high temperature. In both methods, each starting material has two identical functional groups.




Nylon 6, trade name **Perlon**, is used to make rope and tire cord.




Armadillo bicycle tires reinforced with Kevlar are hard to pierce with sharp objects, so a cyclist rarely gets a flat tire.

Kevlar is a polyamide formed from terephthalic acid and 1,4-diaminobenzene. The aromatic rings of the polymer backbone make the chains less flexible, resulting in a very strong material. Kevlar is light in weight compared to other materials that are similar in strength, so it is used in many products, such as bulletproof vests, army helmets, and the protective clothing used by firefighters.




Problem 30.15

What polyamide is formed from each monomer or pair of monomers?




30.6B Polyesters


Polyesters are formed by step-growth polymerization using nucleophilic acyl substitution reactions, as we learned in Section 22.16B. For example, the reaction of terephthalic acid and ethylene glycol forms **polyethylene terephthalate (PET)**, the chapter-opening molecule.



Although PET is a very stable material, some polyesters are more readily hydrolyzed to carboxylic acids and alcohols in aqueous medium, making them suited for applications in which slow degradation is useful. For example, copolymerization of glycolic acid and lactic acid forms a copolymer used by surgeons in dissolving sutures. Within weeks, the copolymer is hydrolyzed to the monomers from which it was prepared, which are metabolized readily by the body. These sutures are used internally to hold tissues together while healing and scar formation occur.




Problem 30.16 Polyethylene terephthalate is also prepared by the transesterification of dimethyl terephthalate with ethylene glycol. Draw the mechanism for this nucleophilic acyl substitution.




Problem 30.17 The first synthetic fibers were prepared by the step-growth polymerization of $\text{HOOC}(\text{CH}_2)_4\text{COOH}$ and $\text{HOCH}_2\text{CH}_2\text{OH}$. Draw the structure of this polymer and suggest reasons why it is less suitable than either nylon 6,6 or PET for use in consumer products.

30.6C Polyurethanes


A **urethane** (also called a **carbamate**) is a compound that contains a carbonyl group bonded to both an OR group and an NHR (or NR_2) group (Section 28.6). Urethanes are prepared by the nucleophilic addition of an alcohol to the carbonyl group of an **isocyanate**, $\text{RN}=\text{C}=\text{O}$.



Polyurethanes are polymers formed by the reaction of a diisocyanate and a diol.




A well-known polyurethane is **spandex**, a strong and flexible polymer that illustrates how the macroscopic properties of a polymer depend on its structure at the molecular level. Spandex was first used in women's corsets, girdles, and support hose, but is now routinely used in both men's and women's active wear. Spandex is strong and lends "support" to the wearer, but it also stretches. Spandex is lighter in weight than many other elastic polymers, and it does not break down when exposed to perspiration and detergents. On the molecular level, it has rigid regions that are joined together by soft, flexible segments. The flexible regions allow the polymer to expand and then recover its original shape. The rigid regions strengthen the polymer.




30.6D Polycarbonates

Although it is not acutely toxic, bisphenol A (BPA) mimics the body's own hormones and disrupts normal endocrine functions. Concern over low-dose exposure by infants has led to a voluntary phase-out of BPA-based polymers in infant formula packaging.


A **carbonate** is a compound that contains a carbonyl group bonded to two OR groups. Carbonates can be prepared by the reaction of phosgene ($\text{Cl}_2\text{C}=\text{O}$) with two equivalents of an alcohol (ROH).



Polycarbonates are formed from phosgene and a diol. The most widely used polycarbonate is **Lexan**, a lightweight, transparent material that is formed from phosgene and bisphenol A, and used in bike helmets, goggles, catcher's masks, and bulletproof glass.




Problem 30.18 Lexan can also be prepared by the acid-catalyzed reaction of diphenyl carbonate with bisphenol A. Draw a stepwise mechanism for this process.




30.6E Epoxy Resins

Epoxy resins represent a class of step-growth polymer familiar to anyone who has used “epoxy” to glue together a broken object. An epoxy resin consists of two components: a fluid **prepolymer** composed of short polymer chains with reactive epoxides on each end, and a **hardener**, usually a diamine or triamine that ring opens the epoxides and cross-links the chains together. The prepolymer is formed by reacting two difunctional monomers, bisphenol A and epichlorohydrin.




Bisphenol A has two nucleophilic OH groups, while epichlorohydrin has polar C–O and C–Cl bonds that can react with two different nucleophiles. The general reaction of epichlorohydrin with nucleophiles is given in the accompanying equation. Nucleophilic attack on the strained epoxide ring affords an alkoxide that displaces chloride by an intramolecular $\text{S}_{\text{N}}2$ reaction, forming a new epoxide. Ring opening with a second nucleophile gives a 2° alcohol.



When bisphenol A is treated with excess epichlorohydrin, this stepwise process continues until all the phenolic OH groups have been used in ring-opening reactions, leaving epoxy groups on both ends of the polymer chains. This constitutes the fluid **prepolymer**, as shown in Figure 30.6.

Figure 30.6


Formation of an epoxy resin from a prepolymer and a hardening agent



When the prepolymer is mixed with a diamine or triamine (the **hardener**), the reactive epoxide rings can be opened by the nucleophilic amino groups to cross-link polymer chains together, causing the polymer to harden. A wide range of epoxy resins is commercially prepared by this process, making them useful for adhesives and coatings. The longer and more extensively cross-linked the polymer chains, the harder the resin.


Problem 30.19

(a) Draw the structure of the prepolymer **A** formed from 1,4-dihydroxybenzene and excess epichlorohydrin. (b) Draw the structure of the cross-linked polymer **B** formed when **A** is treated with $\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ as the hardening agent.

**30.7 Polymer Structure and Properties**

While the chemistry of polymer synthesis can be explained by the usual themes of organic reactions, the large size of polymer molecules gives them some unique physical properties compared to small organic molecules.

Linear and branched polymers do not form crystalline solids because their long chains prevent efficient packing in a crystal lattice. Most polymer chains have **crystalline regions** and **amorphous regions**:



- **Ordered crystalline regions**, called **crystallites**, are places where sections of the polymer chain lie in close proximity and are held together by intermolecular interactions. Ordered regions of polyethylene, $-\text{[CH}_2\text{CH}_2\text{]}_n-$, are held together by van der Waals interactions, whereas ordered regions of nylon chains are held together by intermolecular hydrogen bonding.
- **Amorphous regions** are places where the polymer chains are randomly arranged, resulting in weak intermolecular interactions.

Crystalline regions impart toughness to a polymer, while amorphous regions impart flexibility. The greater the crystallinity of a polymer—that is, the larger the percentage of ordered regions—the harder the polymer. Branched polymers are generally more amorphous and, since branching prevents chains from packing closely, they are softer, too.

Two temperatures, T_g and T_m , often characterize a polymer's behavior on heating:

- T_g , the glass transition temperature, is the temperature at which a hard amorphous polymer becomes soft.
- T_m , the melt transition temperature, is the temperature at which the crystalline regions of the polymer melt to become amorphous. More ordered polymers have higher T_m values.

Thermoplastics are polymers that can be melted and then molded into shapes that are retained when the polymer is cooled. Although they have high T_g values and are hard at room temperature, heating causes individual polymer chains to slip past each other, causing the material to soften. Polyethylene terephthalate and polystyrene are thermoplastic polymers.

Thermosetting polymers are complex networks of cross-linked polymers. Thermosetting polymers are formed by chemical reactions that occur when monomers are heated together to form a network of covalent bonds. Thermosetting polymers cannot be re-melted to form a liquid phase, because covalent bonds hold the network together. **Bakelite**, a thermosetting polymer prepared from phenol (PhOH) and formaldehyde ($\text{H}_2\text{C}=\text{O}$) in the presence of a Lewis acid, is formed by electrophilic aromatic substitution reactions. Since formaldehyde is a reactive electrophile and phenol contains a strongly electron-donating OH group, substitution occurs at all ortho and para positions to the OH group, resulting in a highly cross-linked polymer, shown in Figure 30.7.


Problem 30.20 Draw a stepwise mechanism for Step [2] in Figure 30.7 using AlCl_3 as the Lewis acid catalyst.

Sometimes a polymer is too stiff and brittle to be useful in many applications. In this case, a low molecular weight compound called a **plasticizer** is added to soften the polymer and give it flexibility. The plasticizer interacts with the polymer chains, replacing some of the intermolecular interactions between the polymer chains. This lowers the crystallinity of the polymer, making it more amorphous and softer.

Dibutyl phthalate is a plasticizer added to the poly(vinyl chloride) used in vinyl upholstery and garden hoses. Since plasticizers are more volatile than the high molecular weight polymers, they


Figure 30.7

The synthesis of Bakelite from phenol and formaldehyde



- Bakelite, the first totally synthetic polymeric material, was patented by Leo Baekeland in 1910. Bowling balls are made of Bakelite.

slowly evaporate with time, making the polymer brittle and easily cracked. Plasticizers like dibutyl phthalate that contain hydrolyzable functional groups are also slowly degraded by chemical reactions.



30.8 Green Polymer Synthesis

One hundred fifty years ago there were no chemical manufacturing plants and no synthetic polymers, and petroleum had little value. Synthetic polymers have transformed the daily lives of many in the modern world, but not without a hefty price. Polymer synthesis and disposal have a tremendous impact on the environment, creating two central issues:

- **Where do polymers come from?** What raw materials are used for polymer synthesis and what environmental consequences result from their manufacture?
- **What happens to polymers once they are used?** How does polymer disposal affect the environment, and what can be done to minimize its negative impact?

30.8A Environmentally Friendly Polymer Synthesis—The Feedstock

In Chapter 12, you were introduced to **green chemistry**, the use of environmentally benign methods to synthesize compounds. Given the billions of pounds of polymers manufactured worldwide each year, there is an obvious need for methods that minimize the environmental impact.

To date, green polymer synthesis has been approached in a variety of ways:

- **Using starting materials that are derived from renewable sources, rather than petroleum. The starting materials for an industrial process are often called the chemical feedstock.**
- **Using safer, less toxic reagents that form fewer by-products.**


Recall from Section 4.7 that 3% of a barrel of crude oil is used as the feedstock for chemical synthesis.

- **Carrying out reactions in the absence of solvent or in aqueous solution (instead of an organic solvent).**

Until recently, **the feedstock for all polymer synthesis has been petroleum**; that is, the monomers for virtually all polymer syntheses are made from crude oil, a nonrenewable raw material. As an example, nylon 6,6 is prepared industrially from adipic acid [$\text{HOOC}(\text{CH}_2)_4\text{COOH}$] and 1,6-diaminohexane [$\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$], both of which originate from benzene, a product of petroleum refining (Figure 30.8).

Besides beginning with a nonrenewable chemical feedstock, adipic acid synthesis has other problems. The use of benzene, a carcinogen and liver toxin, is undesirable, especially in a large-scale reaction. Moreover, oxidation with HNO_3 in Step [3] produces N_2O as a by-product. N_2O depletes ozone in the stratosphere in much the same way as the CFCs discussed in Chapter 15. In addition, N_2O also absorbs thermal energy from the earth's surface like CO_2 , and may therefore contribute to global warming, as discussed in Section 4.14.


As a result, several research groups are working to develop new methods of monomer synthesis that begin with renewable, more environmentally friendly raw materials and produce fewer hazardous by-products. As an example, chemists at Michigan State University have devised a two-step synthesis of adipic acid from D-glucose, a monosaccharide available from plant sources. The synthesis uses a genetically altered *E. coli* strain (called a **biocatalyst**) to convert D-glucose to (2Z,4Z)-2,4-hexadienoic acid, which is then hydrogenated to adipic acid. Methods such as this, which avoid starting materials derived from petroleum, are receiving a great deal of attention in the chemical community.



Sorona(R), DuPont's trade name for **poly(trimethylene terephthalate)**, is a large-volume polymer that can now be made at least in part from glucose derived from a renewable plant source such as corn. A biocatalyst converts D-glucose to 1,3-propanediol, which forms poly(trimethylene terephthalate) (PTT) on reaction with terephthalic acid, as shown in Figure 30.9.


In related chemistry, poly(lactic acid) (PLA) is a polymer used in bottles and packaging, and it can also be made into a synthetic fiber (trade name Ingeo) used in clothing and carpets.

Figure 30.8
Synthesis of adipic acid and
1,6-diaminohexane for
nylon 6,6 synthesis




- The synthesis of both monomers needed for nylon 6,6 synthesis begins with benzene, a petroleum product.

Figure 30.9 A swimsuit made (in part) from corn—The synthesis of poly(trimethylene terephthalate) from 1,3-propanediol derived from corn




- Poly(trimethylene terephthalate), sold as Sorona(R) by the DuPont Corporation, is made into fibers used in clothing and other materials. Although 1,3-propanediol, one of the monomers needed for its synthesis, has been prepared from petroleum feedstocks in the past, it is now available from a renewable plant source such as corn.

Poly(lactic acid) is prepared on a large scale by the fermentation of carbohydrates obtained from corn. Fermentation initially yields a cyclic lactone called lactide, derived from two molecules of lactic acid [$\text{CH}_3\text{CH}(\text{OH})\text{CO}_2\text{H}$]. Heating lactide with acid forms poly(lactic acid). PLA is an especially attractive polymer choice, because it readily degrades in a landfill.




30.8B Polymer Synthesis with Less Hazardous Reagents

Other approaches to green polymer synthesis have concentrated on using less hazardous reagents and avoiding solvents. For example, Lexan can now be prepared by the reaction of bisphenol A with diphenyl carbonate [$(\text{PhO})_2\text{C}=\text{O}$] in the absence of solvent. This process avoids the use of phosgene ($\text{Cl}_2\text{C}=\text{O}$, Section 30.6D), an acutely toxic reagent that must be handled with extreme care, as well as the large volume of CH_2Cl_2 typically used as the solvent for the polymerization process.



Problem 30.21 Thermosetting resins similar to Bakelite (Section 30.7) have also been prepared from renewable feedstocks. One method uses cardinol, the major constituent of the liquid obtained from roasted cashew nutshells. What polymer is obtained when cardinol is treated with formaldehyde ($\text{H}_2\text{C}=\text{O}$) in the presence of a proton source?



30.9 Polymer Recycling and Disposal

The same desirable characteristics that make polymers popular materials for consumer products—durability, strength, and lack of reactivity—also contribute to environmental problems. Polymers do not degrade readily, and as a result, billions of pounds of polymers end up in landfills every year. Estimates suggest that synthetic polymers comprise 11% of solid municipal waste, 30% of which comes from packaging materials.

Two solutions to address the waste problem created by polymers are recycling existing polymer types to make new materials, and using biodegradable polymers that will decompose in a finite and limited time span.

30.9A Polymer Recycling

Although thousands of different synthetic polymers have now been prepared, six compounds, called the “**Big Six**,” account for 76% of the synthetic polymers produced in the United States each year. Each polymer is assigned a recycling code (1–6) that indicates its ease of recycling; **the lower the number, the easier to recycle**. Table 30.1 lists these six most common polymers, as well as the type of products made from each recycled polymer.


Recycling begins with sorting plastics by type, shredding the plastics into small chips, and washing the chips to remove adhesives and labels. After the chips are dried and any metal caps or rings are removed, the polymer chips are melted and molded for reuse.

Table 30.1 Recyclable Polymers


Recycling code	Polymer name	Structure	Recycled product
1	PET Polyethylene terephthalate		fleece jackets carpeting plastic bottles
2	HDPE High-density polyethylene		Tyvek insulation sports clothing
3	PVC Poly(vinyl chloride)		floor mats
4	LDPE Low-density polyethylene		trash bags
5	PP Polypropylene		furniture
6	PS Polystyrene		molded trays trash cans

Of the Big Six, only the polyethylene terephthalate (PET) in soft drink bottles and the high-density polyethylene (HDPE) in milk jugs and juice bottles are recycled to any great extent. Since recycled polymers are often still contaminated with small amounts of adhesives and other materials, these recycled polymers are generally not used for storing food or drink products. Recycled HDPE is converted to Tyvek, an insulating wrap used in new housing construction, and recycled PET is used to make fibers for fleece clothing and carpeting. Currently about 23% of all plastics are recycled in the United States.

An alternative recycling process is to re-convert polymers back to the monomers from which they were made, a process that has been successful with acyl compounds that contain C–O or C–N bonds in the polymer backbone. For example, heating polyethylene terephthalate with CH_3OH cleaves the esters of the polymer chain to give ethylene glycol ($\text{HOCH}_2\text{CH}_2\text{OH}$) and dimethyl terephthalate. These monomers then serve as starting materials for more PET. This chemical recycling process is a transesterification reaction that occurs by nucleophilic acyl substitution, as discussed in Chapter 22.



Similarly, treatment of discarded nylon 6 polymer with NH_3 cleaves the polyamide backbone, forming ϵ -caprolactam, which can be purified and re-converted to nylon 6.



Problem 30.22 Why can't chemical recycling—that is, the conversion of polymer to monomers and re-conversion of monomers to polymer—be done easily with HDPE and LDPE?


Problem 30.23 Organic polymers can also be incinerated as a means of disposal. (a) What products are formed on combustion of polyethylene? (b) What products are formed on combustion of polyethylene terephthalate? (c) Are these reactions exothermic or endothermic? (See Sections 6.4 and 29.3 for related reactions.) (d) Propose a reason why HDPE and PET must be separated from poly(vinyl chloride) prior to incineration.

30.9B Biodegradable Polymers

Another solution to the accumulation of waste polymers in landfills is to design and use polymers that are biodegradable.


- **Biodegradable polymers are polymers that can be degraded by microorganisms—bacteria, fungi, or algae—naturally present in the environment.**

Several biodegradable polyesters have now been developed. For example, the **polyhydroxyalkanoates (PHAs)** are polymers of 3-hydroxy carboxylic acids, such as 3-hydroxybutyric acid or 3-hydroxyvaleric acid.



PHAs are bioplastics. PHAs are not synthesized from petroleum products, and they are degraded by soil microorganisms to CO₂ and H₂O.


The two most common PHAs are **polyhydroxybutyrate (PHB)** and a copolymer of **polyhydroxybutyrate** and **polyhydroxyvalerate (PHBV)**. PHAs can be used as films, fibers, and coatings for hot beverage cups made of paper.




Bacteria in the soil readily degrade PHAs, and in the presence of oxygen, the final degradation products are CO₂ and H₂O. The rate of degradation depends on moisture, temperature, and pH. Degradation is slower in enclosed landfills that are lined and covered.

An additional advantage of the polyhydroxyalkanoates is that the polymers can be produced by fermentation. Certain types of bacteria produce PHAs for energy storage when they are grown in glucose solution in the absence of specific nutrients. The polymer forms as discrete granules within the bacterial cell, and it is then removed by extraction to give a white powder that can be melted and modified into a variety of different products.

Biodegradable polyamides have also been prepared from amino acids. For example, aspartic acid can be converted to polyaspartate, abbreviated as **TPA** (thermal polyaspartate). TPA is commonly used as an alternative to poly(acrylic acid), which is used to line the pumps and boilers of wastewater treatment facilities.



Problem 30.24 What polymers are formed from each monomer?




KEY CONCEPTS

Synthetic Polymers

Chain-Growth Polymers—Addition Polymers

[1] Chain-growth polymers with alkene starting materials (30.2)


- General reaction:



- Mechanism—three possibilities, depending in part on the identity of Z:


Type	Identity of Z	Initiator	Comments
[1] radical polymerization	Z stabilizes a radical. Z = R, Ph, Cl, etc.	A source of radicals (ROOR)	Termination occurs by radical coupling or disproportionation. Chain branching occurs.
[2] cationic polymerization	Z stabilizes a carbocation. Z = R, Ph, OR, etc.	H-A or a Lewis acid (BF ₃ + H ₂ O)	Termination occurs by loss of a proton.
[3] anionic polymerization	Z stabilizes a carbanion. Z = Ph, COOR, COR, CN, etc.	An organolithium reagent (R-Li)	Termination occurs only when an acid or other electrophile is added.

[2] Chain-growth polymers with epoxide starting materials (30.3)



- The mechanism is S_N2.
- Ring opening occurs at the less substituted carbon of the epoxide.

Examples of Step-Growth Polymers—Condensation Polymers (30.6)



[Key functional groups are indicated in red.]


Structure and Properties

- Polymers prepared from monomers having the general structure $\text{CH}_2=\text{CHZ}$ can be isotactic, syndiotactic, or atactic depending on the identity of Z and the method of preparation (30.4).
- Ziegler–Natta catalysts form polymers without significant branching. Polymers can be isotactic, syndiotactic, or atactic depending on the catalyst. Polymers prepared from 1,3-dienes have the *E* or *Z* configuration depending on the catalyst (30.4, 30.5).
- Most polymers contain ordered crystalline regions and less ordered amorphous regions (30.7). The greater the crystallinity, the harder the polymer.
- Elastomers are polymers that stretch and can return to their original shape (30.5).
- Thermoplastics are polymers that can be molded, shaped, and cooled such that the new form is preserved (30.7).
- Thermosetting polymers are composed of complex networks of covalent bonds so they cannot be re-melted to form a liquid phase (30.7).


PROBLEMS

Polymer Structure and Properties


30.25 Draw the structure of the polymer formed by chain-growth polymerization of each monomer.



30.26 Draw the structure of the alternating copolymer formed from each pair of monomers.



30.27 What monomer(s) are used to prepare each polymer or copolymer?








30.28 Draw each polymer in Problem 30.27 using the shorthand representation shown in Figure 30.2.

30.29 Draw a short segment of each polymer: (a) isotactic poly(vinyl chloride); (b) syndiotactic polyacrylonitrile; (c) atactic polystyrene.



30.30 Draw the structure of the polymer that results from anionic polymerization of *p*-trichloromethylstyrene ($\text{CCl}_3\text{C}_6\text{H}_4\text{CH}=\text{CH}_2$) using ethylene oxide as the electrophile to terminate the chain.

30.31 Draw the structure of the polymer formed by step-growth polymerization of each monomer or pair of monomers.




30.32 **ABS**, a widely produced copolymer used for high-impact applications like car bumpers and crash helmets, is formed from three monomers—acrylonitrile ($\text{CH}_2=\text{CHCN}$), 1,3-butadiene ($\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$), and styrene ($\text{CH}_2=\text{CHPh}$). Draw a possible structure for ABS.

30.33 Draw the structures of **Quiana** and **Nomex**, two commercially available step-growth polymers formed from the given monomers. Nomex is a strong polymer used in aircraft tires and microwave transformers. Quiana has been used to make wrinkle-resistant fabrics.




30.34 Kevlar (Section 30.6A) is a very stiff polymer because its backbone contains many aromatic rings and its polymer chains are extensively hydrogen bonded to each other. Draw a short segment of two Kevlar chains, and indicate how the chains are hydrogen bonded to each other.

30.35 Explain the differences observed in the T_g and T_m values for each pair of polymers: (a) polyester **A** and PET; (b) polyester **A** and nylon 6,6. (c) How would you expect the T_m value for Kevlar (Section 30.6A) to compare with the T_m value for nylon 6,6? Explain your prediction.




30.36 Explain why diester **A** is now often used as a plasticizer in place of dibutyl phthalate.




Mechanism

30.37 Draw a stepwise mechanism for the polymerization of isoprene to gutta-percha using $(\text{CH}_3)_3\text{CO}-\text{OC}(\text{CH}_3)_3$ as the initiator.



30.38 Cationic polymerization of 3-phenylpropene ($\text{CH}_2=\text{CHCH}_2\text{Ph}$) affords **A** as the major product rather than **B**. Draw a stepwise mechanism to account for this observation.




30.39 Explain why acrylonitrile ($\text{CH}_2=\text{CHCN}$) undergoes cationic polymerization more slowly than 3-butenitrile ($\text{CH}_2=\text{CHCH}_2\text{CN}$).

30.40 Draw a stepwise mechanism for the anionic polymerization of styrene ($\text{CH}_2=\text{CHPh}$) to form polystyrene $-\text{[CH}_2\text{CHPh]}_n-$ using BuLi as the initiator. Use CO_2 as the electrophile that terminates the chain mechanism.


30.41 Although styrene undergoes both cationic and anionic polymerization equally well, one method is often preferred with substituted styrenes. Which method is preferred with each compound? Explain.




30.42 Rank the following compounds in order of increasing ability to undergo anionic chain-growth polymerization.




30.43 In the presence of H_3O^+ , 2-methylpropene oxide undergoes chain-growth polymerization such that nucleophilic attack occurs at the more substituted end of the epoxide. Draw a stepwise mechanism for this process, and explain this regioselectivity.



30.44 Draw a stepwise mechanism for the conversion of dihalide **A** and 1,4-cyclohexanediol to polyether **B** in the presence of AlCl_3 .




30.45 Draw a stepwise mechanism for the reaction of an alcohol with an isocyanate to form a urethane.




Reactions and Synthesis

30.46 Draw the products of each reaction.




30.47 Explain why aqueous NaOH solution can be stored indefinitely in polyethylene bottles, but spilling aqueous base on a polyester shirt or nylon stockings quickly makes a hole.

30.48 What epoxy resin is formed by the following reaction sequence?




30.49 (a) Explain why poly(vinyl alcohol) cannot be prepared by the radical polymerization of vinyl alcohol ($\text{CH}_2=\text{CHOH}$). (b) Devise a stepwise synthesis of poly(vinyl alcohol) from vinyl acetate ($\text{CH}_2=\text{CHOCOCH}_3$). (c) How can poly(vinyl alcohol) be converted to poly(vinyl butyral), a polymer used in windshield safety glass?




30.50 Although 1,3-propanediol ($\text{HOCH}_2\text{CH}_2\text{CH}_2\text{OH}$) can now be prepared from carbohydrate feedstocks (Section 30.8), it can also be prepared from petroleum feedstocks. Devise a synthesis of $\text{HOCH}_2\text{CH}_2\text{CH}_2\text{OH}$ from $\text{CH}_3\text{CH}=\text{CH}_2$, a product of petroleum refining.

30.51 Devise a synthesis of terephthalic acid and ethylene glycol, the two monomers needed for polyethylene terephthalate synthesis, from the given starting materials.





30.52 The reaction of *p*-cresol with $\text{CH}_2=\text{O}$ resembles the reaction of phenol (PhOH) with $\text{CH}_2=\text{O}$, except that the resulting polymer is thermoplastic but not thermosetting. Draw the structure of the polymer formed, and explain why the properties of these two polymers are so different.




Biological Applications


30.53 In addition to glycolic and lactic acids (Section 30.6B), dissolving sutures can also be prepared from each of the following lactone monomers. Draw the structure of the polymer formed from each monomer.



30.54 Compound **A** is a novel poly(ester amide) copolymer that can be used as a bioabsorbable coating for the controlled release of drugs. **A** is a copolymer of four monomers, two of which are amino acids or amino acid derivatives. The body's enzymes recognize the naturally occurring amino acids in the polymer backbone, allowing for controlled enzymatic breakdown of the polymer and steady release of an encapsulated drug. Identify the four monomers used to synthesize **A**; then use Figure 28.2 to name the two amino acids.




30.55 Researchers at Rutgers University have developed biocompatible polymers that degrade into nonsteroidal anti-inflammatory drugs. For example, the reaction of two equivalents of benzyl salicylate and one equivalent of sebacoyl chloride forms a poly(anhydride ester) called PolyAspirin, which hydrolyzes to salicylic acid (an anti-inflammatory agent) and sebacic acid, which is excreted. This technology can perhaps be used for localized drug delivery at specific sites of injury. What is the structure of PolyAspirin?




Challenge Questions

30.56 Melmac, a thermosetting polymer formed from melamine and formaldehyde ($\text{CH}_2=\text{O}$), is used to make dishes and countertops. Draw a stepwise mechanism for the condensation of one mole of formaldehyde with two moles of melamine, which begins the synthesis of Melmac.



- 30.57** Although chain branching in radical polymerizations can occur by intermolecular H abstraction as shown in Mechanism 30.2, chain branching can also occur by intramolecular H abstraction to form branched polyethylene that contains butyl groups as branches.




















- Draw a stepwise mechanism that illustrates which H must be intramolecularly abstracted to form butyl substituents.
 - Suggest a reason why the abstraction of this H is more facile than the abstraction of other H's.
- 30.58** The reaction of urea $[(\text{NH}_2)_2\text{C}=\text{O}]$ and formaldehyde $(\text{CH}_2=\text{O})$ forms a highly cross-linked polymer used in foams. Suggest a structure for this polymer. [Hint: Examine the structures of Bakelite (Figure 30.7) and Melmac (Problem 30.56).]












This page intentionally left blank

Appendix

A

pK_a Values for Selected Compounds

Compound	pK _a	Compound	pK _a
HI	-10		4.0
HBr	-9		4.2
H ₂ SO ₄	-9		4.3
	-7.3		4.5
	-7		4.6
HCl	-7	CH ₃ COOH	4.8
[(CH ₃) ₂ OH] ⁺	-3.8	(CH ₃) ₃ CCOOH	5.0
[CH ₃ OH ₂] ⁺	-2.5		5.1
H ₃ O ⁺	-1.7		5.2
CH ₃ SO ₃ H	-1.2	CH ₃ O- 	5.3
	0.0	H ₂ CO ₃	6.4
CF ₃ COOH	0.2	H ₂ S	7.0
CCl ₃ COOH	0.6		7.1
	1.0		7.8
Cl ₂ CHCOOH	1.3		8.9
H ₃ PO ₄	2.1		
FCH ₂ COOH	2.7		
ClCH ₂ COOH	2.8		
BrCH ₂ COOH	2.9		
ICH ₂ COOH	3.2		
HF	3.2		
	3.4		
HCOOH	3.8		
	3.9		

Compound	pK _a	Compound	pK _a
HC≡N	9.1	CH ₃ OH	15.5
	9.4	H ₂ O	15.7
NH ₄ ⁺	9.4	CH ₃ CH ₂ OH	16
H ₃ ⁺ NCH ₂ COO ⁻	9.8	CH ₃ CONH ₂	16
	10.0	CH ₃ CHO	17
	10.2	(CH ₃) ₃ COH	18
HCO ₃ ⁻	10.2	(CH ₃) ₂ C=O	19.2
CH ₃ NO ₂	10.2	CH ₃ CO ₂ CH ₂ CH ₃	24.5
	10.3	HC≡CH	25
CH ₃ CH ₂ SH	10.5	CH ₃ C≡N	25
[(CH ₃) ₃ NH] ⁺	10.6	CHCl ₃	25
	10.7	CH ₃ CON(CH ₃) ₂	30
[CH ₃ NH ₃] ⁺	10.7	H ₂	35
	10.7	NH ₃	38
[(CH ₃) ₂ NH ₂] ⁺	10.7	CH ₃ NH ₂	40
CF ₃ CH ₂ OH	12.4		41
	13.3		43
	15	CH ₂ =CHCH ₃	43
		CH ₂ =CH ₂	44
			46
		CH ₄	50
		CH ₃ CH ₃	50

Nomenclature

Although the basic principles of nomenclature are presented in the body of this text, additional information is often needed to name many complex organic compounds. Appendix B concentrates on three topics:


- Naming alkyl substituents that contain branching
- Naming polyfunctional compounds
- Naming bicyclic compounds

Naming Alkyl Substituents That Contain Branching


Alkyl groups that contain any number of carbons and no branches are named as described in Section 4.4A: change the *-ane* ending of the parent alkane to the suffix *-yl*. Thus the seven-carbon alkyl group $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ is called *heptyl*.

When an alkyl substituent also contains branching, follow a stepwise procedure:

- [1] Identify the longest carbon chain of the alkyl group that begins at the point of attachment to the parent. Begin numbering at the point of attachment and use the suffix *-yl* to indicate an alkyl group.





4 C's in the chain ----> butyl group




5 C's in the chain ----> pentyl group


- [2] Name all branches off the main alkyl chain and use the numbers from Step [1] to designate their location.



- [3] Set the entire name of the substituent in parentheses, and alphabetize this substituent name by the first letter of the complete name.



(3-methylbutyl)cyclohexane



1-(1,3-dimethylpentyl)-2-methylcyclohexane

- Alphabetize the **d** of **dimethylpentyl** before the **m** of **methyl**.
- Number the ring to give the lower number to the first substituent alphabetically: place the dimethylpentyl group at C1.

Naming Polyfunctional Compounds

Many organic compounds contain more than one functional group. When one of those functional groups is halo (X^-) or alkoxy (RO^-), these groups are named as substituents as described in Sections 7.2 and 9.3B. To name other polyfunctional compounds, we must learn which functional group is assigned a higher priority in the rules of nomenclature. Two steps are usually needed:

- [1] **Name a compound using the suffix of the highest priority group**, and name other functional groups as *substituents*. Table B.1 lists the common functional groups in order of decreasing priority, as well as the prefixes needed when a functional group must be named as a substituent.
- [2] Number the carbon chain to give the lower number to the highest priority functional group, and then follow all other rules of nomenclature. Examples are shown in Figure B.1.

Polyfunctional compounds that contain C–C double and triple bonds have characteristic suffixes to identify them, as shown in Table B.2. The higher priority functional group is assigned the lower number.


Table B.1 Summary of Functional Group Nomenclature

Functional group	Suffix	Substituent name (prefix)
Carboxylic acid	-oic acid	carboxy
Ester	-oate	alkoxycarbonyl
Amide	-amide	amido
Nitrile	-nitrile	cyano
Aldehyde	-al	oxo (= O) or formyl (–CHO)
Ketone	-one	oxo
Alcohol	-ol	hydroxy
Amine	-amine	amino
Alkene	-ene	alkenyl
Alkyne	-yne	alkynyl
Alkane	-ane	alkyl
Ether	–	alkoxy
Halide	–	halo

Increasing priority


Figure B.1

Examples of nomenclature of polyfunctional compounds




3-amino-2-hydroxybutanal

[Name as a derivative of an aldehyde since CHO is the highest priority functional group.]




o-cyanobenzoic acid

[Name as a derivative of benzoic acid since COOH is the higher priority functional group.]



methyl 4-oxohexanoate

[Name as a derivative of an ester since COOR is the higher priority functional group.]



4-formyl-3-methoxycyclohexanecarboxamide

[Name as a derivative of an amide since CONH₂ is the highest priority functional group.]

Table B.2 Naming Polyfunctional Compounds with C–C Double and Triple Bonds


Functional groups	Suffix	Example
C=C and OH	enol	<p>Start numbering here.</p> <p>5-methyl-4-hexen-1-ol</p>
C=C + C=O (ketone)	enone	<p>Start numbering here.</p> <p>(4E)-4-hepten-3-one</p>
C=C + C≡C	enyne	<p>Start numbering here.</p> <p>HC≡CCH₂CH₂CH=CH₂</p> <p>1-hexen-5-yne</p>

Naming Bicyclic Compounds


Bicyclic ring systems—compounds that contain two rings that share one or two carbon atoms—can be bridged, fused, or spiro.



bridged ring







fused ring




spiro ring

- A bridged ring system contains two rings that share two non-adjacent carbons.
- A fused ring system contains two rings that share a common carbon–carbon bond.
- A spiro ring system contains two rings that share one carbon atom.


Fused and bridged ring systems are named as bicyclo[x.y.z]alkanes, where the parent alkane corresponds to the total number of carbons in both rings. The numbers *x*, *y*, and *z* refer to the number of carbons that join the shared carbons together, written in order of *decreasing* size. For a fused ring system, *z* always equals zero, because the two shared carbons are directly joined together. The shared carbons in a bridged ring system are called the **bridgehead carbons**.



Rings are numbered beginning at a shared carbon, and continuing around the *longest* bridge first, then the next longest, and so forth.





3,3-dimethylbicyclo[3.2.1]octane



7,7-dimethylbicyclo[2.2.1]heptane

Spiro ring systems are named as spiro[x.y]alkanes where the parent alkane corresponds to the total number of carbons in both rings, and *x* and *y* refer to the number of carbons that join the shared carbon (the spiro carbon), written in order of *increasing* size. When substituents are present, the rings are numbered beginning with a carbon *adjacent* to the spiro carbon in the *smaller* ring.



Bond Dissociation Energies for Some Common Bonds

[A-B → A• + •B]

Bond	ΔH° kJ/mol	(kcal/mol)
H - Z bonds		
H - F	569	(136)
H - Cl	431	(103)
H - Br	368	(88)
H - I	297	(71)
H - OH	498	(119)
Z - Z bonds		
H - H	435	(104)
F - F	159	(38)
Cl - Cl	242	(58)
Br - Br	192	(46)
I - I	151	(36)
HO - OH	213	(51)
R - H bonds		
CH ₃ - H	435	(104)
CH ₃ CH ₂ - H	410	(98)
CH ₃ CH ₂ CH ₂ - H	410	(98)
(CH ₃) ₂ CH - H	397	(95)
(CH ₃) ₃ C - H	381	(91)
CH ₂ = CH - H	435	(104)
HC ≡ C - H	523	(125)
CH ₂ = CHCH ₂ - H	364	(87)
C ₆ H ₅ - H	460	(110)
C ₆ H ₅ CH ₂ - H	356	(85)
R - R bonds		
CH ₃ - CH ₃	368	(88)
CH ₃ - CH ₂ CH ₃	356	(85)
CH ₃ - CH = CH ₂	385	(92)
CH ₃ - C ≡ CH	489	(117)

Bond	ΔH° kJ/mol	(kcal/mol)
R - X bonds		
CH ₃ - F	456	(109)
CH ₃ - Cl	351	(84)
CH ₃ - Br	293	(70)
CH ₃ - I	234	(56)
CH ₃ CH ₂ - F	448	(107)
CH ₃ CH ₂ - Cl	339	(81)
CH ₃ CH ₂ - Br	285	(68)
CH ₃ CH ₂ - I	222	(53)
(CH ₃) ₂ CH - F	444	(106)
(CH ₃) ₂ CH - Cl	335	(80)
(CH ₃) ₂ CH - Br	285	(68)
(CH ₃) ₂ CH - I	222	(53)
(CH ₃) ₃ C - F	444	(106)
(CH ₃) ₃ C - Cl	331	(79)
(CH ₃) ₃ C - Br	272	(65)
(CH ₃) ₃ C - I	209	(50)
R - OH bonds		
CH ₃ - OH	389	(93)
CH ₃ CH ₂ - OH	393	(94)
CH ₃ CH ₂ CH ₂ - OH	385	(92)
(CH ₃) ₂ CH - OH	401	(96)
(CH ₃) ₃ C - OH	401	(96)
Other bonds		
CH ₂ = CH ₂	635	(152)
HC ≡ CH	837	(200)
O = C = O	535	(128)
O ₂	497	(119)

Reactions That Form Carbon–Carbon Bonds

Section	Reaction
11.11A	S_N2 reaction of an alkyl halide with an acetylide anion, $\text{C}\equiv\text{CR}$
11.11B	Opening of an epoxide ring with an acetylide anion, $\text{C}\equiv\text{CR}$
15.14	Radical polymerization of an alkene
16.12	Diels–Alder reaction
18.5	Friedel–Crafts alkylation
18.5	Friedel–Crafts acylation
20.10	Reaction of an aldehyde or ketone with a Grignard or organolithium reagent
20.13A	Reaction of an acid chloride with a Grignard or organolithium reagent
20.13A	Reaction of an ester with a Grignard or organolithium reagent
20.13B	Reaction of an acid chloride with an organocuprate reagent
20.14A	Reaction of a Grignard reagent with CO_2
20.14B	Reaction of an epoxide with an organometallic reagent
20.15	Reaction of an α,β -unsaturated carbonyl compound with an organocuprate reagent
21.9	Cyanohydrin formation
21.10	Wittig reaction to form an alkene
22.18	S_N2 reaction of an alkyl halide with NaCN
22.18C	Reaction of a nitrile with a Grignard or organolithium reagent
23.8	Direct enolate alkylation using LDA and an alkyl halide
23.9	Malonic ester synthesis to form a carboxylic acid
23.10	Acetoacetic ester synthesis to form a ketone
24.1	Aldol reaction to form a β -hydroxy carbonyl compound or an α,β -unsaturated carbonyl compound
24.2	Crossed aldol reaction
24.3	Directed aldol reaction
24.5	Claisen reaction to form a β -keto ester
24.6	Crossed Claisen reaction to form a β -dicarbonyl compound
24.7	Dieckmann reaction to form a five- or six-membered ring
24.8	Michael reaction to form a 1,5-dicarbonyl compound
24.9	Robinson annulation to form a 2-cyclohexenone
25.14	Reaction of a diazonium salt with CuCN
26.1	Coupling of an organocuprate reagent (R_2CuLi) with an organic halide ($\text{R}'\text{X}$)
26.2	The palladium-catalyzed Suzuki reaction of an organic halide with an organoborane
26.3	The palladium-catalyzed Heck reaction of a vinyl or aryl halide with an alkene
26.4	Addition of a dihalocarbene to an alkene to form a cyclopropane
26.5	Simmons–Smith reaction of an alkene with CH_2I_2 and $\text{Zn}(\text{Cu})$ to form a cyclopropane
26.6	Olefin metathesis
27.10B	Kiliani–Fischer synthesis of an aldose
28.2B	Alkylation of diethyl acetamidomalonate to form an amino acid
28.2C	Strecker synthesis of an amino acid
30.2	Chain-growth polymerization
30.4	Polymerization using Ziegler–Natta catalysts

Characteristic IR Absorption Frequencies

Bond	Functional group	Wavenumber (cm ⁻¹)	Comment
O-H	• ROH	3600–3200	broad, strong
	• RCOOH	3500–2500	very broad, strong
N-H	• RNH ₂	3500–3300	two peaks
	• R ₂ NH	3500–3300	one peak
	• RCONH ₂ , RCONHR	3400–3200	one or two peaks; N-H bending also observed at 1640 cm ⁻¹
C-H	• C _{sp} -H	3300	sharp, often strong
	• C _{sp²} -H	3150–3000	medium
	• C _{sp³} -H	3000–2850	strong
	• C _{sp²} -H of RCHO	2830–2700	one or two peaks
C≡C		2250	medium
C≡N		2250	medium
C=O			strong
	• RCOCl	1800	
	• (RCO) ₂ O	1800, 1760	two peaks
	• RCOOR	1745–1735	increasing $\bar{\nu}$ with decreasing ring size
	• RCHO	1730	
	• R ₂ CO	1715	increasing $\bar{\nu}$ with decreasing ring size
	• R ₂ CO, conjugated	1680	
	• RCOOH	1710	
• RCONH ₂ , RCONHR, RCONR ₂	1680–1630	increasing $\bar{\nu}$ with decreasing ring size	
C=C	• Alkene	1650	medium
	• Arene	1600, 1500	medium
C=N		1650	medium

Characteristic NMR Absorptions

¹H NMR Absorptions

Compound type	Chemical shift (ppm)
Alcohol	
R-O-H	1-5
$\begin{array}{c} \text{H} \\ \\ \text{R}-\text{C}-\text{O}- \\ \end{array}$	3.4-4.0
Aldehyde	
$\begin{array}{c} \text{O} \\ \\ \text{R}-\text{C}-\text{H} \end{array}$	9-10
Alkane	
RCH ₃	~0.9
R ₂ CH ₂	~1.3
R ₃ CH	~1.7
Alkene	
$\begin{array}{c} \text{H} \\ \\ \text{C}=\text{C} \\ \quad \end{array} \quad sp^2 \text{ C-H}$	4.5-6.0
$\begin{array}{c} \quad \\ \quad \text{C}-\text{H} \\ \quad \\ \text{C}=\text{C} \\ \quad \end{array} \quad \text{allylic } sp^3 \text{ C-H}$	1.5-2.5
Alkyl halide	
$\begin{array}{c} \text{H} \\ \\ \text{R}-\text{C}-\text{F} \\ \end{array}$	4.0-4.5
$\begin{array}{c} \text{H} \\ \\ \text{R}-\text{C}-\text{Cl} \\ \end{array}$	3.0-4.0
$\begin{array}{c} \text{H} \\ \\ \text{R}-\text{C}-\text{Br} \\ \end{array}$	2.7-4.0
$\begin{array}{c} \text{H} \\ \\ \text{R}-\text{C}-\text{I} \\ \end{array}$	2.2-4.0
Alkyne	
-C≡C-H	~2.5

Compound type		Chemical shift (ppm)
Amide		7.5–8.5
Amine		0.5–5.0
		2.3–3.0
Aromatic compound	sp^2 C-H	6.5–8
	benzylic sp^3 C-H	1.5–2.5
Carbonyl compound	sp^3 C-H on the α carbon	2.0–2.5
Carboxylic acid		10–12
Ether		3.4–4.0




^{13}C NMR Absorptions

Carbon type	Structure	Chemical shift (ppm)
Alkyl, sp^3 hybridized C		5–45
Alkyl, sp^3 hybridized C bonded to N, O, or X	 Z = N, O, X	30–80
Alkynyl, sp hybridized C		65–100
Alkenyl, sp^2 hybridized C		100–140
Aryl, sp^2 hybridized C		120–150
Carbonyl C		160–210

General Types of Organic Reactions


Substitution Reactions

[1] Nucleophilic substitution at an sp^3 hybridized carbon atom




[2] Nucleophilic acyl substitution at an sp^2 hybridized carbon atom

Carboxylic acids and their derivatives (Chapter 22)




[3] Radical substitution at an sp^3 hybridized C-H bond

Alkanes (Section 15.3)



[4] Electrophilic aromatic substitution


Aromatic compounds (Chapter 18)




Elimination Reactions

β Elimination at an sp^3 hybridized carbon atom

- a. Alkyl halides
(Chapter 8)




- b. Alcohols
(Section 9.8)




Addition Reactions

[1] Electrophilic addition to carbon-carbon multiple bonds

- a. Alkenes
(Chapter 10)




- b. Alkynes
(Section 11.6)



[2] Nucleophilic addition to carbon-oxygen multiple bonds

- Aldehydes and ketones
(Chapter 21)



How to Synthesize Particular Functional Groups

Acetals

- Reaction of an aldehyde or ketone with two equivalents of an alcohol (21.14)

Acid chlorides

- Reaction of a carboxylic acid with thionyl chloride (22.10)

Alcohols

- Nucleophilic substitution of an alkyl halide with ^-OH or H_2O (9.6)
- Hydration of an alkene (10.12)
- Hydroboration–oxidation of an alkene (10.16)
- Reduction of an epoxide with LiAlH_4 (12.6)
- Reduction of an aldehyde or ketone (20.4)
- Hydrogenation of an α,β -unsaturated carbonyl compound with $\text{H}_2 + \text{Pd-C}$ (20.4C)
- Enantioselective reduction of an aldehyde or ketone with the chiral CBS reagent (20.6)
- Reduction of an acid chloride with LiAlH_4 (20.7)
- Reduction of an ester with LiAlH_4 (20.7)
- Reduction of a carboxylic acid with LiAlH_4 (20.7)
- Reaction of an aldehyde or ketone with a Grignard or organolithium reagent (20.10)
- Reaction of an acid chloride with a Grignard or organolithium reagent (20.13)
- Reaction of an ester with a Grignard or organolithium reagent (20.13)
- Reaction of an organometallic reagent with an epoxide (20.14B)

Aldehydes

- Hydroboration–oxidation of a terminal alkyne (11.10)
- Oxidative cleavage of an alkene with O_3 followed by Zn or $(\text{CH}_3)_2\text{S}$ (12.10)
- Oxidation of a 1° alcohol with PCC (12.12)
- Oxidation of a 1° alcohol with HCrO_4^- , Amberlyst A-26 resin (12.13)
- Reduction of an acid chloride with $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]_3$ (20.7)
- Reduction of an ester with DIBAL-H (20.7)
- Hydrolysis of an acetal (21.14B)
- Hydrolysis of an imine or enamine (21.12B)
- Reduction of a nitrile (22.18B)

Alkanes

- Catalytic hydrogenation of an alkene with $\text{H}_2 + \text{Pd-C}$ (12.3)
- Catalytic hydrogenation of an alkyne with two equivalents of $\text{H}_2 + \text{Pd-C}$ (12.5A)
- Reduction of an alkyl halide with LiAlH_4 (12.6)

- Reduction of a ketone to a methylene group (CH_2)—the Wolff–Kishner or Clemmensen reaction (18.14B)
- Protonation of an organometallic reagent with H_2O , ROH , or acid (20.9)
- Coupling of an organocuprate reagent (R_2CuLi) with an alkyl halide, $\text{R}'\text{X}$ (26.1)
- Simmons–Smith reaction of an alkene with CH_2I_2 and $\text{Zn}(\text{Cu})$ to form a cyclopropane (26.5)

Alkenes

- Dehydrohalogenation of an alkyl halide with base (8.3)
- Dehydration of an alcohol with acid (9.8)
- Dehydration of an alcohol using POCl_3 and pyridine (9.10)
- β Elimination of an alkyl tosylate with base (9.13)
- Catalytic hydrogenation of an alkyne with H_2 + Lindlar catalyst to form a cis alkene (12.5B)
- Dissolving metal reduction of an alkyne with Na , NH_3 to form a trans alkene (12.5C)
- Wittig reaction (21.10)
- β Elimination of an α -halo carbonyl compound with Li_2CO_3 , LiBr , and DMF (23.7C)
- Hofmann elimination of an amine (25.12)
- Coupling of an organocuprate reagent (R_2CuLi) with an organic halide, $\text{R}'\text{X}$ (26.1)
- The palladium-catalyzed Suzuki reaction of a vinyl or aryl halide with a vinyl- or arylborane (26.2)
- The palladium-catalyzed Heck reaction of a vinyl or aryl halide with an alkene (26.3)
- Olefin metathesis (26.6)

Alkyl halides

- Reaction of an alcohol with HX (9.11)
- Reaction of an alcohol with SOCl_2 or PBr_3 (9.12)
- Cleavage of an ether with HBr or HI (9.14)
- Hydrohalogenation of an alkene with HX (10.9)
- Halogenation of an alkene with X_2 (10.13)
- Hydrohalogenation of an alkyne with two equivalents of HX (11.7)
- Halogenation of an alkyne with two equivalents of X_2 (11.8)
- Radical halogenation of an alkane (15.3)
- Radical halogenation at an allylic carbon (15.10)
- Radical addition of HBr to an alkene (15.13)
- Electrophilic addition of HX to a 1,3-diene (16.10)
- Radical halogenation of an alkyl benzene (18.13)
- Halogenation α to a carbonyl group (23.7)
- Addition of a dihalocarbene to an alkene to form a dihalocyclopropane (26.4)

Alkynes

- Dehydrohalogenation of an alkyl dihalide with base (11.5)
- $\text{S}_{\text{N}}2$ reaction of an alkyl halide with an acetylide anion, $\text{C}\equiv\text{CR}$ (11.11)

Amides

- Reaction of an acid chloride with NH_3 or an amine (22.8)
- Reaction of an anhydride with NH_3 or an amine (22.9)
- Reaction of a carboxylic acid with NH_3 or an amine and DCC (22.10)
- Reaction of an ester with NH_3 or an amine (22.11)

Amines

- Reduction of a nitro group (18.14C)
- Reduction of an amide with LiAlH_4 (20.7B)
- Reduction of a nitrile (22.18B)
- $\text{S}_{\text{N}}2$ reaction using NH_3 or an amine (25.7A)
- Gabriel synthesis (25.7A)
- Reductive amination of an aldehyde or ketone (25.7C)

Amino acids

- $\text{S}_{\text{N}}2$ reaction of an α -halo carboxylic acid with excess NH_3 (28.2A)
- Alkylation of diethyl acetamidomalonate (28.2B)
- Strecker synthesis (28.2C)
- Enantioselective hydrogenation using a chiral catalyst (28.4)

Anhydrides

- Reaction of an acid chloride with a carboxylate anion (22.8)
- Dehydration of a dicarboxylic acid (22.10)

Aryl halides

- Halogenation of benzene with $\text{X}_2 + \text{FeX}_3$ (18.3)
- Reaction of a diazonium salt with CuCl , CuBr , HBF_4 , NaI , or KI (25.14A)

Carboxylic acids

- Oxidative cleavage of an alkyne with ozone (12.11)
- Oxidation of a 1° alcohol with CrO_3 (or a similar Cr^{6+} reagent), H_2O , H_2SO_4 (12.12B)
- Oxidation of an alkyl benzene with KMnO_4 (18.14A)
- Oxidation of an aldehyde (20.8)
- Reaction of a Grignard reagent with CO_2 (20.14A)
- Hydrolysis of a cyanohydrin (21.9)
- Hydrolysis of an acid chloride (22.8)
- Hydrolysis of an anhydride (22.9)
- Hydrolysis of an ester (22.11)
- Hydrolysis of an amide (22.13)
- Hydrolysis of a nitrile (22.18A)
- Malonic ester synthesis (23.9)

Cyanohydrins

- Addition of HCN to an aldehyde or ketone (21.9)

1,2-Diols

- Anti dihydroxylation of an alkene with a peroxyacid, followed by ring opening with OH^- or H_2O (12.9A)
- Syn dihydroxylation of an alkene with KMnO_4 or OsO_4 (12.9B)

Enamines

- Reaction of an aldehyde or ketone with a 2° amine (21.12)

Epoxydes

- Intramolecular $\text{S}_{\text{N}}2$ reaction of a halohydrin using base (9.6)
- Epoxidation of an alkene with $m\text{CPBA}$ (12.8)
- Enantioselective epoxidation of an allylic alcohol with the Sharpless reagent (12.15)

Esters

- S_N2 reaction of an alkyl halide with a carboxylate anion, RCOO^- (7.19)
- Reaction of an acid chloride with an alcohol (22.8)
- Reaction of an anhydride with an alcohol (22.9)
- Fischer esterification of a carboxylic acid with an alcohol (22.10)

Ethers

- Williamson ether synthesis— S_N2 reaction of an alkyl halide with an alkoxide, ^-OR (9.6)
- Reaction of an alkyl tosylate with an alkoxide, ^-OR (9.13)
- Addition of an alcohol to an alkene in the presence of acid (10.12)
- Anionic polymerization of epoxides to form polyethers (30.3)

Halohydrins

- Reaction of an epoxide with HX (9.15)
- Addition of X and OH to an alkene (10.15)

Imine

- Reaction of an aldehyde or ketone with a 1° amine (21.11)

Ketones

- Hydration of an alkyne with H_2O , H_2SO_4 , and HgSO_4 (11.9)
- Oxidative cleavage of an alkene with O_3 followed by Zn or $(\text{CH}_3)_2\text{S}$ (12.10)
- Oxidation of a 2° alcohol with any Cr^{6+} reagent (12.12, 12.13)
- Friedel–Crafts acylation (18.5)
- Reaction of an acid chloride with an organocuprate reagent (20.13)
- Hydrolysis of an imine or enamine (21.12B)
- Hydrolysis of an acetal (21.14B)
- Reaction of a nitrile with a Grignard or organolithium reagent (22.18C)
- Acetoacetic ester synthesis (23.10)

Nitriles

- S_N2 reaction of an alkyl halide with NaCN (7.19, 22.18)
- Reaction of an aryl diazonium salt with CuCN (25.14A)

Phenols

- Reaction of an aryl diazonium salt with H_2O (25.14A)

Glossary

A

Absolute configuration (Section 27.11): The exact three-dimensional arrangement of the stereogenic centers in a molecule.

Acetal (Section 21.14): A compound having the general structure $R_2C(OR')_2$, where R = H, alkyl, or aryl. Acetals are used as protecting groups for aldehydes and ketones.

Acetoacetic ester synthesis (Section 23.10): A stepwise method that converts ethyl acetoacetate into a ketone having one or two carbons bonded to the α carbon.

Acetylation (Section 22.9): A reaction that transfers an acetyl group (CH_3CO-) from one atom to another.

Acetyl coenzyme A (Section 22.17): A biochemical thioester that acts as an acetylating reagent. Acetyl coenzyme A is often referred to as acetyl CoA.

Acetyl group (Section 21.2E): A substituent having the structure $-COCH_3$.

Acetylide anion (Sections 11.11, 20.9B): An anion formed by treating a terminal alkyne with a strong base. Acetylide anions have the general structure $R-C\equiv C^-$.

Achiral molecule (Section 5.3): A molecule that is superimposable upon its mirror image. An achiral molecule is not chiral.

Acid chloride (Sections 20.1, 22.1): A compound having the general structure $RCOCl$.

Acidity constant (Section 2.3): A value symbolized by K_a that represents the strength of an acid (HA). The larger the K_a , the stronger the acid.

$$K_a = \frac{[H_3O^+][A^-]}{[H-A]}$$

Active site (Section 6.11): The region of an enzyme that binds the substrate.

Acyclic alkane (Section 4.1): A compound with the general formula C_nH_{2n+2} . Acyclic alkanes are also called saturated hydrocarbons because they contain the maximum number of hydrogen atoms per carbon.

Acylation (Sections 18.5A, 22.17): A reaction that transfers an acyl group from one atom to another.

Acyl chloride (Section 18.5A): A compound having the general structure $RCOCl$. Acyl chlorides are also called acid chlorides.

Acyl group (Section 18.5A): A substituent having the general structure $RCO-$.

Acylium ion (Section 18.5B): A positively charged electrophile having the general structure $(R-C\equiv O)^+$, formed when the Lewis acid $AlCl_3$ ionizes the carbon-halogen bond of an acid chloride.

Acyl transfer reaction (Section 22.17): A reaction that transfers an acyl group from one atom to another.

1,2-Addition (Sections 16.10, 20.15): An addition reaction to a conjugated system that adds groups across two adjacent atoms.

1,4-Addition (Sections 16.10, 20.15): An addition reaction that adds groups to the atoms in the 1 and 4 positions of a conjugated system. 1,4-Addition is also called conjugate addition.

Addition polymer (Section 30.1): A polymer prepared by a chain reaction that adds a monomer to the growing end of a polymer chain. Addition polymers are also called chain-growth polymers.

Addition reaction (Sections 6.2C, 10.8): A reaction in which elements are added to a starting material. In an addition reaction, a π bond is broken and two σ bonds are formed.

Aglycon (Section 27.7C): The alcohol formed from hydrolysis of a glycoside.

Alcohol (Section 9.1): A compound having the general structure ROH. An alcohol contains a hydroxy group (OH group) bonded to an sp^3 hybridized carbon atom.

Aldaric acid (Section 27.9B): The dicarboxylic acid formed by the oxidation of the aldehyde and the primary alcohol of an aldose.

Aldehyde (Section 11.10): A compound having the general structure RCHO, where R = H, alkyl, or aryl.

Alditol (Section 27.9A): A compound formed by the reduction of the aldehyde of an aldose to a primary alcohol.

Aldol condensation (Section 24.1B): An aldol reaction in which the initially formed β -hydroxy carbonyl compound loses water by dehydration.

Aldol reaction (Section 24.1A): A reaction in which two molecules of an aldehyde or ketone react with each other in the presence of base to form a β -hydroxy carbonyl compound.

Aldonic acid (Section 27.9B): A compound formed by the oxidation of the aldehyde of an aldose to a carboxylic acid.

Aldose (Section 27.2): A monosaccharide comprised of a polyhydroxy aldehyde.

Aliphatic (Section 3.2A): A compound or portion of a compound made up of C-C σ and π bonds but not aromatic bonds.

Alkaloid (Section 25.6A): A basic, nitrogen-containing compound isolated from a plant source.

Alkane (Section 4.1): An aliphatic hydrocarbon having only C-C and C-H σ bonds.

Alkene (Section 8.2A): An aliphatic hydrocarbon that contains a carbon-carbon double bond.

Alkoxide (Sections 8.1, 9.6): An anion having the general structure RO^- , formed by deprotonating an alcohol with a base.

Alkoxy group (Section 9.3B): A substituent containing an alkyl group bonded to an oxygen (RO group).

Alkylation (Section 23.8): A reaction that transfers an alkyl group from one atom to another.

Alkyl group (Section 4.4A): A group formed by removing one hydrogen from an alkane. Alkyl groups are named by replacing the suffix *-ane* of the parent alkane with *-yl*.

Alkyl halide (Section 7.1): A compound containing a halogen atom bonded to an sp^3 hybridized carbon atom. Alkyl halides have the general molecular formula $C_nH_{2n+1}X$.

1,2-Alkyl shift (Section 9.9): The rearrangement of a less stable carbocation to a more stable carbocation by the shift of an alkyl group from one carbon atom to an adjacent carbon atom.

Alkyl tosylate (Section 9.13): A compound having the general structure $ROSO_2C_6H_4CH_3$. Alkyl tosylates are also called tosylates and are abbreviated as ROTs.

Alkyne (Section 8.10): An aliphatic hydrocarbon that contains a carbon–carbon triple bond.

Allyl carbocation (Section 16.1B): A carbocation that has a positive charge on the atom adjacent to a carbon–carbon double bond. An allyl carbocation is resonance stabilized.

Allyl group (Section 10.3C): A substituent having the structure $-\text{CH}_2-\text{CH}=\text{CH}_2$.

Allylic bromination (Section 15.10A): A radical substitution reaction in which bromine replaces a hydrogen atom on the carbon adjacent to a carbon–carbon double bond.

Allylic carbon (Section 15.10): A carbon atom bonded to a carbon–carbon double bond.

Allylic halide (Section 7.1): A molecule containing a halogen atom bonded to the carbon atom adjacent to a carbon–carbon double bond.

Allyl radical (Section 15.10): A radical that has an unpaired electron on the carbon adjacent to a carbon–carbon double bond. An allyl radical is resonance stabilized.

Alpha (α) carbon (Sections 8.1, 19.2B): In an elimination reaction, the carbon that is bonded to the leaving group. In a carbonyl compound, the carbon that is bonded to the carbonyl carbon.

Ambident nucleophile (Section 23.3C): A nucleophile that has two reactive sites.

Amide (Sections 20.1, 22.1): A compound having the general structure RCONR'_2 , where $\text{R}' = \text{H}$ or alkyl.

Amide base (Sections 8.10, 23.3B): A nitrogen-containing base formed by deprotonating an amine or ammonia.

Amine (Sections 21.11, 25.1): A basic organic nitrogen compound having the general structure RNH_2 , R_2NH , or R_3N . An amine has a nonbonded pair of electrons on the nitrogen atom.

α -Amino acid (Sections 19.14A, 28.1): A compound having the general structure $\text{RCH}(\text{NH}_2)\text{COOH}$. α -Amino acids are the building blocks of proteins.

Amino acid residue (Section 28.5): The individual amino acids in peptides and proteins.

Amino group (Section 25.3D): A substituent having the structure $-\text{NH}_2$.

α -Amino nitrile (Section 28.2C): A compound having the general structure $\text{RCH}(\text{NH}_2)\text{C}\equiv\text{N}$.

Amino sugar (Section 27.14A): A carbohydrate that contains an NH_2 group instead of a hydroxy group at a non-anomeric carbon.

Ammonium salt (Section 25.1): A compound containing a positively charged nitrogen with four σ bonds; for example, $\text{R}_4\text{N}^+\text{X}^-$.

Angle strain (Section 4.11): An increase in the energy of a molecule resulting when the bond angles of the sp^3 hybridized atoms deviate from the optimum tetrahedral angle of 109.5° .

Angular methyl group (Section 29.8A): A methyl group located at the ring junction of two fused rings of the steroid skeleton.

Anhydride (Section 22.1): A compound having the general structure $(\text{RCO})_2\text{O}$.

Aniline (Section 25.3C): A compound having the structure $\text{C}_6\text{H}_5\text{NH}_2$.

Anion (Section 1.2): A negatively charged ion that results from a neutral atom gaining one or more electrons.

Anionic polymerization (Section 30.2C): Chain-growth polymerization of alkenes substituted by electron-withdrawing groups that stabilize intermediate anions.

Annulation (Section 24.9): A reaction that forms a new ring.

Annulene (Section 17.8A): A hydrocarbon containing a single ring with alternating double and single bonds.

α Anomer (Section 27.6): The stereoisomer of a cyclic monosaccharide in which the anomeric OH and the CH_2OH groups are trans.

In a D monosaccharide, the hydroxy group on the anomeric carbon is drawn down.

β Anomer (Section 27.6): The stereoisomer of a cyclic monosaccharide in which the anomeric OH and the CH_2OH groups are cis. In a D monosaccharide, the hydroxy group on the anomeric carbon is drawn up.

Anomeric carbon (Section 27.6): The stereogenic center at the hemiacetal carbon of a cyclic monosaccharide.

Anti addition (Section 10.8): An addition reaction in which the two parts of a reagent are added from opposite sides of a double bond.

Antiaromatic compound (Section 17.7): An organic compound that is cyclic, planar, completely conjugated, and has $4n$ π electrons.

Antibonding molecular orbital (Section 17.9A): A high-energy molecular orbital formed when two atomic orbitals of opposite phase overlap.

Anti conformation (Section 4.10): A staggered conformation in which the two larger groups on adjacent carbon atoms have a dihedral angle of 180° .



Anti dihydroxylation (Section 12.9A): The addition of two hydroxy groups to opposite faces of a double bond.

Antioxidant (Section 15.12): A compound that stops an oxidation from occurring.

Anti periplanar (Section 8.8A): In an elimination reaction, a geometry where the β hydrogen and the leaving group are on opposite sides of the molecule.

Aromatic compound (Section 17.1): A planar, cyclic organic compound that has p orbitals on all ring atoms and a total of $4n + 2$ π electrons in the orbitals.

Aryl group (Section 17.3D): A substituent formed by removing one hydrogen atom from an aromatic ring.

Aryl halide (Sections 7.1, 18.3): A molecule such as $\text{C}_6\text{H}_5\text{X}$, containing a halogen atom X bonded to an aromatic ring.

Asymmetric carbon (Section 5.3): A carbon atom that is bonded to four different groups. An asymmetric carbon is also called a stereogenic center, a chiral center, or a chirality center.


Asymmetric reaction (Sections 12.15, 20.6A, 28.4): A reaction that converts an achiral starting material into predominantly one enantiomer.

Atactic polymer (Section 30.4): A polymer having the substituents randomly oriented along the carbon backbone of an elongated polymer chain.

Atomic number (Section 1.1): The number of protons in the nucleus of an element.

Atomic weight (Section 1.1): The weighted average of the mass of all isotopes of a particular element. The atomic weight is reported in atomic mass units (amu).

Axial bonds (Section 4.12A): Bonds located above or below and perpendicular to the plane of the chair conformation of cyclohexane. Three axial bonds point upwards (on the up carbons) and three axial bonds point downwards (on the down carbons).



Azo compound (Section 25.15): A compound having the general structure $\text{RN}=\text{NR}'$.

B

Backside attack (Section 7.11C): Approach of a nucleophile from the side opposite the leaving group.

Barrier to rotation (Section 4.10): The energy difference between the lowest and highest energy conformations of a molecule.

Base peak (Section 13.1): The peak in the mass spectrum having the greatest abundance value.

Basicity (Section 7.8): A measure of how readily an atom donates its electron pair to a proton.

Benedict's reagent (Section 27.9B): A reagent for oxidizing aldehydes to carboxylic acids using a Cu^{2+} salt, forming brick-red Cu_2O as a side product.

Benzoyl group (Section 21.2E): A substituent having the structure $-\text{COC}_6\text{H}_5$.

Benzyl group (Section 17.3D): A substituent having the structure $\text{C}_6\text{H}_5\text{CH}_2-$.

Benzylic halide (Sections 7.1, 18.13): A compound such as $\text{C}_6\text{H}_5\text{CH}_2\text{X}$, containing a halogen atom X bonded to a carbon that is bonded to a benzene ring.

Beta (β) carbon (Sections 8.1, 19.2B): In an elimination reaction, the carbon adjacent to the carbon with the leaving group. In a carbonyl compound, the carbon located two carbons from the carbonyl carbon.

Bimolecular reaction (Sections 6.9B, 7.10, 7.13A): A reaction in which the concentration of both reactants affects the reaction rate and both terms appear in the rate equation. In a bimolecular reaction, two reactants are involved in the only step or the rate-determining step.

Biodegradable polymer (Section 30.9B): A polymer that can be degraded by microorganisms naturally present in the environment.

Biomolecule (Section 3.9): An organic compound found in a biological system.

Boat conformation of cyclohexane (Section 4.12B): An unstable conformation adopted by cyclohexane that resembles a boat. The instability of the boat conformation results from torsional strain and steric strain. The boat conformation of cyclohexane is 30 kJ/mol less stable than the chair conformation.



Boiling point (Section 3.4A): The temperature at which molecules in the liquid phase are converted to the gas phase. Molecules with stronger intermolecular forces have higher boiling points. Boiling point is abbreviated as bp.

Bond dissociation energy (Section 6.4): The amount of energy needed to homolytically cleave a covalent bond.

Bonding (Section 1.2): The joining of two atoms in a stable arrangement. Bonding is a favorable process that leads to lowered energy and increased stability.

Bonding molecular orbital (Section 17.9A): A low-energy molecular orbital formed when two atomic orbitals of similar phase overlap.

Bond length (Section 1.6A): The average distance between the centers of two bonded nuclei. Bond lengths are reported in picometers (pm).

Branched-chain alkane (Section 4.1A): An acyclic alkane that has alkyl substituents bonded to the parent carbon chain.

Bridged ring system (Section 16.13D): A bicyclic ring system in which the two rings share non-adjacent carbon atoms.

Bromination (Sections 10.13, 15.6, 18.3): The reaction of a compound with bromine.

Bromohydrin (Section 10.15): A compound having a bromine and a hydroxy group on adjacent carbon atoms.

Brønsted–Lowry acid (Section 2.1): A proton donor, symbolized by HA. A Brønsted–Lowry acid must contain a hydrogen atom.

Brønsted–Lowry base (Section 2.1): A proton acceptor, symbolized by :B. A Brønsted–Lowry base must be able to form a bond to a proton by donating an available electron pair.

C

^{13}C NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the type of carbon atoms in a molecule.

Cahn–Ingold–Prelog system of nomenclature (Section 5.6): The system of designating a stereogenic center as either *R* or *S* according to the arrangement of the four groups attached to the center.

Carbamate (Sections 28.7, 30.6): A functional group containing a carbonyl group bonded to both an oxygen and a nitrogen atom. A carbamate is also called a urethane.

Carbanion (Section 2.5D): An ion with a negative charge on a carbon atom.

Carbene (Section 26.4): A neutral reactive intermediate having the general structure $:\text{CR}_2$. A carbene contains a divalent carbon surrounded by six electrons, making it a highly reactive electrophile that adds to C–C double bonds.

Carbinolamine (Section 21.7B): An unstable intermediate having a hydroxy group and an amine group on the same carbon. A carbinolamine is formed during the addition of an amine to a carbonyl group.

Carbocation (Section 7.13C): A positively charged carbon atom. A carbocation is sp^2 hybridized and trigonal planar, and contains a vacant *p* orbital.

Carbohydrate (Sections 21.17, 27.1): A polyhydroxy aldehyde or ketone or a compound that can be hydrolyzed to a polyhydroxy aldehyde or ketone.

Carbonate (Section 30.6D): A compound having the general structure $(\text{RO})_2\text{C}=\text{O}$.

Carbon backbone (Section 3.1): The C–C and C–H σ bond framework that makes up the skeleton of an organic molecule.

Carbon NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the type of carbon atoms in a molecule.

Carbonyl group (Sections 3.2C, 11.9, 20.1): A functional group that contains a carbon–oxygen double bond (C=O). The polar carbon–oxygen bond makes the carbonyl carbon electrophilic.

Carboxy group (Section 19.1): A functional group having the structure COOH.

Carboxylate anion (Section 19.2C): An anion having the general structure RCOO^- , formed by deprotonating a carboxylic acid with a Brønsted–Lowry base.

Carboxylation (Section 20.14): The reaction of an organometallic reagent with CO_2 to form a carboxylic acid after protonation.

Carboxylic acid (Section 19.1): A compound having the general structure RCOOH.

Carboxylic acid derivatives (Section 20.1): Compounds having the general structure RCOZ, which can be synthesized from carboxylic acids. Common carboxylic acid derivatives include acid chlorides, anhydrides, esters, and amides.

Catalyst (Section 6.10): A substance that speeds up the rate of a reaction, but is recovered unchanged at the end of the reaction and does not appear in the product.

Catalytic hydrogenation (Section 12.3): A reduction reaction involving the addition of H_2 to a π bond in the presence of a metal catalyst.

- Cation** (Section 1.2): A positively charged ion that results from a neutral atom losing one or more electrons.
- Cationic polymerization** (Section 30.2C): Chain-growth polymerization of alkene monomers involving carbocation intermediates.
- CBS reagent** (Section 20.6A): A chiral reducing agent formed by reacting an oxazaborolidine with BH_3 . CBS reagents predictably give one enantiomer as the major product of ketone reduction.
- Cephalin** (Section 29.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-\text{CH}_2\text{CH}_2\text{NH}_3^+$. Cephalins are also called phosphatidylethanolamines.
- Chain-growth polymer** (Section 30.1): A polymer prepared by a chain reaction that adds a monomer to the growing end of a polymer chain. Chain-growth polymers are also called addition polymers.
- Chain mechanism** (Section 15.4A): A reaction mechanism that involves repeating steps.
- Chair conformation of cyclohexane** (Section 4.12A): A stable conformation adopted by cyclohexane that resembles a chair. The stability of the chair conformation results from the elimination of angle strain (all C–C–C bond angles are 109.5°) and torsional strain (all groups on adjacent carbon atoms are staggered).



- Chemical shift** (Section 14.1B): The position of an absorption signal on the x axis in an NMR spectrum relative to the reference signal of tetramethylsilane.
- Chirality center** (Section 5.3): A carbon atom bonded to four different groups. A chirality center is also called a chiral center, a stereogenic center, and an asymmetric center.
- Chiral molecule** (Section 5.3): A molecule that is not superimposable upon its mirror image.
- Chlorination** (Sections 10.14, 15.5, 18.3): The reaction of a compound with chlorine.
- Chlorofluorocarbons** (Sections 7.4, 15.9): Synthetic alkyl halides having the general molecular formula $\text{CF}_x\text{Cl}_{4-x}$. Chlorofluorocarbons, abbreviated as CFCs, were used as refrigerants and aerosol propellants and contribute to the destruction of the ozone layer.
- Chlorohydrin** (Section 10.15): A compound having a chlorine and a hydroxy group on adjacent carbon atoms.
- Chromate ester** (Section 12.12A): An intermediate in the chromium-mediated oxidation of an alcohol having the general structure $\text{R}-\text{O}-\text{CrO}_3\text{H}$.
- s*-Cis** (Sections 16.6, 28.5B): The conformation of a 1,3-diene that has the two double bonds on the same side of the single bond that joins them.
- Cis isomer** (Sections 4.13B, 8.2B): An isomer of a ring or double bond that has two groups on the same side of the ring or double bond.
- Claisen reaction** (Section 24.5): A reaction between two molecules of an ester in the presence of base to form a β -keto ester.
- α Cleavage** (Section 13.3B): A fragmentation in mass spectrometry that results in cleavage of a carbon–carbon bond. With aldehydes and ketones, α cleavage results in breaking the bond between the carbonyl carbon and the carbon adjacent to it. With alcohols, α cleavage occurs by breaking a bond between an alkyl group and the carbon that bears the OH group.
- Clemmensen reduction** (Section 18.14B): A method to reduce aryl ketones to alkyl benzenes using $\text{Zn}(\text{Hg})$ in the presence of a strong acid.
- Coenzyme** (Section 12.13): A compound that acts with an enzyme to carry out a biochemical process.
- Combustion** (Section 4.14B): An oxidation–reduction reaction, in which an alkane or other organic compound reacts with oxygen to form CO_2 and H_2O , releasing energy.
- Common name** (Section 4.6): The name of a molecule that was adopted prior to and therefore does not follow the IUPAC system of nomenclature.
- Compound** (Section 1.2): The structure that results when two or more elements are joined together in a stable arrangement.
- Concerted reaction** (Sections 6.3, 7.11B): A reaction in which all bond forming and bond breaking occurs in one step.
- Condensation polymer** (Sections 22.16A, 30.1): A polymer formed when monomers containing two functional groups come together with loss of a small molecule such as water or HCl . Condensation polymers are also called step-growth polymers.
- Condensation reaction** (Section 24.1B): A reaction in which a small molecule, often water, is eliminated during the reaction process.
- Condensed structure** (Section 1.7A): A shorthand representation of the structure of a compound in which all atoms are drawn in but bonds and lone pairs are usually omitted. Parentheses are used to denote similar groups bonded to the same atom.
- Configuration** (Section 5.2): A particular three-dimensional arrangement of atoms.
- Conformations** (Section 4.9): The different arrangements of atoms that are interconverted by rotation about single bonds.
- Conjugate acid** (Section 2.2): The compound that results when a base gains a proton in a proton transfer reaction.
- Conjugate addition** (Sections 16.10, 20.15): An addition reaction that adds groups to the atoms in the 1 and 4 positions of a conjugated system. Conjugate addition is also called 1,4-addition.
- Conjugate base** (Section 2.2): The compound that results when an acid loses a proton in a proton transfer reaction.
- Conjugated diene** (Section 16.1A): A compound that contains two carbon–carbon double bonds joined by a single σ bond. Pi (π) electrons are delocalized over both double bonds. Conjugated dienes are also called 1,3-dienes.
- Conjugated protein** (Section 28.10C): A structure composed of a protein unit and a non-protein molecule.
- Conjugation** (Section 16.1): The overlap of p orbitals on three or more adjacent atoms.
- Constitutional isomers** (Sections 1.4A, 4.1A, 5.2): Two compounds that have the same molecular formula, but differ in the way the atoms are connected to each other. Constitutional isomers are also called structural isomers.
- Coordination polymerization** (Section 30.4): A polymerization reaction that uses a homogeneous catalyst that is soluble in the reaction solvents typically used.
- Copolymer** (Section 30.2D): A polymer prepared by joining two or more different monomers together.
- Core electrons** (Section 1.1): The electrons in the inner shells of orbitals. Core electrons are not usually involved in the chemistry of a particular element.
- Counterion** (Section 2.1): An ion that does not take part in a reaction and is opposite in charge to the ion that does take part in the reaction. A counterion is also called a spectator ion.
- Coupling constant** (Section 14.6A): The frequency difference, measured in Hz, between the peaks in a split NMR signal.
- Coupling reaction** (Section 25.15): A reaction that forms a bond between two discrete molecules.
- Covalent bond** (Section 1.2): A bond that results from the sharing of electrons between two nuclei. A covalent bond is a two-electron bond.

Crossed aldol reaction (Section 24.2): An aldol reaction in which the two reacting carbonyl compounds are different. A crossed aldol reaction is also called a mixed aldol reaction.

Crossed Claisen reaction (Section 24.6): A Claisen reaction in which the two reacting esters are different.

Crown ether (Section 3.7B): A cyclic ether containing multiple oxygen atoms. Crown ethers bind specific cations depending on the size of their central cavity.

Curved arrow notation (Section 1.5A): A convention that shows the movement of an electron pair. The tail of the arrow begins at the electron pair and the head points to where the electron pair moves.

Cyanide anion (Section 21.9A): An anion having the structure $\text{C}\equiv\text{N}^-$.

Cyano group (Section 22.1): A functional group consisting of a carbon–nitrogen triple bond ($\text{C}\equiv\text{N}$).

Cyanohydrin (Section 21.9): A compound having the general structure $\text{RCH}(\text{OH})\text{C}\equiv\text{N}$. A cyanohydrin results from the addition of HCN across the carbonyl of an aldehyde or a ketone.


Cycloalkane (Sections 4.1, 4.2): A compound that contains carbons joined in one or more rings. Cycloalkanes with one ring have the general formula C_nH_{2n} .

Cyclopropanation (Section 26.4): An addition reaction to a carbon–carbon double bond that forms a cyclopropane.

D

D-Sugar (Section 27.2C): A sugar with the hydroxy group on the stereogenic center farthest from the carbonyl on the right side in the Fischer projection formula.

Decalin (Section 29.8A): Two fused six-membered rings. *cis*-Decalin has the hydrogen atoms at the ring fusion on the same side of the rings, whereas *trans*-decalin has the hydrogen atoms at the ring fusion on opposite sides of the rings.



Decarboxylation (Section 23.9A): Loss of CO_2 through cleavage of a carbon–carbon bond.

Degenerate orbitals (Section 17.9B): Orbitals (either atomic or molecular) having the same energy.

Degree of unsaturation (Section 10.2): A ring or a π bond in a molecule. The number of degrees of unsaturation compares the number of hydrogens in a compound to that of a saturated hydrocarbon containing the same number of carbons.

Dehydration (Sections 9.8, 22.10B): A reaction that results in the loss of the elements of water from the reaction components.

Dehydrohalogenation (Section 8.1): An elimination reaction in which the elements of hydrogen and halogen are lost from a starting material.

Delta (δ) scale (Section 14.1B): A common scale of chemical shifts used in NMR spectroscopy in which the absorption due to tetramethylsilane (TMS) occurs at zero parts per million.

Deoxy (Section 27.14B): A prefix that means without oxygen.

Deoxyribonucleoside (Section 27.14B): An *N*-glycoside formed by the reaction of D-2-deoxyribose with certain amine heterocycles.

Deoxyribonucleotide (Section 27.14B): A DNA building block having a deoxyribose and either a purine or pyrimidine base joined

together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.

Deprotection (Section 20.12): A reaction that removes a protecting group, regenerating a functional group.

Deshielding effects (Section 14.3A): An effect in NMR caused by a decrease in electron density, thus increasing the strength of the magnetic field felt by the nucleus. Deshielding shifts an absorption downfield.

Dextrorotatory (Section 5.12A): Rotating plane-polarized light in the clockwise direction. The rotation is labeled *d* or (+).

1,3-Diacid (Section 23.9A): A compound containing two carboxylic acids separated by a single carbon atom. 1,3-Diacids are also called β -diacids.

Dialkylamide (Section 23.3B): An amide base having the general structure R_2N^- .

Diastereomers (Section 5.7): Stereoisomers that are not mirror images of each other. Diastereomers have the same *R,S* designation for at least one stereogenic center and the opposite *R,S* designation for at least one of the other stereogenic centers.

Diastereotopic protons (Section 14.2C): Two hydrogen atoms on the same carbon such that substitution of either hydrogen with a group *Z* forms diastereomers. The two hydrogen atoms are not equivalent and give two NMR signals.

1,3-Diaxial interaction (Section 4.13A): A steric interaction between two axial substituents of the chair form of cyclohexane. Larger axial substituents create unfavorable 1,3-diaxial interactions, destabilizing a cyclohexane conformation.

Diazonium salt (Section 25.13A): An ionic salt having the general structure $(\text{R}-\text{N}\equiv\text{N})^+\text{Cl}^-$.

Diazotization reaction (Section 25.13A): A reaction that converts 1° alkylamines and arylamines to diazonium salts.

1,3-Dicarbonyl compound (Section 23.2): A compound containing two carbonyl groups separated by a single carbon atom.

1,4-Dicarbonyl compound (Section 24.4): A dicarbonyl compound in which the carbonyl groups are separated by three single bonds. 1,4-Dicarbonyl compounds can undergo intramolecular reactions to form five-membered rings.

1,5-Dicarbonyl compound (Section 24.4): A dicarbonyl compound in which the carbonyl groups are separated by four single bonds. 1,5-Dicarbonyl compounds can undergo intramolecular reactions to form six-membered rings.

Dieckmann reaction (Section 24.7): An intramolecular Claisen reaction of a diester to form a ring, typically a five- or six-membered ring.

Diels–Alder reaction (Section 16.12): An addition reaction between a 1,3-diene and a dienophile to form a cyclohexene ring.

1,3-Diene (Section 16.1A): A compound containing two carbon–carbon double bonds joined by a single σ bond. Pi (π) electrons are delocalized over both double bonds. 1,3-Dienes are also called conjugated dienes.

Dienophile (Section 16.12): The alkene component in a Diels–Alder reaction that reacts with a 1,3-diene.

Dihedral angle (Section 4.9): The angle that separates a bond on one atom from a bond on an adjacent atom.

Dihydroxylation (Section 12.9): Addition of two hydroxy groups to a double bond to form a 1,2-diol.

Diol (Section 9.3A): A compound possessing two hydroxy groups. Diols are also called glycols.

Dipeptide (Section 28.5): Two amino acids joined together by one amide bond.

Diphosphate (Section 29.7B): A good leaving group that is often used in biological systems. Diphosphate is abbreviated as OPP.

Dipole (Section 1.11): A separation of electronic charge.

Dipole–dipole interaction (Section 3.3B): An attractive intermolecular interaction between the permanent dipoles of polar molecules. The dipoles of adjacent molecules align so that the partial positive and partial negative charges are in close proximity.

Directed aldol reaction (Section 24.3): A crossed aldol reaction in which the enolate of one carbonyl compound is formed, followed by addition of the second carbonyl compound.

Disaccharide (Section 27.12): A carbohydrate containing two monosaccharide units joined together by a glycosidic linkage.

Disproportionation (Section 30.2): A method of chain termination in radical polymerization involving the transfer of a hydrogen atom from one polymer radical to another, forming a new C–H bond on one polymer chain and a new double bond on the other.

Dissolving metal reduction (Section 12.2): A reduction reaction using alkali metals as a source of electrons and liquid ammonia as a source of protons.

Disubstituted alkene (Section 8.2A): An alkene that has two alkyl groups and two hydrogens bonded to the carbons of the double bond ($R_2C=CH_2$ or $RCH=CHR$).

Disulfide (Section 28.5C): A compound having the general structure RSSR', often formed between the side chain of two cysteine residues.

Diterpene (Section 29.7A): A terpene that contains 20 carbons and four isoprene units.

Doublet (Section 14.6): An NMR signal that is split into two peaks of equal area, caused by one nearby nonequivalent proton.

Doublet of doublets (Section 14.8): A splitting pattern of four peaks observed when a signal is split by two different nonequivalent protons.

Downfield shift (Section 14.1B): In an NMR spectrum, a term used to describe the relative location of an absorption signal. A downfield shift means the signal is shifted to the left in the spectrum to higher chemical shift on the δ scale.

E

***E,Z* System of nomenclature** (Section 10.3B): A system for unambiguously naming alkene stereoisomers by assigning priorities to the two groups on each carbon of the double bond. The *E* isomer has the two higher priority groups on opposite sides of the double bond, and the *Z* isomer has them on the same side.

E1 mechanism (Sections 8.3, 8.6): An elimination mechanism that goes by a two-step process involving a carbocation intermediate. E1 is an abbreviation for “Elimination Unimolecular.”

E1cB mechanism (Section 24.1B): A two-step elimination mechanism that goes by a carbanion intermediate. E1cB stands for “Elimination Unimolecular, Conjugate Base.”

E2 mechanism (Sections 8.3, 8.4): An elimination mechanism that goes by a one-step concerted process, in which both reactants are involved in the transition state. E2 is an abbreviation for “Elimination Bimolecular.”

Eclipsed conformation (Section 4.9): A conformation of a molecule where the bonds on one carbon are directly aligned with the bonds on the adjacent carbon.



Edman degradation (Section 28.6B): A procedure used in peptide sequencing in which amino acids are cleaved one at a time from

the N-terminal end, the identity of the amino acid determined, and the process repeated until the entire sequence is known.

Eicosanoids (Section 29.6): A group of biologically active compounds containing 20 carbon atoms derived from arachidonic acid.

Elastomer (Section 30.5): A polymer that stretches when stressed but then returns to its original shape.

Electromagnetic radiation (Section 13.5): Radiant energy having dual properties of both waves and particles. The electromagnetic spectrum contains the complete range of electromagnetic radiation, arbitrarily divided into different regions.

Electron-donating inductive effect (Section 7.14A): An inductive effect in which an electropositive atom or polarizable group donates electron density through σ bonds to another atom.

Electronegativity (Section 1.11): A measure of an atom's attraction for electrons in a bond. Electronegativity indicates how much a particular atom “wants” electrons.

Electron-withdrawing inductive effect (Sections 2.5, 7.14A): An inductive effect in which a nearby electronegative atom pulls electron density towards itself through σ bonds.

Electrophile (Section 2.8): An electron-deficient compound, often symbolized by E^+ , which can accept a pair of electrons from an electron-rich compound, forming a covalent bond. Lewis acids are electrophiles.

Electrophilic addition reaction (Section 10.9): An addition reaction in which the first step of the mechanism involves addition of the electrophilic end of a reagent to a π bond.

Electrophilic aromatic substitution (Section 18.1): A characteristic reaction of benzene in which a hydrogen atom on the ring is replaced by an electrophile.

Electrospray ionization (Section 13.4C): A method for ionizing large biomolecules in a mass spectrometer. Electrospray ionization is abbreviated as ESI.

Electrostatic potential map (Section 1.11): A color-coded map that illustrates the distribution of electron density in a molecule. Electron-rich regions are indicated in red and electron-deficient regions are indicated in blue. Regions of intermediate electron density are shown in orange, yellow, and green.

α Elimination (Section 26.4): An elimination reaction involving the loss of two elements from the same atom.

β Elimination (Section 8.1): An elimination reaction involving the loss of elements from two adjacent atoms.

Elimination reaction (Sections 6.2B, 8.1): A chemical reaction in which elements of the starting material are “lost” and a π bond is formed.

Enamine (Section 21.12): A compound having an amine nitrogen atom bonded to a carbon–carbon double bond [$R_2C=CH(NR'_2)$].

Enantiomeric excess (Section 5.12D): A measurement of how much one enantiomer is present in excess of the racemic mixture. Enantiomeric excess (*ee*) is also called optical purity; *ee* = % of one enantiomer – % of the other enantiomer.

Enantiomers (Section 5.3): Stereoisomers that are mirror images but are not superimposable upon each other. Enantiomers have the exact opposite *R,S* designation at every stereogenic center.

Enantioselective reaction (Sections 12.15, 20.6A, 28.4): A reaction that affords predominantly or exclusively one enantiomer. Enantioselective reactions are also called asymmetric reactions.

Enantiotopic protons (Section 14.2C): Two hydrogen atoms on the same carbon such that substitution of either hydrogen with a group *Z* forms enantiomers. The two hydrogen atoms are equivalent and give a single NMR signal.

Endo position (Section 16.13D): A position of a substituent on a bridged bicyclic compound in which the substituent is closer to the longer bridge that joins the two carbons common to both rings.

Endothermic reaction (Section 6.4): A reaction in which the energy of the products is higher than the energy of the reactants. In an endothermic reaction, energy is absorbed and the ΔH° is a positive value.

Energy diagram (Section 6.7): A schematic representation of the energy changes that take place as reactants are converted to products. An energy diagram indicates how readily a reaction proceeds, how many steps are involved, and how the energies of the reactants, products, and intermediates compare.

Energy of activation (Section 6.7): The energy difference between the transition state and the starting material. The energy of activation, symbolized by E_{a} , is the minimum amount of energy needed to break bonds in the reactants.

Enolate (Sections 20.15, 23.3): A resonance-stabilized anion formed when a base removes an α hydrogen from the α carbon to a carbonyl group.

Enol tautomer (Sections 9.1, 11.9, 20.15): A compound having a hydroxy group bonded to a carbon-carbon double bond. An enol tautomer [such as $\text{CH}_2=\text{C}(\text{OH})\text{CH}_3$] is in equilibrium with its keto tautomer [$(\text{CH}_3)_2\text{C}=\text{O}$].

Enthalpy change (Section 6.4): The energy absorbed or released in a reaction. Enthalpy change is symbolized by ΔH° and is also called the heat of reaction.

Entropy (Section 6.6): A measure of the randomness in a system. The more freedom of motion or the more disorder present, the higher the entropy. Entropy is denoted by the symbol S° .

Entropy change (Section 6.6): The change in the amount of disorder between reactants and products in a reaction. The entropy change is denoted by the symbol ΔS° . $\Delta S^\circ = S^\circ_{\text{products}} - S^\circ_{\text{reactants}}$.

Enzyme (Section 6.11): A biochemical catalyst composed of at least one chain of amino acids held together in a very specific three-dimensional shape.

Enzyme-substrate complex (Section 6.11): A structure having a substrate bonded to the active site of an enzyme.

Epoxidation (Section 12.8): Addition of a single oxygen atom to an alkene to form an epoxide.

Epoxide (Section 9.1): A cyclic ether having the oxygen atom as part of a three-membered ring. Epoxides are also called oxiranes.

Epoxy resin (Section 30.6E): A step-growth polymer formed from a fluid prepolymer and a hardener that cross-links polymer chains together.

Equatorial bonds (Section 4.12A): Bonds located in the plane of the chair conformation of cyclohexane (around the equator). Three equatorial bonds point slightly upward (on the down carbons) and three equatorial bonds point slightly downward (on the up carbons).



Equilibrium constant (Section 6.5A): A mathematical expression, denoted by the symbol K_{eq} , which relates the amount of starting material and product at equilibrium. $K_{\text{eq}} = [\text{products}]/[\text{starting materials}]$.

Essential oil (Section 29.7): A class of terpenes isolated from plant sources by distillation.

Ester (Sections 20.1, 22.1): A compound having the general structure RCOOR' .

Esterification (Section 22.10C): A reaction that converts a carboxylic acid or a derivative of a carboxylic acid to an ester.

Ether (Section 9.1): A functional group having the general structure ROR' .

Ethynyl group (Section 11.2): An alkynyl substituent having the structure $-\text{C}\equiv\text{C}-\text{H}$.

Excited state (Sections 1.8B, 16.15A): A high-energy electronic state in which one or more electrons have been promoted to a higher energy orbital by absorption of energy.

Exo position (Section 16.13D): A position of a substituent on a bridged bicyclic compound in which the substituent is closer to the shorter bridge that joins the two carbons common to both rings.

Exothermic reaction (Section 6.4): A reaction in which the energy of the products is lower than the energy of the reactants. In an exothermic reaction, energy is released and the ΔH° is a negative value.

Extraction (Section 19.12): A laboratory method to separate and purify a mixture of compounds using solubility differences and acid-base principles.

F

Fat (Sections 10.6B, 29.3): A triacylglycerol that is solid at room temperature and composed of fatty acid side chains with a high degree of saturation.

Fatty acid (Sections 10.6A, 19.6): A long-chain carboxylic acid having between 12 and 20 carbon atoms.

Fehling's reagent (Section 27.9B): A reagent for oxidizing aldehydes to carboxylic acids using a Cu^{2+} salt as an oxidizing agent, forming brick-red Cu_2O as a by-product.


Fibrous proteins (Section 28.10): Long linear polypeptide chains that are bundled together to form rods or sheets.

Fingerprint region (Section 13.6B): The region in an IR spectrum at $< 1500 \text{ cm}^{-1}$. The region often contains a complex set of peaks and is unique for every compound.

First-order rate equation (Sections 6.9B, 7.10): A rate equation in which the reaction rate depends on the concentration of only one reactant.

Fischer esterification (Section 22.10C): An acid-catalyzed esterification reaction between a carboxylic acid and an alcohol to form an ester.

Fischer projection formula (Section 27.2A): A method for representing stereogenic centers with the stereogenic carbon at the intersection of vertical and horizontal lines. Fischer projections are also called cross formulas.



Fishhook (Section 6.3B): A half-headed curved arrow used in a reaction mechanism to denote the movement of a single electron.

Flagpole hydrogens (Section 4.12B): Hydrogens in the boat conformation of cyclohexane that are on either end of the "boat" and are forced into close proximity to each other.

Formal charge (Section 1.3C): The electronic charge assigned to individual atoms in a Lewis structure. The formal charge is calculated by subtracting an atom's unshared electrons and half of its shared electrons from the number of valence electrons that a neutral atom would possess.

Formyl group (Section 21.2E): A substituent having the structure $-\text{CHO}$.

Four-centered transition state (Section 10.16): A transition state that involves four atoms.

Fragment (Section 13.1): Radicals and cations formed by the decomposition of the molecular ion in a mass spectrometer.

Freons (Sections 7.4, 15.9): Chlorofluorocarbons consisting of simple halogen-containing organic compounds that were once commonly used as refrigerants.

Frequency (Section 13.5): The number of waves passing a point per unit time. Frequency is reported in cycles per second (s^{-1}), which

is also called hertz (Hz). Frequency is abbreviated with the Greek letter nu (ν).

Friedel–Crafts acylation (Section 18.5A): An electrophilic aromatic substitution reaction in which benzene reacts with an acid chloride in the presence of a Lewis acid to give a ketone.

Friedel–Crafts alkylation (Section 18.5A): An electrophilic aromatic substitution reaction in which benzene reacts with an alkyl halide in the presence of a Lewis acid to give an alkyl benzene.

Frontside attack (Section 7.11C): Approach of a nucleophile from the same side as the leaving group.

Full-headed curved arrow (Section 6.3B): An arrow used in a reaction mechanism to denote the movement of a pair of electrons.

Functional group (Section 3.1): An atom or group of atoms with characteristic chemical and physical properties. The functional group is the reactive part of the molecule.

Functional group interconversion (Section 11.12): A reaction that converts one functional group into another.

Functional group region (Section 13.6B): The region in an IR spectrum at $\geq 1500\text{ cm}^{-1}$. Common functional groups show one or two peaks in this region, at a characteristic frequency.

Furanose (Section 27.6): A cyclic five-membered ring of a monosaccharide containing an oxygen atom.

Fused ring system (Section 16.13D): A bicyclic ring system in which the two rings share one bond and two adjacent atoms.

G

Gabriel synthesis (Section 25.7A): A two-step method that converts an alkyl halide into a primary amine using a nucleophile derived from phthalimide.

Gas chromatography (Section 13.4B): An analytical technique that separates the components of a mixture based on their boiling points and the rate at which their vapors travel through a column.

Gauche conformation (Section 4.10): A staggered conformation in which the two larger groups on adjacent carbon atoms have a dihedral angle of 60° .



GC–MS (Section 13.4B): An analytical instrument that combines a gas chromatograph (GC) and a mass spectrometer (MS) in sequence.

gem-Diol (Section 21.13): A compound having the general structure $\text{R}_2\text{C}(\text{OH})_2$. gem-Diols are also called hydrates.

Geminal dihalide (Section 8.10): A compound that has two halogen atoms on the same carbon atom.

Gibbs free energy (Section 6.5A): The free energy of a molecule. Gibbs free energy is denoted by the symbol G° .

Gibbs free energy change (Section 6.5A): The overall energy difference between reactants and products. The Gibbs free energy change is denoted by the symbol ΔG° . $\Delta G^\circ = G^\circ_{\text{products}} - G^\circ_{\text{reactants}}$.

Globular proteins (Section 28.10): Polypeptide chains that are coiled into compact shapes with hydrophilic outer surfaces that make them water soluble.

Glycol (Section 9.3A): A compound possessing two hydroxy groups. Glycols are also called diols.

Glycosidase (Section 27.13B): An enzyme that hydrolyzes glycosidic linkages. An α -glycosidase hydrolyzes only α -glycosidic linkages.

Glycoside (Section 27.7A): A monosaccharide with an alkoxy group bonded to the anomeric carbon.

N-Glycoside (Section 27.14B): A monosaccharide containing a nitrogen bonded to the anomeric carbon.

Glycosidic linkage (Section 27.12): An acetal linkage formed between an OH group on one monosaccharide and the anomeric carbon on a second monosaccharide.

Green chemistry (Sections 12.13, 30.8): The use of environmentally benign methods to synthesize compounds.

Grignard reagent (Section 20.9): An organometallic reagent having the general structure RMgX .

Ground state (Sections 1.8B, 16.15A): The lowest energy arrangement of electrons for an atom.

Group number (Section 1.1): The number above a particular column in the periodic table. Group numbers are represented by either an Arabic (1 to 8) or Roman (I to VIII) numeral followed by the letter A or B. The group number of a second-row element is equal to the number of valence electrons in that element.

Grubbs catalyst (Section 26.6): A widely used ruthenium catalyst for olefin metathesis that has the structure $\text{Cl}_2(\text{Cy}_3\text{P})_2\text{Ru}=\text{CHPh}$.

Guest molecule (Section 9.5B): A small molecule that can bind to a larger host molecule.

H

^1H NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the number and type of hydrogen atoms in a molecule. ^1H NMR is also called proton NMR spectroscopy.

Half-headed curved arrow (Section 6.3B): An arrow used in a reaction mechanism to denote the movement of a single electron. A half-headed curved arrow is also called a fishhook.

α -Halo aldehyde or ketone (Section 23.7): An aldehyde or ketone with a halogen atom bonded to the α carbon.

Haloform reaction (Section 23.7B): A halogenation reaction of a methyl ketone (RCOCH_3) with excess halogen, which results in formation of RCOO^- and CHX_3 (haloform).

Halogenation (Sections 10.13, 15.3, 18.3): The reaction of a compound with a halogen.

Halohydrin (Sections 9.6, 10.15): A compound that has a hydroxy group and a halogen atom on adjacent carbon atoms.

Halonium ion (Section 10.13): A positively charged halogen atom. A bridged halonium ion contains a three-membered ring and is formed in the addition of a halogen (X_2) to an alkene.

Hammond postulate (Section 7.15): A postulate that states that the transition state of a reaction resembles the structure of the species (reactant or product) to which it is closer in energy.

Haworth projection (Section 27.6A): A representation of the cyclic form of a monosaccharide in which the ring is drawn flat.

Head-to-tail polymerization (Section 15.14B): A mechanism of radical polymerization in which the more substituted radical of the growing polymer chain always adds to the less substituted end of the new monomer.

Heat of hydrogenation (Section 12.3A): The ΔH° of a catalytic hydrogenation reaction equal to the amount of energy released by hydrogenating a π bond.

Heat of reaction (Section 6.4): The energy absorbed or released in a reaction. Heat of reaction is symbolized by ΔH° and is also called the change in enthalpy.

Heck reaction (Section 26.3): The palladium-catalyzed coupling of a vinyl or aryl halide with an alkene to form a more highly substituted alkene with a new carbon–carbon bond.

α -Helix (Section 28.9B): A secondary structure of a protein formed when a peptide chain twists into a right-handed or clockwise spiral.

Heme (Section 28.10C): A complex organic compound containing an Fe^{2+} ion coordinated with a porphyrin.

- Hemiacetal** (Section 21.14A): A compound that contains an alkoxy group and a hydroxy group bonded to the same carbon atom.
- Hertz** (Section 13.5): A unit of frequency measuring the number of waves passing a point per second.
- Heteroatom** (Section 3.1): An atom other than carbon or hydrogen. Common heteroatoms in organic chemistry are nitrogen, oxygen, sulfur, phosphorus, and the halogens.
- Heterocycle** (Section 9.3B): A cyclic compound containing a heteroatom as part of the ring.
- Heterolysis** (Section 6.3A): The breaking of a covalent bond by unequally dividing the electrons between the two atoms in the bond. Heterolysis generates charged intermediates. Heterolysis is also called heterolytic cleavage.
- Hexose** (Section 27.2): A monosaccharide containing six carbons.
- Highest occupied molecular orbital** (Section 17.4B): The molecular orbital with the highest energy that also contains electrons. The highest occupied molecular orbital is abbreviated as HOMO.
- High-resolution mass spectrometer** (Section 13.4A): A mass spectrometer that can measure mass-to-charge ratios to four or more decimal places. High-resolution mass spectra are used to determine the molecular formula of a compound.
- Hofmann elimination** (Section 25.12): An E2 elimination reaction that converts an amine into a quaternary ammonium salt as the leaving group. The Hofmann elimination gives the less substituted alkene as the major product.
- Homologous series** (Section 4.1B): A group of compounds that differ by only a CH₂ group in the chain.
- Homolysis** (Section 6.3A): The breaking of a covalent bond by equally dividing the electrons between the two atoms in the bond. Homolysis generates uncharged radical intermediates. Homolysis is also called homolytic cleavage.
- Homopolymer** (Section 30.2D): A polymer prepared from a single monomer.
- Hooke's law** (Section 13.7): A physical law that can be used to calculate the frequency of a bond vibration from the strength of the bond and the masses of the atoms attached to it.
- Host-guest complex** (Section 9.5B): The complex that is formed when a small guest molecule binds to a larger host molecule.
- Host molecule** (Section 9.5B): A large molecule that can bind a smaller guest molecule.
- Hückel's rule** (Section 17.7): A principle that states for a compound to be aromatic, it must be cyclic, planar, completely conjugated, and have $4n + 2\pi$ electrons.
- Hybridization** (Section 1.8B): The mathematical combination of two or more atomic orbitals (having different shapes) to form the same number of hybrid orbitals (all having the same shape).
- Hybrid orbital** (Section 1.8B): A new orbital that results from the mathematical combination of two or more atomic orbitals. The hybrid orbital is intermediate in energy compared to the atomic orbitals that were combined to form it.
- Hydrate** (Sections 12.12B, 21.13): A compound having the general structure R₂C(OH)₂. Hydrates are also called *gem*-diols.
- Hydration** (Sections 10.12, 21.9A): Addition of the elements of water to a molecule.
- Hydride** (Section 12.2): A negatively charged hydrogen ion (H⁻).
- 1,2-Hydride shift** (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a hydrogen atom from one carbon atom to an adjacent carbon atom.
- Hydroboration** (Section 10.16): The addition of the elements of borane (BH₃) to an alkene or alkyne.
- Hydrocarbon** (Sections 3.2A, 4.1): A compound made up of only the elements of carbon and hydrogen.
- Hydrogen bonding** (Section 3.3B): An attractive intermolecular interaction that occurs when a hydrogen atom bonded to an O, N, or F atom is electrostatically attracted to a lone pair of electrons on an O, N, or F atom in another molecule.
- Hydrogenolysis** (Section 28.7): A reaction that cleaves a σ bond using H₂ in the presence of a metal catalyst.
- α Hydrogens** (Section 23.1): The hydrogen atoms on the carbon bonded to the carbonyl carbon atom (the α carbon).
- Hydrohalogenation** (Section 10.9): An electrophilic addition of hydrogen halide (HX) to an alkene or alkyne.
- Hydrolysis** (Section 21.9A): A cleavage reaction with water.
- Hydroperoxide** (Section 15.11): An organic compound having the general structure ROOH.
- Hydrophilic** (Section 3.4C): Attracted to water. The polar portion of a molecule that interacts with polar water molecules is hydrophilic.
- Hydrophobic** (Section 3.4C): Not attracted to water. The nonpolar portion of a molecule that is not attracted to polar water molecules is hydrophobic.
- β -Hydroxy carbonyl compound** (Section 24.1A): An organic compound having a hydroxy group on the carbon β to the carbonyl group.
- Hydroxy group** (Section 9.1): The OH functional group.
- Hyperconjugation** (Section 7.14B): The overlap of an empty p orbital with an adjacent σ bond.
- Imide** (Section 25.7A): A compound having a nitrogen atom between two carbonyl groups.
- Imine** (Sections 21.7B, 21.11A): A compound with the general structure R₂C=NR'. Imines are also called Schiff bases.
- Iminium ion** (Section 21.11A): A resonance-stabilized cation having the general structure (R₂C=NR'₂)⁺, where R' = H or alkyl.
- Inductive effect** (Sections 2.5B, 7.14A): The pull of electron density through σ bonds caused by electronegativity differences of atoms.
- Infrared (IR) spectroscopy** (Section 13.6): An analytical technique used to identify the functional groups in a molecule based on their absorption of electromagnetic radiation in the infrared region.
- Initiation** (Section 15.4A): The initial step in a chain mechanism that forms a reactive intermediate by cleavage of a bond.
- Inscribed polygon method** (Section 17.10): A method to predict the relative energies of cyclic, completely conjugated compounds to determine which molecular orbitals are filled or empty. The inscribed polygon is also called a Frost circle.
- Integration** (Section 14.5): The area under an NMR signal that is proportional to the number of absorbing nuclei that give rise to the signal.
- Intermolecular forces** (Section 3.3): The types of interactions that exist between molecules. Functional groups determine the type and strength of these forces. Intermolecular forces are also called noncovalent interactions or nonbonded interactions.
- Internal alkene** (Section 10.1): An alkene that has at least one carbon atom bonded to each end of the double bond.
- Internal alkyne** (Section 11.1): An alkyne that has one carbon atom bonded to each end of the triple bond.
- Inversion of configuration** (Section 7.11C): The opposite relative stereochemistry of a stereogenic center in the starting material and product of a chemical reaction. In a nucleophilic substitution reaction, inversion results when the nucleophile and leaving group are in the opposite position relative to the three other groups on carbon.
- Iodoform test** (Section 23.7B): A test for the presence of methyl ketones, indicated by the formation of the yellow precipitate, CHI₃, via the haloform reaction.

Ionic bond (Section 1.2): A bond that results from the transfer of electrons from one element to another. Ionic bonds result from strong electrostatic interactions between ions with opposite charges. The transfer of electrons forms stable salts composed of cations and anions.

Ionophore (Section 3.7B): An organic molecule that can form a complex with cations so they may be transported across a cell membrane. Ionophores have a hydrophobic exterior and a hydrophilic central cavity that complexes the cation.

Isocyanate (Section 30.6C): A compound having the general structure $RN=C=O$.

Isoelectric point (Sections 19.14C, 28.1A): The pH at which an amino acid exists primarily in its neutral zwitterionic form. Isoelectric point is abbreviated as pI .

Isolated diene (Section 16.1A): A compound containing two carbon-carbon double bonds joined by more than one σ bond.

Isomers (Sections 1.4A, 4.1A, 5.1): Two different compounds that have the same molecular formula.

Isoprene unit (Section 29.7): A five-carbon unit with four carbons in a row and a one-carbon branch on one of the middle carbons.

Isotactic polymer (Section 30.4): A polymer having all the substituents on the same side of the carbon backbone of an elongated polymer chain.

Isotope (Section 1.1): Two or more atoms of the same element having the same number of protons in the nucleus but a different number of neutrons. Isotopes have the same atomic number but different mass numbers.

IUPAC system of nomenclature (Section 4.3): A systematic method for naming compounds developed by the International Union of Pure and Applied Chemistry.

K

K_a (Section 2.3): The symbol that represents the acidity constant of an acid HA. The larger the K_a , the stronger the acid.

$$K_a = \frac{[H_3O^+][A^-]}{[H-A]}$$

K_{eq} (Section 2.3): The equilibrium constant. $K_{eq} = [\text{products}]/[\text{starting materials}]$.

Kekulé structures (Section 17.1): Two equilibrating structures for benzene. Each structure contains a six-membered ring and three π bonds alternating with σ bonds around the ring.

Ketal (Section 21.14): A compound having the general structure $R_2C(OR')_2$, where R = alkyl or aryl. Ketals are derived from ketones and constitute a subclass of acetals.

β -Keto ester (Section 23.10): A compound containing a ketone carbonyl on the carbon β to the ester carbonyl group.

Ketone (Section 11.9): A compound with two alkyl groups bonded to the $C=O$ carbon atom, having the general structures $R_2C=O$ or $RCOR'$.

Ketose (Section 27.2): A monosaccharide comprised of a polyhydroxy ketone.

Keto tautomer (Section 11.9): A tautomer of a ketone that has a $C=O$ and a hydrogen bonded to the α carbon. The keto tautomer is in equilibrium with the enol tautomer.

Kiliani-Fischer synthesis (Section 27.10B): A reaction that lengthens the carbon chain of an aldose by adding one carbon to the carbonyl end.

Kinetic enolate (Section 23.4): The enolate that is formed the fastest—generally the less substituted enolate.

Kinetic product (Section 16.11): In a reaction that can give more than one product, the product that is formed the fastest.

Kinetic resolution (Section 28.3B): The separation of two enantiomers by a chemical reaction that selectively occurs for only one of the enantiomers.

Kinetics (Section 6.5): The study of chemical reaction rates.

L

L-Sugar (Section 27.2C): A sugar with the hydroxy group on the stereogenic center farthest from the carbonyl on the left side in the Fischer projection formula.

Lactam (Section 22.1): A cyclic amide in which the carbonyl carbon-nitrogen σ bond is part of a ring. A β -lactam contains the carbon-nitrogen σ bond in a four-membered ring.

Lactol (Section 21.16): A cyclic hemiacetal.

Lactone (Section 22.1): A cyclic ester in which the carbonyl carbon-oxygen σ bond is part of a ring.

Le Châtelier's principle (Section 9.8D): The principle that a system at equilibrium will react to counteract any disturbance to the equilibrium.

Leaving group (Section 7.6): An atom or group of atoms (Z) that is able to accept the electron density of the $C-Z$ bond during a substitution or elimination reaction.

Leaving group ability (Section 7.7): A measure of how readily a leaving group (Z) can accept the electron density of the $C-Z$ bond during a substitution or elimination reaction.

Lecithin (Section 29.4A): A phosphoglycerol in which the phosphodiester alkyl group is $-CH_2CH_2N(CH_3)_3^+$. Lecithins are also called phosphatidylcholines.

Leukotriene (Section 9.16): An unstable and potent biomolecule synthesized in cells by the oxidation of arachidonic acid. Leukotrienes are responsible for biological conditions such as asthma.

Levorotatory (Section 5.12A): Rotating plane-polarized light in the counterclockwise direction. The rotation is labeled l or $(-)$.

Lewis acid (Section 2.8): An electron pair acceptor.

Lewis acid-base reaction (Section 2.8): A reaction that results when a Lewis base donates an electron pair to a Lewis acid.

Lewis base (Section 2.8): An electron pair donor.

Lewis structure (Section 1.3): A representation of a molecule that shows the position of covalent bonds and nonbonding electrons. In Lewis structures, unshared electrons are represented by dots and a two-electron covalent bond is represented by a solid line. Lewis structures are also called electron dot structures.

Ligand (Section 26.2A): A group coordinated to a metal, which donates electron density to or sometimes withdraws electron density from the metal.

"Like dissolves like" (Section 3.4C): The principle that compounds dissolve in solvents having similar kinds of intermolecular forces; that is, polar compounds dissolve in polar solvents and nonpolar compounds dissolve in nonpolar solvents.

Lindlar catalyst (Section 12.5B): A catalyst for the hydrogenation of an alkyne to a cis alkene. The Lindlar catalyst is Pd adsorbed onto $CaCO_3$ with lead(II) acetate and quinoline.

Lipid (Sections 4.15, 29.1): A biomolecule with a large number of $C-C$ and $C-H$ σ bonds that is soluble in organic solvents and insoluble in water.

Lone pair of electrons (Section 1.2): A pair of valence electrons that is not shared with another atom in a covalent bond. Lone pairs are also called unshared or nonbonded pairs of electrons.

Lowest unoccupied molecular orbital (Section 17.9B): The molecular orbital with the lowest energy that does not contain electrons. The lowest unoccupied molecular orbital is abbreviated as the LUMO.

M

M peak (Section 13.1): The peak in the mass spectrum that corresponds to the mass of the molecular ion. The M peak is also called the molecular ion peak or the parent peak.

M + 1 peak (Section 13.1): The peak in the mass spectrum that corresponds to the mass of the molecular ion plus one. The M + 1 peak is caused by the presence of isotopes that increase the mass of the molecular ion.

M + 2 peak (Section 13.2): The peak in the mass spectrum that corresponds to the mass of the molecular ion plus two. The M + 2 peak is caused by the presence of isotopes, typically of a chlorine or a bromine atom.

Macrocyclic lactone (Section 22.6A): A cyclic ester contained in a large ring. Macrocyclic lactones are also called macrolides.

Macrolide (Section 22.6A): A cyclic ester contained in a large ring. Macrolides are also called macrocyclic lactones.

Magnetic resonance imaging (MRI) (Section 14.12): A form of NMR spectroscopy used in medicine.

Malonic ester synthesis (Section 23.9A): A stepwise method that converts diethyl malonate into a carboxylic acid having one or two carbons bonded to the α carbon.

Markovnikov's rule (Section 10.10): The rule that states in the addition of HX to an unsymmetrical alkene, the H atom bonds to the less substituted carbon atom.

Mass number (Section 1.1): The total number of protons and neutrons in the nucleus of a particular atom.

Mass spectrometry (Section 13.1): An analytical technique used for measuring the molecular weight and determining the molecular formula of an organic molecule.

Mass-to-charge ratio (Section 13.1): A ratio of the mass to the charge of a molecular ion or fragment. Mass-to-charge ratio is abbreviated as m/z .

Megahertz (Section 14.1A): A unit used for the frequency of the RF radiation in NMR spectroscopy. Megahertz is abbreviated as MHz; 1 MHz = 10^6 Hz.

Melting point (Section 3.4B): The temperature at which molecules in the solid phase are converted to the liquid phase. Molecules with stronger intermolecular forces and higher symmetry have higher melting points. Melting point is abbreviated as mp.

Merrifield method (Section 28.8): A method for synthesizing polypeptides using insoluble polymer supports.

Meso compound (Section 5.8): An achiral compound that contains two or more tetrahedral stereogenic centers.

Meta director (Section 18.7): A substituent on a benzene ring that directs a new group to the meta position during electrophilic aromatic substitution.

Meta isomer (Section 17.3B): A 1,3-disubstituted benzene ring. Meta substitution is abbreviated as $m-$.

Metal hydride reagent (Section 12.2): A reagent containing a polar metal-hydrogen bond that places a partial negative charge on the hydrogen and acts as a source of hydride ions (H^-).

Metathesis (Section 26.6): A reaction between two alkene molecules that results in the interchange of the carbons of their double bonds.

Methylation (Section 7.12): A reaction in which a CH_3 group is transferred from one compound to another.

Methylene group (Sections 4.1B, 10.3C): A CH_2 group bonded to a carbon chain ($-CH_2-$) or part of a double bond ($CH_2=$).

1,2-Methyl shift (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a methyl group from one carbon atom to an adjacent carbon atom.

Micelles (Section 3.6): Spherical droplets formed by soap molecules having the ionic heads on the surface and the nonpolar tails packed together in the interior. Grease and oil dissolve in the interior nonpolar region.

Michael acceptor (Section 24.8): The α,β -unsaturated carbonyl compound in a Michael reaction.

Michael reaction (Section 24.8): A reaction in which a resonance-stabilized carbanion (usually an enolate) adds to the β carbon of an α,β -unsaturated carbonyl compound.

Mixed aldol reaction (Section 24.2): An aldol reaction between two different carbonyl compounds. A mixed aldol reaction is also called a crossed aldol reaction.

Mixed anhydride (Section 22.1): An anhydride with two different alkyl groups bonded to the carbonyl carbon atoms.

Molecular ion (Section 13.1): The radical cation having the general structure $M^{+\cdot}$, formed by the removal of an electron from an organic molecule. The molecular ion is also called the parent ion.

Molecular orbital theory (Section 17.9A): A theory that describes bonds as the mathematical combination of atomic orbitals to form a new set of orbitals called molecular orbitals. Molecular orbital theory is also called MO theory.

Molecular recognition (Section 9.5B): The ability of a host molecule to recognize and bind specific guest molecules.

Molecule (Section 1.2): A compound containing two or more atoms bonded together with covalent bonds.

Monomers (Sections 5.1, 15.14): Small organic compounds that can be covalently bonded to each other (polymerized) in a repeating pattern.

Monosaccharide (Section 27.2): A simple sugar having three to seven carbon atoms.

Monosubstituted alkene (Section 8.2A): An alkene that has one alkyl group and three hydrogens bonded to the carbons of the double bond ($RCH=CH_2$).

Monoterpene (Section 29.7A): A terpene that contains 10 carbons and two isoprene units.

Multiplet (Section 14.6C): An NMR signal that is split into more than seven peaks.


Mutarotation (Section 27.6A): The process by which a pure anomer of a monosaccharide equilibrates to a mixture of both anomers when placed in solution.

N

$n + 1$ rule (Section 14.6C): The rule that an NMR signal for a proton with n nearby nonequivalent protons will be split into $n + 1$ peaks.

Natural product (Section 7.19): A compound isolated from a natural source.

Newman projection (Section 4.9): An end-on representation of the conformation of a molecule. The Newman projection shows the three groups bonded to each carbon atom in a particular C–C bond, as well as the dihedral angle that separates the groups on each carbon.



Nitration (Section 18.4): An electrophilic aromatic substitution reaction in which benzene reacts with $^+NO_2$ to give nitrobenzene, $C_6H_5NO_2$.

Nitrile (Sections 22.1, 22.18): A compound having the general structure $RC\equiv N$.

Nitronium ion (Section 18.4): An electrophile having the structure ${}^+\text{NO}_2$.

***N*-Nitrosamine** (Sections 7.16, 25.13B): A compound having the general structure $\text{R}_2\text{N}-\text{N}=\text{O}$. Nitrosamines are formed by the reaction of a secondary amine with ${}^+\text{NO}$.

Nitrosonium ion (Section 25.13): An electrophile having the structure ${}^+\text{NO}$.

NMR peak (Section 14.6A): The individual absorptions in a split NMR signal due to nonequivalent nearby protons.

NMR signal (Section 14.6A): The entire absorption due to a particular kind of proton in an NMR spectrum.

NMR spectrometer (Section 14.1A): An analytical instrument that measures the absorption of RF radiation by certain atomic nuclei when placed in a strong magnetic field.

Nonbonded pair of electrons (Section 1.2): A pair of valence electrons that is not shared with another atom in a covalent bond. Nonbonded electrons are also called unshared or lone pairs of electrons.

Nonbonding molecular orbital (Section 17.10): A molecular orbital having the same energy as the atomic orbitals that formed it.

Nonnucleophilic base (Section 7.8B): A base that is a poor nucleophile due to steric hindrance resulting from the presence of bulky groups.

Nonpolar bond (Section 1.11): A covalent bond in which the electrons are equally shared between the two atoms.

Nonpolar molecule (Section 1.12): A molecule that has no net dipole. A nonpolar molecule has either no polar bonds or multiple polar bonds whose dipoles cancel.

Nonreducing sugar (Section 27.9B): A carbohydrate that cannot be oxidized by Tollens, Benedict's, or Fehling's reagent.

Normal alkane (Section 4.1A): An acyclic alkane that has all of its carbons in a row. A normal alkane is an "*n*-alkane" or a straight-chain alkane.

Nuclear magnetic resonance spectroscopy (Section 14.1): A powerful analytical tool that can help identify the carbon and hydrogen framework of an organic molecule.

Nucleophile (Sections 2.8, 7.6): An electron-rich compound, symbolized by $:\text{Nu}^-$, which donates a pair of electrons to an electron-deficient compound, forming a covalent bond. Lewis bases are nucleophiles.

Nucleophilic acyl substitution (Sections 20.2B, 22.1): Substitution of a leaving group by a nucleophile at a carbonyl carbon.

Nucleophilic addition (Section 20.2A): Addition of a nucleophile to the electrophilic carbon of a carbonyl group followed by protonation of the oxygen.

Nucleophilicity (Section 7.8A): A measure of how readily an atom donates an electron pair to other atoms.

Nucleophilic substitution (Section 7.6): A reaction in which a nucleophile replaces the leaving group in a molecule.

Nucleoside (Section 27.14B): A biomolecule having a sugar and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage.

Nucleotide (Section 27.14B): A biomolecule having a sugar and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.



Observed rotation (Section 5.12A): The angle that a sample of an optically active compound rotates plane-polarized light. The angle is denoted by the symbol α and is measured in degrees ($^\circ$).

Octet rule (Section 1.2): The general rule governing the bonding process for second-row elements. Through bonding, second-row elements attain a complete outer shell of eight valence electrons.

Oil (Sections 10.6B, 29.3): A triacylglycerol that is liquid at room temperature and composed of fatty acid side chains with a high degree of unsaturation.

Olefin (Section 10.1): An alkene; a compound possessing a carbon-carbon double bond.

Optically active (Section 5.12A): Able to rotate the plane of plane-polarized light as it passes through a solution of a compound.

Optically inactive (Section 5.12A): Not able to rotate the plane of plane-polarized light as it passes through a solution of a compound.

Optical purity (Section 5.12D): A measurement of how much one enantiomer is present in excess of the racemic mixture. Optical purity is also called enantiomeric excess (*ee*); *ee* = % of one enantiomer - % of the other enantiomer.

Orbital (Section 1.1): A region of space around the nucleus of an atom that is high in electron density. There are four different kinds of orbitals, called *s*, *p*, *d*, and *f*.

Order of a rate equation (Section 6.9B): The sum of the exponents of the concentration terms in the rate equation of a reaction.

Organoborane (Section 10.16): A compound that contains a carbon-boron bond. Organoboranes have the general structure RBH_2 , R_2BH , or R_3B .

Organocopper reagent (Section 20.9): An organometallic reagent having the general structure R_2CuLi . Organocopper reagents are also called organocuprates.

Organolithium reagent (Section 20.9): An organometallic reagent having the general structure RLi .

Organomagnesium reagent (Section 20.9): An organometallic reagent having the general structure RMgX . Organomagnesium reagents are also called Grignard reagents.

Organometallic reagent (Section 20.9): A reagent that contains a carbon atom bonded to a metal.

Organopalladium compound (Section 26.2): An organometallic compound that contains a carbon-palladium bond.

Organophosphorus reagent (Section 21.10A): A reagent that contains a carbon-phosphorus bond.

Ortho isomer (Section 17.3B): A 1,2-disubstituted benzene ring. Ortho substitution is abbreviated as *o*-.

Ortho, para director (Section 18.7): A substituent on a benzene ring that directs a new group to the ortho and para positions during electrophilic aromatic substitution.

Oxaphosphetane (Section 21.10B): An intermediate in the Wittig reaction consisting of a four-membered ring containing a phosphorus-oxygen bond.

Oxazaborolidine (Section 20.6A): A heterocycle possessing a boron, a nitrogen, and an oxygen. An oxazaborolidine can be used to form a chiral reducing agent.

Oxidation (Sections 4.14A, 12.1): A process that results in a loss of electrons. For organic compounds, oxidation results in an increase in the number of C-Z bonds or a decrease in the number of C-H bonds; Z = an element more electronegative than carbon.

Oxidative addition (Section 26.2A): The addition of a reagent to a metal, often increasing the number of groups around the metal by two.

Oxidative cleavage (Section 12.10): An oxidation reaction that breaks both the σ and π bonds of a multiple bond to form two oxidized products.

Oxime (Section 27.10A): A compound having the general structure $\text{R}_2\text{C}=\text{NOH}$.

Oxirane (Section 9.1): A cyclic ether having the oxygen atom as part of a three-membered ring. Oxiranes are also called epoxides.

Ozonolysis (Section 12.10): An oxidative cleavage reaction in which a multiple bond reacts with ozone (O_3) as the oxidant.

P

Para isomer (Section 17.3B): A 1,4-disubstituted benzene ring. Para substitution is abbreviated as *p*-.

Parent ion (Section 13.1): The radical cation having the general structure $M^{+•}$, formed by the removal of an electron from an organic molecule. The parent ion is also called the molecular ion.

Parent name (Section 4.4): The portion of the IUPAC name of an organic compound that indicates the number of carbons in the longest continuous chain in the molecule.

Pentose (Section 27.2): A monosaccharide containing five carbons.

Peptide bond (Section 28.5): The amide bond in peptides and proteins.

Peptides (Sections 22.6B, 28.5): Low molecular weight polymers of less than 40 amino acids joined together by amide linkages.

Percent *s*-character (Section 1.10B): The fraction of a hybrid orbital due to the *s* orbital used to form it. As the percent *s*-character increases, a bond becomes shorter and stronger.

Percent transmittance (Section 13.6B): A measure of how much electromagnetic radiation passes through a sample of a compound and how much is absorbed.

Peroxide (Section 15.2): A reactive organic compound with the general structure ROOR. Peroxides are used as radical initiators by homolysis of the weak O–O bond.

Peroxyacid (Section 12.7): An oxidizing agent having the general structure RCO_3H .

Peroxy radical (Section 15.11): A radical having the general structure ROO·.

Petroleum (Section 4.7): A fossil fuel containing a complex mixture of compounds, primarily hydrocarbons with 1 to 40 carbon atoms.

Phenol (Sections 9.1, 15.12): A compound such as C_6H_5OH , which contains a hydroxy group bonded to a benzene ring.

Phenyl group (Section 17.3D): A group formed by removal of one hydrogen from benzene, abbreviated as C_6H_5- or $Ph-$.

Pheromone (Section 4.1): A chemical substance used for communication in an animal or insect species.

Phosphatidylcholine (Section 29.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2N(CH_3)_3^+$. Phosphatidylcholines are also called lecithins.

Phosphatidylethanolamine (Section 29.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2NH_3^+$. Phosphatidylethanolamines are also called cephalins.

Phosphoacylglycerols (Section 29.4A): A lipid having a glycerol backbone with two of the hydroxy groups esterified with fatty acids and the third hydroxy group as part of a phosphodiester.

Phosphodiester (Section 29.4): A functional group having the general formula $ROPO_2OR'$ formed by replacing two of the H atoms in phosphoric acid (H_3PO_4) with alkyl groups.

Phospholipid (Sections 3.7A, 29.4): A hydrolyzable lipid that contains a phosphorus atom.

Phosphonium salt (Section 21.10A): An organophosphorus reagent with a positively charged phosphorus and a suitable counterion; for example, $R_4P^+X^-$. Phosphonium salts are converted to ylides upon treatment with a strong base.

Phosphorane (Section 21.10A): A phosphorus ylide; for example, $Ph_3P=CR_2$.

Photon (Section 13.5): A particle of electromagnetic radiation.

Pi (π) bond (Section 1.9B): A bond formed by side-by-side overlap of two *p* orbitals where electron density is not concentrated on the

axis joining the two nuclei. Pi (π) bonds are generally weaker than σ bonds.

pK_a (Section 2.3): A logarithmic scale of acid strength. $pK_a = -\log K_a$. The smaller the pK_a , the stronger the acid.

Plane of symmetry (Section 5.3): A mirror plane that cuts a molecule in half, so that one half of the molecule is the mirror reflection of the other half.

Plane-polarized light (Section 5.12A): Light that has an electric vector that oscillates in a single plane. Plane-polarized light, also called polarized light, arises from passing ordinary light through a polarizer.

Plasticizer (Section 30.7): A low molecular weight compound added to a polymer to give it flexibility.

β -Pleated sheet (Section 28.9B): A secondary structure of a protein formed when two or more peptide chains line up side by side.

Poisoned catalyst (Section 12.5B): A hydrogenation catalyst with reduced activity that allows selective reactions to occur. The Lindlar catalyst is a poisoned Pd catalyst that converts alkynes to cis alkenes.

Polar aprotic solvent (Section 7.8C): A polar solvent that is incapable of intermolecular hydrogen bonding because it does not contain an O–H or N–H bond.

Polar bond (Section 1.11): A covalent bond in which the electrons are unequally shared between the two atoms. Unequal sharing of electrons results from bonding between atoms of different electronegativity values, usually with a difference of ≥ 0.5 units.

Polarimeter (Section 5.12A): An instrument that measures the degree that a compound rotates plane-polarized light.

Polarity (Section 1.11): A characteristic that results from a dipole. The polarity of a bond is indicated by an arrow with the head of the arrow pointing toward the negative end of the dipole and the tail with a perpendicular line through it at the positive end of the dipole. The polarity of a bond can also be indicated by the symbols δ^+ and δ^- .

Polarizability (Section 3.3B): A measure of how the electron cloud around an atom responds to changes in its electronic environment.

Polar molecule (Section 1.12): A molecule that has a net dipole. A polar molecule has either one polar bond or multiple polar bonds whose dipoles reinforce.

Polar protic solvent (Section 7.8C): A polar solvent that is capable of intermolecular hydrogen bonding because it contains an O–H or N–H bond.

Polyamide (Sections 22.16A, 30.6A): A step-growth polymer that contains many amide bonds. Nylon 6,6 and nylon 6 are polyamides.

Polycarbonate (Section 30.6C): A step-growth polymer that contains many $-OC(=O)O-$ bonds in its backbone, often formed by reaction of $Cl_2C=O$ with a diol.

Polycyclic aromatic hydrocarbon (Sections 9.17, 17.5): An aromatic hydrocarbon containing two or more benzene rings that share carbon–carbon bonds. Polycyclic aromatic hydrocarbons are abbreviated as PAHs.

Polyene (Section 16.7): A compound that contains three or more double bonds.


Polyester (Sections 22.16B, 30.6B): A step-growth polymer consisting of many ester bonds between diols and dicarboxylic acids.

Polyether (Sections 9.5B, 30.3): A compound that contains two or more ether linkages.

Polymer (Sections 5.1, 15.14): A large molecule composed of smaller monomer units covalently bonded to each other in a repeating pattern.

Polymerization (Section 15.14A): The chemical process that joins together monomers to make polymers.

- Polysaccharide** (Section 27.13): A carbohydrate containing three or more monosaccharide units joined together by glycosidic linkages.
- Polyurethane** (Section 30.6C): A step-growth polymer that contains many $-\text{NHC}(=\text{O})\text{O}-$ bonds in its backbone, formed by reaction of a diisocyanate and a diol.
- Porphyrin** (Section 28.10C): A nitrogen-containing heterocycle that can complex metal ions.
- Primary (1°) alcohol** (Section 9.1): An alcohol having the general structure RCH_2OH .
- Primary (1°) alkyl halide** (Section 7.1): An alkyl halide having the general structure RCH_2X .
- Primary (1°) amide** (Section 22.1): An amide having the general structure RCONH_2 .
- Primary (1°) amine** (Sections 21.11, 25.1): An amine having the general structure RNH_2 .
- Primary (1°) carbocation** (Section 7.14): A carbocation having the general structure RCH_2^+ .
- Primary (1°) carbon** (Section 4.1A): A carbon atom that is bonded to one other carbon atom.
- Primary (1°) hydrogen** (Section 4.1A): A hydrogen that is bonded to a 1° carbon.
- Primary protein structure** (Section 28.9A): The particular sequence of amino acids joined together by peptide bonds.
- Primary (1°) radical** (Section 15.1): A radical having the general structure $\text{RCH}_2\cdot$.
- Propagation** (Section 15.4A): The middle part of a chain mechanism in which one reactive particle is consumed and another is generated. Propagation repeats until a termination step occurs.
- Prostaglandin** (Section 4.15): A class of lipids containing 20 carbons, a five-membered ring, and a COOH group. Prostaglandins possess a wide range of biological activities.
- Prosthetic group** (Section 28.10C): The non-protein unit of a conjugated protein.
- Protecting group** (Section 20.12): A blocking group that renders a reactive functional group unreactive so that it does not interfere with another reaction.
- Protection** (Section 20.12): The reaction that blocks a reactive functional group with a protecting group.
- Proteins** (Sections 22.6B, 28.5): High molecular weight polymers of 40 or more amino acids joined together by amide linkages.
- Proton** (Section 2.1): A positively charged hydrogen ion (H^+).
- Proton NMR spectroscopy** (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the number and type of hydrogen atoms in a molecule.
- Proton transfer reaction** (Section 2.2): A Brønsted–Lowry acid–base reaction; a reaction that results in the transfer of a proton from an acid to a base.
- Purine** (Section 27.14B): A bicyclic aromatic heterocycle having two nitrogens in each of the rings.



- Pyranose** (Section 27.6): A cyclic six-membered ring of a monosaccharide containing an oxygen atom.
- Pyrimidine** (Section 27.14B): A six-membered aromatic heterocycle having two nitrogens in the ring.



Q

- Quantum** (Section 13.5): The discrete amount of energy associated with a particle of electromagnetic radiation (i.e., a photon).
- Quartet** (Section 14.6C): An NMR signal that is split into four peaks having a relative area of 1:3:3:1, caused by three nearby nonequivalent protons.
- Quaternary (4°) carbon** (Section 4.1A): A carbon atom that is bonded to four other carbon atoms.
- Quaternary protein structure** (Section 28.9C): The shape adopted when two or more folded polypeptide chains aggregate into one protein complex.
- Quintet** (Section 14.6C): An NMR signal that is split into five peaks caused by four nearby nonequivalent protons.

R

- R,S System of nomenclature** (Section 5.6): A system of nomenclature that distinguishes the stereochemistry at a tetrahedral stereogenic center by assigning a priority to each group connected to the stereogenic center. *R* indicates a clockwise orientation of the three highest priority groups and *S* indicates a counterclockwise orientation of the three highest groups. The system is also called the Cahn–Ingold–Prelog system.
- Racemic mixture** (Section 5.12B): An equal mixture of two enantiomers. A racemic mixture, also called a racemate, is optically inactive.
- Racemization** (Section 7.13C): The formation of equal amounts of two enantiomers from an enantiomerically pure starting material.
- Radical** (Sections 6.3B, 15.1): A reactive intermediate with a single unpaired electron, formed by homolysis of a covalent bond.
- Radical anion** (Section 12.5C): A reactive intermediate containing both a negative charge and an unpaired electron.
- Radical cation** (Section 13.1): A species with an unpaired electron and a positive charge, formed in a mass spectrometer by the bombardment of a molecule with an electron beam.
- Radical inhibitor** (Section 15.2): A compound that prevents radical reactions from occurring. Radical inhibitors are also called radical scavengers.
- Radical initiator** (Section 15.2): A compound that contains an especially weak bond that serves as a source of radicals.
- Radical polymerization** (Section 15.14B): A radical chain reaction involving the polymerization of alkene monomers by adding a radical to a π bond.
- Radical scavenger** (Section 15.2): A compound that prevents radical reactions from occurring. Radical scavengers are also called radical inhibitors.
- Rate constant** (Section 6.9B): A constant that is a fundamental characteristic of a reaction. The rate constant, symbolized by k , is a complex mathematical term that takes into account the dependence of a reaction rate on temperature and the energy of activation.
- Rate-determining step** (Section 6.8): In a multistep reaction mechanism, the step with the highest energy transition state.
- Rate equation** (Section 6.9B): An equation that shows the relationship between the rate of a reaction and the concentration of the reactants. The rate equation depends on the mechanism of the reaction and is also called the rate law.
- Reaction coordinate** (Section 6.7): The x axis in an energy diagram that represents the progress of a reaction as it proceeds from reactant to product.
- Reaction mechanism** (Section 6.3): A detailed description of how bonds are broken and formed as a starting material is converted to a product.

- Reactive intermediate** (Sections 6.3, 10.18): A high-energy unstable intermediate formed during the conversion of a stable starting material to a stable product.
- Reactivity–selectivity principle** (Section 15.6): The chemical principle that less reactive reagents are generally more selective, typically yielding one major product.
- Reciprocal centimeter** (Section 13.6A): The unit for wavenumber, which is used to report frequency in IR spectroscopy.
- Reducing sugar** (Section 27.9B): A carbohydrate that can be oxidized by Tollens, Benedict's, or Fehling's reagent.
- Reduction** (Sections 4.14A, 12.1): A process that results in the gain of electrons. For organic compounds, reduction results in a decrease in the number of C–Z bonds or an increase in the number of C–H bonds; Z = an element more electronegative than carbon.
- Reductive amination** (Section 25.7C): A two-step method that converts aldehydes and ketones into amines.
- Reductive elimination** (Section 26.2A): The elimination of two groups that surround a metal, often forming new carbon–hydrogen or carbon–carbon bonds.
- Regioselective reaction** (Section 8.5): A reaction that yields predominantly or exclusively one constitutional isomer when more than one constitutional isomer is possible.
- Resolution** (Section 28.3): The separation of a racemic mixture into its component enantiomers.
- Resonance** (Section 14.1A): In NMR spectroscopy, when an atomic nucleus absorbs RF radiation and spin flips to a higher energy state.
- Resonance hybrid** (Sections 1.5C, 16.4): A structure that is a weighted composite of all possible resonance structures. The resonance hybrid shows the delocalization of electron density due to the different locations of electrons in individual resonance structures.
- Resonance structures** (Sections 1.5, 16.2): Two or more structures of a molecule that differ in the placement of π bonds and nonbonded electrons. The placement of atoms and σ bonds stays the same.
- Retention of configuration** (Section 7.11C): The same relative stereochemistry of a stereogenic center in the reactant and the product of a chemical reaction.
- Retention time** (Section 13.4B): The length of time required for a component of a mixture to travel through a chromatography column.
- Retro Diels–Alder reaction** (Section 16.14B): The reverse of a Diels–Alder reaction in which a cyclohexene is cleaved to give a 1,3-diene and an alkene.
- Retrosynthetic analysis** (Section 10.18): Working backwards from a product to determine the starting material from which it is made.
- RF radiation** (Section 14.1A): Radiation in the radiofrequency region of the electromagnetic spectrum, characterized by long wavelength and low frequency and energy.
- Ribonucleoside** (Section 27.14B): An *N*-glycoside formed by the reaction of D-ribose with certain amine heterocycles.
- Ribonucleotide** (Section 27.14B): An RNA building block having a ribose and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.
- Ring-closing metathesis** (Section 26.6): An intramolecular olefin metathesis reaction using a diene starting material, which results in ring closure.
- Ring current** (Section 14.4): A circulation of π electrons in an aromatic ring caused by the presence of an external magnetic field.
- Ring flipping** (Section 4.12B): A stepwise process in which one chair conformation of cyclohexane interconverts with a second chair conformation.
- Ring-opening metathesis polymerization** (Problem 26.29): An olefin metathesis reaction that forms a high molecular weight polymer from certain cyclic alkenes.
- Robinson annulation** (Section 24.9): A ring-forming reaction that combines a Michael reaction with an intramolecular aldol reaction to form a 2-cyclohexenone.
- Rule of endo addition** (Section 16.13D): The rule that the endo product is preferred in a Diels–Alder reaction.

S

- Sandmeyer reaction** (Section 25.14A): A reaction between an aryl diazonium salt and a copper(I) halide to form an aryl halide (C_6H_5Cl or C_6H_5Br).
- Saponification** (Section 22.11B): Basic hydrolysis of an ester to form an alcohol and a carboxylate anion.
- Saturated fatty acid** (Section 10.6A): A fatty acid having no carbon–carbon double bonds in its long hydrocarbon chain.
- Saturated hydrocarbon** (Section 4.1): A compound that contains only C–C and C–H σ bonds and no rings, thus having the maximum number of hydrogen atoms per carbon.
- Schiff base** (Section 21.11A): A compound having the general structure $R_2C=NR'$. A Schiff base is also called an imine.
- Secondary (2°) alcohol** (Section 9.1): An alcohol having the general structure R_2CHOH .
- Secondary (2°) alkyl halide** (Section 7.1): An alkyl halide having the general structure R_2CHX .
- Secondary (2°) amide** (Section 22.1): An amide having the general structure $RCONHR'$.
- Secondary (2°) amine** (Sections 21.11, 25.1): An amine having the general structure R_2NH .
- Secondary (2°) carbocation** (Section 7.14): A carbocation having the general structure R_2CH^+ .
- Secondary (2°) carbon** (Section 4.1A): A carbon atom that is bonded to two other carbon atoms.
- Secondary (2°) hydrogen** (Section 4.1A): A hydrogen that is attached to a 2° carbon.
- Secondary protein structure** (Section 28.9B): The three-dimensional conformations of localized regions of a protein.
- Secondary (2°) radical** (Section 15.1): A radical having the general structure $R_2CH\cdot$.
- Second-order rate equation** (Sections 6.9B, 7.10): A rate equation in which the reaction rate depends on the concentration of two reactants.
- Separatory funnel** (Section 19.12): An item of laboratory glassware used for extractions.
- Septet** (Section 14.6C): An NMR signal that is split into seven peaks caused by six nearby nonequivalent protons.
- Sesquiterpene** (Section 29.7A): A terpene that contains 15 carbons and three isoprene units.
- Sesterterpene** (Section 29.7A): A terpene that contains 25 carbons and five isoprene units.
- Sextet** (Section 14.6C): An NMR signal that is split into six peaks caused by five nearby nonequivalent protons.
- Sharpless asymmetric epoxidation** (Section 12.15): An enantioselective oxidation reaction that converts the double bond of an allylic alcohol to a predictable enantiomerically enriched epoxide.
- Sharpless reagent** (Section 12.15): The reagent used in the Sharpless asymmetric epoxidation. The Sharpless reagent consists of *tert*-butyl hydroperoxide, a titanium catalyst, and one enantiomer of diethyl tartrate.
- Shielding effects** (Section 14.3A): An effect in NMR caused by small induced magnetic fields of electrons in the opposite direction to

the applied magnetic field. Shielding decreases the strength of the magnetic field felt by the nucleus and shifts an absorption upfield.

1,2-Shift (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a hydrogen atom or an alkyl group from one carbon atom to an adjacent carbon atom.

Sigma (σ) bond (Section 1.8A): A cylindrically symmetrical bond that concentrates the electron density on the axis that joins two nuclei. All single bonds are σ bonds.

Silyl ether (Section 20.12): A common protecting group for an alcohol in which the O–H bond is replaced by an O–Si bond.

Simmons–Smith reaction (Section 26.5): Reaction of an alkene with CH_2I_2 and $\text{Zn}(\text{Cu})$ to form a cyclopropane.

Singlet (Section 14.6A): An NMR signal that occurs as a single peak.

Skeletal structure (Section 1.7B): A shorthand representation of the structure of an organic compound in which carbon atoms and the hydrogen atoms bonded to them are omitted. All heteroatoms and the hydrogens bonded to them are drawn in. Carbon atoms are assumed to be at the junction of any two lines or at the end of a line.

$\text{S}_{\text{N}}1$ mechanism (Sections 7.10, 7.13): A nucleophilic substitution mechanism that goes by a two-step process involving a carbocation intermediate. $\text{S}_{\text{N}}1$ is an abbreviation for “Substitution Nucleophilic Unimolecular.”

$\text{S}_{\text{N}}2$ mechanism (Sections 7.10, 7.11): A nucleophilic substitution mechanism that goes by a one-step concerted process, where both reactants are involved in the transition state. $\text{S}_{\text{N}}2$ is an abbreviation for “Substitution Nucleophilic Bimolecular.”

Soap (Sections 3.6, 22.12B): The carboxylate salts of long-chain fatty acids prepared by the basic hydrolysis or saponification of a triacylglycerol.

Solubility (Section 3.4C): A measure of the extent to which a compound dissolves in a liquid.

Solute (Section 3.4C): The compound that is dissolved in a liquid solvent.

Solvent (Section 3.4C): The liquid component into which the solute is dissolved.

Specific rotation (Section 5.12C): A standardized physical constant for the amount that a chiral compound rotates plane-polarized light. Specific rotation is denoted by the symbol $[\alpha]$ and defined using a specific sample tube length (l in dm), concentration (c in g/mL), temperature (25 °C), and wavelength (589 nm). $[\alpha] = \alpha/(l \times c)$

Spectator ion (Section 2.1): An ion that does not take part in a reaction and is opposite in charge to the ion that does take part in a reaction. A spectator ion is also called a counterion.

Spectroscopy (Section 13.1): An analytical method using the interaction of electromagnetic radiation with molecules to determine molecular structure.

Sphingomyelin (Section 29.4B): A hydrolyzable phospholipid derived from sphingosine.

Spin flip (Section 14.1A): In NMR spectroscopy, when an atomic nucleus absorbs RF radiation and its magnetic field flips relative to the external magnetic field.

Spin–spin splitting (Section 14.6): Splitting of an NMR signal into peaks caused by nonequivalent protons on the same carbon or adjacent carbons.

Spiro ring system (Problem 23.63, Appendix B): A compound having two rings that share a single carbon atom.

Staggered conformation (Section 4.9): A conformation of a molecule in which the bonds on one carbon bisect the R–C–R bond angle on the adjacent carbon.



Step-growth polymer (Sections 22.16A, 30.1): A polymer formed when monomers containing two functional groups come together with loss of a small molecule such as water or HCl. Step-growth polymers are also called condensation polymers.

Stereochemistry (Sections 4.9, 5.1): The three-dimensional structure of molecules.

Stereogenic center (Section 5.3): A site in a molecule at which the interchange of two groups forms a stereoisomer. A carbon bonded to four different groups is a tetrahedral stereogenic center. A tetrahedral stereogenic center is also called a chirality center, a chiral center, or an asymmetric center.

Stereoisomers (Sections 4.13B, 5.1): Two isomers that differ only in the way the atoms are oriented in space.


Stereoselective reaction (Section 8.5): A reaction that yields predominantly or exclusively one stereoisomer when two or more stereoisomers are possible.

Stereospecific reaction (Section 10.14): A reaction in which each of two stereoisomers of a starting material yields a particular stereoisomer of a product.

Steric hindrance (Section 7.8B): A decrease in reactivity resulting from the presence of bulky groups at the site of a reaction.

Steric strain (Section 4.10): An increase in energy resulting when atoms in a molecule are forced too close to one another.

Steroid (Sections 16.14C, 29.8): A tetracyclic lipid composed of three six-membered rings and one five-membered ring.



Straight-chain alkane (Section 4.1A): An acyclic alkane that has all of its carbons in a row. Straight-chain alkanes are also called normal alkanes.

Strecker amino acid synthesis (Section 28.2C): A reaction that converts an aldehyde into an α -amino acid by way of an α -amino nitrile.

Structural isomers (Sections 4.1A, 5.2): Two compounds that have the same molecular formula but differ in the way the atoms are connected to each other. Structural isomers are also called constitutional isomers.

Substituent (Section 4.4): A group or branch attached to the longest continuous chain of carbons in an organic molecule.

Substitution reaction (Section 6.2A): A reaction in which an atom or a group of atoms is replaced by another atom or group of atoms. Substitution reactions involve σ bonds: one σ bond breaks and another is formed at the same atom.

Substrate (Section 6.11): An organic molecule that is transformed by the action of an enzyme.

Sulfonate anion (Section 19.13): An anion having the general structure RSO_3^- , formed by deprotonating a sulfonic acid with a Brønsted–Lowry base.

Sulfonation (Section 18.4): An electrophilic aromatic substitution reaction in which benzene reacts with $^+\text{SO}_3\text{H}$ to give a benzenesulfonic acid, $\text{C}_6\text{H}_5\text{SO}_3\text{H}$.

Sulfonic acid (Section 19.13): A compound having the general structure RSO_3H .

Suzuki reaction (Section 26.2): The palladium-catalyzed coupling of an organic halide (RX) with an organoborane (RBY_2) to form a product $\text{R-R}'$.

Symmetrical anhydride (Section 22.1): An anhydride that has two identical alkyl groups bonded to the carbonyl carbon atoms.

Symmetrical ether (Section 9.1): An ether with two identical alkyl groups bonded to the oxygen.

Syn addition (Section 10.8): An addition reaction in which two parts of a reagent are added from the same side of a double bond.

Syn dihydroxylation (Section 12.9B): The addition of two hydroxy groups to the same face of a double bond.

Syndiotactic polymer (Section 30.4): A polymer having the substituents alternating from one side of the backbone of an elongated polymer chain to the other.

Syn periplanar (Section 8.8): In an elimination reaction, a geometry in which the β hydrogen and the leaving group are on the same side of the molecule.

Systematic name (Section 4.3): The name of a molecule indicating the compound's chemical structure. The systematic name is also called the IUPAC name.

T

Target compound (Section 11.12): The final product of a synthetic scheme.

Tautomerization (Sections 11.9, 23.2A): The process of converting one tautomer into another.

Tautomers (Section 11.9): Constitutional isomers that are in equilibrium and differ in the location of a double bond and a hydrogen atom.

Terminal alkene (Section 10.1): An alkene that has the double bond at the end of the carbon chain.

Terminal alkyne (Section 11.1): An alkyne that has the triple bond at the end of the carbon chain.

C-Terminal amino acid (Section 28.5A): The amino acid at the end of a peptide chain with a free carboxy group.

N-Terminal amino acid (Section 28.5A): The amino acid at the end of a peptide chain with a free amino group.

Termination (Section 15.4A): The final step of a chain reaction. In a radical chain mechanism, two radicals combine to form a stable bond.

Terpene (Section 29.7): A lipid composed of repeating five-carbon isoprene units.

Tertiary (3°) alcohol (Section 9.1): An alcohol having the general structure R_3COH .

Tertiary (3°) alkyl halide (Section 7.1): An alkyl halide having the general structure R_3CX .

Tertiary (3°) amide (Section 22.1): An amide having the general structure RCONR'_2 .

Tertiary (3°) amine (Sections 21.11, 25.1): An amine having the general structure R_3N .

Tertiary (3°) carbocation (Section 7.14): A carbocation having the general structure R_3C^+ .

Tertiary (3°) carbon (Section 4.1A): A carbon atom that is bonded to three other carbon atoms.

Tertiary (3°) hydrogen (Section 4.1A): A hydrogen that is attached to a 3° carbon.

Tertiary protein structure (Section 28.9C): The three-dimensional shape adopted by an entire peptide chain.

Tertiary (3°) radical (Section 15.1): A radical having the general structure $\text{R}_3\text{C}\cdot$.

Tesla (Section 14.1A): A unit used to measure the strength of a magnetic field. Tesla is denoted with the symbol "T."

Tetramethylsilane (Section 14.1B): An internal standard used as a reference in NMR spectroscopy. The tetramethylsilane (TMS) reference peak occurs at 0 ppm on the δ scale.

Tetrasubstituted alkene (Section 8.2A): An alkene that has four alkyl groups and no hydrogens bonded to the carbons of the double bond ($\text{R}_2\text{C}=\text{CR}_2$).

Tetraterpene (Section 29.7A): A terpene that contains 40 carbons and eight isoprene units.

Tetrose (Section 27.2): A monosaccharide containing four carbons.

Thermodynamic enolate (Section 23.4): The enolate that is lower in energy—generally the more substituted enolate.

Thermodynamic product (Section 16.11): In a reaction that can give more than one product, the product that predominates at equilibrium.

Thermodynamics (Section 6.5): A study of the energy and equilibrium of a chemical reaction.

Thermoplastics (Section 30.7): Polymers that can be melted and then molded into shapes that are retained when the polymer is cooled.

Thermosetting polymer (Section 30.7): A complex network of cross-linked polymer chains that cannot be re-melted to form a liquid phase.

Thioester (Section 22.17): A compound with the general structure RCOSR' .

Tollens reagent (Sections 20.8, 27.9B): A reagent that oxidizes aldehydes, and consists of silver(I) oxide in aqueous ammonium hydroxide. A Tollens test is used to detect the presence of an aldehyde.

***p*-Toluenesulfonate** (Section 9.13): A very good leaving group having the general structure $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3^-$ and abbreviated as TsO^- . Compounds containing a *p*-toluenesulfonate leaving group are called alkyl tosylates and are abbreviated ROTs.

Torsional energy (Section 4.9): The energy difference between the staggered and eclipsed conformations of a molecule.

Torsional strain (Section 4.9): An increase in the energy of a molecule caused by eclipsing interactions between groups attached to adjacent carbon atoms.

Tosylate (Section 9.13): A very good leaving group having the general structure $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3^-$, and abbreviated as TsO^- .

***s*-Trans** (Sections 16.6, 28.5B): The conformation of a 1,3-diene that has the two double bonds on opposite sides of the single bond that joins them.

Trans diaxial (Section 8.8B): In an elimination reaction of a cyclohexane, a geometry in which the β hydrogen and the leaving group are trans with both in the axial position.

Trans isomer (Sections 4.13B, 8.3B): An isomer of a ring or double bond that has two groups on opposite sides of the ring or double bond.

Transition state (Section 6.7): An unstable energy maximum as a chemical reaction proceeds from reactants to products. The transition state is at the top of an energy "hill" and can never be isolated.

Triacylglycerol (Sections 10.6, 22.12A, 29.3): A lipid consisting of the triester of glycerol with three long-chain fatty acids. Triacylglycerols are the lipids that comprise animal fats and vegetable oils. Triacylglycerols are also called triglycerides.

Triose (Section 27.2): A monosaccharide containing three carbons.

Triplet (Section 14.6): An NMR signal that is split into three peaks having a relative area of 1:2:1, caused by two nearby nonequivalent protons.

Trisubstituted alkene (Section 8.2A): An alkene that has three alkyl groups and one hydrogen bonded to the carbons of the double bond ($\text{R}_2\text{C}=\text{CHR}$).

Triterpene (Section 29.7A): A terpene that contains 30 carbons and six isoprene units.

U

Ultraviolet (UV) light (Section 16.15): Electromagnetic radiation with a wavelength from 200–400 nm.

Unimolecular reaction (Sections 6.9B, 7.10, 7.13A): A reaction that has only one reactant involved in the rate-determining step, so the concentration of only one reactant appears in the rate equation.

α,β -Unsaturated carbonyl compound (Section 20.15): A conjugated compound containing a carbonyl group and a carbon–carbon double bond separated by a single σ bond.

Unsaturated fatty acid (Section 10.6A): A fatty acid having one or more carbon–carbon double bonds in its hydrocarbon chain. In natural fatty acids, the double bonds generally have the Z configuration.

Unsaturated hydrocarbon (Section 10.2): A hydrocarbon that has fewer than the maximum number of hydrogen atoms per carbon atom. Hydrocarbons with π bonds or rings are unsaturated.

Unsymmetrical ether (Section 9.1): An ether in which the two alkyl groups bonded to the oxygen are different.

Upfield shift (Section 14.1B): In an NMR spectrum, a term used to describe the relative location of an absorption signal. An upfield shift means a signal is shifted to the right in the spectrum to lower chemical shift.

Urethane (Section 30.6C): A compound that contains a carbonyl group bonded to both an OR group and an NHR (or NR₂) group. A urethane is also called a carbamate.

V

Valence bond theory (Section 17.9A): A theory that describes covalent bonding as the overlap of two atomic orbitals with the electron pair in the resulting bond being shared by both atoms.

Valence electrons (Section 1.1): The electrons in the outermost shell of orbitals. Valence electrons determine the properties of a given element. Valence electrons are more loosely held than the core electrons and thus participate in chemical reactions.

van der Waals forces (Section 3.3B): Very weak intermolecular interactions caused by momentary changes in electron density in molecules. The changes in electron density cause temporary dipoles, which are attracted to temporary dipoles in adjacent molecules. van der Waals forces are also called London forces.

Vicinal dihalide (Section 8.10): A compound that has two halogen atoms on adjacent carbon atoms.

Vinyl group (Section 10.3C): An alkene substituent having the structure $-\text{CH}=\text{CH}_2$.

Vinyl halide (Section 7.1): A molecule containing a halogen atom bonded to the sp^2 hybridized carbon of a carbon–carbon double bond.

Vitamins (Sections 3.5, 29.5): Organic compounds needed in small amounts by biological systems for normal cell function.

VSEPR theory (Section 1.6B): Valence shell electron pair repulsion theory. A theory that determines the three-dimensional shape of a molecule by the number of groups surrounding a central atom. The most stable arrangement keeps the groups as far away from each other as possible.

W

Walden inversion (Section 7.11C): The inversion of a stereogenic center involved in an S_N2 reaction.

Wavelength (Section 13.5): The distance from one point of a wave to the same point on the adjacent wave. Wavelength is abbreviated with the Greek letter lambda (λ).

Wavenumber (Section 13.6A): A unit for the frequency of electromagnetic radiation that is inversely proportional to wavelength. Wavenumber, reported in reciprocal centimeters (cm^{-1}), is used for frequency in IR spectroscopy.

Wax (Sections 4.15, 29.2): A hydrolyzable lipid consisting of an ester formed from a high molecular weight alcohol and a fatty acid.

Williamson ether synthesis (Section 9.6): A method for preparing ethers by reacting an alkoxide (RO⁻) with a methyl or primary alkyl halide.

Wittig reaction (Section 21.10): A reaction of a carbonyl group and an organophosphorus reagent that forms an alkene.

Wittig reagent (Section 21.10A): An organophosphorus reagent having the general structure $\text{Ph}_3\text{P}=\text{CR}_2$.

Wohl degradation (Section 27.10A): A reaction that shortens the carbon chain of an aldose by removing one carbon from the aldehyde end.

Wolff–Kishner reduction (Section 18.14B): A method to reduce aryl ketones to alkyl benzenes using hydrazine (NH_2NH_2) and strong base (KOH).

Y

Ylide (Section 21.10A): A chemical species that contains two oppositely charged atoms bonded to each other, and both atoms have octets of electrons.

Z

Zaitsev rule (Section 8.5): In a β elimination reaction, a rule that states that the major product is the alkene with the most substituted double bond.

Ziegler–Natta catalysts (Section 30.4): Polymerization catalysts prepared from an organoaluminum compound and a Lewis acid such as TiCl_4 , which afford polymer chains without significant branching and with controlled stereochemistry.

Zwitterion (Sections 19.14B, 28.1B): A neutral compound that contains both a positive and negative charge.

Credits

Prologue

Figure 1a: © Comstock/Punchstock RF; **1b:** © Corbis Premium RF/Alamy; **1c:** © Steven E. Frischling; **1d:** © Layne Kennedy/Corbis; **2 (left):** © Jill Braaten; **2 (right):** Daniel C. Smith.

Chapter 1

Opener: © Organics Image Library/Alamy; **p. 10:** © Jill Braaten; **p. 18 (both):** © Steve Gschmeissner/Photo Researchers, Inc.; **p. 27:** Courtesy John Somerville; **p. 30:** © The McGraw-Hill Companies, Inc./Elite Images; **p. 36:** © Corbis/RF; **p. 37:** © PhotoDisc/Getty RF; **p. 38:** © PhotoDisc/Getty RF.

Chapter 2

Opener: © The McGraw-Hill Companies, Inc./Elite Images; **p. 56:** © Fancy Photography/Veer; **pp. 65, 66:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 70:** Wilmer Stratton; **p. 71:** © National Geographic/Getty Images.

Chapter 3

Opener: © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 84:** Courtesy of Gebauer Company, Cleveland, Ohio; **p. 88 (top):** © Andrew Syred/Photo Researchers, Inc.; **p. 88 (bottom):** © Martin Harvey/Corbis; **p. 97:** © Comstock/PunchStock RF; **p. 98 (left):** © Banana Stock/Photo Stock RF; **p. 98 (right):** © The McGraw-Hill Companies, Inc./John Flournoy, photographer.

Chapter 4

Opener: © Bettmann/Corbis; **p. 114 (top):** © God of Insects; **p. 114 (bottom):** © Douglas Peebles/Corbis; **p. 119:** © PhotoDisc RF; **p. 128:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **pp. 129, 150:** © PhotoDisc RF.

Chapter 5

Opener: © The McGraw-Hill Companies, Inc./Elite Images; **5.1 (starch):** © Jim Erickson/Corbis; **5.1 (wheat):** © PhotoDisc/Getty RF; **5.1 (cotton):** © Corbis/RF; **5.1 (fabric):** Daniel C. Smith; **p. 163:** Daniel C. Smith; **p. 166 (top):** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 166 (bottom):** © Henriette Kress; **p. 170:** © PhotoDisc RF; **p. 179:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 188 (caraway):** © The McGraw-Hill Companies, Inc./Elite Images; **p. 188 (spearmint):** © David Sieren/Visuals Unlimited.

Chapter 6

Opener: © Jim Erickson; **p. 216:** © Jack Star/PhotoLink/Getty RF.

Chapter 7

Opener: © David Stoecklein/Corbis; **p. 232:** David L. Ballantine, Dept. Marine Sciences, Univ. Puerto Rico; **p. 233 (bottom):** Time Magazine June 30, 1947 in the collection of Beverly Naidus, www.artsforchange.org; **pp. 236, 250:** © Jill Braaten; **p. 261:** © The McGraw-Hill Companies, Inc./Elite Images.

Chapter 8

Opener: © Comstock/Punchstock RF; **p. 281:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **8.5:** Photo by Forest & Kim Starr.

Chapter 9

Opener: Photo collection of Dr. James P. McVey, NOAA Sea Grant Program; **9.4 (corn):** © Jupiterimages RF; **9.4 (ethanol):** Photo by Mark Wilson/Getty Images; **9.5b (painting):** The Boston Medical Library in the Francis A. Countway Library of Medicine; **9.7:** © Henriette Kress; **p. 347:** © The McGraw-Hill Companies, Inc./Jill Braaten, Photographer; **p. 349:** The McGraw-Hill Companies, Inc./John Thoeming, photographer.

Chapter 10

Opener: © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **10.5 (carrots):** © Corbis—All Rights reserved; **10.5 (ginger):** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **10.5 (apples):** © PhotoDisc RF; **10.5 (oranges):** © Iconotec/Alamer and Cali, photographers; **p. 369:** © Jill Braaten; **p. 380:** © 1998 Richard Megna, Fundamental Photographs, NYC; **10.17:** © Henriette Kress.

Chapter 11

Opener: © The McGraw-Hill Companies, Inc./Chris Kerrigan, photographer; **11.3:** © Arachnokulture; **p. 416:** Michael G. Moyer.

Chapter 12

Opener: John A. Rizzo/Getty Images; **p. 432:** © The McGraw-Hill Companies, Inc./Elite Images; **12.4 (both):** © The McGraw-Hill Companies, Inc./Elite Images; **p. 435:** PhotoLink/Getty Images RF; **p. 436:** © 1998 Richard Megna, Fundamental Photographs, NYC; **p. 441:** USDA APHIS PPQ Archives, www.forestryimages.org; **p. 449:** © The McGraw-Hill Companies, Inc./Elite Images; **p. 452:** PhotoDisc/Getty RF.

Chapter 13

Opener: Library of Congress; **13.7:** © Ted Kinsman/Photo Researchers, Inc.

Chapter 14

Opener: © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **14.15a:** James Rosato, RF (R) (MR) (CT) MRI Applications Specialist Brigham and Women's Hospital, Boston MA; **14.15b:** With permission, Daniel C. Smith. Spectra courtesy of the Chemistry Department at Rutgers University.

Chapter 15

Opener: © Jill Braaten; **p. 551:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 557 (top):** © The McGraw-Hill Companies, Inc./Elite Images; **p. 557 (bottom):** © Royalty-Free/Corbis; **p. 561:** The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 562 (PVC blood bags):** © Image Source/Getty RF; **(syringes):** © Royalty-Free/Corbis; **p. 562 (dental floss):** © Jill Braaten.

Chapter 16

Opener: © C Squared Studios/Getty Images; **p. 581:** © Franz-Marc-Frei/Corbis; **16.8:** Stephen Frink/Getty Images; **p. 599:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer.

Chapter 17

Opener: The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **17.3 (left):** PhotoAlto/PictureQuest; **17.3 (right):** Photo by Bob Nichols, USDA Natural Resources Conservation Service; **p. 622:** Daniel C. Smith; **p. 624:** Photo by Forest and Kim Starr; **p. 632 (soccer ball):** © Getty Images RF; **p. 632 (diamonds):** © Charles O'Rear/Corbis; **p. 632 (graphite):** © Lester V. Bergman/Corbis.

Chapter 18

Opener: © Rene Dulhoste/Photo Researchers, Inc.; **p. 646:** © Time Life Pictures/Getty Images; **p. 653:** Bill Aron/PhotoEdit Inc.; **p. 673:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer.

Chapter 19

Opener: Dr. Steven J. Wolf; **p. 694:** Ted Nelson/Dembinsky Photo Associates; **p. 696 (spinach):** © Ryman Cabannes/Photocuisine/Corbis; **p. 696 (meadowsweet):** © Biopix.dk <http://www.biopix.dk>; **p. 707:** © Photodisc Green/Getty RF; **p. 711:** John A. Raineri.

Chapter 20

Opener: © George Grall/National Geographic/Getty Images; **p. 723:** Gerald D. Carr, PhD; **p. 730:** © M.K. Ranjitsinh/Photo Researchers, Inc.; **p. 733 (soybeans):** PhotoDisc/Getty RF; **p. 733 (cereals):** © PhotoDisc/Getty RF; **p. 736:** © Digital Vision/Getty RF; **p. 738:** © The McGraw-Hill Companies, Inc./photographer Joe Franek; **p. 747:** © Iconotec.com/Pepiera Tom Canabi Hugo.

Chapter 21

Opener: Richo Cech, Horizon Herbs, LLC; **21.6:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 791:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 792:** Daniel C. Smith; **p. 808:** Bill Beatty/Visuals Unlimited, Inc.

Chapter 22

Opener: © Royalty-Free/Corbis; **p. 836 (top):** © Jill Braaten; **p. 836 (bottom):** © blickwinkel/Alamy; **p. 842 (top):** © The McGraw-Hill Companies, Inc./Joe Franek, photographer; **p. 842 (bottom):** Photo by Scott Bauer/USDA; **p. 844:** Coby Schal/NC State University; **p. 845:** © Stockbyte Silver/Getty RF; **p. 854:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 855:** © Jill Braaten; **p. 857:** © Andrew McClenaghan/Photo Researchers, Inc.; **p. 859:** Image courtesy of The Advertising Archives.

Chapter 23

Opener: © Colin Young-Wolff/Alamy; **pp. 887, 894:** © The McGraw-Hill Companies, Inc./Joe Franek, photographer; **p. 899:** Sergio Piu-matti; **p. 905:** © Margareta Taute/Visuals Unlimited.

Chapter 24

Opener: © Jill Braaten; **p. 925:** © William Weber/Visuals Unlimited; **p. 926 (both):** © Jill Braaten; **p. 931:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer.

Chapter 25

Opener: © Inga Spence/Visuals Unlimited; **25.5 (nightshade):** Werner Arnold; **25.5 (tobacco):** Creatas Images RF; **25.5 (hemlock):** © Steven P. Lynch; **p. 958:** Daniel C. Smith; **p. 968:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 974:** James Gathany/CDC; **p. 988 (indigo plant):** © Kirsten Soderlind/Corbis; **p. 988 (shell):** © SuperStock; **p. 988 (madder):** Photograph by Paul Redearn/Ozarks Regional Herbarium/Southwest Missouri State University; **p. 989:** © Science & Society Picture Library; **p. 990:** John Fischer, About.com Hawaii Travel.

Chapter 26

Opener: © Degginger E.R./Animals Animals/Earth Scenes; **p. 1012:** © The McGraw-Hill Companies, Inc./John Thoeming, photographer.

Chapter 27

Opener: © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 1029 (top):** © Jill Braaten; **p. 1029 (bottom):** © The McGraw-Hill Companies, Inc./Elite Images; **p. 1042:** © Brand X Pictures/PunchStock RF; **p. 1045 (top):** © Nature Picture Library/Alamy; **p. 1045 (bottom):** © Inga Spence/Alamy; **p. 1057 (top):** © Vol. 3/PhotoDisc/Getty RF; **p. 1057 (bottom):** © The McGraw-Hill Companies, Inc./Elite Images; **27.10 (sugar):** © The McGraw-Hill Companies, Inc./Elite Images; **27.10 (sugarcane):** © David Muench/Corbis; **27.11:** © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **p. 1062:** © Comstock Images/PictureQuest RF.

Chapter 28

Opener: © Royalty-Free/Corbis; **28.8:** Courtesy Protein Technologies, Inc.; **p. 1110:** © Eye of Science/Photo Researchers, Inc.

Chapter 29

Opener: © Jill Braaten; **p. 1121:** Daniel C. Smith; **p. 1123 (palm trees):** © Royalty-Free/Corbis; **p. 1123 (vegetable oils):** © The McGraw-Hill Companies, Inc./Elite Images; **p. 1123 (fish):** © Rick Price/Corbis; **p. 1125 (top):** © AP/Wide World Photos; **(bottom):** © PhotoDisc/Getty RF; **p. 1133 (top):** © Henriette Kress; **p. 1133 (bottom):** © PhotoDisc/Getty RF; **p. 1142:** © Comstock/JupiterImages RF.

Chapter 30

Opener: © The McGraw-Hill Companies, Inc./Jill Braaten, photographer; **30.1:** © Getty Images; **30.4 (balls):** © Dynamicgraphics/JupiterImages RF; **30.4 (paint):** © Royalty-Free/Corbis; **30.4 (blankets):** © Fernando Bengoe/Corbis; **30.4 (super glue):** © The McGraw-Hill Companies, Inc./John Thoeming, photographer; **p. 1159 (top):** DSM Dyneema; **p. 1159 (bottom):** © Bob Krist/Corbis; **p. 1161:** © The McGraw-Hill Companies, Inc./John Thoeming, photographer; **p. 1166:** © Rim Light/PhotoLink/Getty RF; **30.9 (corn):** © Royalty-Free/Corbis; **30.9 (swimsuit):** DuPont; **p. 1171 (both):** Ecoflex/www.basf.de/ecoflex.

Spectra Art

Chapters 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 25: Spectra courtesy of the Chemistry Department at Rutgers University.

Index

Page numbers followed by f indicate figures; those followed by t indicate tables. An A— before page numbers indicates appendix pages.

- ABS, 1174
Acebutolol, 110
Acetal(s)
 conversion to hemiacetals, 807–808
 cyclic, 804
 in carbohydrates, 812–813
 conversion of cyclic hemiacetal to, 811–812
 glycosides, 1043–1045
 hydrolysis of, 807–808
 as protecting groups, 808–809
 structure of, 1042
 synthesis of, 820
 addition of alcohols to aldehydes and ketones, 804–807
 from hemiacetals, 805–806, 811–812
 terminology of, 804
Acetaldehyde, 820
 aldol reaction of, 917, 919, 922
 ethanol oxidized to, 451
 structure of, 777
Acetamide, structure of, 826, 831
Acetaminophen, 845, 856, 874, 999, 1132
Acetanilide, 717
Acetate
 formation from acetaldehyde oxidation, 451
 as nucleophile, 243–245, 252
 resonance structures of, 67–68, 67f, 575–576, 701, 703f
Acetate (fabric), 1060
Acetic acid, 107, 695, 716, 872
 acidity of, 66–67, 70, 701, 704
 as Brønsted–Lowry acid, 55f
 electrostatic potential plot of, 67f, 689, 689f
 in ethyl acetate formation, 218–219
 glacial, 695
 halogenation of carbonyl compounds in, 893
 melting point of, 93
 NMR spectrum of, 523
 pK_a of, 701
 as polar protic solvent, 240f
 resonance structures, 578
 structure of, 691t, 826
Acetic anhydride, 826, 830, 1046
Acetic benzoic anhydride, 830
Acetoacetic ester synthesis, 900, 903–905, 932
Acetone, 566
 acidity of, 67
 aldol reactions of, 922
 as Brønsted–Lowry base, 55f, 56
 dipoles in adjacent molecules, 89
 odor of, 783
 as polar aprotic solvent, 241, 241f
 solubility of, 94–95
 structure and bonding of, 39, 778, 783
 surface area and boiling point of, 91f
 uses of, 783
Acetone enolate, 885, 891f
Acetonitrile
 acidity of, 67
 deprotonation of, 62
 as polar aprotic solvent, 241f
 structure of, 832f
p-Acetophenetidin, 874
Acetophenone, 731, 913, 946, 996
 aldol reaction of, 919
 structure of, 778
N-Acetyl amino acids, 1082
Acetylation, 845
Acetyl chloride, 226, 1046
 structure of, 826, 830
Acetylcholine, 861
Acetyl CoA, 860–862, 1134, 1140
1-Acetylcyclohexene, 909
1-Acetylcyclopentene, 945
Acetylene
 acetylide anion formation from, 418–419
 in acid–base reactions, 61–62
 acidity of, 67–68, 69f
 bond dissociation energies of, 400–401
 bonds in, 38, 39, 39f, 40f
 combustion of, 402
 deprotonation of, 418
 disparlure synthesis from, 441, 442f
 electrostatic potential map of, 405, 405f
 functional group in, 83, 84t
 Lewis structure, 15
 molecular shape of, 24, 26f
 nomenclature, 402
N-Acetyl-D-glucosamine (NAG), 862, 1061–1062
Acetyl group, 1046
 abbreviation for, 1006
 structure, 779
Acetylide anions, 61–62
 conversion of terminal alkynes to, 414
 formation by deprotonation of terminal alkynes, 405–406
 as organometallic compounds, 740
 reactions with alkyl halides, 414–416
 reactions with epoxides, 416–417
 retrosynthetic analysis with, 418–419
 synthesis from acetylene, 418–419
 in synthesis of disparlure, 441, 442f
Acetylsalicylic acid, 54, 696, 856. *See also* Aspirin
Achiral carbocation, 376
Achiral enolate, 891
Achiral epoxides, 443
Achiral intermediate, 381
Achiral meso compound, 178, 382, 382f
Achiral molecules, 164–166, 165f
 halogenation of, 549–550
 optical inactivity of, 182–183
Achiral product, 376
Achiral reactant, 344, 376
Acid(s)
 amino. *See* Amino acids
 Brønsted–Lowry
 carboxylic acids, 699–703
 description of, 55–56
 reactions of, 56–58
 carboxylic. *See* Carboxylic acid(s)
 as catalysts, 218–219
 common, 70
 conjugate, 56–58, 236–237, 237t
 dehydration of alcohols to alkenes in strong, 325–330
 as electron pair acceptors, 55, 58, 72
 in epoxide ring opening reactions, 345–347
 Lewis, 72–74
 as proton donors, 55
 reaction of ethers with strong, 341–343
 strength, 58–70
 sulfonic, 709–710
Acid–base reactions
 Brønsted–Lowry, 56–58, 254
 description, 56–58
 Lewis, 73–74, 235
 organometallic reagents and, 741
 predicting outcome of, 61–62
Acid chlorides
 acidity/basicity of, 829, 829t
 boiling point of, 834t
 functional group in, 86t
 IR spectra of, 835, 835t
 leaving group, 725
 melting point of, 834t
 NMR spectra of, 836
 nomenclature of, 830, 833, 833t
 nucleophilic substitution reactions, 725
 odor of, 842
 physical properties of, 834t
 pK_a , 829t
 reactions of, 842–843, 858t
 with ammonia or amines to form amides, 975–976
 conversion to amides, 842
 conversion to anhydrides, 842, 843
 conversion to carboxylic acids, 842, 843
 conversion to esters, 842
 Friedel–Crafts acylation reactions, 648
 with organometallic reagents, 750–753
 reduction of, 734
 solubility of, 834t
 stability of, 829
 structure and bonding of, 722, 826, 828–829
 synthesis from carboxylic acids, 845–847, 846f
Acidity
 basicity, inverse relationship with, 59, 59t
 of carbon–hydrogen bonds, 64
 of carboxylic acids, 689, 699, 700–705, 703f
 determining relative acidity of protons, 69–70
 electronegativity and, 63–66
 factors affecting, 62–70
 element effects, 63–65, 69f
 hybridization effects, 67–68, 68f, 69f
 inductive effects, 65–66, 66f, 69f
 resonance effects, 66–67, 67f, 69f
 periodic trends in, 63–65
 Acidity constant (K_a), 58–59
 Acid rain, 70
 Acrolein, 591f
 Acrylic acid, 561
 Acrylonitrile, 1155f
 Activation energy. *See* Energy of activation (E_a)
 Active methylene compounds, 923–924, 934
 Active site, enzyme, 219, 220f
 Acyclic alkanes
 branched-chain, 115
 conformations of, 129–137
 butane, 134–137, 134f, 136f
 ethane, 129–134, 132f, 133f
 constitutional isomers, 115–117, 118t
 having more than five carbon atoms, 117–118, 118t
 having one to five carbon atoms, 114–117, 118t
 homologous series, 117, 118t
 molecular formula, 114, 118, 118t
 normal (*n*) (straight-chain), 117–118, 118t
 structure, 114–115
 torsional and steric strain energies in, 136t
Acyases, 1084
Acylation
 definition of, 648
 Friedel–Crafts. *See* Friedel–Crafts acylation
 acylation
 Acyl derivatives, 722
 Acyl group, 648, 830
 naming, 779
 Acylium ion, 471–472, 649–650
 Acyl transfer reactions, 860–862, 861f
 Addition polymers. *See* Chain-growth polymers
 Addition reactions
 in alkenes, 370–390, A–14
 anti addition, 371, 377, 378t, 380, 382, 384
 halogenation, 371f, 379–383, 382f
 halohydrin formation, 371f, 383–385, 385f, 385t
 hydration, 371f, 378–379, 390
 hydroboration–oxidation, 371f, 385–390, 387f, 388f, 389f, 389t
 hydrohalogenation, 371–378, 371f, 372f, 373f, 374f, 377f, 378t
 overview of, 370–371, 371f
 stereochemistry of, 371, 376–378, 378t
 syn addition, 371, 377, 378t, 386, 388–389
 allylic bromination, 554
 anti additions, 434, 436, 442, 443
 carbene addition, 1013
 conjugate, 934
 in conjugated dienes
 Diels–Alder reaction, 588–597, 589f
 electrophilic, 584–586
 kinetic *versus* thermodynamic product, 586–588
 description of, 199
 dihydroxylation, 442–444
 epoxidation, 439–442, 451–454
 exothermic reactions, 372, 372f
 oxidative, 1006, 1008, 1011
 radical reactions, 554, 558–560
 syn additions, 434, 440, 442, 442f, 443–444, 1013
Adenine, 1064, 1065f
S-Adenosylmethionine (SAM), 250
ADHD, 191. *See also* Attention deficit hyperactivity disorder
-adiene (suffix), 362
Adipic acid, 875, 1160
Adipoyl chloride, 859
Adrenal cortical steroids, 1141, 1142

- Adrenaline, 228
 structure of, 958
 synthesis of, 228, 251, 251f
- Advair. *See* Fluticasone
- Advil. *See* Ibuprofen
- Aerosol propellant, 233, 551
- Agent Orange, 646f
- Aglycon, 1045
- al (suffix), 776
- Alanine, 105
 abbreviation for, 711t
 acid–base reactions of, 712, 712f
 in dipeptide synthesis, 1094–1098
 enantiomers of, 168f
N-acetyl, 1082–1084
 isoelectric point of, 713, 1078t
 in peptide formation, 1086–1087
 p*K*_a values for, 1078t
 structure of, 104f, 1076f
- Albuterol, 167, 347, 347f, 819, 1001
- Alcohol dehydrogenase, 451
- Alcoholic beverages, 319, 426
- Alcohols, 312–357, 449–450, 449f
 alkene preparation by acid-catalyzed
 dehydration, 369–370
 allylic, epoxidation of, 452–453
 boiling point of, 318t
 bonding in, 314
 classification of, 313
 dehydration reactions of, 324–331
 alkene synthesis in, 324–331
 carbocation rearrangements in,
 328–330
 dehydration to alkenes, 324–331,
 327f, 331f
 E1 mechanism in, 326
 E2 mechanism in, 327, 330–331
 enthalpy change in, 327, 327f
 of β -hydroxy carbonyl compounds,
 919–920
 Le Châtelier's principle in,
 327–328
 in phosphorus oxychloride and
 pyridine, 330–331
 in strong acid, 325–330
 electrostatic potential maps, 314f
 elimination reactions, 341
 ether formation from, 378, 379
 fragmentation pattern in mass
 spectrum, 471
 Friedel–Crafts alkylation of, 652
 functional group, 83, 85t
 β -hydroxy carbonyl compounds,
 917–928
 infrared (IR) spectra of, 482
 interesting examples, 319, 320f
 leaving groups of, 324, 330–332,
 335–336, 338–339
 melting point of, 318t
 NMR spectra of, 518, 519
 nomenclature, 314–316
 common names, 315–316
 for cyclic alcohols, 315, 315f
 IUPAC system, 314–316
 nucleophilicity of, 263
 oxidation of, 439f, 447–450, 451
 patchouli, 331f
 physical properties of, 318–319, 318t
 as polar protic solvents, 240, 240f, 264
 primary (1°), 390
 classification as, 313
 conversion to alkyl halides with
 HX, 332–334, 337t
 conversion to alkyl halides with
 phosphorus tribromide, 336,
 337t
 conversion to alkyl halides with
 thionyl chloride, 335–336, 337t
 dehydration by E2 mechanism, 327
 hydrogen bonding extent, 318
 from nucleophilic addition to
 aldehydes, 735
 oxidation to carboxylic acids, 447,
 448–450, 698
 from reactions of aldehydes
 with organometallic reagents,
 742–743
 from reduction of acid chlorides,
 734
 from reduction of aldehydes, 432,
 727, 728, 729
 from reduction of carboxylic acid
 derivatives, 727
 from reduction of carboxylic acids,
 735
 from reduction of esters, 734
 from reduction of monosaccharides,
 1047
 reactions of
 conversion to alkyl halides with
 HX, 331–335, 337t
 conversion to alkyl halides with
 thionyl chloride and phosphorus
 tribromide, 335–337, 337t
 conversion to alkyl tosylates,
 338–340
 elimination reactions, 323–331
 general features, 323–324
 nucleophilic substitution reactions,
 323–324, 331–338
 summary of, 341, 341f
 secondary (2°), 390
 classification as, 313
 conversion to alkyl halides with
 HX, 332–334, 337t
 conversion to alkyl halides with
 phosphorus tribromide, 336,
 337t
 conversion to alkyl halides with
 thionyl chloride, 335–336, 337t
 dehydration by E1 mechanism,
 326
 hydrogen bonding extent, 318
 oxidation to ketones, 447, 448, 450
 from reactions of aldehydes
 with organometallic reagents,
 742–743
 from reduction of ketones, 432,
 727, 728, 729
 solubility, 318t
 structure of, 313, 314
 substitution reactions, 341
 synthesis of, 321–322
 from alkyl halides, 321–322
 hydration of alkenes, 378–379
 hydroboration–oxidation of alkenes,
 385–390
 by nucleophilic substitution
 reaction, 321–322
 reactions of organometallic reagents
 with aldehydes and ketones,
 742–743, 789–790
 reactions of organometallic reagents
 with epoxides, 754–755
 reduction of epoxides, 437–438,
 438f
- tertiary (3°)
 classification as, 313
 conversion to alkyl halides with
 HX, 332–334, 337t
 dehydration by E1 mechanism, 326
 hydrogen bonding extent, 318
 from reaction of esters and acid
 chlorides with organometallic
 reagents, 750–752
 from reaction of ketones with
 organometallic reagents,
 742–743
- Aldaric acids, 1047, 1049, 1054–1056
D-aldaric acids, 1052
- aldehyde (suffix), 777
- Aldehyde dehydrogenase, 451
- Aldehyde group, as electron-withdrawing
 group, 655–656
- Aldehydes. *See also* Carbonyl
 compounds
 aldol reactions of, 917–919, 922–923,
 925–926
 boiling point of, 780t
 carbohydrates, 812–813
 electrostatic potential map of, 723,
 723f
 fragmentation pattern in mass
 spectrum, 471
 functional group in, 86t
 hydration of, 802–804
 kinetics of hydrate formation,
 803–804
 thermodynamics of hydrate
 formation, 802–803
- hydroxy
 cyclization of, 810, 813
 β -hydroxy, synthesis in aldol
 reactions, 917–928
 intramolecular cyclization of, 810,
 813
 melting point of, 780t
 nomenclature, 776–777, 779
 common names, 777, 777f
 IUPAC system, 776, 777f
 nucleophilic addition reactions, 724,
 774–813
 acid-catalyzed, 787
 carbanion addition, 789–790
 cyanide addition, 790–791
 hydride addition, 789
 mechanism of, 786–787
 nucleophiles in, 787–788, 788f
 primary amine addition, 797–799,
 801f
 secondary amine addition, 800–801,
 801f
 odors of, 783, 783f
 oxidation reactions, 726–727,
 738–739
 to carboxylic acids, 1047–1048
 physical properties of, 779, 780t
 p*K*_a, 886t
 polyhydroxy, 812–813
 protecting groups for, 808–809
 reactions of
 with amines, 975
 at α carbon, 785–786
 at carbonyl carbon, 785, 786
 halogenation at α carbon, 892–895,
 895f
 with organometallic reagents,
 742–745, 745f, 789–790
 reactivity of, 724
 reduction of, 789–790
 to alcohols, 432, 727, 728, 729
 reductive amination to amines,
 963–966
 solubility of, 780t
 structure of, 722, 775
 synthesis of, 784
 alcohol oxidation, 447, 448–449,
 450
 carboxylic acid reduction, 727
 hydroboration–oxidation of alkynes,
 412–413, 784
 hydrolysis of acetals, 807–808
 hydrolysis of imines and enamines,
 801
 oxidation of primary alcohols, 784
 oxidative cleavage of alkenes,
 444–446, 785
 reduction of acid chlorides, 734
 reduction of acid chlorides and
 esters, 784
 reduction of esters, 734
 reduction of nitriles, 865
 unreactivity to nucleophilic
 substitution, 725–726
 Wittig reaction, 792–797
- Alditol, 1047
- Aldohexoses, 1053–1056
D-aldohexoses, 1034–1035, 1035f,
 1054f
 Haworth projections of, 1039–1040
 pyranose ring of, 1037
 stereogenic centers of, 1032, 1034
- Aldol reactions
 crossed, 921–924
 synthetically useful reactions,
 922–924, 923f
 useful transformations of aldol
 products, 924, 924f
 dehydration of aldol product, 919–920
 directed, 925–926, 925f
 features of, 917–919, 918f
 intramolecular, 926–928, 928f
 in Robinson annulation, 936–939
 mechanism of, 917–918
 retrosynthetic analysis using, 920–921
- Aldonic acid, 1047–1048
- D*-Aldopentoses, 1034, 1035f, 1053, 1054f
- Aldoses
D-aldose family, 1034–1035, 1035f
 in Kiliani–Fischer synthesis, 1049,
 1051–1056
 oxidation of, 1047–1049
 reduction of, 1047
 structure of, 1028–1029
 in Wohl degradation, 1049, 1050
- Aldosterone, 1142
- Aldotetroses
D-aldotetroses, 1034, 1034f, 1035f,
 1052
 stereoisomers of, 1034, 1034f
- Aldrin, 604
- Alendronic acid (Fosamax), 18
- Aleve. *See* Naproxen
- Aliphatic carboxylic acids, 703–704
- Aliphatic hydrocarbons, 83, 114. *See also*
 Alkanes; Alkenes; Alkynes
- Aliskiren, 167
- Alizarin, 988
- Alizarine yellow R, 989
- Alkaloids, 957, 957f
- Alkanes, 113–158
 acyclic
 branched-chain, 115
 conformations of, 129–137, 132f,
 133f, 134f, 136f
 constitutional isomers, 115–117,
 118t
 having more than five carbon atoms,
 117–118, 118t
 having one to five carbon atoms,
 114–117, 118t
 homologous series, 117, 118t
 molecular formula, 114, 118, 118t
 normal (*n*) (straight-chain), 117–
 118, 118t
 structure of, 114–115
 torsional and steric strain energies
 in, 136t
 boiling point of, 129, 130t
 combustion of, 148–149
 cycloalkanes
 molecular formula, 114, 118
 naming, 125–127, 126f, 127f
 structure, 114, 118
 substituted, 141–147, 143f, 146f
- fossil fuels, 128–129
- halogenation, 541–551
 of achiral starting material,
 549–550, 549t
 bromination, 546–547, 547f

- Alkanes (*continued*)
 halogenation (*continued*)
 of chiral starting material, 549t, 550
 chlorination, 544–548, 544f, 545f, 548f, 549–551
 energy changes during, 544, 544f, 545f
 mechanism of, 542–544
 in organic synthesis, 548–549
 stereochemistry of, 549–551, 549t
 infrared (IR) spectra of, 481
 melting point of, 129, 130t
 nomenclature, 119–128, 123f
 common names, 127–128, 128f
 IUPAC system, 121–127, 123f
 substituents, 120–121
 nonreactivity of, 84
 oxidation of, 147–149
 radical reactions, 541
 reduction of an alkyne to, 434–435
 solubility of, 129, 130t
 synthesis of
 reduction of alkenes, 427, 428–432, 428f
 reduction of alkyl halides, 437–438, 438f, 741
 reduction of alkynes, 428f, 434–435
- Alkene oxides, 317
- Alkenes, 358–398
 addition reactions, 370–390
 anti addition, 371, 377, 378t, 380, 382, 384
 halogenation, 371f, 379–383, 382f
 halohydrin formation, 371f, 383–385, 385f, 385t
 hydration, 371f, 378–379, 390
 hydroboration–oxidation, 371f, 385–390, 387f, 388f, 389f, 389t
 hydrohalogenation, 371–378, 371f, 372f, 373f, 374f, 377f, 378t
 overview of, 370–371, 371f
 stereochemistry of, 371, 376–378, 378t
 syn addition, 371, 377, 378t, 386, 388–389
 amino acid synthesis from, 1085–1086
 boiling point of, 365–366
 classification of, 281–282, 281f
 constitutional isomers of, 325
 conversion to alkynes, 404–405
 conversion to cyclopropanes, 1012–1014
 in coupling reactions with organic halides (Heck reaction), 1003, 1009–1011
 cyclopropane synthesis in Simmons–Smith reaction, 1014–1015
 degrees of unsaturation, calculating, 360–361, 431–432
 dehydration of alcohols to, 324–331
 diastereoisomers of, 282–283
 dihydroxylation of, 442–444
 disubstituted, 281, 281f, 288, 289–290, 293, 297–298
 double bonds in, 281–283, 359, 360t
 restricted rotation around, 282–283, 282f
 electrophiles, reactions with, 103
 electrostatic potential plots, 370, 370f
 epoxidation of, 439–442, 439f
 fatty acids, 358, 367–369, 368f, 368t
 Friedel–Crafts alkylation with, 652
 functional group in, 83, 84t
 hybrid orbitals in, 281, 284
 hydrogenation of, 429–434
 alkene stability and, 429–430, 430f
 degrees of unsaturation, determination of, 431–432
 cis and trans isomers compared, 429–430
 mechanism of catalytic hydrogenation, 430–431
 infrared (IR) spectra of, 481
 interesting examples, 366, 366f, 367f
 internal, 359
 lipids, 366–369
 melting point of, 365, 368, 368t
 metathesis, 1015–1019, 1016f, 1018f
 molecular formula of, 360
 monomers in polymerization, 561–563, 562t
 monosubstituted, 281, 281f, 288
 NMR spectra of, 499–500, 516–517, 516f, 517f
 nomenclature, 362–365
 common names, 365, 365f
 IUPAC system, 362–363
 for stereoisomers, 363–364
 organic synthesis, 391–392
 oxidative cleavage of, 444–446, 785
 physical properties of, 365–366, 368t
 polymerization of, 1150–1156, 1153, 1157–1158
 radical reactions, 541, 558–560
 reduction of, 427, 428–432, 428f
 solubility of, 365
 stability of, 283–284, 429–430, 430f
 stereoisomers of, 282–284
 alkyne reduction and, 434–436
 synthesis of, 369–370
 amine conversion, 977–980
 comparison of methods, 796–797
 dehydration of alcohols, 369–370
 dehydrohalogenation of alkyl halides with strong base, 369, 370
 by elimination reactions, 279–280, 369–370
 by Hofmann elimination reaction, 977–980
 reduction of alkynes to cis alkenes, 434, 435
 reduction of alkynes to trans alkenes, 434, 436
 substituted alkene in Heck reaction, 1009–1010
 Wittig reaction, 792–797
 terminal, 359
 tetrasubstituted, 281
 trisubstituted, 281, 281f, 289–290, 293, 297–298
- Alkenols, 363
- Alkoxides, 280
 electrostatic potential plot, 885, 885t
 in ether synthesis, 322
 protonation of, 730, 733, 743, 751, 754–755
- Alkoxy group, 85t, 316
- Alkylamines
 basicity of, 969–972, 973t, 974t
 reaction with nitrous acid, 980–981
- Alkylation reactions
 in acetoacetic ester synthesis, 900, 903–905
 of diethyl malonate derivative, 1079–1080, 1081f
 of enolates, 897–899
 general features of, 897–899
 tamoxifen synthesis, 899
 of unsymmetrical ketones, 898–899
 Friedel–Crafts. *See* Friedel–Crafts alkylation
 in malonic ester synthesis, 900–903
 polyalkylation, 666
- Alkyl benzenes
 electron density of, 655
 halogenation of, 669–671
 oxidation of, 671–672, 698
 synthesis of
 Friedel–Crafts alkylation, 643f, 647–648
 by reduction of aryl ketones, 672–673
- Alkylboranes
 formation by hydroboration, 385–387, 387f
 oxidation of, 388–389
- Alkyl bromides
 mass spectrum of, 469, 469f
 synthesis of, 558–560
- Alkyl chlorides
 in Friedel–Crafts alkylation reactions, 648–649, 651
 mass spectrum of, 468, 469f
- Alkyl diazonium salts, 981
- Alkyl group, 120, 121, 231, 235, 365, 647, 830
 in alkyl tosylates, 338
 carbon radical stability and, 539–540, 540f
 as electron-donating group, 655, 656
 in ethers, 313
 nomenclature
 in alcohols, 315
 for branched substituents, A–3–4
 in ethers, 316
 stabilization of carbonyl group by, 802
- Alkyl halides
 acetylide anion reactions with, 414–416
 alkane formation from, 741
 allylic halides, 229, 229f
 aryl halides, 229, 229f, 267
 benzylic halides, 229, 229f
 classification of, 229–230, 229f, 301
 conversion to alcohols to, 331–337
 with HX, 331–335, 337t
 importance of, 337–338
 with thionyl chloride and phosphorus tribromide, 331–335, 337t
 conversion to carboxylic acids, 754
 conversion to primary amines, 962
 coupling reaction with organocuprates, 1003
 dehydrohalogenation with strong base, 369, 370
 in direct enolate alkylation, 897–899
 elimination reactions, 234, 278–311
 E1 mechanism, 300–304
 E2 mechanism, 287, 288t, 295, 300–304, 979–980, 979f
 examples, 229f, 232–234, 233f
 functional group, 85t
 mass spectrometry of, 468, 469, 469f
 molecular formula of, 229
 nomenclature, 230–231, 231f
 common names, 231
 IUPAC, 230, 231f
 nucleophiles, reactions with, 103
 nucleophilic substitution reactions, 234–269
 alcohol synthesis, 321–322
 alkyl halide structure and, 248, 255, 262, 262f
 with ammonia or amines to form amines, 960
 ether synthesis, 321–322
 S_N1 mechanism, 300–304
 S_N2 mechanism, 300–304
 physical properties, 231–232, 232t
 polar carbon–halogen bond, 234
 primary (1°)
 acetylide anion reactions with, 414–415
 classification as, 229–230
 E2 elimination reactions, 287, 288, 302f
 example of, 229f
 S_N2 substitution reactions, 248, 262, 262f, 302f
 synthesis of, 332, 335, 336
 reaction mechanism, determining type of, 300–304
 reduction of, 437–438, 438f
 secondary (2°)
 acetylide anion reactions with, 414–415
 classification as, 229–230
 E1 elimination reactions, 292, 303f
 E2 elimination reactions, 287, 302f–303f
 example of, 229f
 S_N1 substitution reactions, 255, 262, 262f, 303f
 S_N2 substitution reactions, 248, 262, 262f, 302f
 synthesis of, 332, 335, 336
 synthesis of
 halogenation of alkanes, 541–549
 hydrohalogenation of alkenes, 371–378, 371f, 372f, 373f, 378t
 tertiary (3°)
 acetylide anion reaction with, 414
 classification as, 229–230
 E1 elimination reactions, 292, 294, 302f
 E2 elimination reactions, 287, 288, 302f
 example of, 229f
 S_N1 substitution reactions, 255, 262, 262f, 302f
 synthesis of, 332, 334–335
 unreactive, 229, 229f
 vinyl halides, 229, 229f, 267
- 1,2-Alkyl shift, 328
- Alkylthio group, 85t
- Alkyl titanium compound, 1157
- Alkyl tosylates, 338–340, 341f, 369
 conversion of alcohols to, 338–339, 340
 reactions of, 339–340
- Alkynes, 399–425
 addition reactions
 halogenation, 406f, 409
 hydration, 406f, 409–412
 hydroboration–oxidation, 406f, 412–413
 hydrohalogenation, 406–408, 406f
 overview of, 405
 boiling point of, 402
 bond dissociation energies of, 400–401
 cycloalkynes, 401
 degrees of unsaturation, 400
 deprotonation of, 405–406, 406t
 functional group in, 83, 84t
 hydroboration of, 1007, 1009
 hydroboration–oxidation of, 406f, 412–413
 infrared (IR) spectra of, 481
 interesting examples, 402–404, 402f–404f
- internal
 definition of, 400
 hydration of, 410
 hydroboration–oxidation of, 412–413
 oxidative cleavage of, 446–447, 698
 synthesis in acetylide anion reactions with alkyl halides, 414–416
 melting point of, 402
 molecular formula of, 400
 nomenclature, 401–402
 oxidative cleavage of, 446–447, 698
 physical properties of, 402
 reduction of, 427, 428f, 434–437
 to alkanes, 428f, 434–435

- to cis alkenes, 434, 435
to trans alkenes, 434, 436
solubility of, 402
structure of, 400–401
synthesis of, 404–405, 417–419
from alkenes, 404–405
by dehydrohalogenation, 299–300, 300f
from dihalides, 404–405
E2 elimination reactions, 279, 299–300, 300f
multistep, 418–419
terminology and conventions, 417–418
- terminal
conversion to acetylide anions, 414–415
definition of, 400
hydration of, 410
hydroboration–oxidation, 412–413
oxidative cleavage of, 446–447, 698
reaction as an acid, 405–406
triple bond of, 400–401
- D-Allaric acid, 1049
Allegra. *See* Fexofenadine
Allenene, 195
Allicin, 119
D-Allonic acid, 1071
D-Allose, 1035f, 1049
Allyl group, 365
Allylic alcohol
epoxidation of, 452–453
synthesis from α,β -unsaturated carbonyl compounds, 756
synthesis from aldol reactions, 924, 924f
- Allylic carbocations, 585
as conjugated system, 572, 573–574
electrostatic potential plot, 574, 574f
NMR spectra of, 575, 575f
resonance of, 574–576
- Allylic carbon
oxidation of, 433, 433f
radical halogenation at, 552–555
bromination, 553–554
product mixtures in, 555
in triacylglycerols, 433, 433f
- Allylic halides, 229, 229f
Allylic substitution, 554
Allyl radical, 552–555
Alpha (α) carbon, 279–280, 691, 694, 710
of aldehydes and ketones, 777, 785–786
substitution reactions of carbonyl compounds at the α carbon, 880–906
- Alpha (α) cleavage, 471–472
Alpha (α) elimination reaction, 1012
Alpha (α) helix, 1101–1104, 1102f, 1105f, 1107f
- Alpha (α) hydrogens, 881, 922, 925
Altoacid. *See* Methoprene
D-Altrose, 1035f, 1069
Aluminum chloride, in Friedel–Crafts alkylations, 647–653, 665–666
Alzheimer's disease, 85, 926, 975
Amberlyst A-26 resin, 450–451
Ambident nucleophile, 888
-amide (suffix), 831–832, 833t
Amide bond in peptides, 1086, 1101
Amides
basicity of, 829t, 971–972, 973t, 974t
boiling point of, 834t
cyclic, 827
functional group in, 86t
hydrolysis of, 855–856
interesting examples, 837–838
IR spectra of, 835, 835t
leaving group, 725
melting point of, 834t
nitrogen bases, 70, 71f
NMR spectra of, 836
nomenclature, 831–832, 833t
nucleophilic substitution reactions, 725
physical properties of, 834, 834f, 834t
 pK_a , 829t, 886t
polyamides, 825, 829
primary (1°), 827, 831, 834–836, 842, 844, 850
tautomers, 863
as protecting group for an amine, 977f
proteins, 837
reduction to amines, 736–737, 963
secondary (2°), 827, 831–832, 834–836, 842, 844, 850
solubility of, 834t
stability of, 829
structure and bonding of, 722, 826, 827, 828–829
synthesis of
from acid chlorides, 842, 975–976
from anhydrides, 844–845, 975–976
from carboxylic acids, 846f, 849–850
from esters, 850
tautomers, 863–864
term usage, 887
tertiary (3°), 827, 831–832, 842, 844, 850
enolate formation from, 885–886
- Amide tautomer, 863–864
Amikacin, 1062
-amine (suffix), 952–953
Amine oxide, 444
Amines, 949–991
aromatic
basicity of heterocyclic, 972, 973t, 974t
nomenclature, 953
basicity of, 950, 966–975
aromaticity effects on, 972, 973t
hybridization effects on, 972–973, 973t
inductive effects on, 969, 973t
 pK_a values, 966, 968–969, 974t
relative to other compounds, 968–975, 974t
resonance effects of, 973t
boiling point of, 954t
classification of, 950
electrostatic potential plots of, 951, 951f
functional group, 85t
general molecular formula for, 955
infrared (IR) spectra of, 483
interesting examples, 956–959
derivatives of 2-phenylethylamine, 958–959, 959f
histamine and antihistamines, 957–958
simple amines and alkaloids, 956–957, 957f
IR spectra of, 955, 955f
mass spectra of, 955, 955f
melting point of, 954t
nitrosamine formation, 261
NMR spectra of, 956, 956f
nomenclature of, 952–954
as nucleophiles, 950, 975–977
odors of, 956–957
physical properties of, 954, 954t
primary (1°)
addition to aldehydes and ketones, 797–799, 799f, 801f
amino acids, 1077
from direct nucleophilic substitution, 960–961
Gabriel synthesis, 961–962
IR spectra of, 955, 955f
nomenclature, 952
reactions with acid chlorides and anhydrides, 975–976
reactions with aldehydes and ketones, 975
reactions with nitrous acid, 980–981
from reduction of amides, 963
from reduction of nitriles, 865, 962
from reduction of nitro compounds, 962
structure of, 950
protecting group for, 976, 977f
reactions of
with acid chlorides, 842, 975–976
with aldehydes and ketones, 975
with anhydrides, 844, 845, 975–976
with carboxylic acids, 846f, 849–850
conversion to alkenes, 977–980
conversion to amides, 975–976
with esters, 850
general features, 966
Hofmann elimination, 977–980
with monosaccharides, 1062
with nitrous acid, 980–982
secondary (2°)
addition to aldehydes and ketones, 800–801, 801f
from direct nucleophilic substitution, 960
IR spectra of, 955, 955f
nomenclature, 952–953
proline, 1077
reactions with acid chlorides and anhydrides, 975–976
reactions with aldehydes and ketones, 975
reaction with nitrous acid, 981–982
from reduction of amides, 963
structure of, 950
separation from other organic compounds, 967–968, 967f
solubility of, 954t
spectroscopic properties of, 955–956, 955f
structure and bonding of, 950–952
synthesis of, 960–966
direct nucleophilic substitution reactions, 960–961
Gabriel synthesis, 961–962
reduction of amides, 736–737, 963
reduction of nitriles, 962
reduction of nitro compounds, 962
reductive amination of aldehydes and ketones, 963–966
tertiary (3°)
from direct nucleophilic substitution, 960
IR spectra of, 955, 955f
nomenclature, 952–953
reactions with acid chlorides and anhydrides, 975–976
from reduction of amides, 963
structure of, 950
- Amino acids, 104–105, 710–713. *See also* Peptides; Proteins
abbreviations for, 711t, 1076f
N-acetyl, 1082–1084
acid–base behavior of, 711–712, 712f, 1077–1078, 1077f, 1078t
 α -amino acids, 710, 1075, 1075f
D-amino acids, 710–711, 1075, 1075f
L-amino acids, 710–711, 1075, 1075f
analysis of peptide composition, 1091
C-terminal, 1087–1088
enantiomers of, 1075, 1075f
separation of, 1081–1084
essential, 711, 1077
general features of, 1075, 1075f, 1077
isoelectric points, 713
naturally occurring, 1075, 1076f
reactions with acid chlorides and anhydrides, 975–976
reactions with aldehydes and ketones, 975
reactions with nitrous acid, 980–981
from reduction of amides, 963
from reduction of nitriles, 865, 962
from reduction of nitro compounds, 962
structure of, 950
protecting group for, 976, 977f
reactions of
with acid chlorides, 842, 975–976
with aldehydes and ketones, 975
with anhydrides, 844, 845, 975–976
with carboxylic acids, 846f, 849–850
conversion to alkenes, 977–980
conversion to amides, 975–976
with esters, 850
general features, 966
Hofmann elimination, 977–980
with monosaccharides, 1062
with nitrous acid, 980–982
secondary (2°)
addition to aldehydes and ketones, 800–801, 801f
from direct nucleophilic substitution, 960
IR spectra of, 955, 955f
nomenclature, 952–953
proline, 1077
reactions with acid chlorides and anhydrides, 975–976
reactions with aldehydes and ketones, 975
reaction with nitrous acid, 981–982
from reduction of amides, 963
structure of, 950
separation from other organic compounds, 967–968, 967f
solubility of, 954t
spectroscopic properties of, 955–956, 955f
structure and bonding of, 950–952
synthesis of, 960–966
direct nucleophilic substitution reactions, 960–961
Gabriel synthesis, 961–962
reduction of amides, 736–737, 963
reduction of nitriles, 962
reduction of nitro compounds, 962
reductive amination of aldehydes and ketones, 963–966
tertiary (3°)
from direct nucleophilic substitution, 960
IR spectra of, 955, 955f
nomenclature, 952–953
reactions with acid chlorides and anhydrides, 975–976
from reduction of amides, 963
structure of, 950
- N-terminal, 1087–1088
peptide synthesis from, 1091–1100, 1101f
 pK_a values, 1077, 1078, 1078t
representative examples, 711t
resolution of
by conversion to diastereomers, 1081–1084, 1082f
kinetic, using enzymes, 1084
separation of, 1081–1084
stereogenic center of, 1075
structure of, 710, 1075, 1075f, 1076f, 1077
synthesis of, 1078–1081
alkylation of diethyl malonate derivative, 1079–1080, 1081f
enantioselective hydrogenation, 1085–1086
reaction of α -halo acids with ammonia, 1078–1079, 1081f
Strecker synthesis, 1080–1081, 1081f
zwitterion form, 711–712, 1077
- Aminobenzene. *See* Aniline
p-Aminobenzoic acid (PABA), 599, 990–991
Aminoglycoside antibiotics, 1062
Amino group, 85t, 665–666, 953
as electron-donating group, 654–656, 656f
inductive effect of, 654
nitro reduction to, 673–674
as ortho, para director, 662–663
resonance effect of, 654
- 6-Aminohexanoic acid, 1161
 α -Amino nitrile, 1080–1081
Amino sugars, 1061–1062
Ammonia
basicity of, 969
boiling point, 436
in dissolving metal reductions, 428, 436
hybrid orbitals in, 35, 35f
molecular shape, 26, 26f
 pK_a of, 60
reactions with
acid chlorides, 842, 975–976
alkyl halides to form amines, 960–961
anhydrides, 844, 975–976
esters, 850
 α -halo acids, 1078–1079
- Ammonium cation, of amino acids, 712
Ammonium salt
in amine extraction process, 967–968
enantiomers, 951
in Hofmann elimination reaction, 977–978
synthesis of, 950–951, 960–961
- Amoxicillin, 3, 52, 192, 838
Amphetamine, 72, 76, 191, 964
Amygdalin, 194, 791–792
Amylopectin, 1060–1061
Amylose, 1060–1061
 β -Amyrin, 1145
Anabolic steroids, 1142
Analgesics, 696, 845, 898, 906
Anastrozole, 862
Androsterone, 1141t, 1147
-ane (suffix), 117, 120, 121, 127, 362
Anesthetics, 84, 233f, 320, 673
Angiotensin, 1116
Angle strain, 137–139
in epoxide, 314
Angstrom (\AA), 23
Angular methyl groups, 1138–1139
Anhydrides
acidity/basicity of, 829t
boiling point of, 834t
cyclic, 826–827, 847

- Anhydrides (*continued*)
 derivation of term, 830
 IR spectra of, 835, 835t
 melting point of, 834t
 mixed, 826–827
 NMR spectra of, 836
 nomenclature, 830, 833t
 physical properties of, 834t
 pK_a , 829t
 reactions of, 844–845, 858t
 acetylation, 845
 with ammonia or amines to form amides, 975–976
 conversion to amides, 844–845
 conversion to carboxylic acids, 844
 conversion to esters, 844–845
 solubility of, 834t
 stability of, 829
 structure and bonding of, 826–827, 828–829
 symmetrical, 826–827, 830
 synthesis of
 from acid chlorides, 842, 843
 dehydration of dicarboxylic acids, 847
- Aniline, 970
 basicity of, 969–970
 electrophilic aromatic substitution reaction, 663
 electrostatic potential plot of, 656f, 971f
 nomenclature and, 612
 polyhalogenation of, 665
 resonance structures, 654
 structure of, 611, 953
 synthesis of
 electrophilic aromatic substitution reaction, 646
 by nitrobenzene reduction, 673
- Anionic polymerization, 1154–1155, 1155f
 of epoxides, 1156–1157
- Anions
 acetate, 66–67, 67f
 carbanion, 68, 201–202, 201f
 cyclopentadienyl, 624–625, 630, 630f
 definition of, 7
 enolate, 21
 naked, 241
 radical, 436
 salt formation, 10–11
 solvation by hydrogen bonding, 240
- Anisole, 343, 492
- Annulation
 origin of word, 936
 Robinson, 936–939
- Annulenes, 620–621
 18-annulene, 537
- Anomeric carbon, 1037–1040, 1042, 1044, 1056–1058, 1063
- Anomers, 1037–1040
- Antabuse, 451
- Anthracene, 621
- Anti addition, 371, 377, 378t, 380, 382, 384, 434, 436, 442, 443
- Antiaromatic compounds, 618, 619, 625, 626
- Antibiotics
 aminoglycoside, 1062
 cephalosporins, 838
 chloramphenicol, 167
 ionophores, 101–102
 β -lactam, 463, 485, 838, 856–857
 penicillin(s), 463, 485, 838, 856–857
 sulfa drugs, 990–991, 991f
- Antibonding molecular orbital, 627, 628f, 629–631
- Anticancer drugs
 aromatase inhibitors, 862
 doxorubicin, 1029f
- epothilone A, 1018f
 illudin-S, 905
 imatinib mesylate, 837
 tamoxifen, 880, 899
- Anti conformation, 134f, 135–137, 136f
- Antidepressant, 250, 646f, 653, 913, 958
- Antifreeze, 319f
- Antihistamines, 610, 646f, 957–958, 968, 975
- Anti-inflammatory drugs
 aspirin, 54, 71–72, 215, 268, 269f, 696, 697, 1131–1132
 fluticasone, 230
 ibuprofen, 119, 150, 187, 673, 730f, 916, 932, 1131–1132
 nabumetone, 906
 naproxen, 159, 187, 898
 nonsteroidal anti-inflammatory drugs (NSAIDs), 1131–1132
- Antimalarial drugs, 289f, 389, 389f, 624, 896, 974, 975
- Antioxidants, 557, 571
- Anti periplanar geometry, 295–298, 295f
- Antiseptics, 236
- Antiviral agents, 86, 1018f
- APC (aspirin, phenacetin, caffeine), 877
- Aprotic solvents, polar, 241, 241f, 264, 265, 265t, 287, 288t, 889–890
- Arabinose, 1054–1055
 D-arabinose, 1034, 1035f, 1051
- Arachidic acid, 1122t
- Arachidonic acid, 568, 570
 leukotriene synthesis by oxidation of, 348
 prostaglandin synthesis from, 697
 structure of, 97, 1122t
 synthesis of eicosanoids from, 1130, 1131f
- Arene, 608
- Arene oxide, 157
- Arginine
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
- Aricept. *See* Donepezil
- Arimidex. *See* Anastrozole
- Aripiprazole, 995
- Aromatase inhibitors, 862
- Aromatic compounds, 607–640
 background on, 608
 bonding and antibonding orbitals, 627–628
 buckminsterfullerene, 632
 charged, 634–626
 electrophilic substitution. *See* Electrophilic aromatic substitution reactions
 examples, 620–626
 charged compounds, 624–626
 heterocycles, 621–624, 623f
 multiple rings, 621
 single ring, 620–621
 Hückel's rule, 617–620
 basis of, 626–629
 interesting, 614–615, 614f
 molecular orbitals, 627–629, 628f
 nomenclature, 610–613
 predicting aromaticity using the inscribed polygon method, 629–631, 630f, 631f
 spectroscopic properties, 613–614, 613f, 613t
- Aromaticity, effect on basicity of amines, 972, 973t
- Arrows
 curved, 20–21, 56, 201, 235
 double-headed, 18, 56–57, 202t
 equilibrium, 56–57
 half-headed, 539
 open, 417
 use in describing organic reactions, 202–203, 202t
- Artemisinin, 194, 389, 389f
- Artificial sweeteners, 1058, 1059f, 1090
- Arylamines
 basicity of, 969–971, 973t, 974t
 reaction with nitrous acid, 980–981
- Arylboranes, 1007
- Aryl bromides, synthesis from aryl diazonium salts, 983
- Aryl chlorides
 examples of biologically active, 646f
 synthesis from aryl diazonium salts, 983
- Aryl diazonium salt, 980–988
- Aryl fluorides, synthesis from aryl diazonium salts, 983
- Aryl groups, 612
- Aryl halides, 229, 229f, 267
 coupling reactions
 with alkenes, 1009–1011
 with organoboranes, 1006–1007
 with organocuprates, 1003
 examples of biologically active, 646f
 reactivity of, 650
 synthesis of
 from aryl diazonium salts, 983
 halogenation of benzene, 643f, 645
- Aryl iodides, synthesis from aryl diazonium salts, 983
- Aryl ketones, reduction to alkyl benzenes, 672–673
- Ascorbic acid. *See* Vitamin C
- Asparagine
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
- Asparagopsis taxiformis, 232
- Aspartame, 109, 872, 1058, 1059f, 1090
- Aspartic acid, 1077, 1171
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
- Aspirin, 54, 71–72, 873
 crossing of cell membrane, 72
 derivation of word, 696
 mechanism of action, 215, 697, 1131–1132
 proton transfer reaction, 71–72
 structure, 696
 synthesis of, 268, 269f, 696
- Asthma, 347–348
- Asymmetric carbon. *See* Stereogenic centers
- Asymmetric reaction, 452
- Asymmetric reduction, 731
- Atactic polymer, 1157–1158
- ate (suffix), 692, 830–831, 833t
- Atenolol, 78, 85
- Atherosclerosis, 861f, 862
- Atmosphere
 carbon dioxide concentration in, 149, 149f
 ozone layer destruction, 233
- Atomic mass unit (amu), 7
- Atomic number, 7
- Atomic symbol, 7
- Atomic weight, 7
- Atom mass, infrared (IR) absorption and, 478–479, 479f
- Atoms
 components of, 7
 excited state, 32
 ground state, 32
- Atorvastatin, 1140, 1141f
- Atropine, 957, 957f
- Attention deficit hyperactivity disorder, 191. *See also* ADHD
- Automated peptide synthesis, 1099–1100, 1101f
- Avobenzene, 30, 931
- Axial bond, 160
- Axial hydrogens, 139–140, 139f, 141f
- Azo compound
 applications of, 988–991
 dyes, 988–991
 sulfa drugs, 990–991, 991f
 formation of, 982, 986–988
- Azo coupling, 982, 986–988
- AZT (azidodeoxythymidine), 4
- Azulene, 635
- Backside attack, in nucleophilic substitution reactions, 245–246
- Baekeland, Leo, 1166
- Baeyer strain theory, 137
- Bakelite, 1165, 1166f
- Ball-and-stick model
 of acyclic alkanes, 114–115
 of molecular shape, 24–26
- Barbituric acid, 119
- Barrier to rotation, 136
- Base(s)
 for alkyne deprotonation, 405–406, 406t
 for alkyne synthesis by dehydrohalogenation, 299–300, 300f
- Brønsted–Lowry
 description of, 55–56
 reactions of, 56–58
 substitution reactions with alkyl halides, 234
- common, 70–71
 conjugate, 56–58
 dehydration of β -hydroxy carbonyl compounds with, 920
 as electron pair donors, 55, 58, 72–73
 in elimination reactions, 279–280, 280t
 in E1 elimination reactions, 291, 293, 293t, 298
 in E2 elimination reactions, 286, 286f, 288t, 298
 enolate formation with, 887–888, 887t
 hydrides, 70, 71f
 Lewis, 72–74
 nitrogen-containing, 70, 71f
 nonnucleophilic, 239
 organometallic reagents, 739, 741
 oxygen-containing, 70, 71f
 as proton acceptors, 55
 strong, 70–71
 used to deprotonate carboxylic acids, 700, 700t
 weak as good leaving group in nucleophilic substitution reactions, 236–238
- Base pairs, 1064, 1065f
- Base peak, in mass spectrum, 465
- Bases, nucleoside, 1063–1065, 1065f
- Basicity
 acidity, inverse relationship with, 59, 59t
 of amines, 950, 966–975
 of carboxylic acid derivatives, 828–829, 829t
 of carboxylic acids, 699
 leaving group ability and, 236–238
 nucleophilicity compared to, 239, 240, 241
 periodic trends in, 63–65
- Basketane, 128f
- Beeswax, 150, 150f
- Benadryl, 968
- Benazepril, 995
- Benedict's reagent, 1048

- Bent molecular shape
 alcohols, 314
 description of, 26
 ethers, 314
 polarity and, 44
- Benzaldehyde, 492
 aldol reactions of, 922, 923, 923f
 electron density of, 656, 656f
 resonance structures of, 655
 secondary alcohol formed from, 743
 structure of, 777
- Benzalkonium chloride, 276
- Benzamide, structure of, 831
- Benzene, 52, 83, 84t, 108, 157, 607–640
 as annulene, 620
 background on, 608
 in BTX mixture, 614
p-chlorostyrene synthesis from, 675–676
 conjugation of, 618
 degrees of unsaturation, 608
 electrostatic potential plot of, 610, 610f, 656f
 Friedel–Crafts acylation of, 643f, 647–653, 672–673
 Friedel–Crafts alkylation of, 643f, 647–653, 672–673
 halogenation of, 643f, 645, 668
 heat of hydrogenation, 616–617, 616f
 hybridization and orbitals, 609–610
 lack of addition reactivity, 608, 617, 642
 molecular orbitals of, 628–630, 628f
 nitration of, 643f, 646–647, 669
p-nitrobenzoic acid synthesis from, 675
 NMR spectrum of, 505, 506f, 506t, 619–620
 pi (π) electrons of, 618, 626, 642
 resonance, 609
 resonance structures, 576
 stability, 615–617, 619
 structure of, 609–610
 substitution reactions, 608, 617, 642–687. *See also* Electrophilic aromatic substitution reactions
 sulfonation of, 643f, 646–647
 synthesis from aryl diazonium salts, 983–984
 synthesis of *m*-bromoaniline from, 674
 synthesis of *p*-bromonitrobenzene from, 668–669
 synthesis of *o*-nitrotoluene from, 669
 synthesis of 1,3,5-tribromobenzene from, 984, 984f
 trisubstituted benzene synthesis from, 676–677
- Benzenecarbaldehyde. *See* Benzaldehyde
- Benzene derivatives
 disubstituted, 666–668
 electrophilic aromatic substitution reactions of, 666–668
 halogenation of, 666–667
 nitration of, 667–668
 synthesis of, 668–669
 drugs containing benzene rings, 615f
 interesting examples, 614–615, 614f
 NMR spectra, 518–519, 519f
 nomenclature, 610–613
 aromatic rings as substituents, 612–613
 disubstituted, 611
 monosubstituted, 611
 polysubstituted, 611–612
 spectroscopic properties, 613–614, 613f, 613t
 substituted
 activating and deactivating substituents, 657–661, 660f
 directing effects of substituents, 657–664, 664f
 electrophilic aromatic substitution, 657–660, 660f, 665–666
 electrostatic potential plot of, 656, 656f
 Friedel–Crafts acylation of, 665–666
 Friedel–Crafts alkylation of, 665–666
 halogenation of, 665
 inductive effects on, 654–656
 multistep synthesis of, 675–677
 orientation effects on, 661–664
 oxidation of, 671–672
 reduction of, 672–674
 resonance effects on, 654–656
 synthesis of, 668–669, 675–677
 with diazonium salts, 984–986
 trisubstituted, synthesis of, 676–677
- Benzene ring
 of arylamines, 969
 NMR spectra of, 505, 506f, 506t, 518–519, 519f
- Benzenesulfonic acid, 643f, 646
- Benzo[*a*]pyrene, 349, 614, 614f
- Benzo[*a*]pyrene derivative, 97
- Benzocaine, 673, 874
- Benzoic acid
 acidity of, 705–707, 706f
 separation from cyclohexanol by extraction procedure, 707–709, 709f
 structure of, 691t
 substituted, 705–707, 706f
 synthesis of
 by oxidation of alkyl benzenes, 671, 698
- Benzonitriles, synthesis from aryl diazonium salts, 492, 983
- Benzophenone, structure of, 778
- Benzoquinone, 591f
- Benzoyle group, 779
- Benzphetamine, 996
- Benzylamine, 994, 996
- Benzyl bromide, 683
- Benzyl ester, 1097
- Benzyl group, 612
- Benzylic bromination, 670–671
- Benzylic C–H bond, 669–670, 698
- Benzylic halides, 229, 229f, 670–671
- Benzylic radical, 669–671, 1151–1152
- Benzyl methyl ether, 683
- Benzyl salicylate, 1177
- Bergstrom, Sune, 697
- Beryllium, octet rule exceptions in, 18
- Beryllium hydride
 hybrid orbitals in, 35
 molecular shape of, 24, 26f
- Beta (β) blocker, 85
- Beta (β) carbon, 279–280, 296, 691
- Beta (β) elimination reactions. *See also* Elimination reactions
 of alcohols, 323–331
 dehydrohalogenation as, 279
 Zaitsev rule, 288–291
- Beta (β) lactam(s), 311, 827, 838, 856–857
- Beta (β) lactam family of antibiotics, 463, 485
- Bextra. *See* Valdecoxib
- BHA (butylated hydroxy anisole), 568
- Biaryls, 1025
- Bicyclic compounds, naming, A–5–6
- Bicyclic product, 592–593
- Bicyclic ring system, bridged, 593, 593f
- “Big Six” synthetic polymers, 1169, 1169f
- Bile acid, 158
- Bile salt, 158
- Bilobalide, 153
- Bimatoprost, 697
- Bimolecular elimination. *See* E2 elimination reactions
- Bimolecular reactions
 elimination reactions, 285
 in nucleophilic substitution reactions, 243–244
 organic reactions, 217
- Biocatalyst, 1167, 1168f
- Biodegradable polymers, 1170–1171
- Biofuels, 1125
- Biomolecules, 104–105, 104f, 1028.
See also Carbohydrates; DNA; Lipids; Proteins
 definition of, 104
 families of, 104
 mass spectra of, 474
- 2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl (BINAP), 1085–1086
- Bisphenol A (BPA), 683, 1163, 1164f, 1168
- Blattellaquinone, 843, 844
- Bloch, Konrad, 1140
- Blood alcohol content, measurement of, 449–450, 449f, 485
- Boat form, of cyclohexane, 141, 141f
- Boc (*tert*-butoxycarbonyl protecting group), 1096–1098
- Boiling point
 of alcohols, 318t
 of aldehydes and ketones, 780t
 of alkanes, 129, 130t
 of alkenes, 365, 366
 of alkyl halides, 232t
 of alkyne, 402
 of amines, 954t
 of carboxylic acids, 693t
 definition of, 90
 of enantiomers, 182, 184, 184t
 of epoxides, 318t
 of ethers, 318t
 intermolecular forces and, 90–92
 polarizability and, 91
 of racemic mixtures, 184, 184t
 separation of liquids based on, 92, 92f
 surface area and, 91
- Bombykol, 1002, 1008, 1008f
- Bond(s) and bonding, 10–12. *See also* specific bond types
 bond angle, 24–27, 26f
 bond length, 23, 23t, 40–42, 41t
 breaking and making in organic reactions, 200–203, 201f
 covalent, 10, 11–12
 definition of, 10
 double, 15
 ionic, 10–11
 Lewis structures, 12–18
 in nucleophilic substitution reactions, 242–243, 252
 octet rule, 10
 π (π), 37–40, 38f, 39f, 40f
 polar, 43, 44, 45f
 rotation, 36
 σ (σ), 32, 33, 36–40, 38f, 39f, 40f
 triple, 15
- Bond angles
 in alkyne, 400
 four groups around an atom, 24, 25–26, 26f, 40f
 molecular shape and, 24–27, 26f
 three groups around an atom, 24–25, 26f, 40f
 two groups around an atom, 24, 26f, 40f
- Bond cleavage
 heterolytic, 200–202, 201f
 homolytic, 200–202, 201f
 in nucleophilic substitution reactions, 242–243, 252
- Bond dissociation energy, 203–206
 in alkyne, 400–401
 for benzylic C–H bond, 670
 bond length and, 205
 bond strength and, 204–206
 carbon radical stability and, 539–540
 of common bonds, 204t, A–7–8
 definition of, 203
 enthalpy change calculation from, 205–206
 limitations of, 206
 periodic table trend in, 204–205
- Bonded (shared) electrons, 12–14, 12f
- Bonding molecular orbital, 627, 628f, 629–631, 630f, 631f
- Bond length
 average for common bonds, 23t
 bond dissociation energy and, 205
 bond strength and, 40–42, 41t
 in conjugated dienes, 581–582
 percent *s*-character and, 41–42, 582
 periodic trends in, 23
- Bond polarity, 43–44, 45f
- Bond strength
 bond dissociation energy and, 204–206
 bond length and, 40–42, 41t
 infrared (IR) absorption and, 478–479, 479f, 480
 percent *s*-character and, 41–42
- Bongkreic acid, 395
- 9-Borabicyclo[3.3.1]nonane (9-BBN), 387
- Borane, 731
 addition to alkene, 386–387, 387f, 388f
 hydroboration–oxidation of alkynes, 412–413
- Boron, octet rule exceptions in, 18
- Boron trifluoride, molecular shape of, 24–25, 26f
 hybrid orbitals in, 35
- BPA (bisphenol A), 683, 1163, 1164f, 1168
- Bradykinin, 1089
- Brevicommin, 823
- Bridged bicyclic ring system, 593, 593f
- Bridged halonium ion, 380–382, 383–384, 385t, 409
- Bridgehead carbons, A–6
- Bromination
 of alkanes, 546–547, 547f
 of alkenes, 558–560
 at allylic C–H bonds, 553–555
 with NBS, 553–555
 product mixtures in, 555
 selective, 553–554
 substitution *versus* addition, 554
 of benzene, 645, 668
 benzylic, 670–671
 chlorination compared, 546–547
 as endothermic reaction, 547, 547f
 polybromination, 665
 of substituted benzenes, 665
 as test for π (π) bond, 380
- Bromine, isotopes of, 469
- α -Bromoacetone, 893
- 3-Bromoacetophenone, 779f
- m*-Bromoacetophenone, 779f
- m*-Bromoaniline, 674
- o*-Bromoaniline, 953
- p*-Bromobenzaldehyde, synthesis of, 985–986
- Bromobenzene, 343, 645
- 4-Bromobutanal, 819
- 1-Bromobutane, 288, 874
- 2-Bromobutane, 280, 550–551
- 3-Bromo-1-butene, 555, 587, 588f
- 1-Bromo-2-butene, 555, 587, 588f
- cis*-1-Bromo-4-*tert*-butylcyclohexane, 309
- trans*-1-Bromo-4-*tert*-butylcyclohexane, 309
- o*-Bromochlorobenzene, 611

- 1-Bromo-1-chloroethylene, NMR spectrum of, 500
- Bromocyclohexane, 308
- Bromocyclohexane, 376
- 2-Bromocyclohexanone, 896
- 1-Bromocyclohexene, 1005
- 1-Bromo-3,3-dimethylbutane, 567
- 2-Bromo-2,3-dimethylbutane, 567
- (5*R*)-6-Bromo-2,5-dimethylnonane, 275
- (6*R*)-6-Bromo-2,6-dimethylnonane, 275
- 1-Bromohexane, 874
- 3-Bromohexane, 166
- enantiomers of, 168f
- 6-Bromo-2-hexanone, 823
- trans*-1-Bromo-1-hexene, 1004
- 1-Bromo-2-hexene, 274
- Bromohydrins, 383–384
- Bromomethane, in nucleophilic substitution reactions, 243–245, 248, 248f, 249f
- 2-Bromo-2-methylbutane, 289–290
- cis*-1-Bromo-4-methylcyclohexane, 263
- 1-Bromo-1-methylcyclopentane, 293
- 2-Bromo-2-methylpropane, 280, 288, 414
- p*-Bromonitrobenzene, 668–669
- Bromonium ion, 554
- 1-Bromopentane, 492
- 2-Bromopentane, in E2 elimination reaction, 979
- 1-Bromopropane
- NMR spectrum of, 514–515, 515f
- synthesis of, 558
- 2-Bromopropane, 310, 322
- mass spectrum of, 469, 469f
- NMR spectrum of, 513, 513f
- synthesis of, 547, 558
- α -Bromopropiophenone, 894
- (*Z*)-2-Bromostyrene, 1009, 1024
- N*-Bromosuccinimide (NBS), 383–384, 553–555
- p*-Bromotoluene, 611
- Brompheniramine, 975
- Bronchodilator, 167, 321, 347, 347f
- Brønsted–Lowry acid
- carboxylic acids, 699–703
- description of, 55–56
- reactions of, 56–58
- Brønsted–Lowry base
- alkyl halide and elimination reactions, 234
- description of, 55–56
- in elimination reactions with alkyl halides, 279
- reactions of, 56–58
- Brucine, 1115
- BTX mixture, 83, 614
- Buckminsterfullerene, 632
- Bufotenin, 958–959
- Bufo* toads, 958
- Bupropion, 646f, 686, 913
- 1,2-Butadiene, 424
- 1,3-Butadiene, 397
- conformations of, 1088
- conjugation in, 572–573, 573f
- electrostatic potential plot for, 573, 573f
- HBr addition to, 584–588, 588f
- polymerization of, 1156, 1160
- resonance structures, 576, 582
- sigma (σ) bond length in, 581–582
- stability of, 619
- ultraviolet absorption, 598
- Butanal, 944
- boiling point of, 91
- intermolecular forces in, 90
- melting point of, 93
- Butane
- alkene conversion to, 429
- anti conformation, 282f
- conformations of, 134–137, 134f, 136f
- constitutional isomers of, 118t
- eclipsed conformation, 282f
- halogenation of, 549–550
- molecular formula of, 118t
- as propellant, 551
- rotation about carbon–carbon single bonds of, 282, 282f
- solubility of, 94–95
- structure of, 115
- Butanedioic acid, 692
- 2,3-Butanediol, 345
- 2,3-Butanedione, addition to margarine of, 432
- Butanenitrile, synthesis from 1-propanol, 337–338
- Butanoic acid, 695
- IR spectrum of, 694f
- synthesis of, 901–902
- 1-Butanol
- boiling point of, 91, 108, 352
- intermolecular forces in, 90
- melting point of, 93
- naming, 315
- 2-Butanol
- (2*R*)-2-Butanol, 873
- (2*S*)-2-butanol, conversion to its tosylate, 338–339
- enantiomers of, 171–173, 729–730
- infrared (IR) spectrum, 482
- stereogenic center of, 168
- synthesis using Grignard reaction, 747
- tert*-Butanol, as polar protic solvent, 240f
- 2-Butanone, 422
- boiling point of, 779
- infrared (IR) spectrum, 482
- reduction of, 729–730
- structure of, 778
- synthesis of, 905
- 1-Butene, 306, 360t, 363f, 1021
- halogenation of, 555
- 2-Butene
- cis*-2-butene, 306, 360t, 365, 382, 382f
- carbene addition to, 1013
- epoxidation of, 440
- hydrogenation of, 429–430
- trans*-2-butene, 360t, 365, 382, 382f, 384
- carbene addition to, 1014
- epoxidation of, 440
- hydrogenation of, 429–430
- stereoisomers of, 282–284, 282f
- tert*-Butoxide, nucleophilicity of, 239
- tert*-Butoxycarbonyl protecting group (Boc), 1096–1098
- tert*-Butoxy group, 316
- Butter, 369, 432
- oxidation of lipids in, 215
- Butylamine, mass spectrum of, 955, 955f
- (*R*)-*sec*-Butylamine, 195
- Butylated hydroxy anisole (BHA), 568
- Butylated hydroxy toluene (BHT), 557
- Butylbenzene, 677
- tert*-Butylbenzene, 611
- Butylcyclohexane, 126f, 460
- tert*-Butylcyclohexane, 143f
- tert*-Butyl cyclohexanecarboxylate, 831
- cis*-4-*tert*-Butylcyclohexanol, 773
- 4-*tert*-Butylcyclohexanone, 773
- tert*-Butylcyclopentane, 125
- tert*-Butyldimethylsilyl ether (TBDMS), 749, 750
- Butyl group, 121
- sec*-butyl group, 121
- tert*-butyl group, 121, 143
- tert*-Butyl hydroperoxide, 438, 452
- tert*-Butyl iodide, 231
- Butyllithium, 71, 423, 794, 888
- Butylmagnesium chloride, 759
- 1-*sec*-Butyl-3-methylcyclohexane, 127f
- sec*-Butyl methyl ether, 316
- tert*-Butyl methyl ether (MTBE), 95
- formation of, 378–379
- as fuel additive, 378, 497
- NMR spectrum of, 497, 499
- 5-*tert*-Butyl-3-methylnonane, 124
- tert*-Butyl pentyl ether, 492
- 1-Butyne, 424
- addition reactions of, 405, 406f
- hydrohalogenation of, 406f, 407
- 2-Butyne, 424
- hydration of, 410f, 411
- infrared (IR) spectrum, 480
- Butyric acid, structure of, 691t
- γ -Butyrolactone, 719, 872
- Cadaverine, 956–957
- Caffeine, 109, 529, 949
- Cahn–Ingold–Prelog system, 170
- Calcitriol, 581, 581f
- Calorie (cal), 40
- Camphene, 1147
- Camphor, 1134
- Cancer. *See also* Anticancer drugs
- nitrosamines and, 261
- polycyclic aromatic hydrocarbons and, 349
- Capnellene, 416, 416f
- Caproic acid, 688, 691t
- ϵ -Caprolactam, 1161, 1170
- ϵ -Caprolactone, 1177
- Capsaicin, 4, 49, 485, 529, 607, 914
- Caraway, odor of, 188
- Carbamate, 870, 1095–1096, 1162
- Carbanions
- addition to aldehydes and ketones, 789
- electrostatic potential plots, 68, 68f
- formation in alkyne reduction, 436
- nucleophilic addition with, 724, 789
- nucleophilic substitution with, 725
- vinyl, 436
- Carbenes
- intermediates in cyclopropane synthesis, 1012–1014
- metal–carbene intermediates in metathesis, 1017
- preparation of dihalocarbenes, 1012–1013
- structure of, 1012
- Carbenoid, 1015
- Carbinolamines, 788, 797, 800, 975
- Carbocation(s), 20
- achiral, 376
- allylic
- as conjugated system, 572, 573–574
- electrostatic potential plot, 574, 574f
- NMR spectra of, 575, 575f
- resonance of, 574–576
- classification, 256
- in electrophilic addition reactions, 372–376
- in electrophilic aromatic substitution reactions, 643–644, 659–660
- electrostatic potential maps, 257, 257f
- energy diagram for formation of, 260f
- formation in Lewis acid–base reactions, 74
- Hammond postulate, 374–375, 374f
- as Lewis acid, 372
- localized *versus* delocalized, 574f
- in nucleophilic substitution reactions, 243
- planar, 254, 263, 377, 1043–1044
- primary (1°), 256
- as reactive intermediate, 330
- resonance-stabilized, 408, 410, 411, 471, 574–575, 574f, 575f, 584–585, 587, 643
- secondary (2°), 256
- in S_N1 substitution reactions, 252–260, 260f
- tertiary (3°), 256, 649, 651–652
- trigonal planar, 253–254, 376
- vinyl, 267, 407–408, 411
- Carbocation intermediates, 558
- Carbocation rearrangements
- conversions of alcohols to alkyl halides, 334–335
- in dehydration of alcohols, 328–330
- Friedel–Crafts alkylation, 651–652
- in hydration reactions, 378
- 1,2-shifts, 328–330
- 1,2-hydride shift, 651
- 1,2-methyl shift, 328–329
- in S_N1 reactions, 334–335
- Carbocation stability
- of allylic carbocations, 573–574
- conjugation and, 573
- in electrophilic aromatic substitution reaction, 659–660
- Hammond postulate and, 258–261
- hyperconjugation and, 257–258
- inductive effects and, 256–257
- 1,2-shifts and, 328
- Carbohydrates, 812–813, 1027–1073
- amino sugars, 1061–1062
- cellulose, 160–162, 161f, 162f, 858, 858f, 1029f, 1059–1060
- disaccharides, 1056–1058, 1058f, 1059f
- energy in, 1028
- examples of, 1028, 1029f
- glycosides, 1042–1046
- N*-glycosides, 1062–1065, 1065f
- molecular formula of, 1028
- monosaccharides. *See* Monosaccharides
- natural fibers, 858, 858f
- nomenclature, 1028, 1029, 1034
- polysaccharides, 1059–1061
- starch, 160–161, 161f, 162f, 319, 320f, 1060–1061
- Carbon
- alpha (α), 279–280, 691, 694, 710
- of aldehydes and ketones, 777, 785–786
- substitution reactions of carbonyl compounds at the α carbon, 880–906
- beta (β), 279–280, 296, 691
- bonding of, 11, 12f, 15, 17t, 32–33
- charged atom, 31
- chiral. *See* Stereogenic centers
- delta (δ), 691
- electron-deficient, 65, 82, 84, 85, 103, 202, 234, 234f
- electronegativity value, 739
- electrophilic, 234, 234f
- elemental forms of, 632
- excited state, 32
- formal charge of, 17t
- gamma (γ), 691
- ground state, 32
- isotopes of, 7, 8f, 472t
- mass of, 472t
- periodic table entry, 8
- primary (1°), 116
- quaternary (4°), 116
- radicals, 539–540, 540f
- secondary (2°), 116
- tertiary (3°), 116
- tetravalence of, 4, 25, 27
- valence electrons of, 32
- Carbonate, 1163
- Carbon backbone (skeleton), 82
- Carbon–carbon bonds
- bond length and bond strength, 40–41, 41t

- double. *See also* Double bonds
in alkenes, 281–283. *See also*
Alkenes
properties of, 360t
radical reactions, 541, 558–560
restricted rotation around,
282–283, 282f
- NMR spectra of, 505, 506f, 506t
nonpolar nature of, 43
reactions forming, 1002–1026, A–9
coupling reactions of
organocuprates, 1003–1005
cyclopropane synthesis,
1012–1014
Heck reaction, 1003, 1009–1011
metathesis, 1015–1019, 1016f,
1018f
Simmons–Smith reaction,
1014–1015
Suzuki reaction, 1002, 1003,
1005–1009, 1008f
rotation around, 129, 131
- Carbon dioxide
atmospheric concentration of, 149,
149f
electrostatic potential plot for, 45f
global warming, role in, 149
as nonpolar molecule, 44
reaction of Grignard reagents with,
753–754
release in combustion reactions, 148
- Carbon–halogen bond, polar, 234, 234f
- Carbon–hydrogen bonds
acidity of, 63, 64
benzylic, 669–670
bond length and bond strength, 41–42
halogenation of, 552–555
nonpolar nature of, 43
radical reaction with, 540–541
- Carbon monoxide, 1110
- Carbon NMR, 495, 522–526
of aldehydes, 782, 782f
of amines, 956
basis of, 522–523
of benzene derivatives, 613–614, 613f,
613t
of carbocations, 575, 575f
of carboxylic acid derivatives, 836
of carboxylic acids, 694, 695f
characteristic absorptions, A–12
chemical shifts in, 523, 524, 525t,
575, 575f
examples of spectra, 526f
of ketones, 782
of nitriles, 836
number of signals in, 523–524
position of signals in, 524–526
reference signal, 523
- Carbon–oxygen bonds, polar nature of,
43
- Carbon skeleton (backbone), 82
- Carbon tetrachloride, 232
as solvent, 94
- carbonyl chloride (suffix), 830, 833t
- Carbonyl compounds. *See also specific
classes*
classes of, 722
condensation reactions, 916–939
conversion of ozonides to, 445–446
electron deficiency of, 723
electrostatic potential map of, 723,
723f
formation by oxidation of alcohols,
447
 β -hydroxy
synthesis of, in aldol reactions,
917–928
useful transformations of, 924, 924f
leaving groups of, 722, 724, 725–726
 pK_a , 884–885, 885t, 886t
- reactions at α carbon, 880–906
acetoacetic ester synthesis, 900,
903–905
enolate alkylation, 897–899
halogenation, 892–895, 895f
malonic ester synthesis, 900–903
racemization, 891, 891f
reactions of, 723–726
nucleophilic addition, 724
nucleophilic substitution, 724,
725–726
with organometallic reagents,
712–745, 748–750, 749f
oxidation, 726–727, 738–739
reduction, 726–738
reduction of, 726–738
biological, 732–733
carboxylic acids and derivatives,
733–738
by catalytic hydrogenation, 729
enantioselective, 731–733, 732f
with metal hydride reagents,
728–729, 730f, 738t
stereochemistry of, 729–731
synthesis by hydroboration–oxidation
of alkynes, 412–413
tautomers, 863–864
 α,β -unsaturated compounds, 755–758
unsymmetrical, enolates of, 889
- Carbonyl condensation reactions,
916–939
aldol reactions, 917–928. *See also*
Aldol reactions
Claisen reaction, 928–932, 929f
Dieckmann reaction, 932–933
Michael reaction, 934–936, 935f
Robinson annulation, 936–939
- Carbonyl groups, 409, 722
conjugated, 780–781
as electron-withdrawing group, 591
features of, 828
formation by oxidative cleavage of
alkenes, 444–446
as functional group, 85, 86t
hydrogenation of, 432
IR absorption of, 693, 694f, 780–781,
781f
of monosaccharides, 1028
nucleophilic addition to, 786–788,
788f
reactivity of, 85
stabilization by alkyl groups, 802
tautomers of, 881–883
- Carboxy, derivation of term, 689
- Carboxy group, 86t, 689
of benzene derivatives, 671–672
- Carboxylate anion
of amino acids, 712
formation of, 700
metal salt of, 692, 693f
synthesis of
hydrolysis of amides, 855–856
hydrolysis of esters, 850, 851–852
hydrolysis of nitriles, 863–864
- Carboxylation, 753–754
- Carboxylic acid(s), 688–709
acidity of, 689, 699, 700–705, 703f,
829t, 845
aliphatic, inductive effects in, 703–704
aspirin as, 696–697
basicity of, 699
benzene derivatives, 671–672
boiling point of, 693t
deprotonation of, 700, 700t
diacids, 692, 900
as dimers, 692, 693f
dipole–dipole interactions of, 692
electrostatic potential plot, 689, 689f
functional group in, 86t
hydrogen bonding, 692, 693f
- interesting examples, 694–696
intermolecular forces, 692, 693f, 693t
IR spectra of, 693–694, 694f
leaving group, 725
melting point of, 693t
NMR spectra of, 694, 695f
nomenclature of, 690–692
common names, 691–692, 691t,
692f
IUPAC system, 690–692
nucleophilic acyl substitution reactions,
845–850, 846f, 858t
conversion to acid chlorides,
845–847, 846f
conversion to amides, 846f,
849–850
conversion to cyclic anhydrides,
846f, 847
conversion to esters, 846f, 847–849
nucleophilic substitution reactions, 725
oxidation of aldehydes, 727, 738–739
physical properties, 692–693, 834, 834t
 pK_a of, 700–704, 829t
protonation of, 699
reactions of
ester synthesis, 846f, 847–849
general features, 699
nucleophilic acyl substitution
reactions, 845–850, 846f, 858t
reduction, 726, 735
resonance stabilization of, 701–702,
702f, 703f
salts of, 696
separation from other organic
compounds, 707–709, 709f
solubility of, 693t
spectroscopic properties, 693–694,
694f
stability of, 829
structure and bonding of, 689, 722,
826, 828–829
substituted benzoic acids, 705–707,
706f
synthesis of, 697–699
conversion of acid chlorides, 842,
843
conversion of anhydrides, 844
hydrolysis of amides, 855–856
hydrolysis of esters, 850–852
hydrolysis of nitriles, 863–864
malonic ester synthesis, 900–903
oxidation of aldehydes, 738
oxidation of alkyl benzenes, 698
oxidation of alkynes, 698
oxidation of primary (1°) alcohols,
447, 448–450, 698
oxidative cleavage of alkynes,
446–447
reaction of organometallic reagents
with carbon dioxide, 753–754
-carboxylic acid (suffix), 860
- Carboxylic acid derivatives. *See also* Acid
chlorides; Amides; Anhydrides;
Esters
acidity/basicity of, 828–829, 829t
boiling point of, 834, 834t
IR spectra of, 835, 835t
leaving groups of, 827, 828, 829t,
838–840
melting point of, 834, 834t
NMR spectra of, 836
nomenclature of, 830–834, 833t
nucleophilic acyl substitution reactions
of, 725, 838–839
mechanism of, 838–839
relative reactivity, 839–841, 859
summary of, 857–858, 858t
physical properties of, 834, 834t
reaction of organometallic reagents
with, 750–753
- reactivity with nucleophiles, 725
reduction of, 727
resonance structures of, 828
solubility of, 834, 834t
stability of, 829
structure and bonding of, 722, 826–829
- Carboxypeptidase, 1093, 1093t
- Carcinogen, 349
- Cardinol, 1169
- Carnauba wax, 1121
- Carotatoxin, 109
- β -Carotene, 97–98, 367f, 795, 795f
addition to margarine, 432
- Carson, Rachel, 233
- Carvacrol, 640
- Carvone, 188, 640, 1145
- Catalysts. *See also specific catalysts*
acids as, 218–219
biocatalyst, 1167, 1168f
chiral, 1085–1086
definition of, 218
energy of activation lowered by, 219,
219f
enzymes, 219–220, 220f
homogeneous, 1159
hydrogenation, 429
Lewis acid, 333
Lindlar, 435
metal, 218–219, 429
palladium, 219, 729. *See also*
Palladium-catalyzed reactions
Ziegler–Natta, 1157–1159, 1160
- Catalytic amount, 219
- Catalytic hydrogenation, 429–431
of aldehydes and ketones, 729
- Catecholborane, 1007, 1009
- Cation(s)
(carbocation. *See* Carbocation(s))
counterions, 55
cyclopentadienyl, 625
definition of, 7
radical, 465
salt formation, 10–11
solvation by ion–dipole interactions,
240, 241, 264–265
spectator ions, 55
propylion, 625–626, 630, 630f
- Cationic polymerization, 1153–1154,
1155f
- CBS reagent, 731–732, 732f
- Cedrol, 1133f
- Celecoxib (Celebrex), 1132
- Cell membrane
cholesterol in, 150, 150f, 1140
functions, 100
phospholipids in, 1126–1128
structure of, 100, 101f
transport across, 100–102, 102f
- Cellobiose, 1058
- Cells, 100
- Cellulose, 858, 858f
hydrolysis of, 160, 161f, 1060
solubility of, 162
structure of, 160–161, 161f, 162f,
1029f, 1059–1060
- Cellulose acetate, 1060
- Cembrene, 601
- Cephalexin, 838
- Cephalin, 1126
- Cephalosporins, 838
- Cetylpyridinium chloride (CPC), 236
- CFCs. *See* Chlorofluorocarbons (CFCs)
- Chain branching, in polymerization,
1152–1153
- Chain-growth polymerization,
1150–1157
anionic, 1154–1155, 1155f,
1156–1157
cationic, 1153–1154, 1155f
chain branching in, 1152–1153

- Chain-growth polymerization (*continued*)
 copolymers, 1156
 of epoxides, 1156–1157
 ionic, 1153–1156, 1155f
 radical, 1151–1152, 1151f, 1153–1156, 1155f
- Chain-growth polymers, 1149, 1150–1157
- Chain mechanism, 544, 551
- Chain termination, 1152
- Chair conformation
 of cyclohexanes, 138–147, 139f
 of glucose, 1041
- Chantix. *See* Varenicline
- Charge
 Brønsted–Lowry acids and bases, 55f, 56
 on carbon atom, 31
 dipole, 43, 44, 45f
 formal, 15–16, 17t
 in resonance hybrid, 22
- Charged ion, 7
- Chauvin, Yves, 1015
- Chemical shift
 in ^{13}C NMR, 523–524, 525t, 575, 575f
 in ^1H NMR
 equation for, 498
 predicting values of, 503–504
 protons on benzene rings, 505, 506t
 protons on carbon–carbon double bonds, 505, 506t
 protons on carbon–carbon triple bonds, 505, 506t
 scale, 497
 values for common bonds, 504t
- Chiral catalysts, 1085–1086
- Chiral drugs, 187
- Chirality
 basic principles of, 165f
 of naturally occurring objects, 166
 in substituted cycloalkanes, 180
 test for, 164
- Chirality center, 165. *See also* Stereogenic centers
- Chiral molecules, 163–166, 165f
 halogenation of, 550–551
 optical activity of, 183
- Chiral reagent, 1084, 1085
- Chiral reducing agent, 731
- Chitin, 1061–1062
- Chloral, 802
- Chloral hydrate, 802, 803
- Chloramphenicol, 167
- Chlorination
 of alkanes, 544–548, 544f, 545f, 548f, 549–551
 of benzene, 645
 bromination compared to, 546–548, 548f
 of ethane, 543–544, 544f, 545f
 as exothermic reaction, 544, 544f, 547–548, 548f
 mechanism of, 543–544
- Chlorine
 isotopes of, 468
 mass spectra of alkyl chlorides, 468, 469f
- 2-Chloroacetamide, 879
- 2-Chloroacetic acid, 704
- Chlorobenzene, 611, 645
- 2-Chloro-1,3-butadiene, polymerization of, 1160
- 1-Chlorobutane, synthesis of, 549–550
- 2-Chlorobutane
 NMR spectrum of, 501
 synthesis of, 549–550
- Chlorocyclohexane, chair conformation of, 296–297
- Chlorocyclopentane, 96
- Chlorocyclopropane, NMR spectrum of, 500
- Chloroethane, 51, 84, 231
 NMR spectrum of, 524
- Chloroethylene, NMR spectrum of, 500
- 4-Chloro-1-ethyl-2-propylbenzene, 612
- Chlorofluorocarbons (CFCs), 3, 233, 551–552, 552f
- Chloroform, 232
 dichlorocarbene preparation from, 1012–1013
- Chlorohydrin, estrone synthesis from, 385f
- Chloromethane, 233f. *See also* Methyl chloride
- 1-Chloro-1-methylcyclohexane, 375
- 1-Chloro-2-methylcyclohexane
cis-1-chloro-2-methylcyclohexane, 296
trans-1-chloro-2-methylcyclohexane, 296
- 1-Chloro-3-methylcyclohexane, 375
- 1-Chloro-1-methylcyclopropane, 307
- 2-Chloro-5-methylheptane, 230
- 3-Chloro-3-methylhexane, 376
- (3*R*)-1-Chloro-3-methylpentane, 310
- 1-Chloro-2-methylpropane, 231f
- meta*-Chloroperoxybenzoic acid, 438f, 439–441
- 2-Chloropropanal, 777f
- 1-Chloropropane, 374, 673
- 2-Chloropropane, 374
 mass spectrum of, 468, 469f
- 2-Chloropropene, 407
- 3-Chloropropenoic acid, NMR spectra of, 516, 516f
- α -Chloropropionaldehyde, 777f
- Chloroquine, 974
- p*-Chlorostyrene, 675–676
- Chlorpheniramine, 646f
- Cholesterol, 596, 861–862, 861f, 1119, 1139–1141
 biosynthesis of, 1140, 1140f
 in cell membranes, 150, 150f, 1140
 cholesterol-lowering drugs, 1140, 1141f
 trans fats and, 433
 solubility of, 95, 150
 specific rotation of, 185
 structure of, 150, 150f, 170, 1139
 synthesis stimulation by saturated fats, 369
- Cholesteryl esters, 861, 861f
- Cholic acid, 158
- Choline, 861
- Chromate ester, 448
- Chromium
 in oxidizing agents, 438–439, 448–450, 738–739
- Chromium(VI) oxide, 438–439
- Churchane, 128f
- Churchill, Winston, 233
- Chymotrypsin, 1093, 1093t
- Ciguatoxin (CTX3C), 735–736, 736f
- Cimetidine, 957, 958
- Cinnamaldehyde, 783f, 922
- Cis-fused, 592
- Cis geometry
 alkenes, 359, 360t, 363
 cycloalkenes, 359
- Cis isomers
 of 2-butene, 282–283, 282f
 of cycloalkanes, 144–147
 definition of, 144, 282, 377
 of disubstituted cycloalkanes, 180
- Cis protons, 516, 517f
- Citral, 1133f
- Citric acid, as Brønsted–Lowry acid, 50, 55f
- Citronellal, 783f
- Citronellol, 493
- Claisen reactions, 928–929, 929f
 crossed, 930–932
 intramolecular, 932–933
- Claritin. *See* Loratadine
- Clemmensen reduction, 672
- Clomiphene, 394
- Clopidogrel, 175, 912
- Clostridium botulinum*, 261
- ^{13}C NMR. *See* Carbon NMR
- Cocaine, 78, 110, 836, 957
- Cocoa butter, 1125
- Coconut oil, 369, 1123, 1124
- Codeine, 716, 959
- Coenzyme, 451
 NADH, 733
- Coffee, decaffeinated, 233f
- Collagen, 81, 1075
 structure of, 1108–1109, 1109f
- Column, in periodic table, 8
- Combustion
 of acetylene, 402
 of alkanes, 148–149
 definition of, 148
 of triacylglycerols, 1125
- Common names
 of alcohols, 315–316
 of aldehydes, 777, 777f
 of alkenes, 127–128, 128f
 of alkenes, 365, 365f
 of alkyl halides, 231
 of amines, 952–953
 of benzene derivatives, 610–612
 of carbohydrates, 1029
 of carboxylic acid derivatives, 830
 of carboxylic acids, 691–692, 691t, 692f
 of epoxides, 317
 of ethers, 316
 of ketones, 778, 779f
 of nitriles, 832, 832f
- Compounds
 examples of, 10
 formation of, 10
 ionic, 11
- Concentration, reaction rate and, 216, 217–218
- Concerted reaction, 200, 285
- Condensation polymer. *See also* Step-growth polymers
 nylon, 859
 polyesters, 859–860
- Condensation reactions
 carbonyl, 916–939
 definition of, 919
- Condensed structures, 27–29, 28f
- Configuration
 inversion of
 in alkyl tosylate reactions, 339–340
 retention of
 in conversion of alcohols to alkyl tosylates, 338–339, 340
 in oxidation of alkylboranes, 388–389
 in $\text{S}_{\text{N}}2$ reactions, 246–247, 247f, 263
 of stereoisomers, 162
- Conformations
 of acyclic alkanes, 129–137
 butane, 134–137, 134f, 136f
 ethane, 129–134, 132f, 133f
 anti, 134f, 135–137, 136f
 of cycloalkanes, 137–147
 disubstituted, 143–147
 monosubstituted, 141–143
 of cyclohexane(s), 138–147
 boat, 141, 141f
 chair, 138–147, 139f
 disubstituted cyclohexane, 144–147, 146f
 monosubstituted cyclohexane, 141–143, 143f
 ring-flipping, 140–141, 141f
 definition of, 131
 eclipsed, 131–137, 132f, 133f, 134f, 136f, 136t, 295f
 free energy change and, 208–209
 gauche, 134f, 135, 136f
 staggered, 131–137, 132f, 133f, 134f, 136f, 295, 295f
- Congo red, 989, 990
- Coniine, 957, 957f
- Conjugate acid, 56–58, 236–237, 237t
- Conjugate addition, 584, 934
- Conjugate base, 56–58
- Conjugated dienes, 580–599
 addition reactions
 Diels–Alder reaction, 588–597, 589f
 electrophilic, 584–586
 kinetic *versus* thermodynamic product, 586–588
 carbon–carbon bond length in, 581–582
 conformation of
s-cis, 580
s-trans, 580
 electrostatic potential plot for, 573, 573f
 hydrogenation of, 583–584
 interesting examples of, 581
 numbering of carbons in, 584
 stability of, 583–584, 583f
 stereoisomers of, 580
 synthesis in Suzuki reaction, 1006–1009
 ultraviolet (UV) light absorption, 597–599, 598f
- Conjugated double bond, 920
- Conjugated proteins, 1109
- Conjugated systems
 allylic carbocation, 572, 573–574
 1,3-dienes, 572–573, 573f
- Conjugation, 571–606
 definition of, 572
 derivation of word, 572
 β -dicarbonyl compounds, 882
 effect on carbonyl group IR absorption, 780–781
 Hückel's rule and, 618
- Constitutional isomers
 of alkanes, 115–117, 118t
 of alkenes, 325
 definition of, 17
 of dipeptides, 1087
 in elimination reactions, 280
 features of, 162
 ketone and aldehyde, 413
 of monosaccharides, 1029
 stereoisomers compared, 163f
 tautomers, 410
- Copolymers, 1156
- Copper
 electronegativity value of, 739
 organocuprate reagents, 739–740, 1003–005
- Copper(I) cyanide, 983
- Core electrons, 9
- Corey, E. J., 5, 348
- Corn
- ethanol production from, 319, 320f
 use in green polymer synthesis, 1167–1168, 1168f
- Coronary artery disease, 150, 175, 369
- Cortisol, 314, 1142
- Cortisone, 783, 1142
 Diels–Alder reactions, 596–597
- Cotton
 binding dyes to, 990
 cellulose in, 1059
 structure of, 858, 858f

- Coumarin, 946
 Counterions, 55
 Coupling constant, 510, 516
 Coupling reactions
 alkenes with organic halides (Heck reaction), 1003, 1009–1011
 of aryl diazonium salts, 982, 986–988
 of organocuprate reagents, 1003–1005
 Covalent bonding
 description of, 10, 11–12
 in Lewis structures, 12–15
 Covalent molecules
 boiling points of, 91
 intermolecular forces, 87–90, 88f, 90t
 solubility of, 96f
 COX-1, 1131–1132, 1131f
 COX-2, 1131–1132, 1131f
 COX-3, 1132
 CPC (cetylpyridinium chloride), 236
 Cracking, 366
 Crafts, James, 643
 CRCs (chlorofluorocarbons), 233, 551–552, 552f
p-Cresol, 686, 1176
 Crossed aldol reactions, 921–924
 synthetically useful reactions, 922–924, 923f
 with two different aldehydes, both having α H atoms, 922
 useful transformations of aldol products, 924, 924f
 Crossed Claisen reactions, 930–932
 Cross formula, 1030
 Crown ethers, 101–102, 320–321, 321f
 Crude oil, 128
 Crutzen, Paul, 551
 Crystallites, 1165
 CTX3C (ciguatoxin), 735–736, 736f
 Cubane, 128f, 138
 Cucumber aldehyde, 779
 Curcumin, 639
 Curved arrow notation, 20–21, 56, 201, 235
 Cyanide
 addition to aldehydes and ketones, 790–791
 addition to imines, 1080
 α -Cyano carbonyl compounds, 923–924
 2-Cyanocyclohexanone, structure of, 832f
 Cyano group, 791, 832
 of nitriles, 827
 Cyanohydrins
 in Kiliani–Fischer synthesis, 1049, 1051
 naturally occurring derivatives, 791–792
 synthesis by cyanide addition to aldehydes and ketones, 790–791
 in Wohl degradation, 1049, 1050
 Cyclic anhydrides, 826–827, 847
 Cyclic compounds
 entropy decrease with formation of, 209
 stereogenic centers in, 168–170
 Cyclic esters, 837
 Cyclic ethers, 317
 Cyclization, of hydroxy aldehyde, 810, 813
 Cyclo- (prefix), 118, 125
 Cycloalkanes
 conformations of, 137–147
 disubstituted, 143–147
 monosubstituted, 141–143
 disubstituted, 180–181
 molecular formula, 114, 118
 naming, 125–127, 126f, 127f
 NMR spectra, 499–500
 stereoisomers, 144–147
 structure of, 114, 118
 substituted, 141–147, 143f, 146f
 Cycloalkenes
 cis and trans geometry of, 359
 nomenclature, 363, 363f
 Cycloalkynes, 401
 Cyclobutadiene, 618–619, 626, 631
 Cyclobutane
 conformation, 137
 structure of, 118, 138f
 1-Cyclobutylhexane, 126f
 Cyclodecane, structure of, 138f
 α -Cyclogeraniol, 398
 Cycloheptane, structure of, 138f
 1,3,5-Cycloheptatriene, 618, 625
 1,3-Cyclohexadiene, 618
 heat of hydrogenation, 616
 stability of, 619
 ultraviolet absorption, 598
 Cyclohexanamine. *See* Cyclohexylamine
 Cyclohexane(s)
 conformations of, 138–147
 boat, 141, 141f
 chair, 138–147, 139f, 296–297
 disubstituted cyclohexane, 144–147, 146f
 free energy change and, 208
 monosubstituted cyclohexane, 141–143, 143f
 ring-flipping, 140–141, 141f
 conversion to cyclohexene, 549
 E2 elimination in, 297
 in mango, 114
 NMR spectrum, 518
 skeletal structures, 29
 structure of, 114, 118, 125
 Cyclohexanecarbonyl chloride, structure of, 830
cis-1,2-Cyclohexanediol, 443
 Cyclohexanediols, 443, 1175
 1,3-Cyclohexanedione, 717
 Cyclohexanol, 819
 cyclohexene formation from, 392
 dehydration of, 330–331, 392
 1,2-dibromocyclohexane synthesis from, 391–392
 separation from benzoic acid by extraction procedure, 707–709, 709f
 separation of cyclohexylamine from, 967–968, 967f
 synthesis from cyclohexene, 378–379
 Cyclohexanone, 566, 760, 821
 aldol reactions of, 924, 924f
 alkylation of, 898
 conversion to alkene, 796–797
 enolate formation from, 884
 halogenation of, 896
 nucleophilic addition reactions, 788, 788f
 tertiary alcohol formed from, 743
 Cyclohexene
 addition reactions of, 371f, 376, 378–379, 380
 bromination of, 554
 1,2-dibromocyclohexane synthesis from, 392
 heat of hydrogenation, 616
 as Lewis base, 74
 stereoisomers, 592
 synthesis
 cyclohexane conversion, 549
 from cyclohexanol, 392
 by dehydration of cyclohexanol, 330–331
 as synthetic intermediate, 392
 2-Cyclohexen-1-ol, 356
 2-Cyclohexenone, 493, 821, 928
 formation in Robinson annulation, 936–939
 reduction of, 729
 synthesis of, 896
 Cyclohexylamine, 952
 ammonium salt of, 967–968, 967f
 in Hofmann elimination reaction, 978
 separation from cyclohexanol, 967–968, 967f
 Cyclononatriene, 637
 Cyclooctane, 187
 Cyclooctatetraene, 618, 637
 NMR spectrum of, 619–620
 Cyclooctene, cis and trans isomers of, 359
 Cyclooctyne, 401
 Cycloooxygenase, 697, 1130, 1131–1132, 1131f
 Cyclopentadiene, 593, 595–596, 625, 636
 Cyclopentadienyl anion, 624–625
 Cyclopentadienyl cation, 625
 Cyclopentadienyl radical, 625
 Cyclopentane, structure of, 118, 138f
 Cyclopentanecarboxylic acid, 902
 1,2-Cyclopentanediol
 cis isomer of, 443
 trans isomer of, 316
 Cyclopentanol, 96
 Cyclopentanone, 422, 819
 Cyclopentene, chlorination of, 381
 Cyclopropane(s)
 conformation, 137
 dihalo to dialkyl conversion, 1014
 NMR spectrum of, 500
 structure of, 118
 synthesis of
 in Simmons–Smith reaction, 1014–1015
 using carbene intermediates, 1012–1014
 Cyclopropenone, 821
 Cysteine, 190
 abbreviation for, 711t
 isoelectric point for, 1078t
 in peptide formation, 1086–1087
 pK_a values for, 1078t
 structure of, 1076f
 Cytidine, 1063
 Cytidine monophosphate, 1064
 Cytosine, 1064, 1065f
 2,4-D (2,4-dichlorophenoxyacetic acid), 646f
 Darvon (propoxyphene), 106, 166–167
 DBN (1,5-diazabicyclo[4.3.0]non-5-ene), 286, 286f, 300
 DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), 286, 300
 DCC. *See* Dicyclohexylcarbodiimide
d (dextrorotatory) compounds, 183
 DDE (dichlorodiphenyldichloroethylene), 278, 280
 DDT (dichlorodiphenyltrichloroethane), 4, 109, 215, 233, 683
 biological effects of, 278
 DDE formed from, 278
 degradation of, 280
 Dean–Stark trap, 804, 805f
 Decalin, 158, 460, 1138, 1139f
 Decamethrin, 1012
 Decane, 113
 constitutional isomers of, 118t
 molecular formula of, 118t
 Decarboxylation, 1079
 of β -diacids, 900
 of β -keto acids, 901
 in malonic ester synthesis, 900–902
 DEET, 842–843
 Dehydrating agent, 849–850
 Dehydration reactions
 of alcohols to alkenes, 324–331, 369–370
 alkene synthesis in, 324–331
 carbocation rearrangements in, 328–330
 E1 mechanism in, 326
 E2 mechanism in, 327, 330–331
 enthalpy change in, 327, 327f
 Le Châtelier's principle in, 327–328
 with phosphorus oxychloride and pyridine, 330–331
 in strong acid, 325–330
 of aldol product, 919–920
 dicarboxylic acid conversion to cyclic anhydrides, 847
 as β -elimination reaction, 325
 fragmentation pattern in mass spectrum, 471
 Dehydrohalogenation
 alkene synthesis by, 279
 of alkyl halides with strong base, 369, 370
 alkyne synthesis by, 299–300, 300f
 bases commonly used in, 280, 280t
 drawing products of, 280
 E1 mechanism of, 291
 E2 mechanism of, 285, 290
 16,17-Dehydroprogesterone, 1146
 Delocalization, resonance, 67
 Delta (δ) carbon, 691
 Delta (δ) scale, 497
 Dementia, 85, 926
 Demerol. *See* Meperidine
Dendrobates histrionicus (poison dart frog), 404, 404f
 Deoxy (prefix), 1063
 2-Deoxyadenosine, 1063
 Deoxyadenosine monophosphate, 104f, 105, 1029f, 1064
 2-Deoxy-D-ribose, 1063
 Deoxyribonucleic acid (DNA), 104f, 105, 1064–1065, 1065f
 Deoxyribonucleosides, 1063–1064
 Deoxyribonucleotides, 1064
 Deprotection, 748
 Deprotonation
 in acetoacetic ester synthesis, 904
 of alkyne, 405–406, 406f
 of carboxylic acids, 700, 700t
 of diethyl acetamidomalonate, 1079
 in direct enolate alkylation, 897
 in glycoside formation, 1043
 in halogenation at α carbon of carbonyl compounds, 893–895
 in malonic ester synthesis, 901
 racemization at α carbon of carbonyl compounds, 891
 Deshielding effects, in NMR, 502–503, 502f, 503f, 506f, 506t
 DET (diethyl tartrate), 452–453
 Detergents, 100
 Deuterium, 7
 Deuteriochloroform, 496f
 Dextropimaric acid, 1145
 Dextrorotatory, 1033
 Dextrorotatory (*d*) compounds, 183
 Di- (prefix), 123
 Diabetes, 783
 Diacids, 692, 900
 formation from monosaccharide oxidation, 1049
 in nylon synthesis, 859, 1160
 Diacon. *See* Methoprene
 Dialkylborane, 387, 387f
 Diamine, in nylon synthesis, 859, 1160
 1,4-Diaminobenzene, 1161
 1,6-Diaminohexane, 1160, 1167, 1167f
 Diamond, 632
 Diastereomers, 175–177, 177f
 alkenes, 360t, 363–364
 of 2-butene, 282–284, 282f
 definition of, 177
 disubstituted cycloalkanes, 180–181
 epimers, 1035

- Diastereomers (*continued*)
of glucose, 1038, 1038f
halogenation reactions and, 550–551
of monosaccharides, 1035
NMR spectrum of, 500–502
nucleophilic substitution reactions
and, 263
physical properties of, 185–186, 186f
separation of amino acid enantiomers
into, 1081–1083, 1082f
- 1,3-Diaxial interactions, 143
Diazepam, structure, 615f
Diazomethane, 1014
Diazonium ions, 261
Diazonium salts
alkyl, 981
aryl, 980–988
coupling reactions of, 982, 986–988
explosiveness of, 981
formation from primary amines,
980–981
substitution reactions of, 982–986
synthesis of aryl chlorides and
bromides, 983
synthesis of aryl fluorides, 983
synthesis of aryl iodides, 983
synthesis of benzene, 983–984
synthesis of benzonitriles, 983
synthesis of phenols, 982
uses in synthesis, 984–986
Diazoization, 980–981, 984
DIBAL-H (diisobutylaluminum hydride),
734, 735, 736f, 738t, 865–866
Dibromobenzene
meta-, 610, 613f
ortho-, 610, 613f
para-, 610, 613f
1,4-Dibromobutane, 902
2,2-Dibromobutane, 407–408, 424
2,3-Dibromobutane, 166, 177–179, 179f,
309
Dibromocarbene, 1014
1,2-Dibromocyclohexane, synthesis from
cyclohexanol, 391–392
1,3-Dibromocyclopentane, 180–181
1,2-Dibromoethane, 395
4,5-Dibromooctane, 396
2,3-Dibromopentane, stereoisomers of,
176–177, 177f, 179
Di-*tert*-butyl dicarbonate, 1096
Dibutyl phthalate, 1165–1166, 1174
Dicarbonyl compounds, 445
aldol reactions with, 926–928
1,4-dicarbonyl compounds, 926
1,5-dicarbonyl compounds, 927, 928f
in Robinson annulation, 936–937
synthesis of, 935, 936–937
 β -dicarbonyl compounds, 923, 923f
 pK_a values, 886, 886f
synthesis in crossed Claisen
reactions, 931
tautomers, 882
2,5-Dichloroaniline, 612
4,4'-Dichlorobiphenyl, 95
Dichlorocarbene, 1012
1,2-Dichlorocyclopentane, 381
Dichlorodifluoromethane, 551
Dichlorodiphenyldichloroethylene. *See*
DDE
Dichlorodiphenyltrichloroethane. *See*
DDT
1,1-Dichloroethylene, NMR spectrum
of, 500
Dichloromethane, 233f, 448
as solvent in extraction procedure,
707–708, 708f, 709f
2,4-Dichlorophenoxyacetic acid herbicide
(2,4-D), 646f
trans-1,3-Dichloropropene, 517
Dictyopterene D', 460
Dicyclohexylcarbodiimide (DCC), 846f,
849–850, 1095, 1098, 1100
Dicyclohexylurea, 849
Dicyclopentadiene, 595–596
Didrex. *See* Benzphetamine
Dieckmann reaction, 932–933
Diels–Alder reaction, 588–597
diene reactivity in, 590–591
dienophile reactivity in, 591
drawing product of, 590
features of, 589
retro, 595–596
retrosynthetic analysis of product, 595
rule of endo addition, 593–594
rules governing, 590–594, 593–594
stereospecificity of, 592
in steroid synthesis, 596–597
tetradotoxin synthesis, 589, 589f
as thermal reaction, 589
Dienes, 362
conjugated. *See* Conjugated dienes
isolated, 572, 573f, 583, 583f, 584
in metathesis, 1017–1019
ozonolysis of, 446
1,3-Dienes, 572–573, 604. *See also*
Conjugated dienes
polymerization of, 1160
synthesis in Suzuki reaction,
1006–1009
Dienophiles
cis, 590–591, 592
common, 594f
cyclic, 590–591, 592
Diels–Alder reaction, 588–596
reactivity of, 591
stereochemistry, 592
trans, 590–591, 592
Diesel fuel, composition of hydrocarbons
in, 128
Diesters
in Dieckmann reaction, 932–933
1,3-diesters, pK_a of, 886f
1,6-diesters, in Dieckmann reaction,
932
1,7-diesters, in Dieckmann reaction,
933
 β -diesters
as active methylene compounds,
924
synthesis in crossed Claisen
reaction, 931–932
Diethyl acetamidomalonate, 1079
Diethylamine, solubility of, 96
Diethyl carbonate, in crossed Claisen
reaction, 931
1,2-Diethylcyclohexane, 155
cis-1,2-Diethylcyclopropane, 1015
Diethyl ether, 84, 108
as anesthetic, 233f, 320
boiling point of, 779
in extraction procedures, 707
infrared (IR) spectrum, 482
naming, 316
solubility of, 96
as solvent, 94, 341, 707
Diethyl maleate, 1026
Diethyl malonate, 910, 923, 932
alkylation of derivatives, 1079–1080,
1081f
conversion to a carboxylic acid,
900–903
Diethyl oxalate, 946
Diethyl tartrate (DET), 452–453
N,N-Diethyl-*m*-toluamide, 842–843
Diethyl tartrate (DET), 452–453
Digoxin, 812
Dihalides, 299
alkyne synthesis from, 404–405
Dihalocarbenes, 1012–1014
Dihedral angle, 131, 132f, 135, 136f, 295
Dihydroxyacetone, 1029, 1035, 1036f
1,4-Dihydroxybenzene, 1164
Dihydroxylation, 442–444
anti, 442, 443
syn, 442, 443–444
1,4-Dihydroxymethylcyclohexane, 860
6,7-Dihydroxy-2-nonanone, 823
Diisobutylaluminum hydride (DIBAL-H),
734, 735, 736f, 738t, 865–866
Diisopropylamine, 888, 952
 β -Diketone, pK_a of, 886f
Dimers, carboxylic acids as, 692, 693f
2,2-Dimethoxy-1,3-cyclopentanedi-
carbaldehyde, 785
5,5-Dimethoxy-2-pentanone, 820
Dimethylacetylene carboxylate, 606
Dimethylallyl diphosphate, 1134–1135,
1135f
4-(*N,N*-Dimethylamino)pyridine, 973
N,N-Dimethylbenzamide, 832
1,3-Dimethylbenzene, 667
2,3-Dimethylbutanal, 776
3,3-Dimethyl-2-butanol, 328, 329
3,3-Dimethyl-2-butanone, 536
2,2-Dimethyl-3-butenal, 779
2,3-Dimethyl-1-butene, 397
3,3-Dimethyl-1-butene, 567
2,3-Dimethyl-2-butene, 397
Dimethyl carbonate, 769
(1*E*,4*R*)-1,4-Dimethylcyclohexene, 395
1,2-Dimethylcyclohexane, 127f
1,3-Dimethylcyclohexane, 126
1,4-Dimethylcyclohexane, 144–146, 146f
2,2-Dimethyl-1,3-cyclohexanedione, 909
5,5-Dimethyl-1,3-cyclohexanedione, 909
2,2-Dimethylcyclohexanone, 898, 899
2,6-Dimethylcyclohexanone, 898
1,2-Dimethylcyclohexene, 377, 377f
1,6-Dimethylcyclohexene, 363f
1,2-Dimethylcyclopentane, 144
cis isomer, 163f
trans isomer, 163f
1,3-Dimethylcyclopentane, 186
3,3-Dimethylcyclopentene, 353
Dimethylcyclopropanes, 194, 1026
Dimethyl ether
isomers of, 17, 78
NMR spectrum of, 524
Dimethylformamide (DMF), 537
boiling point of, 92
as solvent, 241f, 896
2,5-Dimethyl-3-heptyne, 402f
4,5-Dimethylhexanoic acid, 690
2,4-Dimethyl-3-hexanone, 760
2,4-Dimethylhexanoyl chloride, 833
6,6-Dimethyl-3-octyne, 402
2,2-Dimethylloxirane, 317, 346
2,3-Dimethylloxirane, 345
2,3-Dimethylpentane, 123f
mass spectrum of, 470–471
2,2-Dimethylpropane, 116
2,2-Dimethylpropanoic acid, 704
Dimethyl sulfide, conversion of ozonide to
carbonyl compounds, 445–446
Dimethyl sulfoxide (DMSO), 266
aqueous, 383–384
as polar aprotic solvent, 241f
as solvent for alkyne synthesis, 299,
300f
Dimethyl terephthalate, 1162, 1170
1,3-Dinitriles
active methylene compounds,
923–924
 pK_a , 886f
Dinoprostone, 1130
-dioic acid (suffix), 692
-diol (suffix), 315
Diols, 315
1,2-diol formation by dihydroxylation,
442–444
gem-diol, 802–804
p-Dioxanone, 1177
Dioxybenzene, 109
Dipeptides
constitutional isomers of, 1087
synthesis of, 1086–1087, 1094–1098
Diphenhydramine, 275, 610, 968
Diphenyl carbonate, 1163, 1168
Diphosphate, as leaving group,
1134–1137
Dipole, 43, 44, 45f
in alkenes, 365
temporary, 87–88
Dipole–dipole interactions, 89, 90t
of aldehydes and ketones, 779
of alkyl halides, 231
of carbonyl compounds, 834
of carboxylic acids, 692
of nitriles, 834
in polar aprotic solvents, 241, 265
Dipole moment, infrared (IR) absorption
and, 480
Directed aldol reactions, 925–926, 925f
Disaccharides, 1056–1058
artificial sweeteners, 1058, 1059f
general features, 1056
lactose, 1027, 1057–1058
maltose, 1057
sucrose, 1058, 1058f
Discodermolide, 195
Disorder, entropy as measure of, 209
Disparlure, synthesis of, 441, 442f
Disproportionation, 1152
Dissolving metal reductions, 428, 436
Distillation, of crude petroleum, 129
Distillation apparatus, 92, 92f
trans-1,2-Disubstituted cycloalkanes, 345
Disulfide bonds
in α -keratins, 1108
in peptides, 1089
in tertiary structure of proteins, 1105,
1106f
in vulcanized rubber, 1159, 1159f
Diyne, 401
DMAP, 973
DMF. *See* Dimethylformamide
DMSO. *See* Dimethyl sulfoxide
DNA, 104f, 105, 1064–1065, 1065f
Dobutamine, 952
Docetaxel, 876
Docosanoic acid, 460
Dodecahedrane, 127, 138, 606
Domagk, Gerhard, 990
Donepezil, 85, 926
L-Dopa (levodopa), 6, 45, 191, 1115
Dopamine, 820, 958, 959f
Double bonds. *See also* Alkenes
in alkenes, 83, 84t
in alkyl halides, 229, 229f
bond length and bond strength, 41, 41t
components of carbon–carbon, 37,
38f
in condensed structures, 28
conjugated, 576, 920
degrees of unsaturation, calculating,
431–432
effect on melting point of fatty acids,
368, 368t
in functional group, 83, 84t
hydrogenation of, 432–433, 433f
Lewis structure, 15
nitrogen–nitrogen, 986
NMR spectra of, 505, 506f, 506t
 π (π) bonds, 82
planar, 376–377, 381, 386, 388
radical reactions with, 541, 558–560
in resonance hybrid, 22
resonance structures
conjugated, 576
major contributor, 577–578, 582

- minor contributors, 577–578, 582
with one atom more electronegative than the other, 576, 577
rotation restriction, 37, 360t
in triacylglycerols, 369, 1122–1124, 1122t
in unsaturated fatty acids, 358, 368
Z configuration, 368
- Double dagger notation, 211
- Double-headed arrow, 18, 56–57, 202t
- Doublet, NMR spectrum, 509–510, 511t, 512, 512t, 513
- Doublet of doublets, NMR spectrum, 516, 516f
- Doxorubicin, 1029f
- Dyes
azo compounds, 987, 988–991
binding to fabric, 989–990
direct, 989
natural and synthetic, 988–989
sulfa drugs, 990–991
- Dyneema, 1159
- E,Z* system of nomenclature, 364
- E1cB elimination reactions, 920
- Eclipsed conformation, 131–137, 132f, 133f, 134f, 136f, 136t, 295f
- Edman degradation, 1091–1092
- E1 elimination reactions
acid-catalyzed dehydration of alcohols, 369–370
base in, 291, 293, 293t, 298
characteristics of, 293t
dehydration of secondary and tertiary alcohols, 326
E2 mechanism compared, 298–299, 298t, 300–304, 302f–303f
energy diagram for, 292, 292f
kinetics of, 291, 293t
leaving group in, 292, 293t
rate of reaction, 292–293, 293t
regioselectivity of, 293
S_N1 substitution reactions compared, 294, 300–304, 302f–303f
S_N2 substitution reactions compared, 300–304, 302f–303f
solvent for, 293t
two-step mechanism, 291–292, 293t
- E2 elimination reactions
acetylide anion reactions with alkyl halides, 414–415
alkyl halide identity in, 287–288, 288t
alkyl tosylates reactions, 339
alkyne synthesis, 299–300, 404–405
anti periplanar geometry in, 295–298, 295f
base in, 286, 286f, 288t, 298
characteristics of, 288t
dehydration of alcohols, 327, 330–331
dehydrohalogenation of alkyl halides, 369, 370
trans diaxial geometry in, 296–298, 296f
- E1 mechanism compared, 298–299, 298t, 300–304, 302f–303f
energy diagram of, 285, 286f
in Hofmann elimination, 978, 979, 979f
kinetics of, 285, 288t
leaving group in, 287, 288t
one-step mechanism, 285–287, 288t
organic synthesis examples, 289f
rate of reaction, 287
regioselectivity of, 290
S_N1 and S_N2 substitution reactions compared, 300–304, 302f–303f
solvent for, 287, 288t
stereochemistry of, 295–298, 295f
stereoselectivity of, 290
- syn periplanar geometry in, 295, 295f
Zaitsev rule, 288–291, 297
- Effexor. *See* Venlafaxine
- E-85 fuel, 319, 320f
- Ehrlich, Paul, 990
- Eicosane
constitutional isomers of, 118t
molecular formula, 118t
- Eicosanoids, 1129–1132
biological activity of, 1130, 1130t
biosynthesis of, 1130–1132, 1131f
derivation of word, 1129
- E* isomer, 364
- Elaidic acid, 395
- Elastomer, 1159
- Electromagnetic radiation, 474–476, 475f
- Electromagnetic spectrum, 475, 475f
- Electron cloud, 7
- Electron-deficient sites/species, 58, 82, 84, 85, 102–103, 202, 723
- Electron density, delocalization, 67
- Electron-donating groups, 256, 654–657, 660, 703–706, 706f
- Electron dot representation. *See* Lewis structures
- Electronegativity
acidity trends and, 63–66
bond polarity and, 43, 44
definition of, 42
inductive effects on substituted benzenes, 654
oxidation and, 147
periodic trends in, 42, 42f
values for common elements, 42f
- Electron pair acceptor, 55, 72
- Electron pair donor, 55, 72–73
- Electron pairs
in Brønsted–Lowry acid–base definition, 55, 56
Lewis acids and bases, 72–74
movement of, 56, 201
- Electron-poor sites/species, 73
- Electron-rich sites/species, 58, 73, 102–103, 370
- Electrons
bonded (shared), 12
core, 9
delocalized, 18, 22, 578–579, 609
density of, 8–9, 32, 43–44, 44f
excited state, 597–598
ground state, 597–598
in Lewis structure, 12–14
lone pairs, 21
nonbonded (unshared), 12–14, 12f
number of, 7, 9
in resonance structures, 18–23
unpaired, 200–201
valence, 10–16, 12–16, 17t
- Electron-withdrawing groups, 256, 591, 654–657, 660–661, 665, 703–707, 706f
- Electrophiles, 85
alkyne reactions with, 405
carbocations as, 201, 202
Lewis acid, 73
meaning of term, 73
radicals as, 202
reactivity of, 102–103
- Electrophilic addition reactions, A–14
of conjugated dienes
kinetic *versus* thermodynamic products, 586–588, 587f, 588f
1,2- *versus* 1,4-addition, 584–586
energy diagram for, 372, 373f
halogenation of alkynes, 409
hydration of alkenes, 378–379, 390
hydrohalogenation of alkenes, 371–378, 372f, 373f, 374f, 377f, 378t
hydrohalogenation of alkynes, 407–408
- Markovnikov's rule on, 374–375, 374f, 378t
stereochemistry of, 376–378, 377f, 378t
- Electrophilic aromatic substitution reactions, 641–687, A–13
azo coupling, 986–987
energy changes in, 644, 644f
examples of, 643f
Friedel–Crafts acylation, 643f, 647–653, 653f, 665–666, 672–673, 677
Friedel–Crafts alkylation, 643f, 647–653, 665–666, 672–673, 675–676
halogenation, 643f, 644–645, 646f, 665, 666–667
mechanism of, 642–644
nitration, 643f, 646–647, 667–668
of substituted benzenes, 657–660, 660f
disubstituted benzenes, 666–668
limitations on reactions, 665–666
rate of reactions, 660, 660f
sulfonation, 643f, 646–647
- Electrophilic carbon, in alkyl halides, 234
- Electrospray ionization (ESI), 474
- Electrostatic interactions, between ions, 10–11
- Electrostatic potential plot
of acetic acid, 67f, 689, 689f
of acetylene, 405, 405f
of an alcohol, 314f
of an alkoxide, 885, 885f
of amines, 951, 951f
of benzene, 610, 610f
of carbanions, 68f
of carbocations, 257, 257f
localized and delocalized, 574, 574f
carbon dioxide, 45f
description of, 43–44
of dienes, 573, 573f
of an enolate, 885, 885f
of an epoxide, 314f
of ethanol, 66f, 67f
of an ether, 314f
of ethylene, 370, 370f
of formaldehyde, 723, 723f
of halomethanes, 234f
of methyl chloride, 44, 44f
of pyridine and pyrrole, 622, 623f
of substituted anilines, 971, 971f
of 2,2,2-trifluoroethanol, 66f
of vitamins A and E, 1128, 1129f
of water, 45f
- Elements, periodic table, 8–10, 8f
- Eleostearic acid, 395
- Elimination reactions, A–14
of alcohols, 323–331
alkene synthesis, 279–280, 369–370
of alkyl halides, 234, 278–311
alkyne synthesis, 279, 404–405
alpha (α), 1012
beta (β), 279
bimolecular, 285. *See also* E2 elimination reactions
description of, 198
elimination, unimolecular, conjugate base (E1cB) mechanism, 920
features of, 279–280
mechanisms of, 285
E1. *See* E1 elimination reactions
E2. *See* E2 elimination reactions
E1 and E2 compared, 298–299, 298t, 300–304, 302f–303f
reductive, 1006, 1008, 1011
stereochemistry, 295–298
unimolecular, 285. *See also* E1 elimination reactions
Zaitsev rule on, 288–291
- Enalapril, 191
- Enals, 779
- Enamines
hydrolysis of, 801
synthesis of
from aldehydes and ketones, 800–801, 801f
amine reaction with aldehydes and ketones, 975
- Enanthotoxin, 25
- Enantiomeric excess, 184–185, 453, 732
- Enantiomers
of amino acids, 710–711, 1075, 1075f
separation of, 1081–1084, 1082f
of an ammonium salt, 951
in carbonyl reductions, 729–730
chemical properties of, 186–188
chiral drugs, 187
of cyclic compounds, 169
definition of, 164
drawing, 168, 168f
in electrophilic addition to alkenes, 376–377, 377f
enantioselective reactions, 452–453
in enolate alkylation, 898
from epoxidation of alkenes, 440, 441
in epoxide ring opening reactions, 344–345
formation in halogenation of alkenes, 380, 381, 382
halogenation reactions and, 549–551
in halohydrin formation, 384
in hydroboration–oxidation reactions, 389
labeling stereogenic centers with *R* or *S*, 170–175, 172f, 175f, 179–180
of monosaccharides, 1030, 1033–1034, 1034f, 1036
in nucleophilic substitution reactions, 246, 254–255
optical purity, 184–185
physical properties, 182–186, 184t, 186f
racemic mixture of, 183–184, 184t, 376
sense of smell and, 187–188, 188f
three-dimensional representations for, 168f
- Enantioselective reactions, 452–453
amino acid synthesis, 1085–1086
biological reduction, 732–733
carbonyl reductions, 731–733, 732f
- Enantiotopic protons, NMR signals of, 500–502
- Enclomiphene, 394
- Endergonic reaction, 210
- Endo addition, 593–594, 594f
- Endothermic reaction
bond breaking and, 203–204
bromination, 547, 547f
dehydration of alcohols, 327f
energy diagram of, 212f
enthalpy change and, 210
reaction rate, 259
transition state in, 258–259, 259f
- ene (suffix), 362
- Energy
nonrenewable, 129
of photons, 474, 475–476
release in bond formation, 202
torsional, 133–134
units of measurement, 40
- Energy barrier, 211, 213–214, 215, 285, 292
- Energy change, during the chlorination of ethane, 544, 544f, 545f
- Energy diagrams, 210–215, 212f, 215f
for carbocation formation, 260f
for chlorination of ethane, 545f
for E1 elimination reactions, 292, 292f
for E2 elimination reactions, 285, 286f

- Energy diagrams (*continued*)
 for electrophilic addition of HBr to 1,3-butadiene, 587, 588f
 for electrophilic additions, 372, 373f
 for electrophilic aromatic substitution, 644, 644f
 rate of electrophilic aromatic substitution of substituted benzenes, 660, 660f
 for S_N1 substitution reactions, 253, 253f, 260f
 for S_N2 substitution reactions, 245, 245f, 249f
 for two-step reaction mechanisms, 213–215, 215f
- Energy of activation (E_a)
 catalysts and, 219, 219f
 definition of, 211
 energy barrier height determined by, 211
 in energy diagrams, 211–215, 212f, 215f
 in exothermic reactions, 259, 260f
 rate constant, relationship to, 217
 reaction rate and, 215–217
 transition state energy effect on, 259
- Enol(s), 881–884
 hydroxy group of, 313
 reaction mechanism, 883–884
 resonance structures, 883–884
 synthesis of, 755
 hydration of alkyne, 409–412
 hydroboration–oxidation of alkynes, 412–413
 tautomers, 881–883
- enol (suffix), A–5
- Enolate(s), 21, 884–891
 achiral, 891
 in aldol reactions, 918, 918f, 920, 922, 923, 923f, 925, 927
 alkylation, 897–899
 general features of, 897–899
 tamoxifen synthesis, 899
 of unsymmetrical ketones, 898–899
 in Claisen reactions, 928–932
 electrostatic potential plot of, 885, 885f
 examples, 885–886
 kinetic, 889–890, 898, 925
 in Michael reaction, 934–936, 935f
 reactions of, 888–889, 892, 897–899
 resonance-stabilized, 920, 929, 935
 resonance structures, 884
 in Robinson annulation, 936–939
 synthesis of, 755
 with bases, 887–888, 887t
 from carbonyl compounds, 884
 from esters, 885–886
 from tertiary amides, 885–886
 thermodynamic, 890–891, 899
 of unsymmetrical carbonyl compounds, 889–891
- Enol tautomers, 410–412, 882–883
- enone (suffix), A–5
- Enones, 779
- Entacapone, 52
- Enthalpy change (ΔH°), 203–206
 calculation from bond dissociation energies, 205–206
 in dehydration of alcohols, 327, 327f
- E1 elimination reactions, 292, 292f
 in energy diagrams, 211–215, 212f, 215f
- free energy change, relation to, 209–210
 negative, 203, 210
 positive, 203, 210
- Entropy (S°), 209
- Entropy change (ΔS°), 209–210
- Environmental Protection Agency, 450
- enyne (suffix), A–5
- Enynes, 401
- Enzymes, 219–220, 220f, 1075
 active site of, 219, 220f
 catalytic action of, 219–220, 220f
 in kinetic resolution of amino acids, 1084
 in oxidation reactions, 451
 in peptide sequencing, 1093, 1093t
- Enzyme–substrate complex, 219–220, 220f
- Ephedrine, 166–167, 192
- Epi-aristolochene, 1147
- Epiclorohydrin, 356, 1163, 1164f
- Epimers
 definition of, 1035
 of monosaccharides, 1051, 1053–1054
- Epinephrine, 228
 structure of, 958
 synthesis of, 228, 251, 251f
- Eplrenone, 321
- Epoxidation, 439–442, 439f
 in anti dihydroxylation, 443
 Sharpless, 451–454
 of squalene, 1140, 1140f
 stereochemistry of, 440
 in synthesis of disparlure, 441, 442f
- Epoxides, 277
 achiral, 443
 anionic polymerization of, 1156–1157
 applications, 347–349
 boiling point of, 318t
 bonding, 314
 electrostatic potential plot of, 314f
 interesting examples, 321
 melting point of, 318t
 nomenclature, 317–318
 physical properties of, 318–319, 318t
 reactions of, 343–347
 with acetylide anions, 416–417
 with acids HZ, 345, 346f, 347
 general features, 324
 with organometallic reagents, 754–755
 reduction, 437–438, 438f
 with strong nucleophiles, 343–345
 reduction of, 437–438, 438f
 ring opening reactions, 1156–1157
 with acids, 345–347
 enantiomers in, 344–345
 nucleophilic substitution reactions, 343–347
 regioselectivity of, 345–346
 solubility of, 318t
 structure of, 313, 314
 synthesis of, 322–323
 from alkenes, 439–441
 from halohydrins, 323
- Epoxy (prefix), 317
- Epoxyalkane, 317
- 1,2-Epoxy-cyclohexane, 317, 344
- 1,2-Epoxy-2-methylpropane, 317
- cis-2,3-Epoxy-pentane, 317
- Epoxy resins, 1163–1164, 1164f
- Equal. *See* Aspartame
- Equations for organic reactions, writing, 197, 197f
- Equatorial bond, 160
- Equatorial hydrogens, 139–140, 139f, 141f
- Equilibrium, 206–208, 207f, 208t
 acid–base reactions, 61–62
 direction of, 216
 nucleophilic substitution and, 238
- Equilibrium arrows, 56–57, 202t
- Equilibrium constant (K_{eq}), 206–209, 207f, 208t, 216
- Ergot, 641
- Erlotinib, 421
- D-Erythrose, 193, 1034, 1034f, 1035f
- L-Erythrose, 1034, 1034f
- D-Erythrulose, 1036f
- ESI (electrospray ionization), 474
- Esmolol, 79
- Essential amino acids, 711, 1076f, 1077
- Essential fatty acids, 368, 369
- Essential oils, 1132
- Esterification, 227
- Esterification, Fischer, 847–848, 859–860
- Esters
 acidity/basicity of, 829t
 boiling point of, 834t
 cholesteryl, 861, 861f
 chromate, 448
 cyclic, 837
 enolates from, 885–886
 functional group in, 86t
 general structure of, 367
 interesting examples, 836–837
 IR spectra of, 835, 835t
 leaving group, 725
 melting point of, 834t
 NMR spectra of, 836
 nomenclature of, 830–831, 833, 833t
 nucleophilic substitution reactions, 725
 odors of, 836
 physical properties of, 834t
 pK_a , 829t, 886t
 polyesters, 859–860
 reactions of, 850–855, 855f, 858t
 with ammonia and amines to form amides, 850
 carboxylic acid preparation by malonic ester synthesis, 900–903
 Claisen reaction, 928–932, 929f
 hydrolysis to carboxylic acids, 850–852
 lipid hydrolysis, 853–855
 methyl ketone preparation by acetoacetic ester synthesis, 900, 903–905
 reaction with organometallic reagents, 750–752
 reduction of, 734
 solubility of, 834t
 stability of, 829
 structure and bonding of, 722, 826, 828–829
 synthesis of
 from acid chlorides, 842
 from anhydrides, 844–845
 from carboxylic acids, 846f, 847–849
 from monosaccharides, 1046
 thioesters, 860–861
 in triacylglycerols, 366, 367
 waxes, 1121
- Estradiol, 770
 analogs of, 402, 403
 function in body, 1141t
 structure of, 49, 157, 403, 1141t
 synthesis of, 288, 289f
- Estragole, 684
- Estrogens, 1141, 1141t
- Estrone
 function in the body, 1141t
 structure of, 117, 750, 935t, 1141t
 synthesis of, 687
 from a chlorohydrin, 385f
 in Diels–Alder reactions, 596–597
 using the Michael reaction, 935, 935f
- Ethambutol, 49
- Ethanal. *See* Acetaldehyde
- Ethane
 acidity of, 67–68, 69f
 bonds in, 36, 40f
 carbon–carbon bond length in, 581
 chlorination of, 543–544, 544f, 545f
 conformations of, 129–134, 132f, 133f
 formation from ethylene and hydrogen, 218–219
 molecular formula, 114, 115, 118t
 nonreactivity of, 82, 83
 structure of, 82, 83, 114
- Ethanedioic acid, 692
- 1,2-Ethanediol, 316
- Ethanol, 110, 227
 in acid–base reactions, 61–62
 acidity of, 61–62, 65, 66, 67, 701, 703f
 in alcoholic beverages, 319, 426
 description, 2–3
 electrostatic potential plot, 66f, 67f
 in ethyl acetate formation, 218–219
 isomers of, 17, 78
 NMR spectrum of, 518, 518f
 oxidation of, 451
 pK_a of, 701, 703f
 as polar protic solvent, 240f
 preparation from ethylene, 366f
 production by fermentation of carbohydrates, 319, 320f
 reactivity of, 82
 as renewable fuel source, 319, 320f
 sodium ethoxide production from, 322
 solubility of, 95
 structure, 82, 83
- Ethene, 365. *See also* Ethylene
- Ethers
 boiling point of, 318t
 bonding of, 314
 crown, 101–102, 320–321, 321f
 cyclic, 317
 electrostatic potential maps, 314f
 formation by alcohol addition to alkenes, 378–379
 functional group, 85t
 infrared (IR) spectra of, 482
 interesting examples, 320–321, 321f
 melting point of, 318t
 nomenclature, 316–317
 physical properties of, 318–319, 318t
 preparation by nucleophilic substitution reaction, 269
 reactions of
 alkyl halide formation, 341–343
 general features, 324
 with strong acid, 341–343
 simple and complex, 316
 solubility of, 318t
 structure of, 313, 314
 symmetrical, 313, 322
 synthesis of
 from alkyl halides, 321–322
 from monosaccharides, 1046
 by nucleophilic substitution reactions, 321–322
 Williamson ether synthesis, 322
 unsymmetrical, 313, 322
- Ethoxide, 61–62, 66, 701, 703f
 nucleophilicity of, 239
- Ethoxy group, 316
- 4-Ethoxyoctane, 317
- Ethyl acetate, 227
 in Claisen reaction, 928
 formation from acetic acid and ethanol, 218–219
 as solvent, 847
 structure of, 831
 synthesis of, 847
- Ethyl acetoacetate, 888, 908, 914
 conversion to ketone, 903–905
 synthesis in Claisen reaction, 928
- Ethyl *p*-aminobenzoate, 673
- N*-Ethylaniline, 953, 996

- Ethylbenzene, 611
 bromination of, 670
 styrene synthesis from, 671
- Ethyl benzoate, in crossed Claisen reactions, 879, 930–931
- Ethyl butanoate, 78, 836
- Ethyl chloride, 84, 231, 524
- Ethyl chloroformate, in crossed Claisen reaction, 931
- Ethyl chrysanthemate, 873
- Ethyl α -cyanoacrylate, 1155f
- Ethylcyclobutane, 127f
- N*-Ethylcyclohexanamine, 965
- 2-Ethylcyclohexanecarbaldehyde, 776
- 2-Ethylcyclohexanone, 898
- 4-Ethyl-3,4-dimethylcyclohexane, 123f
- 5-Ethyl-2,6-dimethylcyclohexane, 123f
- Ethylcyclopentane, 489
- Ethylene, 365
 acidity of, 67–68, 69f
 bond dissociation energies, 400–401
 bonds of, 36–38, 38f, 40f, 359
 bromine addition to, 608
 as Brønsted–Lowry base, 55f, 56
 electrostatic potential plot, 370f
 ethane formation from, 218–219
 functional group in, 83, 84f
 in Heck reaction, 1009–1010
 as industrial starting material, 366, 366f
 Lewis structure, 14–15
 molecular shape of, 24–25, 26f
 monomers in polymerization, 561
 pK_a of, 60
 polymerization of, 1153, 1158
 preparation from petroleum, 366
 structure of, 1151f
 synthesis from alkyl tosylates, 339
- Ethylene glycol, 155, 451, 807, 1176
 acetal formation from, 804
 in antifreeze, 319f
 formation from PET recycling, 1170
 naming, 316
 in polyester synthesis, 860
 in polyethylene terephthalate (PET), 1148, 1150f, 1161–1162
 in polyurethane synthesis, 1162
 preparation from ethylene, 366f
 structure of, 319f
- Ethylene oxide, 317
 polymerization of, 1156–1157
 ring opening of, 1156–1157
- 1-Ethyl-2-fluorocyclopentane, 231f
- N*-Ethylformamide, 832
- Ethyl formate, in crossed Claisen reactions, 930–931
- Ethyl group, 121
- 2-Ethyl-1-heptene, 362
- Ethyl 2,4-hexadienoate, 946
- 1-Ethyl-3-isopropylcyclohexane, 155
- Ethyl isopropyl ketone, 779f
- 1-Ethyl-3-methylcyclohexane, 126
- N*-Ethyl-*N*-methylcyclopentanamine, 953
- N*-Ethyl-*N*-methylcyclopentylamine, 953
- Ethyl methyl ether, synthesis from alkyl tosylates, 339
- 3-Ethyl-3-methylheptane, 489
- Ethyl methyl ketone, 778
- 4-Ethyl-5-methylcyclohexane, 123f
- Ethyl *p*-nitrobenzoate, 673
- 3-Ethylpentane, bromination of, 546
- 2-Ethyl-1-pentene, 376
- Ethyl phenylacetate, 875
- Ethyl tosylate, 339
- Ethyne, 402. *See also* Acetylene
- Ethynylcyclohexane, 402f, 1021
- Ethynylestradiol, 399f, 402–403, 403f, 745, 745f, 750, 1142
- Ethynyl group, 402
- Etoside, 817
- Excited state
 of atom, 32
 of electron, 597–598
- Exergonic reaction, 210
- Exo product, 593–594, 594f
- Exothermic reactions
 addition reaction, 372, 372f
 bond formation and, 203–204, 206
 chlorination, 544, 544f, 547–548, 548f
 energy diagram of, 211, 212f
 hydrogenation, 429
 reaction rate, 259
 transition state in, 258, 259f, 260f
- Exponent, 59
- Extraction, 707–709, 708f, 709f
 for separation of amines, 967–968, 967f
- Fabric, binding of dyes to, 989–990
- Farnesene, 306, 367f, 601
- Farnesol, 1133f, 1136
- Farnesyl diphosphate, 1135–1136, 1135f, 1147
- Fats, 369, 432–433, 853, 854
 body content of, 1125
 degree of unsaturation, 432
 fatty acid composition, 1123, 1123t
 melting point of, 369, 1122–1123
 saturated, 369
 trans, 433
- Fat-soluble vitamins, 97, 1128, 1129f
- Fatty acids
 characteristics of, 1122
 composition in fats and oils, 1123, 1123t
 essential, 368, 369, 1122
 melting point of, 368, 368f
 omega-3, 368, 369
 oxidation of, 853
 saturated, 358, 368, 433, 1122, 1122t, 1123, 1123t
 synthesis in lipid hydrolysis, 853, 854–855
 thioesters of, 861
 trans, 433
 triacylglycerol formation and, 105
 in triacylglycerols, 367, 369, 1122–1125, 1122t
 unsaturated, 358, 368, 1122, 1122t, 1123, 1123t, 1124
 hydrogenation of, 1124
 oxidation of, 1124
- Feedstock, 1166
- Fehling's reagent, 1048
- Femara. *See* Letrozole
- Fenfluramine, 76
- Fenn, Dr. John, 474
- Fermentation, alcohol production by, 319, 320f
- Ferulic acid, 605
- Fexofenadine, 49, 957, 958
- Fibers
 natural, 858, 858f
 synthetic, 825, 858–860
- Fibrinogen, 1075
- Fibrous proteins, 1106
- Fingerprint, of IR spectrum, 477, 478, 478f, 479, 479f
- First-order kinetics, 217
 of E1 elimination reactions, 291, 293t
 of nucleophilic substitution reactions, 243–244
 of S_N1 substitution reactions, 252
- First-order rate equation, 217, 243, 244
- Fischer, Emil, 1053
- Fischer esterification, 847–848, 859–860
- Fischer projection formulas, 1030–1033, 1033f
- Fischer proof of the structure of glucose, 1053–1056, 1054f
- Fishhook notation, 201
- Five-membered rings, synthesis of
 in Dieckmann reaction, 932–933
 in intramolecular aldol reactions, 926–928
- FK506, 837
- Flagpole hydrogens, 141, 141f
- Flat helical ribbon, shorthand for α -helix, 1103–1104
- Flat wide arrow, shorthand for β -pleated sheet, 1104
- Fleming, Sir Alexander, 463, 857
- Flexibilene, 1146
- Flonase. *See* Fluticasone
- Flosal, 923
- 9-Fluorenylmethoxycarbonyl protecting group (Fmoc), 1096, 1099–1100
- Fluorine
 polarizability, 88–89
 valence electrons of, 11
- Fluoromethane, polarizability and boiling point of, 91f
- m*-Fluoronitrobenzene, 611
- Fluoxetine, 3, 76, 187, 250, 250f, 732, 958
- Fluticasone, 230
- Fmoc (9-fluorenylmethoxycarbonyl protecting group), 1096, 1099–1100
- Folic acid, 990–991
- Formal charge, 15–16, 17t
- Formaldehyde
 aldol reactions of, 922
 Bakelite preparation from, 1165
 electrostatic potential map of, 723, 723f
 formation from methanol oxidation, 451
 hydration of, 802
 primary alcohol formed from, 743
 structure of, 777, 783
 uses of, 783
- Formaldehyde hydrate, 802
- Formalin, 783
- Formic acid, 60, 119, 451, 691t, 694
- Formyl group, 779
- Fosamax (alendronic acid), 18
- Fossil fuels, 70, 128–129, 149
- Four-centered transition state, 386
- Fragmentation, 465, 469–472, 470f
- Free energy change (ΔG°), 206–209, 207f, 208t
- Frequency
 of electromagnetic radiation, 474–476, 475f
 of infrared radiation, 476–477
- Friedel, Charles, 643
- Friedel–Crafts acylation, 643f, 647–653, 677
 alkyl benzene synthesis, 672–673, 672f
 of benzene, 643f
 electrophilic aromatic substitution reaction, 643f
 general features of, 647–648
 intramolecular, 653, 653f
 ketone synthesis, 643f, 648, 672–673
 limitations of, 665–666
 mechanism of, 649–650
- Friedel–Crafts alkylation, 643f, 647–653, 675–676
 of alcohols, 652
 of alkenes, 652
 in alkyl benzene synthesis, 647–648, 672–673, 672f
 of benzene, 643f
 carbocations and rearrangements, 650–652
 electrophilic aromatic substitution reaction, 643f
- general features of, 647–648
 intramolecular reactions, 653
 limitations in, 665–666
 mechanism of, 648–649
 polyalkylation, 666
 unreactive halides, 650
- Frontalin, 454, 454f
- Frontside attack, in nucleophilic substitution reactions, 245–246
- Frost circle, 629
- Fructofuranose ring, 1058
- Fructose
 D-fructose, 1029, 1035, 1036f
 stereogenic centers, 166–167
 in sucrose, 1058
- Fuels, renewable, 319, 320f
- Full-headed curved arrow, 201, 202t
- Fumaric acid, 112, 592
- Functional groups, 82–87. *See also specific functional groups*
 compounds containing a C=O group, 84, 85t
 compounds containing a C–Z σ bond, 84, 85t
 definition of, 82
 drawing connection to carbon skeleton, 82
 fragmentation pattern in mass spectrum, 471–472
 grouping organic compounds by, 83
 hydrocarbons, 83–84, 84t
 importance of, 86
 infrared (IR) spectroscopy and, 477, 478, 478f, 479, 479f, 480t
 intermolecular forces and, 87
 nomenclature of, A–4
 nucleophilic substitutions and, 267, 268t
 reactivity and, 102–104
 synthesis of, A–15–18
- Funk, Casimir, 97
- Funnel, separatory, 707, 708f
- Furan, 636, 1037
- Furanose rings, 1037, 1041–1042
- Fused ring system, 592–593, 593f
- Gabapentin, 170
- Gabriel synthesis, 961–962
- Galactose, 156, 1057
 D-galactose, 1035f, 1050
 in lactose, 219, 220f
 Wohl degradation, 1050
- Gamma (γ) carbon, 691
- Garlic, 119
- Gas chromatography–mass spectrometry (GC–MS), 473–474, 473f, 474f
- Gasohol, 319, 320f
- Gasoline
 additives
 benzene, 83
 BTX mixture, 614
 ethanol, 319
 high-octane, 497, 614
 MTBE, 95
 combustion of, 148
 composition of hydrocarbons in, 128, 129
 isooctane in, 196, 216
 octane rating, 83
 oxidation of isooctane, 196
- Gauche conformation, 134f, 135, 136f
- GC–MS (gas chromatography–mass spectrometry), 473–474, 473f, 474f
- Geckos, 389
- Geminal dichloride, 404
- Geminal dihalide, 299
 alkyne synthesis from, 404–405
 synthesis in hydrohalogenation of alkynes, 406

- Geminal protons, 516, 517f
 Generic name, 119
 Geranial, 306, 783f
 Geraniol, 454, 1134, 1136
 Geranyl diphosphate, 1134–1137, 1135f, 1136–1137
 GHB (γ -hydroxybutyric acid), 696, 872
 Gibberellic acid, 945
 Gibbs free energy (G°), 207
Ginkgo biloba, 4–5, 688
 Ginkgolide B, 4–5
 Glacial acetic acid, 695
 Gleevec, 837
 Global warming, 149
 Globular proteins, 1106
 D-Glucaric acid, 1049
 Glucitol, 1047
 Glucopyranose
 α -D-glucopyranose, 1038, 1048f
 β -D-glucopyranose, 1038, 1062
 Glucosamine, 862
 D-glucosamine, 1061
 Glucose, 156, 812–813
 acyclic aldehyde, 1038–139, 1038f
 anomers of, 1038, 1038f, 1040, 1041f
 in complex carbohydrates, 105
 conversion to ester, 1046
 conversion to ether, 1046
 diastereomers of, 1038, 1038f
 drawing as cyclic hemiacetal, 1037–139, 1038f
 energy in, 1028
 Fischer projection of, 1032, 1033f
 Fischer proof of structure of, 1053–1056, 1054f
 D-glucose, 1029, 1032, 1033, 1033f, 1035, 1035f, 1037–1038, 1041f, 1047, 1049–1051, 1053, 1071, 1167
 α -D-glucose, 813, 1038, 1038f, 1043–1044, 1046, 1057, 1060
 β -D-glucose, 812–813, 823, 1029f, 1038, 1038f, 1041, 1043–1044, 1046, 1057, 1060
 L-glucose, 1033
 glycoside formation from, 1043–1044
 in honey, 1042
 from hydrolysis
 of cellulose, 1060
 of glycosides, 1044
 of maltose, 1057
 of starch, 160, 1061
 in lactose, 219, 220f
 oxidation of, 196, 206, 1048–1049
 reduction of carbonyl group, 1047
 structure, 104f, 161f
 synthesis and metabolism of, 1028
 three-dimensional representations for
 D-glucose, 1041, 1041f
 Wohl degradation, 1050
 Glucosidase
 α -glucosidase, 1061
 β -glucosidase, 1060
 Glutamic acid, 720
 abbreviation for, 711t
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
 Glutamine
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
 Glutathione, 1090
 Glutaric acid, 683
 Glyceraldehyde
 enantiomers of, 1030–1031, 1033
 D-glyceraldehyde, 1029, 1033, 1034
 L-glyceraldehyde, 1033
 (R)-glyceraldehyde, 1030, 1033
 (S)-glyceraldehyde, 1030, 1033
 optical activity, 183
 Glycerol
 naming, 316
 synthesis in lipid hydrolysis, 853, 854–855
 in triacylglycerols, 367
 Glycine, 80, 710–711
 abbreviation for, 711t
 in dipeptide synthesis, 1094–1098
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1075f, 1076f
 Glycogen, 1061
 Glycolic acid, 66
 copolymerization with lactic acid, 1161
 Glycols, 315
 formation by dihydroxylation, 442
 Glycopeptide transpeptidase, 857
 Glycoside(s)
 formation, 1043–1044
 α glycoside, 1043–1044
 β glycoside, 1043–1044
 hydrolysis of, 1044–1045
 naturally occurring, 1045
 N-Glycoside(s), 1062–1065, 1065f
 α -N-glycoside, 1062
 β -N-glycoside, 1062
 Glycosidic bond/linkage
 in disaccharides, 1056–1058
 α -glycosidic linkage, 1058, 1060–1061
 1→4- α -glycosidic bond, 1057, 1060–1061
 1→6- α -glycosidic bond, 1060–1061
 β -glycosidic linkage, 1063
 1→4- β -glycosidic bond, 1057, 1059, 1062
 in polysaccharides, 1059–1061
 Goodyear, Charles, 1159
 Grandisol, 1134
 Graphite, 632
 Green chemistry
 Amberlyst A-26 resin and, 450–451
 definition of, 450
 green polymer synthesis, 946, 1166–1169, 1167f, 1168f
 Grignard products, retrosynthetic analysis of, 746–748
 Grignard reagents, 739–740
 addition to nitriles, 866
 organic synthesis with, 759–761
 preparation of, 740
 reaction as base, 741
 reaction with α,β -unsaturated carbonyl compounds, 757
 reaction with aldehydes and ketones, 742–745, 789
 reaction with carbon dioxide, 753–754
 reaction with epoxides, 754–755
 reaction with esters and acid chlorides, 750–753
 Grignard, Victor, 739
 Ground state
 of atom, 32
 of electron, 597–598
 Group, in periodic table, 8
 Group number, 8, 10
 Grubbs catalyst, 1015–1019
 Grubbs, Robert, 1015
 Guanine, 1064, 1065f
 D-Gulose, 1035f, 1069
 Gutta-percha, 1159, 1160, 1175
 Gypsy moth, 441
 Half-headed curved arrow, 201, 202t, 539
 Halides. *See also* Alkyl halides; Aryl halides; Hydrogen halides; Vinyl halides
 allylic, 229, 229f
 methylation
 coupling reaction with organocuprates, 1003
 S_N2 substitution reactions, 248, 262
 organic
 coupling reaction with organocuprate reagents, 1003–1005
 coupling with alkenes (Heck reaction), 1003, 1009–1011
 coupling with organoboron reagents (Suzuki reaction), 1003, 1005–1009
 unreactive, 650
 α -Halo aldehyde or ketone, synthesis of, 892–895
 Halo alkane, 230. *See also* Alkyl halides
 α -Halo carbonyl compounds
 reactions of, 896
 synthesis of, 892–895
 α -Halo carboxylic acids, amino acid synthesis from, 1078–1079, 1081f
 Halocyclohexanes, E2 elimination in, 296–298
 Haloform, 894–895, 895f
 Haloform reaction, 894–895, 895f
 Halogenation
 of alkanes, 541–551
 of achiral starting material, 549–550, 549t
 bromination, 546–547, 547f
 of chiral starting material, 549t, 550
 chlorination, 544–548, 544f, 545f, 548f, 549–551
 energy changes during, 544, 544f, 545f
 mechanism of, 542–544
 in organic synthesis, 548–549
 stereochemistry of, 549–551, 549t
 of alkenes, 371f, 379–383, 382f
 of alkyl benzenes, 669–671
 of alkynes, 409
 of benzene, 643f, 645, 668
 of carbonyl compounds, at α carbon, 892–895, 895f
 in acid, 892–893, 895f
 in base, 892, 893–895, 895f
 electrophilic aromatic substitution reaction, 643f, 644–645, 646f, 665, 666–667
 haloform reaction, 894–895, 895f
 monohalogenation, 542
 polyhalogenation, 665
 stereochemistry of, 381–383
 transition states in, 546–548, 547f, 548
 Halogens
 in alkyl halides. *See* Alkyl halides
 as electron-withdrawing groups, 655–656, 661
 polarizable, 380
 Halo group, 85t
 Halohydrin
 epoxide synthesis from, 323
 synthesis from alkenes, 371f, 383–385, 385f, 385t
 mechanism of, 383–384, 385t
 regioselectivity of, 384–385, 385t
 stereochemistry of, 384–385, 385t
 use in organic synthesis, 385
 Halomethanes, electrostatic potential map of, 234f
 Halonium ion, bridged, 380–382, 383–384, 385t, 409
 Halothane, 49, 233f, 320
 Hammond postulate, 258–261, 293, 374–375, 374f, 408, 546–547
 electrophilic aromatic substitution reaction, 659
 features of, 258–259
 S_N1 substitution reactions, 258, 260–261
 Handedness, 160, 163, 166
 Hansen's disease (leprosy), 169, 990
 Hardener, 1163–1164, 1164f
 Hardening, 432, 1124
 Haworth projections, 1038, 1039–1041
 HCFCs (hydrochlorofluorocarbons), 552
 HDL particles (high density lipoproteins), 862
 HDPE (high-density polyethylene), 561, 1152, 1169t, 1170
 Heat of hydrogenation, 429–430, 583–584
 benzene, 616–617, 616f
 Heat of reaction, 203. *See also* Enthalpy change
 Heck reaction, 1003, 1009–1011
 Helicene, 614–615
 Helminthosporal, 947
 Heme, 1109, 1110
 Hemiacetal(s)
 in carbohydrates, 812–813
 conversion to acetals, 805–806
 conversion to carbonyl compounds, 808
 cyclic
 conversion to cyclic acetals, 811–812
 structure, 809
 synthesis of, 810–811
 monosaccharides, 1036–1046
 structure of, 809
 synthesis of, 805–806, 807
 Hemibrevetoxin B, 84
 Hemoglobin, 1075
 sickle cell, 1110
 structure, 1105, 1109–1110, 1109f
 Heptalene, 635
 Heptanal, 923
 Heptane
 constitutional isomers of, 118t
 molecular formula, 118t
 2,6-Heptanedione, 926
 2-Heptanone, synthesis of, 905
trans-2-Heptene, 1004
 Herbicides, 646, 646f
 Heroin, 192, 490, 845
 Heteroatom(s), 4
 in condensed structures, 27–28
 definition of, 82
 in functional groups, 82, 102–103
 in skeletal structures, 29, 30
 stereogenic centers, 167
 Heterocycle(s), 317
 aromatic, 621–624, 623f
 nitrogen, 953
 Heterocyclic aromatic amines, basicity of, 972, 973t, 974t
 Heterogeneous reaction mixture, 429
 Heterolysis (heterolytic cleavage)
 carbocation formation, 252, 260
 description, 200
 in E1 elimination reactions, 291
 reactive intermediates of, 201–202, 201f
 in two-step reaction mechanism, 213
 Hexachloroethane, 187
 2,4-Hexadiene, 580, 619
 (2Z,4Z)-2,4-Hexadienoic acid, 1167
 1,3-Hexadiyne, 402f
 Hexamethylenediamine, 859, 875
 Hexamethylphosphoramide (HMPA), 241f
 Hexane, 113
 constitutional isomers of, 118t
 infrared (IR) spectrum, 481
 mass spectrum of, 466, 466f, 469–470, 470f

- molecular formula, 118t
skeletal structures, 29
as solvent, 94
solvent in extraction procedure, 707
- 2,5-Hexanedione, aldol reactions of, 926–927
- Hexanoic acid, 688, 717
- 2-Hexanol, conversion to other compounds, 759
- 3-Hexanone, 425
- 1,3,5-Hexatriene, 617, 619
- 1-Hexene, infrared (IR) spectrum of, 481
- 3-Hexene
cis-3-hexene, 363, 396, 1015
trans-3-hexene, 363, 396
- 1-Hexyne, 1009
infrared (IR) spectrum, 481
- HFCs (hydrofluorocarbons), 552
- High density lipoproteins (HDLs), 862
- High-density polyethylene (HDPE), 561, 1152, 1158, 1169t, 1170
- Highest occupied molecular orbital (HOMO), 629
- High-performance liquid chromatography (HPLC), 1091
- Hinckley, Robert, 320
- Histamine, 622–623, 957–958, 995
- Histidine
isoelectric point for, 1078t
 pK_a values for, 1078t
structure of, 1076f
- Histrionicotoxin, 106, 404, 404f
- HMPA (hexamethylphosphoramide), 241f
- ¹H NMR. *See* Proton NMR
- Hoffman, Felix, 696
- Hofmann elimination, 977–980
- Homogeneous catalysts, 1159
- Homologous series, 117, 118t
- Homolysis (homolytic cleavage)
of allylic C–H bond, 552–553
bond dissociation energy for, 203
chlorination of alkanes, 543, 545
endothermic nature of, 203
entropy increase with, 209
- Honey, 1042
- Honeycomb, 150
- Hooke's law, 478, 479f
- Hormones. *See also specific hormones*
local mediators distinct from, 1130
peptide, 1089, 1090f
sex, 1141–1142, 1141t
- Host–guest complex, 320
- Housane, 128f
- 5-HPETE (5-hydroperoxyeicosatetraenoic acid), 348, 568, 573
- Hückel, Erich, 618
- Hückel's rule, 617–620
basis of, 626–629
- Humulene, 1008, 1008f, 1145
- Hybridization, 32–40, 609–610
conjugated systems and, 578–579
definition of, 33
- Hybridization effects
on acidity, 67–68, 68f, 69f
on basicity of amines, 972–973, 973t
- Hybrid orbitals, 32–40
percent *s*-character, 41–42, 68, 284
sp orbitals, 34–35, 34f, 38–39, 40f
in alkynes, 400, 401, 405
percent *s*-character, 68
*sp*² orbitals, 34–35, 34f, 37, 39f, 40f
in alkenes, 281, 283–284
in aromatic heterocycles, 621–623, 623f
carbon radical, 539
carboxy group, 689
percent *s*-character, 68
*sp*³ orbitals, 33–34, 33f, 35, 35f, 36, 36f, 39f, 40f
in acyclic alkanes, 114
in alkenes, 283–284
percent *s*-character, 68
superscripts, meaning of, 34
- Hydrates
oxidation of, 449
synthesis from aldehydes and ketones, 802–804
kinetics, 803–804
thermodynamics, 802–803
- Hydration
of aldehydes and ketones, 802–804
kinetics, 803–804
thermodynamics, 802–803
of alkenes, 371f, 378–379, 390
definition of, 791
hydroboration–oxidation compared, 390
- Hydrazine, in Wolff–Kishner reduction, 672
- Hydrides
as base, 70, 71f
metal hydride reagents
reducing agents, 428, 727–731, 733–738, 738t
reduction of aldehydes and ketones, 727–729
reduction of nitriles, 865
nucleophilic addition with, 724, 728
nucleophilic substitution with, 725
reduction of ketones and aldehydes, 728–729, 730f, 738t, 789
- 1,2-Hydride shift, 328–330, 651
- Hydroboration
of alkenes, 386–387, 387f, 388f
of alkynes, 1007, 1009
definition of, 386
mechanism of, 386–387
regioselectivity of, 387
- Hydroboration–oxidation reactions
of alkenes, 371f, 385–390
mechanism of, 385–387
regioselectivity of, 387, 389t
stereochemistry of, 388–389, 389t
hydration compared, 390
synthesis of artemisinin, 389, 389f
- Hydrocarbons
aliphatic, 83, 114
aromatic, 83
definition of, 83
functional groups in, 83–84, 84t
infrared (IR) absorptions in, 480
oxidation and reduction of, 427, 428f
polycyclic aromatic (PAH), 349
saturated, 114
synthesis in organocuprate couplings, 1004–1005
unsaturated, 360. *See also* Alkenes
- Hydrochloric acid, 70
elimination in nylon synthesis, 859
in epoxide ring opening reactions, 345, 346
as Lewis acid, 74
for removal of Boc protecting group, 1096
 pK_a of, 62
- Hydrochloride salts, 968
- Hydrochlorofluorocarbons (HCFCs), 552
- Hydrofluorocarbons (HFCs), 552
- Hydrogen
covalent bond formation by, 11–12
isotopes of, 7, 472t
mass of, 472t
octet rule exceptions, 17
primary (1°), 116–117
as reducing agent, 428. *See also* Hydrogenation
secondary (2°), 116–117
sigma (σ) bond in molecular hydrogen, 32
tertiary (3°), 116–117
- Hydrogenation
of aldehydes and ketones, 729
of alkenes, 429–434
alkene stability and, 429–430, 430f
cis and *trans* isomers compared, 429–430
degrees of unsaturation, determination of, 431–432
mechanism of catalytic hydrogenation, 430–431
alkene stability and, 429–430, 430f
catalytic, 429–431
of conjugated dienes, 583–584
enantioselective synthesis of amino acids, 1085–1086
as exothermic reactions, 429
heat of, 429–430, 583–584, 616–617, 616f
of oils, 432–434, 433f
of unsaturated fatty acids, 1124
- Hydrogen bonding
in alcohols, 318
in amides, 834, 834f
in amines, 954
anion solvation by, 240
in carboxylic acids, 692, 693f
in DNA, 104f, 105, 1064, 1065f
intramolecular in β -dicarbonyl compounds, 882
in polar protic solvents, 240, 265
solubility and, 94, 95
strength of, 89, 90t
in water, 89
- Hydrogen bromide, for removal of Boc protecting group, 1096
- Hydrogen chloride. *See* Hydrochloric acid
- Hydrogen cyanide, 792
- Hydrogen fluoride, 12
- Hydrogen halides
addition to alkenes, 371–378, 371f, 372f, 373f, 378t
addition to alkynes, 406–408, 406f
alcohol conversion to alkyl halides with, 331–335
reactivity of, 333
- Hydrogen ions (H⁺), Brønsted–Lowry acid definition and, 55
- Hydrogenolysis
of benzyl esters, 1097
in peptide synthesis, 1097–1098
- Hydrogen peroxide, as oxidizing agent, 438
- Hydrohalogenation
of alkenes, 371–378
Markovnikov's rule, 374–375, 374f, 378t
mechanism of, 371–373, 372f, 373f, 378t
stereochemistry of, 376–377, 377f, 378t
- of alkynes, 406–408, 406f
Markovnikov's rule, 407, 408
- Hydrolysis
of acetals, 807–808
in acetoacetic ester synthesis, 904, 905
of amides, 855–856
in amino acid synthesis, 1079
base-promoted, 852
definition of, 791
of esters, 850–855
acid-catalyzed, 851
applications of, 853–855
base-promoted, 851–852
glucose formation by, 1044, 1057, 1060, 1061
of imines and enamines, 801
of lipids, 853–855
in malonic ester synthesis, 901
of nitriles, 863–864
partial, of peptides, 1092–1093
of starch, 319, 320f
of triacylglycerols, 1124
- Hydronium ion, formal charge on, 16
- Hydroperoxide, 1124
- 5-Hydroperoxyeicosatetraenoic acid (5-HPETE), 348, 573
- Hydrophilic, 95, 97
- Hydrophobic, 95, 97, 150, 1121
- Hydroxide
nucleophile in alcohol synthesis, 322
oxygen bases, 70, 71f
- α -Hydroxy acids, 66
- Hydroxy aldehyde
cyclization of, 810, 813
 β -Hydroxy, synthesis in aldol reactions, 917–928
intramolecular cyclization of, 810, 813
- o*-hydroxybenzaldehyde, 946
- Hydroxybenzene. *See* Phenol
- o*-Hydroxybenzoic acid, 720
- p*-Hydroxybenzoic acid, 720
- 3-Hydroxybutanal, 917
- 4-Hydroxybutanal, 810
- 2-Hydroxybutanedioic acid, 720
- 4-Hydroxybutanoic acid, 696
- 3-Hydroxy-2-butanone, addition to margarine, 432
- 3-Hydroxybutyric acid, 1170–1171
- γ -Hydroxybutyric acid (GHB), 696
- β -Hydroxy carbonyl compounds
synthesis of, in aldol reactions, 917–928
useful transformations of, 924, 924f
- 3-Hydroxy carboxylic acids, 1170–1171
- Hydroxy group, 85t
of alcohols, 313, 324
of enols, 313
of ethanol, 82–83
hydrophilic nature of, 95
IR absorption, 482, 693, 694f
of monosaccharides, 1028, 1046–1047
NMR absorption, 694, 695f
OH proton, NMR spectra of, 517–518, 518f
of phenols, 313
- 6-Hydroxy-2-hexanone, 808
- β -Hydroxy ketone, synthesis of, 919
- Hydroxylamine, 1050
- 5-Hydroxypentanal, 810, 813
- 5-Hydroxy-2-pentanone, 748, 749f, 750
- N*-Hydroxysuccinimide, 1117
- 3-Hydroxyvaleric acid, 1170–1171
- Hyperconjugation, 257–258
- Hypertension, 85
- Hypophosphorous acid, 983
- Ibuprofen
anti-inflammatory action of, 150, 916
enantiomers, 187, 909
mode of action, 1131–1132
structure of, 78, 106
synthesis of, 673, 730f, 911, 916, 932
systematic name of, 119
- ic acid (suffix), 691, 831, 832, 860
- ide (suffix), 231
- D*-Idose, 1035f
- Illudin-S, 905
- Imatinib mesylate, 837
- Imidazole, 749, 750, 870
- Imides, in Gabriel synthesis of primary amines, 961
- Imidic acid tautomer, 863–864
- Imines
hydrolysis of, 801
reduction to amines, 963
synthesis of
from aldehydes and ketones, 797–799, 799f, 801f, 975, 1080

- Imines (*continued*)
 synthesis of (*continued*)
 from nucleophilic addition reactions, 788, 788f
 reduction of nitriles, 865–866
 Iminium ion, 798, 800–801, 801f
 Immunosuppressants, 837
 Indene, 636
 Indigo, 988
 Inductive effects
 on acidity, 65–66, 66f, 69f
 in aliphatic carboxylic acids, 703–704
 on basicity of amines, 969, 973t
 in carbocation stability, 256–257
 definition of, 65
 electron-donating, 654–656
 electron-withdrawing, 65, 654–656
 on substituted benzenes, 654–657
 -ine (suffix), 230–231
 Influenza, 86
 Infrared (IR) absorptions/spectra
 of aldehydes, 780–781, 781f
 of amines, 955, 955f
 atom mass and, 478–479, 479f
 of benzene derivatives, 613
 bond strength and, 478–479, 479f, 480
 of carboxylic acid derivatives, 835, 835t
 of carboxylic acids, 693–694, 694f
 characteristic frequencies, A–10
 functional groups and, 478, 479, 479f, 480t, 481–485
 of hydrocarbons, 481
 important, 479, 480t
 of ketones, 780–781
 of nitriles, 835
 of nitrogen-containing compounds, 483
 of oxygen-containing compounds, 482
 percent *s*-character and, 480
 vibration of bonds and, 476–477, 478–480
 wavenumber scale of, 476
 Infrared (IR) spectroscopy, 464, 474, 476–487
 background on, 476–477
 functional group identification, 476, 477
 IR inactive vibration, 480
 isomers distinguished by, 484
 structure determination in, 485–487
 vibration modes, 476–477
 wavenumber scale, 476
 Infrared (IR) spectrum
 characteristics of, 477–478, 478f
 fingerprint region, 477, 478, 478f, 479, 479f
 functional group region, 477, 478, 478f, 479, 479f, 480t
 Inorganic compounds, 1
 Inscribed polygon method for predicting aromaticity, 629–631, 630f, 631f
 Insecticide, 233, 1012
 Inspira. *See* Eplerenone
 Insulin, 1075, 1105, 1106f
 Integral, NMR spectrum, 507–508
 Intermolecular forces, 87–90
 of aldehydes and ketones, 779, 780t
 boiling point and, 90–92, 91f
 in carboxylic acids, 692, 693f, 693t
 dipole–dipole interactions, 89, 90t
 hydrogen bonding, 89, 90t, 954
 ion–ion interactions, 87, 90t
 melting point and, 92–93
 solubility and, 93–96, 94f, 96f
 van der Waals forces, 87–89, 88f, 90t
 International Union of Pure and Applied Chemistry, 119. *See also* IUPAC nomenclature
 Intramolecular cyclization, of hydroxy aldehydes, 810, 813
 Intramolecular hydrogen bonding, in β -dicarbonyl compounds, 882
 Intramolecular malonic ester synthesis, 902
 Intramolecular reactions
 aldol reactions, 926–928, 928f
 Robinson annulation, 936–939
 Friedel–Crafts reactions, 652–653, 653f
 intramolecular Claisen (Diekmann reaction), 932–933
 Inversion of configuration, 263
 in alkyl tosylate reactions, 339–340
 in S_N2 substitution reaction, 246–247, 247f, 263
 Invirase. *See* Saquinavir
 Iodine, polarizability of, 88–89
 Iodoform, 894–895
 Iodoform test, 894
 Iodomethane, polarizability and boiling point of, 91f
 1-Iodo-1-methylcyclohexane, 289
 Ion(s)
 anions
 acetate, 66–67, 67f
 carbanion, 68, 201–202, 201f
 cyclopentadienyl, 624–625, 630, 630f
 definition of, 7
 enolate, 21
 naked, 241
 radical, 436
 salt formation, 10–11
 solvation by hydrogen bonding, 240
 bond formation by, 202
 cations
 carbocation. *See* Carbocation(s)
 counterions, 55
 cyclopentadienyl, 625
 definition of, 7
 radical, 465
 salt formation, 10–11
 solvation by ion–dipole interactions, 240, 241, 264–265
 spectator ions, 55
 tropylium, 625–626, 630, 630f
 counterions, 55
 polyatomic, 16
 spectator ions, 55
 transport across cell membranes, 100–102, 102f
 Ion–dipole interactions
 cation solvation by, 240, 241, 264–265
 description of, 94, 94f
 Ionic bonding, 10–11
 Ionic compounds
 boiling points of, 91
 formation of, 10–11
 intermolecular forces, 87, 90t
 melting points of, 93
 solubility of, 94, 96f
 Ionic intermediates, 554
 Ionic polymerization, 1153–1156, 1155f
 Ion–ion interactions, 87, 90t
 Ionophores, 100–102, 102f
 IR spectroscopy. *See* Infrared (IR) spectroscopy
 Iso- (prefix), 120
 Isoamyl acetate, 836
 Isobutane, 115
 p-Isobutylbenzaldehyde, 911
 Isobutyl chloride, 231f
 Isobutylene, 1155
 Isobutyl group, 121
 Isocyanate, 1162
 Isoelectric point (pI), for amino acids, 713, 1078, 1078t
 Isoleucine
 isoelectric point for, 1078t
 p*K*_a values for, 1078t
 structure of, 1076f, 1077
 Isomers
 constitutional. *See* Constitutional isomers
 definition of, 17, 162
 infrared (IR) spectroscopy of, 484
 Lewis structures, 17
 ortho and para, 668–669
 resonance structures compared, 19
 stereoisomers. *See* Stereoisomers
 types of, 181, 181f
 Isoniazid, 995
 Isooctane
 combustion of, 148
 oxidation of, 196, 206, 216
 Isopentane, 127
 melting point of, 93
 structure, 116
 Isopentenyl diphosphate, 1134–1136, 1135f, 1138
 Isophorone, 948
 Isoprene, 581, 605, 1145, 1159
 Isoprene unit, 1132–1138
 Isopropyl alcohol, 315
 Isopropylamine, 356, 952, 963
 Isopropylbenzene
 oxidation to benzoic acid, 671
 structure of, 673
 synthesis from benzene, 673
 Isopropylcyclopentane, 761
 Isopropyl group, 120
 3-Isopropyl-4-methylcyclohexanone, 778
 Isopropyl methyl ether, 322
 Isopropyl 2-methylpentanoate, 833
 (*R*)-Isoproterenol, 768
 Isopulegone, 493
 Isotactic polymer, 1157–1158
 Isotopes, 7, 8f
 of bromine, 469
 of carbon, 7, 8f, 472t
 of chlorine, 468
 definition of, 7
 masses of common, 472t
 of nitrogen, 472t
 of oxygen, 472t
 IUPAC nomenclature
 of alcohols, 314–316, 315f
 of aldehydes, 776, 777f
 of alkanes, 121–127, 123f, 126f, 127f
 of alkenes, 362–363
 of alkyl halides, 230–231, 231f
 of alkynes, 401–402
 of benzene derivatives, 610–613
 of carboxylic acids, 690–692
 of cycloalkanes, 125–127, 126f, 127f
 of epoxides, 317–318
 of ethers, 316–317
 of ketones, 777–778, 779f
 Jack-o'-lantern mushroom, 905
 Januvia. *See* Sitagliptin
 cis-Jasmone, 435
 Johnson, W. S., 928
 Joule (J), 40
 Juvenile hormones, 721, 745, 745f, 1004
 mimics of, 745
 Kavain, 364
 Keflex. *See* Cephalexin
 Kekulé, August, 608
 Kekulé structures, 608, 609
 Kelsey, Frances Oldham, 170
 Keratins, 1075
 α -keratins, 1107–1108, 1108f
 Kerosene, composition of hydrocarbons in, 128
 Ketal, 804
 Ketamine, 191
 Ketene, 50
 Keto–enol tautomers, 410, 881–883
 β -Keto esters
 as active methylene compounds, 924
 conversion to ketones, 903–905
 p*K*_a, 886t
 synthesis of
 Claisen reaction, 928–932, 929f
 Michael reaction, 935, 936–937
D-Ketohexoses, 1035, 1036f
 Ketones. *See also* Carbonyl compounds
 aldol reactions of, 917–919, 922–923, 925–926
 boiling point of, 780t
 carbohydrates, 812–813
 in crossed Claisen reactions, 931
 fragmentation pattern in mass spectrum, 471
 functional group in, 86t
 hydration of, 802–804
 kinetics of hydrate formation, 803–804
 thermodynamics of hydrate formation, 802–803
 interesting examples, 783
 iodoform test of, 894
 IR spectra of, 482, 780–781
 melting point of, 780t
 NMR spectra of, 782, 782f
 nomenclature, 777–778, 779
 common names, 778, 779f
 IUPAC system, 777–778, 779f
 nucleophilic addition reactions, 724, 774–813
 acid catalyzed, 787
 carbanion addition, 789–790
 cyanide addition, 790–791
 hydride addition, 789
 mechanism of, 786–787
 nucleophiles in, 787–788, 788f
 primary amine addition, 797–799, 801f
 secondary amine addition, 800–801, 801f
 physical properties, 779, 780t
 p*K*_a, 886t
 protecting groups for, 808–809
 reactions of
 aldol reactions, 917–919, 922–923, 925–926
 with amines, 975
 at α carbon, 785–786
 at carbonyl carbon, 785, 786
 crossed Claisen reactions, 931
 halogenation at α carbon, 892–895, 895f
 hydration, 802–804
 nucleophilic addition reactions, 724, 774–813
 with organometallic reagents, 742–745, 745f, 789–790
 reactivity of, 724
 reduction of, 789–790
 to alcohols, 727, 728, 729
 of aryl ketones to alkyl benzenes, 672–673
 reductive amination to amines, 963–966
 to secondary alcohols, 432
 solubility of, 780t
 structure of, 722, 775
 synthesis of
 acetotacetic ester synthesis, 900, 903–905
 alcohol oxidation, 447, 448, 450
 Friedel–Crafts acylation, 643f, 648, 672–673, 784
 hydration of alkynes, 409–412, 784

- hydroboration–oxidation of alkynes, 412–413
- hydrolysis of acetals, 807–808
- hydrolysis of imines and enamines, 801
- oxidation of secondary alcohols, 784
- oxidative cleavage of alkenes, 444–446, 785
- reactions of acid chlorides with organocuprates, 784
- tautomers, 864
- unreactivity to nucleophilic substitution, 725
- unsymmetrical, alkylation of, 898–899
- Wittig reaction, 792–797
- Ketoprofen, 70, 191
- Ketoses
- D-ketose family, 1035–1036, 1036f
 - structure, 1028–1029
- Keto tautomers, 882–883
- Kevlar, 1161, 1174
- Kiliani–Fischer synthesis, 1049, 1051–1056
- Kinetic enolates, 889–890, 898
- Kinetic product, in electrophilic addition reactions of conjugated dienes, 586–588, 587f, 588f
- Kinetic resolution, of amino acids using enzymes, 1084
- Kinetics
- definition of, 206, 215
 - E1 elimination reactions, 291, 293t
 - E2 elimination reactions, 285
 - first-order, 217
 - rate equations, 216–218
 - second-order, 217
 - S_N1 substitution reactions, 252, 256t
 - S_N2 substitution reactions, 244, 249t
- Lactams, 827
- β-Lactam(s), 311, 827, 838, 856–857
- β-Lactam family of antibiotics, 463, 485
- Lactase, 219, 220f, 1027, 1058
- Lactate dehydrogenase, 733
- Lactic acid, 29, 696, 733
- copolymerization with glycolic acid, 1161
 - optical activity of, 183
 - specific rotation of, 185
- Lactide, 1168
- Lactols, 809, 810
- Lactone(s), 398, 827
- macrocyclic, 837
 - synthesis by esterification of carboxylic acids, 848
- Lactose, 812, 1027, 1057–1058
- breakdown of, 219, 220f
- Lactose intolerance, 1027, 1058
- Laetrile, 792
- Lanolin, 1121
- Lanosterol, 1140, 1140f
- Lard, 369
- Latanoprost, 697
- Latex, 1159, 1160
- Lauric acid, 1122t, 1125
- Lavandulol, 747
- l* (levorotatory) compounds, 183, 1033
- LDA. *See* Lithium diisopropylamide (LDA)
- LDL particles (low density lipoproteins), 862
- L-Dopa (levodopa), 6, 45, 191, 1115
- LDPE (low-density polyethylene), 561, 1152
- Leaving group
- of alcohols, 324, 330–332, 335–336, 338–339
 - of carbonyl compounds, 722, 724, 725–726
 - of carboxylic acid derivatives, 827, 828, 829t, 838–840
 - in E2 elimination reactions, 287, 288t
 - in nucleophilic acyl substitution reactions, 827, 838–840
 - in nucleophilic substitution reactions, 235–236, 264
 - good, 237t
 - periodic trends in ability, 236–237
 - poor, 237t
 - reaction rate and, 264
 - weak bases, 236–238 - in terpene biosynthesis, 1134–1137
- Le Châtelier's principle, 327–328, 1042
- Lecithin, 1126, 1127f
- Length, units used to report wavelength, 474
- Letrozole, 862
- Leucine, 191
- isoelectric point for, 1078t
 - pK_a values for, 1078t
 - structure of, 1076f
- Leukotrienes, 1129–1130, 1130t, 1131f
- biological activity of, 348
 - leukotriene A₄, 348
 - leukotriene C₄, 348, 568
 - synthesis of, 348
- Levonorgestrel, 403
- Lemorotatory (*l*) compounds, 183, 1033
- Lewis acid, 72–74
- alkenes as, 372
 - borane as, 386
 - carbocation as, 372
 - catalyst, 333
- Lewis acid–base reactions, 73–74, 235. *See also* Nucleophilic substitution reactions
- electrophilic addition, 372
 - S_N1 substitution reactions, 252
- Lewis base, 72–74
- nucleophile as, 238
 - tetrahydrofuran (THF) as, 386
- Lewis structures, 12–18
- of acyclic alkanes, 114–115
 - condensed formula conversion to, 28–29
 - drawing of, 12–14
 - formal charge, 15–16, 17t
 - isomers, 17
 - molecular shape determination, 23–27
 - multiple bonds in, 14–15
 - octet rule exceptions, 17–18
 - resonance structures, 18–23
 - skeletal structure conversion to, 29–31
- Lexan, 1149f, 1163, 1168
- Ligands, in palladium-catalyzed reactions, 1005–1006
- Light
- polarized, 182–183
 - speed of, 475
- Limonene, 50, 446, 1137
- (*R*)-limonene, 367f
 - (*S*)-limonene, 366
- Linalool, 747
- Linalyl diphosphate, 1137
- Linamarin, 791–792
- Lindlar catalyst, 435
- Linear molecule, 24, 26f, 44
- alkynes, 400
- Linen, 858, 858f
- Linoleic acid, 459, 1122, 1122t
- as essential fatty acid, 368, 369
 - melting point, 368t
 - structure, 368f, 368t
- Linolenic acid, 1122, 1122t
- as essential fatty acid, 368
 - melting point, 368t
 - as omega-3 fatty acid, 368
 - sources of, 369
 - structure, 368f, 368t
- Lipases, 853
- Lipid bilayer, 100, 101f, 102, 102f, 1126–1127, 1128
- Lipids, 149–151, 1119–1147
- bonding in, 1120
 - definition of, 149
 - eicosanoids, 1129–1132, 1130t, 1131f
 - energy in, 151, 1125
 - examples of, 150, 150f, 1120f
 - fat-soluble vitamins, 1128, 1129f
 - hydrolysis of, 853–855, 1121
 - hydrolyzable, 1121
 - metabolism of, 1125
 - nonhydrolyzable, 1121
 - origin of word, 1120
 - oxidation of, 215, 556–557, 556f
 - phospholipids. *See* Phospholipids
 - solubility of, 149, 1120
 - steroids. *See* Steroids
 - structure of, 150f
 - terpenes, 1132–1138, 1133f, 1133t, 1135f
 - triacylglycerols. *See* Triacylglycerols
 - unsaturated, oxidation of, 556–557, 556f
 - waxes, 1121
- Lipitor. *See* Atorvastatin
- Lipoprotein, 861f, 862
- Lipoxygenase, 348, 1130
- Lithium
- electronegativity value of, 739
 - valence electrons of, 11
- Lithium acetylide, 740–741, 745
- Lithium aluminum hydride (LiAlH₄), 727–730, 733–738, 738t
- as reducing agent, 428
 - reduction of aldehydes and ketones, 789
 - reduction of alkyl halides, 437–438, 438f
 - reduction of a nitrile, 865
 - reduction of epoxides, 437–438, 438f
 - reduction of nitriles to primary amines, 962
- Lithium diisopropylamide (LDA), 898, 925
- enolate formation with, 887–888, 889–890
 - pK_a, 887t
 - preparation of, 888
- Lithium dimethylcuprate, 492, 1014
- Lithium fluoride, 11
- Lithium tri-*tert*-butoxyaluminum hydride, 734, 735, 738t
- Lithium tri-*sec*-butylborohydride, 773
- Living polymerization, 1155
- Local anesthetic, 673
- Local mediators, 1130
- Log, 59, 60
- London forces, 87. *See also* van der Waals forces
- Lone pairs
- in aromatic heterocycles, 621–623
 - Brønsted–Lowry bases and, 55–56, 55f
 - in carbanions, 202
 - in heteroatoms, 82, 103
 - in Lewis structures, 12–15
 - of negatively charged carbon atoms, 31
 - number of, 31
 - in resonance structures, 19, 21, 67, 576
 - second-row elements and, 12
- Long, Dr. Crawford, 320
- Loratadine, 615f
- Lotensin. *See* Benazepril
- Low density lipoproteins (LDLs), 862
- Low-density polyethylene (LDPE), 1152, 1169t
- Lowest unoccupied molecular orbital (LUMO), 629
- LSD, 606, 641
- intramolecular Friedel–Crafts acylation in the synthesis of, 653, 653f
 - structure of, 959
 - synthesis of, 641
- LTC₄, 1130f
- LUMO (lowest unoccupied molecular orbital), 629
- Lycopene, 571, 581, 598, 598f, 1145
- Lycra. *See* Spandex
- Lyrica. *See* Pregabalin
- Lynen, Feodor, 1140
- Lysergic acid diethyl amide. *See* LSD
- Lysine, 720
- abbreviation for, 711t
 - ionizable side chain of, 1077, 1078t
 - isoelectric point for, 1078t
 - pK_a values for, 1078t
 - structure of, 1076f
- Lysozyme, 1104, 1104f
- D-Lyxose, 1034, 1035f, 1050
- Macrocyclic lactone, 837
- Macrolide, 837
- Magnesium, electronegativity value of, 739
- Magnesium monoperoxyphthalate (MMPP), 438f
- Magnetic resonance imaging (MRI), 527, 527f
- Ma huang, 166, 192
- Malaria, 289, 974
- Maleic acid, 112, 592, 603
- Maleic anhydride, 591f
- Malonic acid, 79, 692
- Malonic acid synthesis
- intramolecular, 902
 - retrosynthetic analysis of, 902–903
 - step in, 901–902
- Malonic ester synthesis, 900–903
- Maltose, 824, 1057
- Mandelic acid, 194, 1115
- Manganese, in oxidizing agents, 438, 439
- Mannose
- in Fischer proof, 1053–1056
 - D-mannose, 1035, 1035f, 1051
 - Haworth projections of, 1039–1040
- Manoalide, 1134
- Margarine, 432–433
- Markovnikov's rule, 374–375, 374f, 378t, 407, 408, 584
- in cationic polymerization, 1153
- Mass number, 7
- Mass spectrometer
- features of, 464
 - high-resolution, 472
 - schematic of, 464f
- Mass spectrometry (MS)
- of alkyl halides, 468–469, 469f
 - of amines, 955, 955f
 - analysis of unknowns using molecular ion, 466–468
 - base peak, 465
 - of biomolecules, 474
 - definition of, 464
 - electrospray ionization (ESI), 474
 - features of, 464–466
 - fragmentation in, 465, 469–472, 470f
 - gas chromatography and, 473–474, 473f, 474f
 - high-resolution, 472
 - M peak, 465, 468–469
 - M + 1 peak, 466
 - M + 2 peak, 468–469
- Mass spectrum, 465
- Mass-to-charge (*m/z*) ratio, 464f, 465
- Mauveine, 988–989
- Maytansine, 1025
- mCPBA (*meta*-chloroperoxybenzoic acid), 438f, 439–441

- MDMA (3,4-methylenedioxy-methamphetamine), 998
- Mefloquine, 194
- Melamine, 1177
- Melanin, 599
- Melatonin, 494, 522
- Melmac, 1177
- Melting point
of alcohols, 318t
of aldehydes and ketones, 780t
of alkanes, 129, 130t
of alkenes, 365, 368, 368t
of alkyl halides, 232t
of alkynes, 402
of amines, 954t
of carboxylic acids, 693t
definition of, 92
of enantiomers, 182, 184, 184t
of epoxides, 318t
of ethers, 318t
of fats, 369, 1122–1123
intermolecular forces and, 92–93
of monosaccharides, 1036
of oils, 369, 1122–1123
of racemic mixtures, 184, 184t
of triacylglycerols, 433f
- Membrane proteins, 1075
- Menthol, 49, 108, 156, 1132–1133
- Menthone, 108
- Meperidine, 950
- Mercapto group, 85t
- Mercury, in Clemmensen reduction, 672–673
- Merrifield method, of peptide synthesis, 1099
- Merrifield, R. Bruce, 1099
- Mescaline, 958, 999
- Mesityl oxide, 773
- Meso compounds, 177–179
achiral compounds, 178, 180–181, 382, 382f, 592, 1013
carbene addition and, 1013
disubstituted cycloalkanes, 180–181
from epoxidation of alkenes, 440
plane of symmetry, 178
- Mestranol, 109, 770
- Meta (prefix), 611
- Metabolite, 157
- Meta director, 657, 658, 661, 663–664, 664f, 665, 666, 668–669, 674
- Metal-carbenes, 1017
- Metal catalyst, 218–219, 429
- Metal hydride reagents
reducing agents, 428, 727–731, 733–738, 738t
reduction of aldehydes and ketones, 727–729
reduction of nitriles, 865
- Metals
as catalysts, 218–219
in oxidizing agents, 438–439
- Metathesis, 1015–1019, 1016f, 1018f
derivation of word, 1015
drawing products of, 1016, 1016f
mechanism, 1017
ring-closing (RCM), 1018–1019, 1018f
- Met-enkephalin, 837
- Methamphetamine, 65, 958, 964, 964f
- Methanal. *See* Formaldehyde
- Methanamine. *See* Methylamine
- Methane
bonding in, 32–33, 33f
combustion of, 148
description, 2
halogenation of, 542f
Lewis structure, 13
mass spectrum of, 465–466
molecular formula, 114, 115, 118t
molecular shape of, 25
in natural gas, 128
oxidation of, 427, 427f
 pK_a of, 62, 63
structure of, 114
whole-number mass of, 465
- Methanesulfonic acid, 710, 837
- Methanethiol, 128
- Methanol
ether formation by addition to alkenes, 378, 379
hybrid orbitals in, 35
Lewis structure, 14
oxidation of, 451
as polar protic solvent, 240f
structure of, 319f
toxicity of, 319f
- Methanolone, 1146
- Methionine
abbreviation for, 711t
isoelectric point for, 1078t
 pK_a values for, 1078t
structure of, 1076f
synthesis of, 1081f
- Methoprene, 745
- Methoxide, as Brønsted–Lowry base, 55f, 56
- p*-Methoxybenzoic acid, 706
- Methoxychlor, 234
- Methoxycyclohexane, conformations of, 209
- Methoxy group, 316
- Methoxy methyl ether, 819
- p*-Methoxyphenyl ester, 1117
- Methyl acetate, 226
formation of, 243
NMR spectrum of, 524, 526f
structure of, 826
- Methyl acrylate, 591f
- Methylamine, 952
as Brønsted–Lowry base, 55f, 56
electrostatic potential plot of, 951, 951f
- N*-Methylaniline, NMR spectrum of, 956, 956f
- p*-Methylaniline
basicity of, 970
electrostatic potential plot of, 971f
- Methyl anthranilate, 875
- Methylation, with SAM, 250
- Methylbenzene. *See* Toluene
- Methyl benzoate, synthesis of, 847
- (*R*)- α -Methylbenzylamine, 1082–1084
- 2-Methyl-1,3-butadiene, 581, 1159
- 2-Methylbutane, 116, 127
- 2-Methylbutanenitrile, structure of, 832f
- 2-Methyl-2-butanol, 325
- 2-Methylbutanoyl chloride, structure of, 830
- (*R*)- α -Methylbutyrophenone, 909
- 2-Methyl-CBS-oxazaborolidine, 731
- Methyl chloride, electrostatic potential plot for, 44, 44f. *See also* Chloromethane
- Methyl *trans*-chrysanthemate, 1023
- 7-Methyl-1,3,5-cycloheptatriene, 636
- 3-Methylcycloheptene, 363f
- Methylcyclohexane, 125, 142, 143f
- 2-Methyl-1,3-cyclohexanedione, 936
- 1-Methylcyclohexanol, 352
- 3-Methylcyclohexanol, 315f
cis-3-methylcyclohexanol, 340
trans-4-methylcyclohexanol, 263
- 2-Methylcyclohexanone, 412, 925
alkylation of, 898–899
enolates formed from, 889
- 1-Methylcyclohexene, 760, 1005
- 3-Methylcyclohexene, 375
stereogenic center, 169
- 5-Methyl-1,3-cyclopentadiene, 636
- 2-Methylcyclopentanamine, in Hofmann reaction, 979
- Methylcyclopentane, 169
- 2-Methylcyclopentanecarboxamide, structure of, 831
- (*3R*)-3-Methylcyclopentanone, 790
- 1-Methylcyclopentene, 363f, 928
- Methyl dimethylphosphonate, 537
- Methylene, 1014
 α -Methylene- γ -butyrolactone, 899
- Methylene chloride, 233f
- Methylenecyclohexane, 365, 796
- 3,4-Methylenedioxy-methamphetamine (MDMA), 998
- Methylene group, 117, 365
- Methyl ester, 1097
- 3-Methylfentanyl, 467–468
- Methyl α -D-glucopyranoside, 1044, 1048f
- Methyl group, 121
angular, 1138–1139
as electron-donating group, 657
inductive effect of, 654
as ortho, para director, 657, 662, 666–668
- Methyl halides
coupling reaction with organocuprates, 1003
 S_N2 substitution reactions, 248, 262
- 6-Methyl-6-hepten-2-ol, 363
- 2-Methylhexanoic acid, synthesis of, 903
- 5-Methyl-3-hexanol, 315
- 5-Methyl-4-hexen-1-yne, 402f
- Methyl 5-hydroxyhexanoate, 808
- N*-Methylisopropylamine, 952
- Methyl ketones, 410
halogenation of, 894–895, 895f
synthesis in acetoacetic ester synthesis, 900, 903–905
- Methyl lithium, 740
- Methylmagnesium bromide, 740
- Methylmagnesium chloride, 748
- Methyl 2-methylbutanoate, 836
- N*-Methylmorpholine *N*-oxide (NMO), 444
- 4-Methyl-2-nonene, 395
- Methyl orange, 987, 989–990
- 3-Methylpentanal, 777f
- 4-Methyl-1-pentanamine, 953
- 2-Methylpentane, 163f
- 3-Methylpentane, 163f
- 4-Methyl-1,4-pentanediol, 748, 749f, 750
- 2-Methyl-3-pentanone, 779f
- 3-Methyl-2-pentanone, 778
- 3-Methyl-2-pentene, 364
(*E*)-3-methyl-2-pentene, 364
(*Z*)-3-methyl-2-pentene, 364
- 4-Methyl-3-penten-2-one, 779
- Methylphenidate, 191
- 1-Methyl-4-phenyl-4-propionoxy-piperidine, 467–468
- N*-Methyl-2-propanamine, 952
- 2-Methyl-1-propanol, 269
- 2-Methylpropene, 396, 686, 1014, 1155
- 2-Methylpropene oxide, 1175
- 2-[4-(2-Methylpropyl)phenyl]propanoic acid, 119
- 4-Methylpyridine, 948
- Methyltriphenylphosphonium bromide, 794
- β -Methylvaleraldehyde, 777f
- Methyl vinyl ketone, 591f, 936
- Metoprolol, 78
- Mevalonic acid, 157
- Micelles, 99
- Michael acceptor, 934
- Michael reaction, 934–936, 935f
- Mifepristone, 403
- Mineral oil, 158
- Mirror images, 160, 163–165, 177
- Miscibility, 95
- Misoprostol, 1131
- Mixed aldol reaction. *See* Crossed aldol reactions
- MMPP (magnesium monoperoxyphthalate), 438f
- Molecular formula
of acyclic alkanes, 114, 118, 118t
of alkenes, 360
of alkyl halides, 229
of alkynes, 400
of amines, 955
of butane, 118t
of carbohydrates, 1028
of cycloalkanes, 114, 118
of decane, 118t
of eicosane, 118t
of ethane, 114, 115, 118t
of heptane, 118t
of hexane, 118t
of methane, 114, 115, 118t
of nonane, 118t
of octane, 118t
of pentane, 118t
of propane, 114, 115, 118t
- Molecular ion, 465, 466–468
- Molecular orbitals (MOs), 627–629, 628f
antibonding, 627, 628f, 629–630
bonding, 627, 628f, 629–631, 630f, 631f
degenerate, 629
highest occupied MO (HOMO), 629
lowest unoccupied MO (LUMO), 629
patterns for cyclic, completely conjugated systems, 631f
prediction/determination of relative energies of, 629–631, 630f, 631f
- Molecular orbital (MO) theory, 627
- Molecular recognition, 320
- Molecular shape
bent, 26, 44
determining
bond angle, 24–27, 26f
bond length, 23, 23t
four groups around an atom, 25–27
linear, 24, 26f, 44
tetrahedral, 25–26, 26f
three groups around an atom, 24–25
trigonal planar, 24–25, 26f
trigonal pyramid, 26
two groups around an atom, 24
- Molecules
definition of, 11
polarity of, 44, 45f
- Molina, Mario, 551
- Molozonide, 445
- Molybdenum, 1015
- Monensin, 812
- Monohalogenation, 542
- Monomers, 1149
- Monosaccharides, 1028–1056
D-aldoses, family of, 1034–1035, 1035f
common names of, 1029, 1034
constitutional isomers of, 1029
conversion to esters, 1046–1047
conversion to ethers, 1046–1047
cyclic forms, 1036–1042
anomers, 1037–1042, 1041, 1041f
chair form, 1040
drawing glucose as a cyclic hemiacetal, 1037–1038, 1038f
furanoses, 1041–1042
Haworth projections, 1038, 1039–1041
three-dimensional representations for D-glucose, 1041, 1041f
determining structure of unknown, 1052–1053

- examples of, 1029
Fischer projection formulas of, 1030–1033, 1033f
Fischer proof of glucose structure, 1053–1056, 1054f
glycosides
 formation, 1043–1044
 hydrolysis, 1044–1045
 naturally occurring, 1045
isomers (D and L) of, 1033–1034
D-ketoses, family of, 1035–1036, 1036f
melting point of, 1036
physical properties of, 1036
reactions at carbonyl group, 1047–1053
 addition or removal of one carbon, 1049–1053
 Kiliani–Fischer synthesis, 1049, 1051–1056
 oxidation, 1047–1049, 1048f
 reduction, 1047
 Wohl degradation, 1049, 1050
reactions at hydroxy groups, 1046–1047
solubility of, 1036, 1047
stereogenic centers in, 1030–1034
 D and L designations, 1033–1034
 multiple, 1032
 structure of, 1028
 sweetness of, 1036
Monosodium glutamate (MSG), specific rotation of, 185
Morphine, 55–56, 597, 845, 957
Morton, Dr. William, 320
Motrin. *See* Ibuprofen
M peak, in mass spectrum, 465, 468–469
M + 1 peak, in mass spectrum, 466
M + 2 peak, in mass spectrum, 468–469
MPPP, 467–468
MRI (magnetic resonance imaging), 527, 527f
MS. *See* Mass spectrometry (MS)
MSG (monosodium glutamate), specific rotation of, 185
MTBE. *See* *tert*-Butyl methyl ether (MTBE)
Multiple bond
 in condensed structures, 27–28
 Lewis structures, 14–15
 position of, 22
Multiplet, NMR spectrum, 511, 511t, 514, 514f
 α -Multistriatin, 454, 454f
Muscalure, 276, 460
Muscone, 730f
Mushroom
 jack-o'-lantern, 905
 Psilocybe, 959
Musk deer, 730
Mutarotation, 1039, 1044, 1057
Mycomycin, 195
Myelin sheath, 1128
Myoglobin, 1074, 1109–1110, 1109f
Myrcene, 49, 1132–1133
Myristic acid, 1122t

Nabumetone, 906, 941
NADH (nicotinamide adenine dinucleotide), 732–733
NAG (*N*-acetyl-D-glucosamine), 862, 1061–1062
Nandrolone, 1142
Naphthalene, 614, 621, 683
1-Naphthol, 356
Naprosyn. *See* Naproxen
Naproxen, 76, 159, 187, 898
Natta, Giulio, 1158
Natural gas, 128
Natural product, 268

NBS (*N*-bromosuccinimide), 383–384
Nelfinavir, 615f
Neopentane
 melting point of, 93
 structure, 116
 van der Waals forces, 88, 88f
Neral, 779, 1145
Nerol, 398
Neryl diphosphate, 1137, 1145
Neurotoxin, 84
Neurotransmitter
 acetylcholine, 861
 dopamine, 958, 959f
 serotonin, 958
Neutrons, 7
Newman projection, 131–132, 132f, 133f
Niacin, 98
Nickel, as catalyst, 219
Nicotinamide adenine dinucleotide (NADH), 732–733
Nicotine, 52, 252, 275, 957, 957f, 973
Niphatoxin B, 416, 416f
Nitration
 of benzene, 643f, 646–647, 669
 electrophilic aromatic substitution reactions, 643f, 646–647, 667–668
Nitriles, 862–866
 acidity of, 885–886
 addition of organometallic reagents, 863, 866
 dipole–dipole interactions of, 834
 hydrolysis of, 863–864, 1081
 infrared (IR) spectra of, 483, 835
 NMR spectra of, 836
 nomenclature, 832, 832f, 833t
 physical properties, 834
 p*K*_a of, 886, 886t
 reduction of, 863, 865–866
 reduction to primary amines, 962
 structure and bonding of, 827, 829, 862–863
 synthesis of, 862
Nitrite preservatives, 261
p-Nitroaniline
 basicity of, 970
 electrostatic potential plot of, 971f
Nitrobenzene, 643f, 646, 647, 657, 663
 reduction to aniline, 673
p-Nitrobenzoic acid, 707
Nitrogen
 in aromatic heterocycles, 621–624, 623f
 bonding patterns and formal charge, 17t
 infrared (IR) absorptions in nitrogen-containing compounds, 483
 isotopes, 472t
 mass of, 472t
Nitrogen heterocycles, 953
Nitrogen–nitrogen double bond, 986
Nitrogen rule, 467
Nitro group
 as electron-withdrawing group, 657
 in Friedel–Crafts reactions, 665
 as meta director, 657, 663–664, 666, 668, 674
 reduction of, 673–674, 962
Nitromethane, 78
Nitronium ion, 646
o-Nitrophenol, 611
Nitrosamines
 in foods, 261
 formation of, 261
 N-nitrosamine from primary amines, 981
 N-nitrosamine from secondary amines, 980, 981–982
N-Nitrosodimethylamine, 261

Nitrosonium ion, 980–982
N-Nitrosopyrrolidine, 261
o-Nitrotoluene, synthesis from benzene, 669
p-Nitrotoluene, 666
Nitrous acid, reaction of amines with, 980–982
NMO (*N*-methylmorpholine *N*-oxide), 444
NMR. *See* Nuclear magnetic resonance (NMR) spectroscopy
NMR spectrometers, 496, 496f
Noble gases, stability of, 10
Node of electron density, 8–9
Nomenclature, A–3–6. *See also* Common names; IUPAC nomenclature
 of acid chlorides, 830, 833, 833t
 of alcohols, 314–316, 315f
 of aldehydes, 776–777, 777f, 779
 of alkanes, 119–128
 common names, 127–128, 128f
 IUPAC system, 121–127, 123f
 substituents, 120–121
 of alkenes, 362–365
 of alkyl halides, 230–231, 231f
 of alkyl substituents that contain branching, A–3–4
 of alkynes, 401–402
 of amides, 831–832, 833t
 of amines, 952–954
 of anhydrides, 830, 833t
 of bicyclic compounds, A–5–6
 of carboxylic acids, 690–692, 691t
 E,Z system, 364
 of epoxides, 317–318
 of esters, 830–831, 833, 833t
 of ethers, 316–317
 generic name, 119
 of ketones, 777–778, 779, 779f
 of monosaccharides, 1028–1029
 of nitriles, 33t, 831–832, 832, 832f, 833t
 of polyfunctional compounds, A–4–5
 systematic name, 119
 trade name, 119
Nomex, 1174
Nonactin, 101, 102
Nonane
 constitutional isomers of, 118t
 molecular formula, 118t
Nonbonded (unshared) electrons, 12
Nonnucleophilic bases, 239
Nonpolar bond, 43
Nonpolar molecule
 description of, 44, 45f
 solubility of, 94
Nonreducing sugars, 1048, 1048f, 1058
Nonsteroidal anti-inflammatory drugs (NSAIDs), 1131–1132
Noradrenaline (norepinephrine)
 structure of, 958
 synthesis of, 251, 251f
Norethindrone, 97, 192, 402–403, 403f, 1142
Novadex. *See* Tamoxifen
Novocain. *See* Procaine
Noyori, Ryoji, 1085
NSAIDs (nonsteroidal anti-inflammatory drugs), 150, 1131–1132
Nuclear magnetic resonance (NMR) spectroscopy, 464, 474, 494–537
 basis of, 495–497
 ¹³C (carbon), 495, 522–526. *See also* Carbon NMR
 ¹H (proton), 495, 497–522. *See also* Proton NMR
 medical uses of, 527, 527f
 nuclear spin in, 495–497
 reference signal, 497, 523

scale, 497
signal *versus* peak, 509
spectrometers, 496, 496f
variables in, 496–497
Nucleophiles, 85, 238–242
 acetylide anions as, 406
 addition to carbonyl groups, 787–788, 788f
 alkyl halide substitution reactions with, 234
 ambident, 888
 attack of electron-deficient atoms by, 238
 biological, 261
 carbanions as, 201, 202
 common, table of, 242t
 enolates as, 888–889, 892
 hydroxide, in alcohol synthesis, 322
 Lewis base, 73
 meaning of term, 73
 negatively charged, 242t
 neutral, 242t
 opening of epoxide rings, 343–345
 organometallic reagents as, 739, 741–742
 pi (π) bonds and, 82, 103
 reactivity of, 102–103
 strength, 239–242
 strong, 263
 in substitution reactions, 235–236. *See also* Nucleophilic substitution reactions
 weak, 263
Nucleophile strength. *See* Nucleophilicity
Nucleophilic acyl substitution reactions, 725, 838–858, A–13
 biological, 860–862, 861f
 general reaction of, 881
 leaving groups in, 827
 mechanism of, 838–839
 polyester synthesis, 1161–1162
 reactions of acid chlorides, 842–843, 858t
 reactions of amides, 855–857, 858t
 reactions of anhydrides, 844–845, 858t
 reactions of carboxylic acids, 845–850, 846f, 858t
 conversion to acid chlorides, 845–847, 846f
 conversion to amides, 846f, 849–850
 conversion to cyclic anhydrides, 846f, 847
 conversion to esters, 846f, 847–849
 reactions of esters, 850–855, 855f, 858t
 reactivity of carboxylic acids and derivatives in, 839–841
 summary of reactions, 857–858, 858t
Nucleophilic addition reactions
 aldehydes and ketones, 724, A–14
 aldol reactions. *See* Aldol reactions
 of carbonyl compounds, 881
 Michael reaction, 934–935
 of nitriles, 863
 organometallic reagents with aldehydes and ketones, 742–745, 745f
 polyurethane synthesis, 1162
 α,β -unsaturated carbonyl compounds, 755–758
Nucleophilicity
 basicity compared to, 239, 240, 241
 periodic trends in, 239, 240, 241
 S_N1 *versus* S_N2 substitution reactions, 262–264
 solvent effects, 240–242
 polar aprotic solvents, 241, 241f
 polar protic solvents, 240, 240f
 steric effects and, 239

- Nucleophilic substitution reactions, A-13
 acetylide anion reactions with alkyl halides, 414-415
 adrenaline synthesis, 228, 251, 251f
 of alcohols, 323-324, 331-338
 alcohol synthesis by, 321-322
 of alkyl halides, 234-269
 amine formation by, 960-962
 bimolecular, 243-244
 of carbonyl compounds, 724, 725-726, 881
 Claisen reaction, 929-932, 929f
 components of, 235
 crown ether use in, 321, 321f
 epoxide ring opening reactions, 343-347
 equilibrium and, 238
 ethers with strong acids, 341-343
 ether synthesis by, 321-322
 features of, 235-236
 first-order rate equation, 243, 244
 leaving groups in, 235-238, 237t, 264
 mechanisms, 242-269
 distinguishing between S_N1 and S_N2 , 262-266, 262f, 265t
 one-step, 243, 244-245, 249t
 order of bond breaking and bond making, 242-243, 252
 S_N1 , 252-261, 253f, 255f, 256f, 260f
 S_N2 , 244-252, 245f, 247f, 248f, 249f, 249t, 250f, 251f, 268t
 two-step, 243, 244, 252-253, 256t
 nucleophile strength (nucleophilicity), 239-242, 262-264
 organic synthesis by, 267-269, 268t, 272f
 in polycarbonate synthesis, 1163
 rate-determining step in, 243, 244
 second-order rate equation, 243, 244
 solvent effects on, 240-242, 241f, 264-265, 265t
 unimolecular, 244, 252
 vinyl halides and, 267
- Nucleotides, 104-105, 104f
 Nucleus, atomic, 7
- Nylon
 history of, 825, 859
 nylon 6, 1161, 1170
 nylon 6,6, 825, 859, 1149f, 1150, 1160, 1167, 1167f
 synthesis of, 859, 1160-1161
- o (suffix), 230
 Occidentalol, 606
- Octane, 1021
 constitutional isomers of, 118t
 molecular formula, 118t
- Octanenitrile, infrared (IR) spectrum of, 483
- Octane rating, of gasoline, 83
 4-Octene, 396
 1-Octen-3-ol, 769
- Octet rule, exceptions to, 17-18
 Octinoxate, 30, 852, 947
- Octylamine, infrared (IR) spectrum of, 483
- Octyl salicylate, 852
 1-Octyne, 741
- Odor(s)
 of acetone, 783
 of acid chlorides, 842
 of aldehydes, 783, 783f
 of amines, 956-957
 of caraway, 188
 of esters, 836
 of spearmint, 188
- OH proton, NMR spectra, 517-518, 518f
- oic acid (suffix), 690, 831
- Oil(s), 369
 crude, 128
 degree of unsaturation, 432
 essential, 1132
 fatty acid composition, 1123, 1123t
 fish, 1123
 hardening, 1124
 hydrogenation of, 432-434, 433f, 1124
 melting points of, 369, 1122-1123
 oxidation of, 556
 partial hydrogenation of, 432-433, 433f
 rancid, 1124
 unsaturated, 369
 vegetable, 835, 854, 855, 1123, 1124
- Oil of ginger, 367f
 Oil refinery, 128f
 Oil spill, 129
- Olefin metathesis, 1015-1019, 1016f, 1018f
- Olefins, 359. *See also* Alkenes
- Oleic acid, 358, 485, 1122t
 melting point, 368t
 structure, 104f, 368f, 368t
 triacylglycerol formation, 105
- Olestra, 854
- Olfactory receptors, 187, 188f
- Olive oil, 369
- Omega carbon, 691
- Omega-3 fatty acids, 368
- one (suffix), 777
- onitrile (suffix), 832, 833t
- Open arrow, 417
- Opium, 845
- Opsin, 799, 799f
- Optical activity, 182-183
 of chiral compounds, 183
 dextrorotatory (*d*) and levorotatory (*l*) compounds, 183
 enantiomeric excess, 184-185
 of racemic mixtures, 183
 specific rotation, 184, 185
- Optical purity, 184-185
- Orajel. *See* Benzocaine
- Oral contraceptives, 399f, 402-403, 403f
- Orbitals. *See also* Hybrid orbitals;
 Molecular orbitals (MOs);
 specific orbitals
 description, 8-9
 number of electrons in, 9
- Order, of a rate equation, 217
- Organic chemistry
 definition of, 1
 importance of, 1, 2f
- Organic halides. *See also* Alkyl halides
 coupling reaction with organocuprate reagents, 1003-1005
 coupling with alkenes (Heck reaction), 1003, 1009-1011
 coupling with organoboron reagents (Suzuki reaction), 1003, 1005-1009
- Organic molecules
 aromatic compounds, 607-640
 common features of, 4
 drawing structures of, 3, 27-31
 condensed structures, 27-29, 28f
 skeletal structures, 27-31, 29f
 examples of, 2-4
 grouping by functional group, 83
 shape. *See* Molecular shape
 solubility of, 94-95
- Organic reactions, 196-227. *See also*
 specific types of reactions
 acid-base, 56-58, 61-62
 arrows used in describing, 202-203, 202t
 asymmetric, 452
 bimolecular, 217
 bond breaking and bond making, 200-203, 201f
 bond dissociation energy, 203-206, 204t
 catalysts, 218-219, 219f
 endergonic, 210
 endothermic, 203-204
 energy diagrams for, 210-215, 212f, 215f
 energy of activation, 211-216, 212f, 215f, 219, 219f
 enthalpy change in, 203-206, 209-210
 entropy change in, 209-210
 enzymes in, 219-220, 220f
 equations for, 197, 197f
 equilibrium constants for, 206-209, 207f, 208t
 exergonic, 210
 exothermic, 203-204, 206
 free energy changes in, 206-209, 207f, 208t
 general types, A-13-14
 kinetics of, 215-218
 learning details of, 390-391
 mechanisms of. *See* Reaction mechanism
 multistep, 214
 oxidation and reduction reactions, 147-148, 148f
 polar, 201, 201f
 radical, 201, 201f
 rate of. *See* Reaction rate
 sequential, 197
 thermodynamics of, 206-209, 207f, 208t
 types of, 198-199. *See also* specific types of reactions
 unimolecular, 217
- Organic synthesis
 alkenes in, 391-392
 of aspirin, 268, 269f
 carbon-carbon bond-forming reactions in, 1002-1026
 definition of, 267
 halogenation reactions, 548-549
 halohydrin use in, 385
 by nucleophilic substitution reactions, 267-269, 268t
 reactions with organometallic reagents, 759-761
 retrosynthetic analysis, 417-419
 sodium borohydride reductions used in, 730f
 of taxol, 268
- Organoaluminum compounds, 1157
- Organoborane
 definition of, 387
 in Suzuki reaction, 1003, 1006-1009
 synthesis of, 1007
- Organocopper reagents. *See*
 Organocuprate reagents
- Organocuprate reagents, 739-740
 coupling reactions of, 1003-1005
 general features of, 1003-1004
 hydrocarbon synthesis in, 1004-1005
 stereospecificity of, 1004
 preparation of, 740
 reactions with α,β -unsaturated carbonyl compounds, 757
 reactions with acid chlorides, 752
- Organolithium reagents, 739-740
 addition to nitriles, 866
 in anionic polymerization, 1154
 arylborane synthesis, 1007
 in ethynylestradiol synthesis, 745
 in lithium diisopropylamide preparation, 888
- reactions
 with aldehydes and ketones, 742, 789
 as base, 741
 with esters, 752
 with α,β -unsaturated carbonyl compounds, 757
- Organomagnesium reagents, 739-740. *See also* Grignard reagents
- Organometallic reagents, 739-761. *See also* specific reagents
 acetylide anions, 740-741
 definition of, 739
 organic synthesis with, 759-761
 preparation of, 740
 protecting groups and, 748-750, 749f
 reactions
 acid-base, 748
 with acid chlorides, 750-753
 with aldehydes, 742-745, 745f, 789-790
 as base, 739, 741
 with carbon dioxide, 753-754
 with carbonyl compounds, 742-745, 748-750, 749f
 with carboxylic acid derivatives, 750-753
 with epoxides, 754-755
 with esters, 750-752
 with ketones, 742-745, 745f, 789-790
 as nucleophile, 739, 741-742
 summary of, 758
 with α,β -unsaturated carbonyl compounds, 755-758
- Organophosphorus reagents, 793-794
- Organosodium reagents, 740
- Ortho (prefix), 611
- Ortho, para activator, 658
- Ortho, para deactivator, 658
- Ortho, para director, 657-658, 661, 662-663, 664f, 666-668, 669, 984
- ose (suffix), 1029
- Oseltamivir, 86, 872
- Osmium tetroxide
 as oxidizing agent, 439
 in syn dihydroxylation, 443-444
- Osprey, 278
- Oxalic acid, 696
- Oxaphosphetane, 794
- Oxazaborolidine, 731
- Oxetane, 772
- Oxidation, 147-148, 148f, 426-462
 of alcohols, 439f, 447-450, 451
 of aldehydes, 726-727, 738-739
 of aldoses, 1047-1049
 of alkanes, 147-149
 of alkenes, 428f, 439f, 444-446
 of alkyl benzenes, 671-672
 of alkylboranes, 388-389
 of alkynes, 428f, 439f, 446-447
 of carbonyl compounds, 726-727
 carboxylic acid synthesis
 from alkyl benzenes, 698
 from alkynes, 698
 from primary alcohols, 698
 definition of, 147, 427
 dihydroxylation, 442-444
 anti, 442, 443
 syn, 442, 443-444
 energy released by, 206
 epoxidation, 439-442, 439f
 Sharpless, 451-454
 stereochemistry of, 440
 in synthesis of disparlure, 441, 442f
 of ethanol, 451
 of fatty acids, 853
 general scheme for, 427f
 green chemistry, 450-451

- hydroboration–oxidation reactions, 371f, 385–390, 387f, 388f, 389f, 389t, 412–413
- of hydrocarbons, 427, 428f
- of lipids, 151, 215
- oxidative cleavage
of alkenes, 444–446
of alkynes, 446–447
in haloform reaction, 894–895, 895f
- of polycyclic aromatic hydrocarbons (PAHs), 349
- reduction, relationship to, 427
- of substituted benzenes, 671–672
- of unsaturated fatty acids, 1124
- of unsaturated lipids, 556–557, 556f
- Oxidation and reduction reactions, 147–148
- Oxidative addition
in Heck reaction, 1011
- in palladium-catalyzed reactions, 1006, 1008, 1011
- in Suzuki reactions, 1008
- Oxidative cleavage
of alkenes, 444–446
of alkynes, 446–447, 698
- in haloform reaction, 894–895, 895f
- Oxidized state, 147
- Oxidizing agents, 438–439. *See also specific agents*
NAD⁺, 733
peroxyacids, 438, 438f
- Oxime, 1050
- Oxiranes, 317. *See also Epoxides*
- 6-Oxoheptanal, 945
- oxy (suffix), 316
- Oxycodone, 490
- Oxygen
bonding patterns and formal charge, 17t
- in condensed structures, 28
- infrared (IR) absorptions in oxygen-containing compounds, 482
- isotopes of, 472t
- mass of, 472t
- as radical scavenger, 541
- Oxytocin, 828, 1089, 1090f, 1105
- oyl (suffix), 779
- Ozone
destruction of atmospheric, 233, 551–552, 552f
- as oxidizing agent, 438, 445–447
- synthesis of, 551
- Ozonide, 445, 447
- PABA (*para*-aminobenzoic acid), 599, 684, 990–991
- Pacific yew tree, 268
- Padimate O, 599
- PAHs. *See* Polycyclic aromatic hydrocarbons (PAHs)
- Pain relievers, 906. *See also Analgesics*
- Palladium-catalyzed reactions, 219
- alkene reduction, 429
- alkyne reduction, 434–435
- general features of, 1005–1006
- Heck reaction, 1003, 1009–1011
- hydrogenation of aldehydes and ketones, 729
- oxidative addition in, 1006, 1008, 1011
- in peptide synthesis, 1097–1098
- reductive elimination in, 1006, 1008, 1011
- Suzuki reactions, 1003, 1005–1009
- Palmitic acid, 1122, 1122t
- Palmitoleic acid, 1122t
- Palm oil, 1123
- Palytoxin, 312, 356
- Para (prefix), 611
- Paracyclophane, 639
- Para red, 989
- Parent ion. *See* Molecular ion
- Parent name, 120
- Parkinson's disease, treatment of, 6, 45, 191, 958, 959f
- Paroxetine (Paxil), 78
- Patchouli alcohol, 331f, 1145
- PCB (polychlorinated biphenyl), 95–96
- PCC (pyridinium chlorochromate), 438–439, 448
- Peanut butter, 432
- PEG (polyethylene glycol), 109, 1157
- Penicillamine, 1113
- Penicillin(s), 838, 856–857
discovery of, 463
penicillin G, 106, 158, 463, 485
- 1,4-Pentadiene, 572, 573f
hydrogenation of, 583
- (3*E*)-1,3-Pentadiene, hydrogenation of, 583
- Pentalene, 635
- 2-Pentanamine, in E2 elimination reaction, 979–980, 979f
- Pentane, 466–467
boiling point, 91
constitutional isomers of, 118t
melting point of, 93
molecular formula, 118t
structure, 116
van der Waals forces, 88, 88f, 90
- 2,4-Pentanedione, 882, 886, 908, 909
- Pentanoic anhydride, 870
- 3-Pentanol, 746
- 2-Pentanone, 910
mass spectrum of, 472
preparation from an acid chloride and an organocuprate reagent, 753
- 3-Pentanone, boiling point of, 91, 91f, 914
- Pentanoyl chloride, 492, 870
- 1-Pentene, 466–467
- 2-Pentylcinnamaldehyde, 923
- 1-Pentyne, 466–467
- Peptide bonds, 1086, 1088–1089
s-cis conformation, 1088
planar geometry of, 1088–1089
s-trans conformation, 1088–1089
- Peptides, 837, 1086–1101
constitutional isomers, 1087
interesting examples, 1089–1090, 1090f
- N- and C-terminal amino acids, 1087
- peptide bond in, 1088–1089
- sequencing, 1090–1094
amino acid analysis, 1091
Edman degradation, 1091–1092
partial hydrolysis of peptide, 1092–1093, 1093t
- simple, 1086–1088
synthesis of, 1094–1101
automated, 1099–1100, 1101f
protecting groups in, 1094–1098
solid phase technique, 1099–1100
- Percent *s*-character, 41, 68
bond length and, 582
effect on basicity of amines, 972–973
hybrid orbital in alkenes, 284
infrared (IR) absorption and, 480
- Percent transmittance, 477
- Perfume, 435
- Period, in periodic table, 8
- Periodic table, 7–10
bond length trends, 23
element location and bonding type, 10
of elements common in organic chemistry, 8f
first-row elements, 8f, 9
organization of, 8
second-row elements, 8f, 9–10
bonding of, 12
exceptions to octet rule, 18
shape arrangements, 24
- third-row elements, 18
- trends
acidity, 63–65
electronegativity, 42, 42f
- Periodic trends
in bond dissociation energies, 204–205
in leaving group ability, 236–237
nucleophilicity, 239, 240, 241
- Periplanone B, 486, 925, 925f, 1145
- Perkin, William Henry, 988–989
- Perlon. *See* Nylon, Nylon 6
- Peroxide, as radical initiator, 540
- Peroxyacetic acid, 438f, 439
- Peroxyacids
in epoxidation, 439–441
as oxidizing agents, 438, 438f
- Peroxy radical, 1151
- Pesticide, 233
- PET. *See* Polyethylene terephthalate (PET)
- Petroleum, 113, 128–129, 129f
alkenes prepared from, 562
ethylene preparation from, 366
as feedstock of polymer synthesis, 1167
- Ph- (abbreviation), 612
- PHAs (polyhydroxyalkanoates), 1170–1171
- PHB (polyhydroxybutyrate), 1171
- PHBV, 1171
- α -Phellandrene, 1133f
- Phenacetin, 877
- Phenanthrene, 621, 636, 640
- Phenol(s), 151, 227, 611, 717
acidity of, 701, 703f
as antioxidants, 557
in aspirin synthesis, 268, 269f
Bakelite preparation from, 1165
conjugate base of, 701–702, 702f, 703f
formation from anisole, 343
hydroxy group of, 313
naming, 612
 pK_a of, 701, 703f
polyhalogenation of, 665
synthesis from aryl diazonium salts, 982
- Phenoxide, 701–702, 702f, 703f
- Phentermine, 76
- Phenylacetaldehyde, 777f, 884
- Phenylacetamide, 871
- Phenylacetic acid, 871
- Phenylacetoneitrile, 871
- Phenylalanine
abbreviation for, 711t
N-acetyl, 1086
isoelectric point for, 1078t
 pK_a values for, 1078t
structure of, 1076f
synthesis of, 1080, 1086
- 4-Phenyl-2-butanone, 998
- Phenylcyclohexane
conformations of, 208
phenyl group in, 83
- Phenylethanal. *See* Phenylacetaldehyde
- 2-Phenylethylamine, 965
derivatives of, 958–959
in Hofmann elimination reaction, 978
- Phenyl group, 83, 84t, 612
(1*E*)-1-Phenyl-1-hexene, 1024
- Phenyl isothiocyanate, 1091–1092
(1*Z*,3*E*)-1-Phenyl-1,3-octadiene, 1009
- 3-Phenyl-1-propanamine, 995
- Phenylpropanoic acid, 195
- N*-Phenylthiohydantoin (PTH), 1091–1092, 1118
- N*-Phenylthiourea, 1091–1092
- Pheromones, 114, 115, 441, 442f, 454, 454f, 843, 844, 925, 925f, 1008
- Phomallenic acid C, 421
- Phosgene, 1163
- Phosphatidylcholine, 1126
- Phosphatidylethanolamine, 1126
- Phosphine, in palladium-catalyzed reactions, 1005
- Phosphoacylglycerols, 1126–1127, 1127f, 1128f
- Phosphodiester, 1126, 1127, 1128f
- Phospholipids
in cell membranes, 100, 101f, 102, 102f, 1126–1128
phosphoacylglycerols, 1126–1127, 1127f, 1128f
sphingomyelins, 1127–1128, 1128f
structure of, 100, 101f, 111, 1126
- Phosphoric acid, 1126
- Phosphoric acid diester, 1126
- Phosphorus, octet rule exceptions in, 18
- Phosphorus oxychloride, dehydration of
alcohols using, 330–331
- Phosphorus tribromide, conversion of
alcohols to alkyl halides with, 335, 336–337
- Photons, 474–476
- Photosynthesis, 1028
- Phthalic acid, 672
- Phthalimide, 961
- Phylloquinone, 98
- Physical properties. *See* Boiling point; Melting point; Solubility
- Pi* (π^*) antibonding molecular orbital, 627, 628f
- Pi (π) bond, 37–40, 38f, 39f, 40f
in alkene double bond, 281
in alkenes, 359, 360, 370–373, 391
in alkynes, 299, 300f, 400–401, 405–406
in aromatic compounds, 83, 608, 609
breaking in addition reactions, 199, 359, 370–373, 371f, 378, 379, 380, 383, 390, 391
bromination as test for, 380
Brønsted–Lowry bases and, 55–56, 55f
in carbonyl group, 85, 722–723, 724
degrees of unsaturation and, 360
delocalized, 583
formation in elimination reactions, 198, 279–280, 291, 294, 299, 300f
in functional groups, 82, 102–103
in Lewis base, 72, 74
in polyenes, 580, 581
radical reactions with, 541, 562
in resonance structure, 67
strength, 359
- Pi (π) bonding molecular orbital, 627, 628f
- Picometer (pm), 23
- Pi (π) electrons
in aromatic compounds, 613, 617, 618–619, 619t
in benzene, 609, 610
Hückel's rule and, 618–619, 619t, 626–629
in NMR spectroscopy, 505–506, 506t
- Pinacol, 357
- Pinacolone, 357
- α -Pinene, 1145
- Piperidine
basicity of, 972–973, 973t
structure, 953
- Pitocin. *See* Oxytocin
- Pivalic acid, 60
- pK_a values, 59–68, A–1–2
acid strength and, 59–60
of amines, 966, 968–969, 974t
for amino acids, 1077, 1078, 1078t
of carbonyl compounds, 884–885, 885t, 886t
definition, 59
of leaving groups in nucleophilic substitution reactions, 237
logarithmic scale of, 60

- pK_a values (*continued*)
of selected organic compounds, 59t
of strong acids, 70
of strong bases, 71
- PLA (polylactic acid), 860
- Planar carbocation, 377, 1043–1044
- Planar double bond, 376–377, 381, 386, 388
- Planar molecule
benzene, 609–610
- Plan B, 403
- Plane of symmetry, 165–166, 165f
in disubstituted cycloalkanes, 180
of meso compound, 178
- Plane-polarized light, 182–183
- Plaque, atherosclerotic, 861f, 862
- Plastic, polyethylene, 84
- Plasticizers, 96, 1165–1166
- Platinum, as catalyst, 219
- Plavix. *See* Clopidogrel
- β -Pleated sheet, 1101–1104, 1103f, 1105f, 1107f
- PMMA [Poly(methyl methacrylate)], 568
- Poison ivy, 707
- Polar aprotic solvents, 241, 241f, 264, 265, 265t, 287, 288t, 889–890
- Polar bond
carbon–halogen, 234, 234f
description of, 43, 44, 45f
- Polarimeter, 182
- Polarity, of carboxylic acids, 699
- Polarizability
boiling point and, 91f
inductive effects on substituted benzenes, 654
van der Waals forces and, 88–89
- Polarized light, 182–183
- Polar molecule
description, 44, 45f
solubility of, 94
- Polar protic solvents, 240, 240f, 264–265, 265t, 293t, 890
- Polar reactions, 201, 201f
- Poly- (prefix), 1155
- Poly(acrylic acid), 561, 1171
- Polyacrylonitrile, 1155f
- Polyalkylation, 666
- Polyamides, 825, 859, 1160–1161
- PolyAspirin, 1177
- Polyatomic ion, 16
- Polycaprolactone, 1177
- Polycarbonates, 1163
- Polychlorinated biphenyl (PCB), 95–96
- Polycyclic aromatic hydrocarbons (PAHs), 349, 614, 614f, 621
- Polydioxanone, 1177
- Polyenes, 580, 581
ultraviolet absorption, 598
- Polyesters, 859–860, 1161–1162
- Polyethers, 320, 1156
- Poly(ethyl acrylate), 568
- Poly(ethyl α -cyanoacrylate), 1155f
- Polyethylene, 84, 366, 366f, 561, 1149f, 1152–1153, 1158, 1159
- Poly(ethylene glycol) (PEG), 109, 1157
- Polyethylene terephthalate (PET), 719, 859–860, 1148, 1150f, 1161–1162, 1169t, 1170, 1174
- Polyfunctional compounds, nomenclature of, A–4–5
- Polyhalogenation, 665
- Polyhydroxyalkanoates (PHAs), 1170–1171
- Polyhydroxybutyrate (PHB), 1171
- Poly(hydroxyethyl methacrylate), 568
- Polyhydroxyvalerate, 1171
- Polyisobutylene, 568, 1155f
- Poly(lactic acid) (PLA), 860, 1167–1168
- Polymerization, 560–563
anionic, 1154–1155, 1155f
of epoxides, 1156–1157
cationic, 1153–1154, 1155f
chain branching in, 1152–1153
coordination, 1159
definition of, 561, 1149
feedstock for, 1166–1167
head-to-tail, 563
ionic, 1153–1156, 1155f
living, 1155
radical, 562–563, 1151–1152, 1151f, 1153
Ziegler–Natta, 1157–1159
- Polymers, 560–562, 560–563, 1148–1178
alternating, 1156
amorphous regions of, 1164–1165
atactic, 1157–1158
“Big Six,” 1169, 1169t
biodegradable, 1170–1171
chain-growth, 1149, 1150–1157
in consumer products, 1149, 1149f
copolymers, 1156
crystalline regions of, 1164–1165
definition of, 160, 1149
derivation of word, 1149
green synthesis of, 1166–1169, 1167f, 1168f
insoluble, 1099
isotactic, 1157–1158
in medicine and dentistry, 561, 562t
molecular weights of, 1150
natural and synthetic fibers, 858–860, 858f
nylon, 825, 859, 1149f, 1150, 1160–1161, 1167, 1167f, 1170
physical properties of, 1164–1166
polyesters, 859–860
random, 1156
recycling of, 1148, 1169–1170, 1169t
shorthand representation of, 1150, 1150f
step-growth, 1150, 1160–1164
stereochemistry of, 1157–1159
structure of, 1149, 1164–1166
syndiotactic, 1157–1158
synthetic, 1148–1178. *See also* Synthetic polymers
- Poly(methyl methacrylate) (PMMA), 568
- Polypropylene, 562t, 1169t
- Polysaccharides, 1059–1061
cellulose, 1059–1060
glycogen, 1061
starch, 1060–1061
- Polystyrene, 96, 538, 562, 1150f, 1169t
preparation from ethylene, 366f
- Polystyrene derivative, 1099
- Polytetrafluoroethylene (Teflon), 562t
- Poly(trimethylene terephthalate) (PTT), 1167, 1168f
- Polytulipalin, 946
- Polyurethanes, 1162
- Poly(vinyl acetate), 562, 1155f
preparation from ethylene, 366f
- Poly(vinyl alcohol), 1176
- Poly(vinyl butyral), 1176
- Poly(vinyl chloride) (PVC), 109, 232, 562t, 1149, 1165, 1169t
preparation from ethylene, 366f
- p Orbitals
description, 8–9
hybridization of, 33–40, 33f–35f, 38f–40f
hyperconjugation, 257
overlap of, 572–573, 578–579, 582, 610, 610f
- $2p$ orbitals, 9, 32–35, 34f
in alkenes, 281
in alkynes, 400, 401, 405
- Potassium *tert*-butoxide, 239
in dehydrohalogenation reactions, 280, 280t
- Potassium dichromate, 438, 448, 449, 449f
- Potassium hydroxide, in dehydrohalogenation reactions, 280t
- Potassium iodide, 10, 983
- Potassium permanganate (KMnO₄)
as oxidizing agent, 71–672, 439, 675, 738
in syn dihydroxylation, 443–444
- Potassium propanoate, 692
- Potassium sorbate, 696
- Precor. *See* Methoprene
- Prednisone, 783
- Prefix, 120
cis, 363
cyclo-, 118, 125
deoxy, 1063
di-, 123
E, 363
epoxy, 317
iso-, 120
meta-, 611
ortho-, 611
para-, 611
R, 170–175, 179–180
S, 170–175, 179–180
sec-, 120
tert-, 120, 121
tetra-, 123
trans, 363
tri-, 123
Z, 363
- Pregabalin, 106
- Preparations, 697
- Prepolymer, 1163–1164, 1164f
- Preservatives, 261
- Primary (1°) alcohols, 390
classification as, 313
conversion to alkyl halides with HX, 332–334, 337t
conversion to alkyl halides with phosphorus tribromide, 336, 337t
conversion to alkyl halides with thionyl chloride, 335–336, 337t
dehydration by E2 mechanism, 327
hydrogen bonding extent, 318
from nucleophilic addition to aldehyde, 735
oxidation to carboxylic acids, 447, 448–450, 698
from reactions of aldehydes with organometallic reagents, 742–743
from reduction of acid chlorides, 734
from reduction of aldehydes, 432, 727, 728, 729
from reduction of carboxylic acid derivatives, 727
from reduction of carboxylic acids, 735
from reduction of esters, 734
from reduction of monosaccharides, 1047
- Primary (1°) alkyl halides
acetylide anion reactions with, 414–415
classification as, 229–230
E2 elimination reactions, 287, 288, 302f
example of, 229f
S_N2 substitution reactions, 248, 262, 262f, 302f
synthesis of, 332, 335, 336
- Primary (1°) amides, 827, 831, 834–836, 842, 844, 850
tautomers, 863
- Primary (1°) amines
addition to aldehydes and ketones, 797–799, 799f, 801f
amino acids, 1077
from direct nucleophilic substitution, 960–961
Gabriel synthesis, 961–962
IR spectra of, 955, 955f
nomenclature, 952
reactions with acid chlorides and anhydrides, 975–976
reactions with aldehydes and ketones, 975
reactions with nitrous acid, 980–981
from reduction of amides, 963
from reduction of nitriles, 962
from reduction of nitro compounds, 962
structure of, 950
synthesis from reduction of nitriles, 865
- Primary (1°) carbocations, 256
- Primary carbon (1°), 116
- Primary hydrogen (1°), 116–117
- Primary structure, of proteins, 1101, 1107f
- Primobolan, 1146
- Procaine, 615f
- Progesterone
analogs of, 402, 403
function in body, 1141t
structure of, 403, 1120f, 1141t
synthesis of, 928
- Progestins, 1141
- Prograf. *See* FK506
- Proline, 719
isoelectric point for, 1078t
pK_a values for, 1078t
structure of, 1076f, 1077
- Prontosil, 990
- Propanal
aldol reaction of, 918–919, 922
enolate formation from, 884
IR spectrum of, 780, 781f
NMR spectra of, 782, 782f
- Propanamide
boiling point of, 92
infrared (IR) spectrum of, 483
- 2-Propanamine, 952
- Propane, 224
bromination of, 546–547, 547f
chlorination of, 546–548, 548f
molecular formula of, 114, 115, 118t
as propellant, 551
structure of, 114–115
torsional energy of, 134
- Propanedioic acid, 692
- 1,2-Propanediol, 352
- 1,3-Propanediol, 352, 1167, 1168f, 1176
- 1,2,3-Propanetriol, 316
- Propanil, 999
- Propanoic acid
NMR spectra of, 695f
pK_a of, 77
- 1-Propanol
conversion to butanenitrile, 337–338
IR spectrum of, 477
NMR spectrum of, 524, 526f
- Proparacaine, 684, 1001
- Propene
borane addition to, 387
hydrohalogenation, 374
monomers in polymerization, 562t
radical halogenation, 552–553
synthesis from propylamine, 977–978
- 2-Propen-1-ol, 363
- Propiolic acid, 77
- Propionic acid, structure of, 691t

- Propiophenone, 732
 halogenation of, 893–894
- Propoxyphene (Darvon), 106, 166–167
- Propranolol, 70, 356
- Propylamine, conversion to propene, 977–978
- Propylbenzene, 227, 673
- Propyl group, 120
- Prostacyclins, 1129–1130, 1130t, 1131f
- Prostaglandins, 150, 150f, 215, 1129–1131
 analogues, 1131
 biological functions, 697
 inhibitors of synthesis, 1131–1132
 PGA₂, synthesis of, 286, 933
 PGE₁, 768, 1131
 PGE₂, 1130
 PGF_{2α}, 150, 150f, 223, 306, 398, 570, 697, 1120f, 1130f, 1144
 PGG₂, 570, 697
 PGI₂, 1130f
 synthesis of, 697, 1130, 1131f
- Protecting groups, 748–750
 acetals as, 808–809
 amide for amines, 977
 in amino acid synthesis, 1094–1098
 general strategy for using, 749f
- Proteins, 1086, 1101–1110. *See also* Peptides
 amide linkages in, 837, 855–856
 conjugated, 1109
 fibrous, 1106
 globular, 1106
 important examples, 1106–1110, 1108f, 1109f
 membrane, 1075
 natural fibers, 858, 858f
 stability of, 855–856
 structure, 1101–1106
 primary, 1101, 1107f
 quaternary, 1105, 1107f
 secondary, 1101–1104, 1102f, 1103f, 1105f, 1107f
 tertiary, 1104–1105, 1107f
 subunits, 1105
- Protic solvents, polar, 240, 240f, 264–265, 265t, 293t, 890
- Proton acceptor, 55
- Protonation
 in acid-catalyzed ester hydrolysis, 851
 in aldehyde/ketone nucleophilic addition reactions, 786–787
 of carboxylic acids, 699
 in glycoside formation, 1043–1044
 in glycoside hydrolysis, 1044
 in nucleophilic addition to aldehydes and ketones, 724
 tautomerization in acid, 410–412
- Proton donor, 55
- Proton NMR, 495, 497–522
 of aldehydes, 782, 782f
 of amines, 956, 956f
 of aromatic compounds, 619–620
 of benzene derivatives, 518–519, 519f, 613–614, 613t
 of carboxylic acid derivatives, 836
 of carboxylic acids, 694, 695f
 characteristic absorptions, A–11–12
 chemical shift
 equation for, 498
 predicting values of, 503–504
 protons on benzene rings, 505, 506t
 protons on carbon–carbon double bonds, 505, 506t
 protons on carbon–carbon triple bonds, 505, 506t
 scale, 497
 values for common bonds, 504t
 coupled protons, 510
 of cyclohexane conformations, 518
 general principles
 chemical shifts, 497–498
 downfield signals, 497–498
 reference signal, 497
 scale, 497
 signal intensity, 497
 upfield signals, 497–498
 identification of an unknown, 519–522
 integrals, 507–508
 intensity of signals, 507–508
 of ketones, 782
 number of signals, 498–502
 enantiotopic and diastereotopic protons, 500–502
 equivalent protons in alkenes and cycloalkanes, 499–500
 general principles, 498–499
 OH proton, 517–518, 518f
 position of signals, 502–504
 chemical shift values, 503–504, 504t
 shielding and deshielding effects, 502–503, 502f, 503f, 506f, 506t
 regions of spectra, 506, 506f
 spin–spin splitting, 508–517
 in alkenes, 516–517, 516f, 517f
 common patterns of, 512t
 complex examples of, 513–515
 how a doublet arises, 509–510
 how a triplet arises, 510
 overview of, 508–509
 rules and examples, 510–513
- Protons
 description of, 7
 relative acidity of, determining, 69–70
- Proton transfer reaction
 aspirin and, 71–72
 description of, 56–58
- Proventil. *See* Albuterol
- Proximity effect, 587
- Prozac. *See* Fluoxetine
- D-Psicose, 1036f
- Pseudoephedrine, 65, 192, 966, 999
- Psilocin, 958–959
- Psilocybe mushroom, 959
- PTH (*N*-phenylthiohydantoin), 1091–1092
- PTT. *See* Poly(trimethylene terephthalate)
- Puffer fish, 589, 589f
- Purification procedures
 extraction, 707–709, 708f, 709f
- Purine, 635, 1063–1064
- Putrescine, 956–957
- PVC. *See* Poly(vinyl chloride)
- Pyran, 1037
- Pyranose ring, 1037, 1056
- Pyrethrin I, 106, 1012, 1023
- Pyridine, 71, 369–370, 621–622, 687, 842
 alcohol conversion to alkyl tosylates in, 338–340, 341f
 basicity of, 972–973, 973t, 974t
 conversion of alcohols to alkyl halides, 335–336
 conversion of monosaccharides to esters using, 1046
 dehydration of alcohols using, 330–331
 electrostatic potential plot of, 622, 623f
 structure of, 953
- Pyridinium chlorochromate (PCC), 438–439, 448
- Pyridoxine, 109
- Pyrimidine, 1063–1064
- α-Pyrone, 635
- Pyrolole, 622, 636, 687, 995
 basicity of, 972, 973t, 974t
 electrostatic potential plot of, 622, 623f
 structure of, 953
- Pyrolidone, structure of, 953, 995
- Pyruvic acid, 733
- Qinghao, 389
- Quantum, 474
- Quartet, NMR spectrum, 511t, 512t, 518
- Quaternary ammonium salt. *See also* Ammonium salt
 in Hofmann elimination reaction, 977–978
 synthesis of, 960–961
- Quaternary carbon (4°), 116
- Quaternary structure, of proteins, 1105, 1107f
- Quiana, 1174
- Quinine, 194
 as natural alkaloid, 957
 source of, 289
 structure of, 624, 975
 synthesis of, 288, 289f, 896, 896f
- Quinoline, 435
- Quintet, NMR spectrum, 511t
- Quinuclidine, 277
- R (prefix), 170–175, 179–180
- Racemic mixture (racemate), 183–184, 184t
 of alcohols from hydroboration–oxidation of alkenes, 389
 alkene halogenation, 381
 of alkyl chlorides, 334
 of amino acids, separation of, 1081–1084, 1082f
 in carbene addition, 1014
 in enolate alkylation, 898
 from epoxidation of alkenes, 440
 in epoxide ring opening reactions, 344
 formation by hydrohalogenation, 376
 halogenation reactions and, 381, 549–550
 in hydride reduction of ketones, 730
 S_N1 substitution reactions and, 254–255, 255f
- Racemic switch, 187
- Racemization
 at α carbon of carbonyl compounds, 891
 definition of, 254
 in S_N1 substitution reactions, 254–255, 255f
- Radiation
 electromagnetic, 474–476, 475f
 infrared, 476–477
 RF (radiofrequency), 495–496, 527
 ultraviolet, 599
- Radical(s)
 bond formation from, 202
 definition of, 539
 as electrophile, 202
 as reactive intermediate, 201–202, 201f
 resonance-stabilized, 555
 stability of, 539–540, 540f
- Radical anion, 436
- Radical cation, 465
- Radical inhibitor, 541
- Radical initiator, 540
- Radical intermediates, 539, 554, 558
- Radical polymerization, 1151–1152
 chain branching in, 1153
 chain termination in, 1152
 monomers used in, 1151, 1151f
- Radical reactions, 201, 201f, 538–570. *See also specific reactions*
 of alkanes, 541–551
 of alkenes, 541, 558–560
 applications of
 antioxidants, 557
 oxidation of unsaturated lipids, 556–557, 556f
 ozone destruction by CFCs, 551–552, 552f
 carbon–hydrogen bond, reaction of radical with, 540–541
 double bonds, addition to, 541, 558–560
 general features of, 540–541
 halogenation at allylic carbon, 552–555
 bromination, 553–554
 product mixtures in, 555
 halogenation of alkanes, 541–551
 of achiral starting material, 549–550, 549t
 bromination, 546–547, 547f
 of chiral starting material, 549t, 550
 chlorination, 544–548, 544f, 545f, 548f, 549–551
 energy changes during, 544, 544f, 545f
 mechanism of, 542–544
 in organic synthesis, 548–549
 stereochemistry of, 549–551, 549t
 halogenation of alkyl benzenes, 669–671
 inhibitors of, 541
 polymerization, 560–563, 562t
 two radicals reacting with each other, 541
- Radical scavenger, 541
- Radical substitution reaction, 541, A–13
- Randomness, entropy as measure of, 209
- Rate constant (*k*), 216–217
- Rate-determining step
 in chlorination of ethane, 544, 545f, 548
 E1 elimination reactions, 292
 electrophilic additions, 373, 373f, 374
 in halogenation reactions, 546–548
 bromination, 547
 chlorination, 548
 halogenation of alkenes, 382
 in a multistep mechanism, 214–215, 217
 in nucleophilic substitution reactions, 243, 244
 rate equation for, 217
 S_N1 substitution reactions, 260
- Rate equation, 216–218
 first-order, 217, 243, 244
 for nucleophilic substitution reactions, 243–244
 second-order, 217, 243, 244
- Rate law, 216–218
- RCM (ring-closing metathesis), 1018–1019, 1018f
- Reaction arrow, 197, 202t
 curved, 201, 202, 202t
 double, 202t
- Reaction coordinate, 210, 211, 212f, 215f
- Reaction mechanism
 bond cleavage, 200–202
 bond formation, 202
 concerted, 200
 definition of, 200
 energy diagram for a two-step, 213–215, 215f
 rate equations and, 216–218
 stepwise, 200, 213–215, 217
- Reaction rate
 catalysts and, 218–219, 219f
 concentration effect on, 216, 217–218
 of endothermic reactions, 212f, 259
 energy of activation (*E*_a) and, 215–217
 of exothermic reactions, 212f, 259
 nucleophilic substitution reactions, 264–265
 rate-determining step, 214–215, 217
 temperature effect on, 216
- Reactions. *See* Organic reactions
- Reactive intermediate
 carbanion, 201–202, 201f
 carbocation, 201–202, 201f, 252–253, 330
 description of, 200
 radicals, 201–202, 201f
 synthetic intermediate compared, 392

- Reactivity, functional groups and, 102–104
- Reactivity–selectivity principle, 546
- Reagent, 197, 197f
- Rebaudioside A, 1045
- Receptors, chiral, 187
- Recycling of polymers, 1148, 1169–1170, 1169t
- Red seaweed (*Asparagopsis taxiformis*), 232
- Reduced state, 147
- Reducing agents, 428. *See also specific agents*
- chiral, 731
- metal hydride reagents, 428, 727–731, 733–738, 738t
- NADH, 732–733
- Reducing sugars, 1048, 1048f, 1057
- Reduction, 147–148, 148f, 426–462
- of acid chlorides and esters, 734–735
- of alkenes, 427, 428–432, 428f
- of alkyl halides, 437–438, 438f
- of alkynes, 427, 428f, 434–437, 437f
- to alkanes, 428f, 434–435
- to cis alkenes, 434, 435
- to trans alkenes, 434, 436
- of amides to amines, 736–737, 963
- of aryl ketones to alkyl benzenes, 672–673
- asymmetric, 731
- of carbonyl compounds, 726–738
- biological, 732–733
- carboxylic acids and derivatives, 733–738
- by catalytic hydrogenation, 729
- enantioselective, 731–733, 732f
- with metal hydride reagents, 728–729, 730f, 738t
- stereochemistry of, 729–731
- of the carbonyl group of monosaccharides, 1047
- Clemmensen, 672
- definition of, 147, 427
- dissolving metal, 428, 436
- of epoxides, 437–438, 438f
- general scheme for, 427f
- of hydrocarbons, 427, 428f
- hydrogenation, 429–434
- of ketones to amines, 963–966
- of nitriles, 863, 865–866
- of nitriles to amines, 962
- of nitro groups, 673–674
- oxidation, relationship to, 427
- of polar C–X sigma (σ) bonds, 437–438
- of triple bonds, 437–437, 437f
- Wolff–Kishner, 672
- Reductive amination, of aldehydes and ketones to amines, 963–966
- Reductive elimination
- in Heck reaction, 1011
- in palladium-catalyzed reactions, 1006, 1008, 1011
- in Suzuki reactions, 1008
- Refining, 128–129, 129f
- Refrigerant, 233
- Regioselectivity
- of alcohol dehydrations, 325
- of E1 elimination reactions, 293
- of E2 elimination reactions, 289, 296
- of electrophilic addition of hydrogen halides, 374
- of epoxide ring opening reactions, 345–346
- of halohydrin formation, 384–385, 385t
- of Hofmann elimination, 979
- of hydroboration, 387
- Reserpine, 879
- Resolution, of amino acids, 1081–1084, 1082f
- Resolving agent, 1082
- Resonance, 18–23
- of allylic carbocations, 574–575, 574f
- allyl type, 575–576
- benzene, 609
- examples of, 575–577
- theory/principles, 18
- usage of term, 574
- Resonance delocalization, 67
- Resonance effects
- on acidity, 66–67, 67f, 69f
- on basicity of amines, 973t
- electron-donating, 654–656
- electron-withdrawing, 654–656
- on substituted benzenes, 654–657
- Resonance hybrid, 18, 22–23
- of benzene, 609
- drawing, 22
- electron delocalization, hybridization, and geometry, 578–579
- major contributor to, 577–578
- minor contributors to, 577–578
- Resonance stabilization, of carboxylic acids, 701–702, 702f, 703f
- Resonance-stabilized allyl carbocation, 574–575, 574f, 575f
- Resonance-stabilized carbocation, 584–585, 587
- in electrophilic aromatic substitution reaction, 643
- in glycoside formation, 1043–1044
- in glycoside hydrolysis, 1044
- Resonance-stabilized radical, 555
- Resonance structures, 18–23, 574–579
- of acetate, 701, 703f
- acidity and, 66–67
- of allyl radical, 553
- of benzylic radical, 669
- of carboxylic acid derivatives, 828
- cations with positive charge adjacent to lone pair, 576
- conjugated allyl carbocation, 574
- of conjugated dienes, 582, 583
- double bond
- conjugated, 576
- major contributor, 577–578, 582
- minor contributors, 577–578, 582
- with one atom more electronegative than the other, 576, 577
- drawing, 19–22
- of enolates, 884
- of enols, 883–884
- isomers compared to, 19
- of naphthalene, 621
- of phenoxide, 701–702, 702f, 703f
- stability of, 577–578
- of substituted benzenes, 654–655
- three atom allyl system, 575–576, 577
- of α,β -unsaturated carbonyl compound, 934
- of Wittig reagent, 793
- Retention of configuration
- in conversion of alcohols to alkyl tosylates, 338–339, 340
- in oxidation of alkylboranes, 388–389
- in S_N2 substitution reactions, 246
- Retention time, in gas chromatography, 473–474
- Retinal, 799, 799f
- Retinol, 97
- Retro Diels–Alder reaction, 595–596
- Retrosynthetic analysis, 392, 417–419
- in acetoacetic ester synthesis, 904
- in benzene derivative synthesis, 985
- of Diels–Alder product, 595
- of disparlure, 441
- of Grignard products, 746–748
- of Heck reaction, 1010
- organometallic reagent reactions, 760–761
- in substituted benzene syntheses, 674, 675–677
- synthesis of *m*-bromoaniline from benzene, 674
- using aldol reaction, 920–921
- using malonic ester synthesis, 902–903
- Wittig reaction, 795–796
- RF radiation, 495–496, 527
- R group, 82, 120
- identity in S_N2 substitution reactions, 248–249, 249t
- Rhodium, 1085
- Rhodopsin, 799, 799f
- Ribbon diagrams, 1104
- α -D-Ribofuranose, 1042, 1062
- β -D-Ribofuranose, 1042
- Ribonucleosides, 1063–1064
- Ribonucleotides, 1064
- D-Ribose, 1034, 1035f, 1063
- D-Ribulose, 1036f
- Rimantadine, 966
- Ring-closing metathesis (RCM), 1018–1019, 1018f
- Ring-flipping, 140–141, 141f
- Ring formation, in Robinson annulation, 936–939
- Ring-opening metathesis polymerization (ROMP), 1022
- Ritalin. *See* Methylphenidate
- RNA, 1064, 1065f
- Robinson, Sir Robert, 936
- Robinson annulation, 936–939
- Rocaltrol. *See* Calcitriol
- Rofecoxib, 1132
- Row, in periodic table, 8
- Rowland, F. Sherwood, 551
- RU 486, 403
- Rubber, 1149f, 1159–1160, 1159f
- Rubbing alcohol, 319f
- Ruthenium, 1015
- S* (prefix), 170–175, 179–180
- Saccharide, origin of word, 1028
- Saccharin, 1058, 1059f
- Safflower oil, 369
- Safrole, 998
- Salicin, 71, 195, 696, 1045
- Salicylamide, 874
- Salicylates, 71
- Salicylic acid, 696, 873
- Salmeterol, 347, 347f, 732f, 772
- Salsolinol, 820
- Salts
- Brønsted–Lowry bases and, 55
- of carboxylic acids, 696
- formation of, 10–11
- negatively charged nucleophile, 235
- SAM (*S*-Adenosylmethionine), 250
- SAME, 250
- Samuelsson, Bengt, 697
- Sandmeyer reaction, 983
- Santalbic acid, 401
- Saponification, 851, 854–855
- Saponins, 98
- Saquinavir, 195
- Saran, 1156
- Saturated fats, 369
- Saturated fatty acid, 358
- Saturated hydrocarbons, 114
- Saytzeff rule, 289
- SBR (styrene–butadiene rubber), polymerization of, 1156
- Sch38516, 1018f
- s*-character, of hybrid orbitals, 68
- Schiff base, 797
- Schrock, Richard, 1015
- s*-cis conformation, 580
- Scombrod fish poisoning, 622
- Seashells, chiral helical, 166
- Seaweed, 232
- Sebacic acid, 1177
- Sebacyl chloride, 1177
- sec*- (prefix), 120
- Secondary (2°) alcohols, 390
- classification as, 313
- conversion to alkyl halides with HX, 332–334, 337t
- conversion to alkyl halides with phosphorus tribromide, 336, 337t
- conversion to alkyl halides with thionyl chloride, 335–336, 337t
- dehydration by E1 mechanism, 326
- hydrogen bonding extent, 318
- oxidation to ketones, 447, 448, 450
- from reactions of aldehydes with organometallic reagents, 742–743
- from reduction of ketones, 432, 727, 728, 729
- Secondary (2°) alkyl halides
- acetylide anion reactions with, 414–415
- classification as, 229–230
- E1 elimination reactions, 292, 303f
- E2 elimination reactions, 287, 302f–303f
- example of, 229f
- S_N1 substitution reactions, 255, 262, 262f, 303f
- S_N2 substitution reactions, 248, 262, 262f, 302f
- synthesis of, 332, 335, 336
- Secondary (2°) amides, 827, 831–832, 834–836, 842, 844, 850
- Secondary (2°) amines
- addition to aldehydes and ketones, 800–801, 801f
- from direct nucleophilic substitution, 960
- IR spectra of, 955, 955f
- nomenclature, 952–953
- proline, 1077
- reactions with acid chlorides and anhydrides, 975–976
- reactions with aldehydes and ketones, 975
- reactions with nitrous acid, 981–982
- from reduction of amides, 963
- structure of, 950
- Secondary carbon (2°), 116
- Secondary hydrogen (2°), 116–117
- Secondary structure, of proteins, 1101–1104, 1102f, 1103f, 1105f, 1107f
- Second-order kinetics, 217
- of E2 elimination reactions, 285
- in nucleophilic substitution reactions, 243–244
- Second-order rate equation, 217, 243, 244
- L-Selectride, 773
- Selective serotonin reuptake inhibitors (SSRIs), 958
- Separation, of amino acids, 1081–1084
- Separator funnel, 707, 708f
- Septet, NMR spectrum, 511t, 513
- Serevent. *See* Salmeterol
- Serine
- abbreviation for, 711t
- isoelectric point for, 1078t
- pK_a values for, 1078t
- structure of, 1076f
- Serotonin, 874, 958
- Sertraline, 615f, 653
- Sex hormones, 1141–1142, 1141t. *See also specific hormones*
- Sex pheromone, 486, 1002
- Sextet, NMR spectrum, 511t, 515f

- Sharpless epoxidation, 451–454, 1085
 Sharpless, K. Barry, 452
 Sharpless reagent, 452
 Shell, of electrons, 8–9
 Shielding effects, in NMR, 502–503, 502f, 503f, 506f, 506t
 1,2-Shifts, 328–330
 Shikimic acid, 86
 Sickle cell anemia, 1110
 Sickle cell hemoglobin, 1110
 Side chain, of amino acid, 1075, 1077, 1078t
 Sigma (σ) bond, 32, 33, 36–40, 38f, 39f, 40f
 in alkanes, 83
 in alkenes, 281, 359, 428–429
 in alkynes, 400–401
 breaking in addition reactions, 199
 carbon–oxygen, 313
 formation in addition reactions, 370, 371, 371f, 372
 formation in elimination reactions, 198
 functional groups that contain C–Z σ bonds, 84, 85t
 hyperconjugation, 257
 inductive effect and, 65, 256
 length in conjugated dienes, 581–582
 radical reactions with, 540
 rotation around, 129
 in substitution reactions, 198, 242
 Sildenafil, 615f
Silent Spring, 233
 Silk, 858, 858f
 binding of dyes to, 989–990
 structure of, 1104, 1105f
 Silver(I) oxide
 in aqueous ammonium hydroxide, 738
 as oxidizing agent, 439
 Silyl ether, 749
 Simmons, H. E., 1014
 Simmons–Smith reaction, 1014–1015
 Simmons–Smith reagent, 1015
 Simvastatin, 170, 581, 581f, 1140, 1141f
 α -Sinensal, 723
 Singlet, NMR spectrum, 508, 511t, 518
 Sirenin, 1024
 Sitagliptin, 624
 Six-membered ring, synthesis of
 in aldol reaction, 928
 in Dieckmann reaction, 933
 in Robinson annulation, 936–939
 Skeletal structures, 29–31, 29f
 with charged carbon atoms, 31
 Smith, R. D., 1014
 S_N1 substitution reactions, 252–261
 alcohol conversion to alkyl halide, 332–334
 alkyl halide identity in, 262, 262f
 applications of, 261
 carbocation stability, 256–258
 Hammond postulate and, 258–261
 hyperconjugation and, 257–258
 inductive effects and, 256–257
 E1 elimination reaction compared, 294, 300–304, 302f–303f
 E2 elimination reaction compared, 300–304, 302f–303f
 energy diagram for, 253, 253f, 260f
 ethers with strong acids, 341–342
 features of, 252
 kinetics of, 252, 256t
 leaving group in, 264
 nucleophile strength in, 262–264
 racemization in, 254–255, 255f
 rate-determining step, 260
 R group in, 255–256, 256t
 S_N2 reactions compared, 262–266, 262f, 265t, 300–304, 302f–303f
 solvent effects on, 264–265
 stereochemistry, 253–255, 255f, 256t
 transition state in, 253, 253f, 264
 two-step mechanism for, 252–253, 256t
 S_N2 substitution reactions, 244–252
 in acetoacetic ester synthesis, 904, 905
 acetylide anion reactions with alkyl halides, 414–415
 acetylide anion reaction with epoxides, 416–417
 alcohol conversion to alkyl halides, 336–337
 with HX, 332–334
 with phosphorus tribromide, 336–337
 with thionyl chloride, 335–336
 alcohol synthesis from alkyl halides, 321–322
 alkyl halide identity in, 262, 262f
 alkyl halide with ammonia or amine, 960–961
 alkyl tosylate reactions, 339–340
 applications of, 250–252, 250f, 251f
 in automated peptide synthesis, 1099
 backside attack in, 245–247, 263
 direct enolate alkylation, 897
 E1 elimination reactions compared, 300–304, 302f–303f
 E2 elimination reactions compared, 300–304, 302f–303f
 energy diagrams for, 245, 245f, 249f
 epoxide ring opening reactions, 343–345
 epoxide synthesis from halohydrin, 323
 epoxy resin synthesis, 1163
 ethers with strong acids, 341–342
 ether synthesis from alkyl halides, 321–322
 features of, 244, 249t
 frontside attack in, 245–246
 of α -halo acids with ammonia, 1078–1079, 1081f
 of α -halo carbonyl compounds, 896, 896f
 kinetics of, 244, 249t
 leaving group in, 264
 in malonic ester synthesis, 901
 nitrile synthesis, 862
 nucleophile strength in, 262–264
 one-step mechanism for, 244–245, 249t
 organic synthesis, 267, 268t, 269
 reduction of alkyl halides, 437–438, 438f
 reduction of epoxides, 437–438, 438f
 R group in, 248–249, 249t
 S_N1 reactions compared, 262–266, 262f, 265t, 300–304, 302f–303f
 solvent effects on, 264–265
 stereochemistry of, 245–248, 247f, 249t
 steric hindrance and, 248–249, 248f, 249t
 transition state, 245, 245f, 246, 247f, 264
 Soap, 98–99, 99f, 851, 854–855
 Sodium, in dissolving metal reductions, 428, 436
 Sodium acetate, 93, 692
 Sodium amide, for alkyne synthesis by dehydrohalogenation, 299, 300f
 Sodium benzoate, 696, 708
 Sodium bisulfite, 443–444
 Sodium borohydride, 428, 727–731, 730f, 738, 738t, 789, 1047
 Sodium chloride, 10–11
 Sodium cyanoborohydride, for reductive amination of aldehydes and ketones, 964
 Sodium dichromate, 438, 448
 Sodium ethoxide, 322
 in dehydrohalogenation reactions, 280t
 Sodium hydroxide
 in dehydrohalogenation reactions, 280, 280t
 use in extraction procedure, 707–708, 709f
 Sodium hypochlorite, 451
 Sodium iodide, 983
 Sodium methoxide, 322
 in dehydrohalogenation reactions, 280t
 Sodium nitrite, 261
 Sodium salicylate, 696
 Sodium trichloroacetate, 1023
 Soft coral (*Capnella imbricata*), 416
 Solanine, 1045
 Solid phase technique, of amino acid synthesis, 1099–1100
 Solubility, 93–96
 of alcohols, 318t
 of aldehydes and ketones, 780t
 of alkanes, 129, 130t
 of alkenes, 365
 of alkynes, 402
 of amines, 954t
 of carboxylic acids, 693t
 definition of, 93
 of enantiomers, 182
 of epoxides, 318t
 of ethers, 318t
 intermolecular forces and, 93–96, 94f
 “like dissolves like,” 94
 of lipids, 149, 1120
 of monosaccharides, 1036, 1047
 Solute, 93
 Solvent effects, on nucleophilic substitution reactions, 240–242, 240f, 241f, 264–265, 265t
 Solvents, 93
 alkyl halides, 232
 in E1 elimination reactions, 293t
 in E2 elimination reactions, 287, 288t
 for extraction procedure, 707–709, 708f, 709f
 liquid, 197
 nonpolar, 94
 nucleophilicity and, 240–242, 265
 organic, 94
 polar, 94
 polar aprotic, 241, 241f, 264, 265, 265t, 287, 288t, 889–890
 polar protic, 240, 240f, 264–265, 265t, 293t, 890
 s Orbitals
 description, 8–9
 hybridization of, 33–40, 33f–35f, 38f–40f
 1s orbital, 9, 32
 2s orbital, 9, 32–34, 34f
 Sorbitol, 179, 180, 1047
 D-Sorbitose, 1036f
 Sorona, 1167, 1168f
 Spam, 261
 Spandex, 1162
 Spearmint, odor of, 188
 Specific rotation, 184, 185
 Spectator ions, 55
 Spectrometer, NMR, 496, 496f
 Spectroscopy, 465
 Spermaceti wax, 1121
 Spermine, 950
 SPF rating (sun protection factor), 599
 Sphingomyelins, 1127–1128, 1128f
 Sphingosine, 1127
 sp hybrid orbitals, 34–35, 34f, 38–39, 40f
 in alkynes, 400, 401, 405
 percent s -character, 41, 68
 sp^2 hybrid orbitals, 34–35, 34f, 37, 39f, 40f, 257, 267, 313, 365
 in alkenes, 281, 283–284, 359
 of alkyl halides, 229, 229f, 267
 in aromatic heterocycles, 621–623, 623f
 carbon radical, 539
 carboxy group, 689
 percent s -character, 41, 68
 sp^3 hybrid orbitals, 33–34, 33f, 35, 35f, 36, 36f, 39f, 40f, 242, 313, 365
 in acyclic alkanes, 114
 in alkenes, 283–284
 of alkyl halides, 229, 229f
 in nucleophilic substitution reactions, 235–236
 percent s -character, 41, 68
 Spiriva. *See* Tiotropium bromide
 Spiro ring system, A–6
 Splenda. *See* Sucralose
 Squalene, 157, 459, 1133f, 1135–1136, 1135f, 1140, 1140f
 Squalene oxide, 1140, 1140f
 Stachyose, 1072
 Staggered conformation, 131–137, 132f, 133f, 134f, 136f, 295, 295f
 Stanazolol, 639, 1142
 Starch
 hydrolysis of, 160, 161, 161f, 319, 320f, 1061
 structure of, 160–161, 161f, 162f, 1060–1061
 Stearic acid, 358, 1122, 1122t
 melting point, 368t
 structure, 368f, 368t
 Stearidonic acid, 368–369, 457
 Step-growth polymers, 1150, 1160–1164
 epoxy resins, 1163–1164, 1164f
 polyamides, 1160–1161
 polycarbonates, 1163
 polyesters, 1161–1162
 polyurethanes, 1162
 Stepwise reaction, 200
 Stereocenter. *See* Stereogenic centers
 Stereochemistry, 129, 159–195
 of addition reactions, 371, 376–378, 377f, 378t
 of carbonyl reduction, 729–731
 chemical properties of enantiomers, 186–188
 chiral and achiral molecules, 163–166, 165f
 diastereomers, 175–177, 185–186, 186f
 of Diels–Alder reaction, 592
 of dienophiles, 592
 of dihydroxylation, 442–444
 of disubstituted cycloalkanes, 180–181
 of E2 elimination reactions, 295–298, 295f
 of electrophilic addition reactions, 376–378, 378t
 of enolate alkylation, 898
 of epoxidation, 440
 of halogenation reactions, 381–383, 549–551, 549t
 of halohydrin formation, 384–385, 385t
 of hydroboration–oxidation reactions, 388–389, 389t
 meso compounds, 177–179, 177f
 physical properties of stereoisomers, 182–186, 184t, 186f
 polymer, 1157–1159
 reactions of organometallic reagents, 744
 of S_N2 substitution reactions, 245–248, 247f, 249t
 of S_N1 substitution reactions, 253–255, 255f, 256t

- Stereochemistry (*continued*)
of starch and cellulose, 160–162, 161f, 162f
stereogenic centers, 166–180, 168f, 172f
- Stereogenic centers, 166–180
in amines, 951–952
in amino acids, 710–711, 1075, 1077
on carbon atoms not part of ring, 166–167
in cyclic compounds, 168–170
definition of, 164
in diastereomers, 175–177
drawing enantiomers, 168, 168f
formation from hydrohalogenation, 376, 377
halogenation reactions and, 549–551
inversion of configuration around, 246–247, 247f
labeling with R or S, 170–175, 172f, 175f, 179–180
in meso compounds, 177–179
in monosaccharides, 1030–1034
number in a molecule, 165
priority assignment, 170–174, 172f
retention of configuration around, 246
tetrahedral, 164–165, 166, 168, 169, 178
- Stereoisomers. *See also* Enantiomers
in alkenes, 360t
chemical properties of, 186–188
cis, 144–147
configuration of, 162
of conjugated dienes, 580
constitutional isomers compared, 163f
of cycloalkanes, 144–147
definition of, 144
diastereomers, 175–177, 177f, 185–186, 186f
disubstituted cycloalkanes, 180–181
enantiomers, 164
of monosaccharides, 1032–1036, 1034f
nomenclature for alkenes, 363–364
number of, maximum, 175
physical properties of, 182–186, 184t, 186f
starch and cellulose, 160–162
trans, 144–147
- Stereoselective reaction
dissolving metal reduction of alkynes, 436
of E2 elimination reactions, 290
- Stereospecific reactions
carbene addition, 1013–1014
coupling reactions with organocuprates, 1004
definition of, 382
epoxidation, 440
halogenation of alkenes, 382
Heck reaction, 1009–1010
Simmons–Smith reaction, 1015
Suzuki reaction, 1006
- Steric effects, nucleophilicity and, 239
- Steric hindrance, 239
in alcohols, 318
in hydroboration reactions, 387
S_N2 substitution reactions and, 248–249, 248f, 249t
- Steric strain, 135, 136t, 138, 239
- Steroids, 1138–1142
adrenal cortical, 1142
anabolic, 1142
cholesterol. *See* Cholesterol
sex hormones, 1141–1142, 1141t
structure of, 1138–1139, 1139f
synthesis in Diels–Alder reaction, 596–597
- Stone, Reverend Edmund, 71
- Strain
angle, 137–139, 314
Baeyer strain theory, 137
steric, 135, 136t, 138, 239
torsional, 133–134, 136t, 137, 138
s-trans conformation, 580
- Strecker synthesis, 1080–1081, 1081f
- Strong electrostatic interactions, in ionic compounds, 87
- Strychnine, 1084
- Styrene, 538, 562, 671, 1016, 1016f, 1021
polymerization of, 1150f, 1151, 1153, 1156
structure of, 1150f, 1151f
- Styrene–butadiene rubber (SBR), 1156
- Substituents, naming, 120–121, 611–612
naming aromatic rings as, 612
- Substitution reactions, A–13
alcohols, 323–324, 331–338
allylic, 554
of aryl diazonium salts, 982–986
at α -carbon of carbonyl compounds, 880–906
description of, 198
electrophilic aromatic. *See* Electrophilic aromatic substitution reactions
Heck reaction, 1003, 1009–1011
nucleophilic, 235–269
radical, 541
- Substrate, 219–220, 220f
- Succinic acid, 692
- Sucralose, 1058, 1059f
- Sucrose, 109, 157, 1058, 1058f
olestra synthesis from, 854
stereogenic centers, 170
structure of, 1058f
- Sudafed, 65. *See also* Pseudoephedrine
- Suffix, 120
-adiene, 362
-al, 776
-aldehyde, 777
-amide, 831–832, 833t
-amine, 952–953
-ane, 117, 120, 121, 127, 362
-ate, 692, 830–831, 833t
-carbonyl chloride, 830, 833t
-carboxylic acid, 830
-dioic acid, 692
-diol, 315
-ene, 362
-enol, A–5
-enone, A–5
-enyne, A–5
-ic acid, 691, 830, 831, 832
-ide, 231
-ine, 230–231
-o, 230
-oic acid, 690, 831
-one, 777
-onitrile, 832, 833t
-ose, 1029
-oxy, 316
-oyl, 779
-yl, 120, 316
-yl chloride, 833, 833t, 860
-ylic acid, 831
-yne, 401
- Sugar, simple, 104–105. *See also* Monosaccharides
- Sugars. *See also* Carbohydrates
amino, 1061–1062
nonreducing, 1048, 1048f, 1058
reducing, 1048, 1048f, 1057
simple. *See* Monosaccharides
- Sulfa drugs, 990–991
- Sulfamethoxazole, 991f
- Sulfanilamide, 990–991
- Sulfide (functional group), 85t
- Sulfisoxazole, 991f
- Sulfonate anion, 710
- Sulfonation
of benzene, 643f, 646–647
electrophilic aromatic substitution reaction, 643f, 646–647
- Sulfonic acids, 709–710
- Sulfur
in fossil fuels, 70
octet rule exceptions in, 18
- Sulfuric acid, 70
dehydration of alcohols in, 325–330, 327f
in epoxide ring opening reactions, 345
in nitration and sulfonation reactions of benzene, 646–647
- Sulfur nucleophile, 348
- Sulfur oxides, 70
- Sulfur trioxide, 646–647
- Sunscreen, 30, 599
- Supercoil, 1108, 1108f
- Superhelix, 1108–1109
- Surface area
boiling point and, 91f
van der Waals forces and, 88, 88f
- Suzuki reaction, 1002, 1003, 1005–1009, 1008f
- Sweeteners, artificial, 1058, 1059f, 1090
- Sweetness, of monosaccharides, 1036
- Sweet'n Low. *See* Saccharin
- Symbol, atomic, 7
- Symmetrical anhydrides, 826–827, 830
- Symmetry
melting points and, 93
plane of, 165–166, 165f, 178
- Syn additions, 371, 377, 378t, 386, 388–389, 434, 440, 442, 442f, 443–444, 1013
- Syndiotactic polymer, 1157–1158
- Syn periplanar geometry, 295, 295f
- Synthetic intermediate, 392
- Synthetic polymers, 560–562, 1148–1178
amorphous regions of, 1164–1165
atactic, 1157–1158
“Big Six,” 1169, 1169t
biodegradable, 1170–1171
chain-growth, 1149, 1150–1157
in consumer products, 1149, 1149f
crystalline regions of, 1164–1165
disposal of, 1169–1171
green synthesis of, 1166–1169, 1167f, 1168f
isotactic, 1157–1158
molecular weights of, 1150
nylon, 825, 859
physical properties of, 1164–1166
plasticizer addition to, 1165–1166
polyesters, 859–860
recycling of, 1148, 1169–1170, 1169t
shorthand representation of, 1150, 1150f
step-growth, 1150, 1160–1164
stereochemistry of, 1157–1159
structure of, 1149, 1164–1166
syndiotactic, 1157–1158
thermoplastics, 1165
thermosetting, 1165
- Systematic name, 119. *See also* IUPAC nomenclature
of amines, 952–953
- 2,4,5-T (2,4,5-trichlorophenoxyacetic acid), 646f
- Tacrine, 975
- Tacrolimus. *See* FK506
- Tagamet. *See* Cimetidine
- D-Tagatose, 1036f
- Talose
D-talose, 1035f, 1050
Wohl degradation, 1050
- Tamiflu, 86
- Tamoxifen, 880, 899
- Tandem ring-opening–ring-closing metathesis, 1026
- Target compound, 417
- Tartaric acid, stereoisomers of, 185–186, 186f
- Tautomerization
in acid, 410–411, 883
of amides, 863–864
in base, 883
of carbonyl compounds, 863–864, 881–883, 893
mechanism of, 883
- Tautomers
definition of, 410
enol, 410–412, 864, 882–883
keto, 410–412, 864, 882–883
- Taxol, 170, 268, 726
- Taxotere, 876
- TBDMS (*tert*-butyldimethylsilyl), 749, 750
- Teflon. *See* Polytetrafluoroethylene (Teflon)
- Temperature, reaction rate and, 216
- Teratogen, 169
- Terephthalic acid, 860, 1148, 1150f, 1161, 1176
- Terpenes, 1132–1138
biosynthesis of, 1134–1138, 1135f
classes of, 1133t
examples of, 1133f
locating isoprene units in, 1132–1134, 1133f
rubber, 1159
- α -Terpinene, 1138
- α -Terpineol, 398, 1137
- tert*- (prefix), 120, 121
- Tertiary (3°) alcohols
classification as, 313
conversion to alkyl halides with HX, 332–334, 337t
dehydration by E1 mechanism, 326
hydrogen bonding extent, 318
from reaction of esters and acid chlorides with organometallic reagents, 750–752
from reaction of ketones with organometallic reagents, 742–743
- Tertiary (3°) alkyl halides
acetylide anion reaction with, 414
classification as, 229–230
E1 elimination reactions, 292, 294, 302f
E2 elimination reactions, 287, 288, 302f
example of, 229f
S_N1 substitution reactions, 255, 262, 262f, 302f
synthesis of, 332, 334–335
- Tertiary (3°) amides, 827, 831–832, 842, 844, 850
enolates formation from, 885–886
- Tertiary (3°) amines
from direct nucleophilic substitution, 960
IR spectra of, 955, 955f
nomenclature, 952–953
reactions with acid chlorides and anhydrides, 975–976
from reduction of amides, 963
structure of, 950
- Tertiary carbon (3°), 116
- Tertiary hydrogen (3°), 116–117
- Tertiary structure, of proteins, 1104–1105, 1107f

- Testosterone, 1141t
Tetra- (prefix), 123
Tetrabutylammonium fluoride, 749
Tetrachloromethane, 232
Tetrafluoroethylene, monomers in polymerization, 562t
Tetrahalide, 409
Tetrahedral arrangement, 25–26, 26f
Tetrahedral geometry, of acyclic alkanes, 114–115
Tetrahedral oxygen, 314
Tetrahedral stereogenic center, 164–165, 166, 169, 178
Tetrahedrane, 138
Tetrahydrocannabinol (THC)
 mass spectrum of, 474, 474f
 structure of, 70
Tetrahydrofuran (THF), 317, 888, 889–890, 898
 complexed with borane, 386
 as solvent, 241f, 341
Tetrahydrogestrinone, 1142
 α -Tetralone, 653
2,3,4,6-Tetra-*O*-methyl-D-galactose, 1073
Tetramethyl α -D-glucopyranose, 1048f
Tetramethylsilane (TMS), 497, 523
Tetravalent carbon, 4, 25, 27
Tetrodotoxin, synthesis of, 589, 589f
Thalidomide, 169
THC. *See* Tetrahydrocannabinol (THC)
Thermodynamic enolates, 890–891, 899
Thermodynamic reaction, in electrophilic addition reactions of conjugated dienes, 586–588, 587f, 588f
Thermodynamics, 206–209, 207f, 208t
 of hydrate formation, 802–803
Thermoplastics, 1165
Thermosetting polymers, 1165
THF. *See* Tetrahydrofuran (THF)
Thiazolinone, 1091–1092, 1118
Thioester, 860–861
Thiol (functional group), 85t
Thionyl chloride
 conversion of alcohols to alkyl halides with, 335–336
 conversion of carboxylic acids to acid chlorides, 846–847, 846f
Three-dimensional representation, of acyclic alkanes, 114–115
Threonine, 719
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f, 1077
Threose, 190
 D-threose, 1034, 1034f, 1035f
 L-threose, 1034, 1034f
Thrombin, 1075
Thromboxanes, 1129–1130, 1130t, 1131f
Thymine, 1063–1064, 1065f
Thymol, 529, 638
Tiotropium bromide, 321
Titanium(IV) isopropoxide, 452
Toads (*Bufo*), 958
Tobacco, 614f
Tobramycin, 1062
Tollens reagent, 738, 1048
Toluene, 108
 in BTX mixture, 614
 electrophilic aromatic substitution reaction, 657, 662
 oxidation to benzoic acid, 671
 structure of, 611
 synthesis by Friedel–Crafts alkylation, 675
Toluene 2,6-diisocyanate, 1162
p-Toluenesulfonate. *See* Tosylate
p-Toluenesulfonic acid (TsOH), 70, 709–710, 804
 dehydration of alcohols to alkenes in, 325, 331f
 p-Toluenesulfonyl chloride (TsCl), 338–340, 340, 341f
 p-Toluidine
 basicity of, 970
 electrostatic potential plot of, 971f
 Torsional energy, 133–134
 Torsional strain, 133–134, 136f, 137, 138
 Tosic acid, 710
 Tosylate, 338–340, 341f
 Tosyl group, 338, 710
 TPA (thermal polyaspartate), 1171
 Trade name, 119
 Trans diaxial geometry, 296–298, 296f
 Trans dihalide, 409
 Trans fats, 433
 Trans geometry
 alkenes, 359, 360t, 363
 cycloalkenes, 359
 Trans isomers
 of 2-butene, 282–283, 282f
 of cycloalkanes, 144–147
 definition of, 144, 282, 377
 of disubstituted cycloalkanes, 180
 Transition state
 definition of, 210–211
 drawing structure of, 211
 of E1 elimination reactions, 292, 292f
 of E2 elimination reactions, 285–286, 286f, 287, 289, 295
 in energy diagrams, 210–215, 215f
 four-centered, 386
 in halogenation reactions, 546–548, 547f, 548f
 Hammond postulate and, 258–261, 259f, 260f, 546–547
 S_N1 substitution reactions, 264
 S_N2 substitution reactions, 264
 in two-step reactions, 213–215, 215f
 Trans protons, 516, 517f
Tri- (prefix), 123
Triacylglycerols, 1122–1125
 combustion of, 1125
 common fatty acids in, 1122t
 degrees of unsaturation in, 432
 energy storage in, 1125
 fats and oils, 369
 formation of, 105
 hydrogenation of, 432–434, 433f, 1124
 hydrolysis of, 367, 853, 854–855, 1124
 melting point of, 1122–1123
 mixed, 1122
 olestra as a substitute for, 854
 oxidation of, 556, 556f, 1124
 physical properties of, 369
 saponification of, 854–855
 saturated, 853, 853f, 1122–1123, 1122t, 1123f
 simple, 1122
 structure of, 366–367, 853, 853f, 1120f, 1122–1123, 1122t, 1123f, 1128f
 unsaturated, 1122–1123, 1122t, 1123f, 1124
Trialkylborane, 387, 387f
Triarylphosphine, 1009
1,3,5-Tribromobenzene, synthesis of, 984, 984f
Tributyltin hydride, 570
Trichlorofluoromethane, 3, 233, 551
Trichloromethane, 232
2,4,5-Trichlorophenoxyacetic acid
 herbicide (2,4,5-T), 646f
Tricyclohexylphosphine, 1005
Trienes, 362
Triethylamine, 71, 277, 952, 1009
1,2,4-Triethylcyclopentane, 127f
Trifluoroacetic acid, for removal of Boc protecting group, 1096
2,2,2-Trifluoroethanol, 65–66, 66f
Trifluoromethanesulfonic acid, 710
Trigonal planar carbocation, 253–254, 376
Trigonal planar carbon radical, 539
Trigonal planar double bond, 281
Trigonal planar molecule, 24–25, 26, 26f
 carbonyl group, 722–723
 carboxy group, 689
Trigonal pyramidal molecule
 amines, 950–951
Trihalomethanes, 1012
Trimethylamine, 956
 electrostatic potential plot of, 951, 951f
Trimethyl borate, 1007
1,3,5-Trimethylcyclohexane, 156
2,5,5-Trimethylcyclohexanecarboxylic acid, 690
2,5,5-Trimethylcyclohexanol, 315f
2,3,5-Trimethyl-2-hexene, 362
1,3,6-Tri-*O*-methyl-D-fructose, 1073
2,3,4-Tri-*O*-methyl-D-glucose, 1073
2,2,4-Trimethylpentane, combustion of, 148
2,3,5-Trimethyl-4-propylheptane, 123f
Triols, 315
Tripeptide, 1086–1088
Triphenylene, 640
Triphenylphosphine, 793, 1005
Triphenylphosphine oxide, 792, 794
Triple bonds
 in alkynes, 83, 84t, 400–401
 bond length and bond strength, 41, 41t
 components of carbon–carbon, 39
 as functional group, 83, 84t
 Lewis structure, 15
 NMR spectra of, 505, 506f, 506t
 reduction of, 434–437, 437f
Triple helix, of collagen, 1108–1109, 1109f
Triplet, NMR spectrum, 509, 510, 511t, 512, 512t, 515f, 518
Tri(*o*-tolyl)phosphine, 1005
Triynes, 401
Tropylium cation, 625–626
Truvia. *See* Rebaudioside A
Trypsin, 1093, 1093t
Tryptophan, 720
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
TsCl (*p*-Toluenesulfonyl chloride), 338–340, 341f
TsOH. *See* *p*-Toluenesulfonic acid (TsOH)
Tuberculosis, 250
Tulipalin A, 946
Tungsten, 1015
ar-Turmerone, 926
Twistoflex, 614–615
Two-step reaction mechanism, 213–215, 215f, 217
TXA₂, 1130f
Tylenol. *See* Acetaminophen
Tyrian purple, 988
Tyrosine
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
Tyvek, 1170
Ultraviolet light
 absorption by conjugated dienes, 597–599, 598f
 skin damage from, 599
Undecane, 114
Unimolecular elimination. *See* E1 elimination reactions
Unimolecular reaction
 in nucleophilic substitution reactions, 244, 252
 organic reactions, 217
Unknown
 analysis using mass spectrometry, 466–468
 using ¹H NMR to identify an, 519–522
 α,β -Unsaturated carbonyl compounds, 755–758
 1,2-addition to, 755–756
 1,4-addition to, 755–756
 in Michael reaction, 934–936, 935f
 resonance structures, 934
 in Robinson annulation, 936–939
 synthesis of
 in dehydration of aldol product, 919–920, 922, 924f
 from α -halo carbonyl compounds, 896
Unsaturated fatty acid. *See* Fatty acids, unsaturated
Unsaturation, degrees of
 in alkenes, 431–432
 benzene, 608
 calculation of, 360–361, 431–432
 hydrogenation data and, 431–432
 in triacylglycerols, 432
Uracil, 1063–1064, 1065f
Urethanes, 1162, 1175
Urushiols, 707
Valdecoxib, 1132
Valence bond theory, 627
Valence electrons, 10–16, 17t, 32
Valence shell electron pair repulsion (VSEPR) theory, 24
Valeric acid, structure of, 691t
Valine
 isoelectric point for, 1078t
 pK_a values for, 1078t
 structure of, 1076f
Valinomycin, 101
Valium. *See* Diazepam
Valproic acid, 78, 910
Vancomycin, 111
van der Waals forces, 87–89, 88f, 90t
 in alkanes, 129
 surface area and, 88, 88f
Vane, John, 697
Vanillin, 30, 529, 783f
Vaniiloids, 4
Varenicline, 995
Vasopressin, 1089, 1090f, 1105
Vasotec. *See* Enalapril
Vegetable oil, 835, 854, 855
 partial hydrogenation of, 432–433, 433f
Venlafaxine, 748, 773
Ventolin. *See* Albuterol
 β -Vetivone, 913, 915, 943
Viagra. *See* Sildenafil
Vibration of bonds, infrared spectroscopy and, 476–477, 478–480
Vicinal dibromide, 404, 405
Vicinal dihalides, 299, 379
 alkyne synthesis from, 404–405
Vinegar, 695
Vinyl acetate
 NMR spectrum of, 517f
 polymerization of, 1152, 1155f
 structure of, 1151f
Vinylboranes, 1007, 1009
 trans vinylborane, 1006–1007
Vinyl bromide, 1009
Vinyl carbanion, 436
Vinyl carbocation, 407, 408








- Vinyl chloride
bond strength, 51
monomers in polymerization, 562t
polymerization of, 1149, 1156
structure of, 1149, 1151f
- 1-Vinylcyclohexene, 365
- Vinyl group, 365
- Vinyl halides, 229, 229f, 267
coupling reactions with
alkenes, 1009–1011
organoboranes, 1006–1007
organocuprates, 1003–1004
reactivity of, 650
synthesis in hydrohalogenation of
alkynes, 406–407
- Vinylidene chloride, polymerization of,
1156
- Vinyl iodide, 1010–1011
- Vioxx. *See* Rofecoxib
- Viracept. *See* Nelfinavir
- Vitamin(s), 97–98
definition of, 97, 1128
fat-soluble, 97, 149, 1128, 1129f
origin of term, 97
water-soluble, 97–98
- Vitamin A, 97–98, 331f
 β -carotene conversion to, 97–98
deficiency, 97, 1129f
electrostatic potential plot of, 1128,
1129f
role in the body, 1129f
solubility, 97
sources, 1128, 1129f
structure, 97, 282, 1129f, 1134
- Vitamin B₃, 98
- Vitamin B₆, 109
- Vitamin C (ascorbic acid), 80, 81, 568
solubility of, 98
sources of, 98
structure of, 98, 836
- Vitamin D₃, 581
deficiency, 1129f
role in the body, 1129f
sources of, 1128
structure of, 282, 1129f
- Vitamin E, 109
as antioxidant, 557
deficiency, 1129f
electrostatic potential plot of, 1128,
1129f
as radical scavenger, 541
role in the body, 1129f
sources of, 1128
structure of, 557, 1129f
- Vitamin K (phylloquinone), 98
deficiency, 1129f
role in the body, 1129f
sources of, 1128
structure of, 1129f
- VSEPR (valence shell electron pair
repulsion) theory, 24
- Vulcanization, 1159, 1159f
- Wald, George, 799
- Walden, Dr. Paul, 246
- Walden inversion, 246
- Wang resin, 1118
- Water
as achiral molecule, 164
electrophilic addition of, 378–379,
390
electrostatic potential plot for, 45f
hybrid orbitals in, 35, 35f
hydration of alkynes, 406f, 409–412
hydrogen bonding, 89
as Lewis base, 73
Markovnikov addition of, 410
molecular shape of, 26, 26f
nucleophilicity of, 73, 263
 pK_a of, 59t, 62, 63
as polar molecule, 44
removing from reaction mixture using
Dean–Stark trap, 804, 805f
as solvent, 94–96, 240, 240f, 264
in extraction procedure, 707, 708f
- Wavelength
definition of, 474
electromagnetic spectrum and,
474–476, 475f
length units for, 474
- Wavenumber, 476
- Waxes, 150, 150f, 1121
- Wedges, in Fischer projection formula,
1032
- Wellbutrin. *See* Bupropion
- Williamson ether synthesis, 322
- Willow tree, 71
- Wittig, Georg, 792
- Wittig reaction
mechanism of, 794
retrosynthetic analysis using, 795–796
- Wittig reagent, 792–797
- Wohl degradation, 1049, 1050
- Wolff–Kishner reduction, 672
- Wood alcohol, 319f. *See also* Methanol
- Wool
binding of dyes to, 989–990
structure of, 858, 858f
- Xalatan. *See* Latanoprost
- X-ray crystallography, 1053
- m*-Xylene, 667
- p*-Xylene, 614
- D-Xylose, 1034, 1035f
- D-Xylulose, 1036f
- yl (suffix), 120, 316, 779
- yl chloride (suffix), 833, 833t, 860
- ylic acid (suffix), 831
- Ylide, 793
- yne (suffix), 401
- Zaitsev rule, 288–291, 293, 297, 980
- Z configuration, of double bonds, 368
- Zidovudine. *See* AZT
(azidodeoxythymidine)
- Ziegler, Karl, 1158
- Ziegler–Natta catalysts, 1157–1158, 1160
- Zigzag skeletal structures, 137
- Zileuton, 348
- Zinc
in Clemmensen reduction, 672–673
conversion of ozonide to carbonyl
compounds, 445
- Zinc chloride, 333
- Zinc–copper couple, 1014–1015
- Zingerone, 50
- Zingiberene, 367f, 459, 601, 1133f
- Z isomer, 364
- Zocor. *See* Simvastatin
- Zolof. *See* Sertraline
- Zuclomiphene, 394
- Zwitterion, 80, 711–712, 1077
- Zyban. *See* Bupropion
- Zyflo. *See* Zileuton

Common Abbreviations, Arrows, and Symbols

Abbreviations

Ac	acetyl, $\text{CH}_3\text{CO}-$
BBN	9-borabicyclo[3.3.1]nonane
BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
Boc	<i>tert</i> -butoxycarbonyl, $(\text{CH}_3)_3\text{COCO}-$
bp	boiling point
Bu	butyl, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2-$
CBS reagent	Corey–Bakshi–Shibata reagent
DBN	1,5-diazabicyclo[4.3.0]non-5-ene
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DCC	dicyclohexylcarbodiimide
DET	diethyl tartrate
DIBAL-H	diisobutylaluminum hydride, $[(\text{CH}_3)_2\text{CHCH}_2]_2\text{AlH}$
DMF	dimethylformamide, $\text{HCON}(\text{CH}_3)_2$
DMSO	dimethyl sulfoxide, $(\text{CH}_3)_2\text{S}=\text{O}$
<i>ee</i>	enantiomeric excess
Et	ethyl, CH_3CH_2-
Fmoc	9-fluorenylmethoxycarbonyl
HMPA	hexamethylphosphoramide, $[(\text{CH}_3)_2\text{N}]_3\text{P}=\text{O}$
HOMO	highest occupied molecular orbital
IR	infrared
LDA	lithium diisopropylamide, $\text{LiN}[\text{CH}(\text{CH}_3)_2]_2$
LUMO	lowest unoccupied molecular orbital
<i>m</i> -	meta
mCPBA	<i>m</i> -chloroperoxybenzoic acid
Me	methyl, CH_3-
MO	molecular orbital
mp	melting point
MS	mass spectrometry
MW	molecular weight
NBS	<i>N</i> -bromosuccinimide
NMO	<i>N</i> -methylmorpholine <i>N</i> -oxide
NMR	nuclear magnetic resonance
<i>o</i> -	ortho
<i>p</i> -	para
PCC	pyridinium chlorochromate
Ph	phenyl, C_6H_5-
ppm	parts per million
Pr	propyl, $\text{CH}_3\text{CH}_2\text{CH}_2-$
RCM	ring-closing metathesis
ROMP	ring-opening metathesis polymerization
TBDMS	<i>tert</i> -butyldimethylsilyl
THF	tetrahydrofuran
TMS	tetramethylsilane, $(\text{CH}_3)_4\text{Si}$
Ts	tosyl, <i>p</i> -toluenesulfonyl, $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_2-$
TsOH	<i>p</i> -toluenesulfonic acid, $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3\text{H}$
UV	ultraviolet

Arrows

	reaction arrow
	equilibrium arrows
	double-headed arrow, used between resonance structures
	full-headed curved arrow, showing the movement of an electron pair
	half-headed curved arrow (fishhook), showing the movement of an electron
	retrosynthetic arrow
	no reaction

Symbols

$\overset{+}{\longrightarrow}$	dipole
$h\nu$	light
Δ	heat
δ^+	partial positive charge
δ^-	partial negative charge
λ	wavelength
ν	frequency
$\tilde{\nu}$	wavenumber
HA	Brønsted–Lowry acid
B:	Brønsted–Lowry base
$:\text{Nu}^-$	nucleophile
E^+	electrophile
X	halogen
\blacktriangleleft	bond oriented forward
\blacktriangleright	bond oriented behind
$---$	partial bond
$[\]^\ddagger$	transition state
[O]	oxidation
[H]	reduction